

MilkChEST, an integrative approach of the secretion pathways in the mammary epithelial cell: molecular dissection of milk fat adaptability

Christine Leroux, Christian Beauvallet, Laurence Bernard, Claudia Bevilacqua, Frédéric Bouvier, Cédric Cabau, Christelle Cebo, Eric Chanat, Isabelle Dugail, Thierry Fassier, et al.

▶ To cite this version:

Christine Leroux, Christian Beauvallet, Laurence Bernard, Claudia Bevilacqua, Frédéric Bouvier, et al.. MilkChEST, an integrative approach of the secretion pathways in the mammary epithelial cell: molecular dissection of milk fat adaptability. European Regional Conference On Goats, Jun 2013, Tromsø, Norway. hal-01194090

HAL Id: hal-01194090

https://hal.science/hal-01194090

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

MilkChEST, an integrative approach of the secretion pathways in the mammary epithelial cell: molecular dissection of milk fat adaptability

Leroux Ch.¹, Beauvallet C.², Bernard L.¹, Bevilacqua C.², BouvierF.³, Cabau C.⁴, Cebo Ch.², Chanat E.⁵, Dugail I.⁶, Fassier T.³, Faulconnier Y.¹, de la Foye A.¹, Huau Ch.⁷, Henry C.⁸, Le Lay S.⁹, Longin Ch.⁵, Miranda G.², Pauloin A.⁵, Robert-Granié Ch.⁷, Truchet S.⁵ and Martin P.²

¹ INRA, UMR1213 Herbivores, F-63122 Saint-Genès Champanelle, France; ² INRA, UMR1313 Gabi, F-78350 Jouy-en-Josas, France; ³ INRA, UE de Bourges, F-18390 Osmoy, France; ⁴ INRA, URA, F-37380 Nouzilly, France; ⁵ INRA, UR1196 GPL, F-78350 Jouy-en-Josas, France; ⁶ INSERM, UMRS 872 INSERM, F-75006 Paris, France; ⁷ INRA UR631 Saga, F-31326 Toulouse, France; ⁸ INRA, UMR 1319 Papso, F-78350 Jouy-en-Josas, France; ⁹ INSERM, U1063 Sopam, F-49933 Nantes, France

During lactation, mammary epithelial cells (MEC) secrete huge quantities of components, the biosynthesis of which are known in general outline. There are some evidences that a close mechanistic relationship exist between the biosynthesis and secretion pathways of the major milk components: casein micelles and milk fat globules (MFG). However, our knowledge of the molecular events occurring during assembly, intracellular transport and secretion of the micelles and MFG remains scarce. Lately there is also a realization that MEC release in milk microparticules, including exosomes, which contain mRNA and microRNA. To go further into the molecular dissection of the biosynthetic and secretion pathways of the major milk components, processes that determine their structure, as well as their biological and technological properties, a large collaborative project involving several French teams has been launched (funded by the French National Agency for Research and Apis-Gene, France).

Our strategy is based on the relevance of the goat model, with a "naturally occurring Knock Out (KO)" at the CSN1S1 locus and on the use of global approaches (including NGS, proteomics and lipidomics analyses) to decipher the multiple effects of this natural KO on milk secretion and quality.

Our main research goals will be: i) to achieve a better definition of the genome fraction which is expressed in goat MEC during lactation, ii) to obtain a better understanding of the role played by 2s1-casein in the export of caseins from the endoplasmic reticulum to the Golgi apparatus, iii) to identify factors that determine the fatty acid profile of milk fat, iiii) to go further into the molecular dissection of the mechanisms driving lipid droplets formation and their interaction with the apical plasma membrane during the course of MFG secretion.

The project includes a comparative study of lipid droplet formation in and microparticules release by MEC versus adipocytes, the later cells producing large lipid droplets but lacking the machinery to secrete them. On the other hand, we will focus on the consequences of the absence of the ②s1-casein on the protein and fat secretory pathways, as well as on the fine composition of milk. The ultimate goal is to identify genes playing key roles in the production of milk fat by MEC on which selection and breeding program might be engaged to modulate the fat composition of milk.