Irrigation, soil organic carbon and N₂O emissions. A review

Benjamin Trost, Annette Prochnow, Katrin Drastig, Andreas Meyer-Aurich,
Frank Ellmer, Michael Baumecker

► To cite this version:

Benjamin Trost, Annette Prochnow, Katrin Drastig, Andreas Meyer-Aurich, Frank Ellmer, et al..
Irrigation, soil organic carbon and N₂O emissions. A review. *Agronomy for Sustainable Development*,
2013, 33 (4), pp.733-749. 10.1007/s13593-013-0134-0 . hal-01201385

HAL Id: hal-01201385

<https://hal.science/hal-01201385>

Submitted on 17 Sep 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Irrigation, soil organic carbon and N₂O emissions. A review

Benjamin Trost · Annette Prochnow · Katrin Drastig ·
Andreas Meyer-Aurich · Frank Ellmer ·
Michael Baumecker

Accepted: 18 January 2013 / Published online: 20 February 2013
© INRA and Springer-Verlag France 2013

Abstract Irrigation has a critical role for crop production worldwide. In particular, irrigation is a major issue due to the growing food demand and climate change. Irrigation affects yields and the emission of greenhouse gases such as CO₂ and N₂O by soils. Here, we review the effect of irrigation on soil organic carbon and N₂O emissions. We

B. Trost (✉) · A. Prochnow · K. Drastig · A. Meyer-Aurich
Leibniz-Institute for Agricultural Engineering Potsdam-Bornim,
Max-Eyth-Allee 100,
14469 Potsdam, Germany
e-mail: btröst@atb-potsdam.de

A. Prochnow
e-mail: aprochnow@atb-potsdam.de

A. Prochnow
e-mail: annette.prochnow@agrar.hu-berlin.de

K. Drastig
e-mail: kdrastig@atb-potsdam.de

A. Meyer-Aurich
e-mail: ameyer@atb-potsdam.de

A. Prochnow
Chair Utilization Strategies for Bioresources, Faculty of
Agriculture and Horticulture, Humboldt-University of Berlin,
Hinter der Reinhardtstr. 8-18,
10115 Berlin, Germany

F. Ellmer
Division of Agronomy and Crop Science, Faculty of Agriculture
and Horticulture, Humboldt-University of Berlin, Albrecht-Thaer-
Weg 5,
14195 Berlin, Germany
e-mail: frank.ellmer@agrar.hu-berlin.de

M. Baumecker
Field Study and Research Station, Faculty of Agriculture and
Horticulture, Humboldt-University of Berlin, Dorfstraße 9,
14974 Thyrow, Germany
e-mail: michael.baumecker@agrar.hu-berlin.de

analysed 22 investigations in various regions of the world. Interactions between irrigation, soil and management factors are described. The main points are: (1) The influence of irrigation is strongly dependent on climate and initial soil organic carbon content. For instance, irrigation of cultivated desert soils led to an average increase of 90 % to over 500 % of soil organic carbon. (2) Irrigation of semiarid regions increases soil organic carbon by 11 % to 35 %. (3) No consistent effects of irrigation were observed in humid regions. In many cases, N₂O emissions increase after precipitation or irrigation. (4) Comparison of N₂O emissions from irrigated and non-irrigated fields shows that availability of reactive nitrogen compounds controls increased N₂O emissions under irrigation, in most cases. Here, increases of about 50 % to 140 % in N₂O emissions were reported.

Keywords Irrigation · Soil organic carbon · Nitrous oxide emissions

1 Introduction

Agriculture today is facing unprecedented challenges. The world population will grow from 6.9 billion people in 2010 to an estimated 9.15 billion people in 2050 (Alexandratos and Bruinsma 2012). At the same time, the individual food energy intake will increase from 2,850 to 3,130 kcal per capita and day (Bruinsma 2009). These trends will apply pressure on resources needed for agricultural production such as land, water and energy, and increase greenhouse gas emissions from agriculture. The situation is expected to be aggravated further by climate change (Alexandratos and Bruinsma 2012). Irrigation is an important means to ensure water supply for crop production and to adapt agriculture to increasing water scarcity due to climate change. On the other hand, irrigation itself might affect climate by altering

the capacity of soils to act as sinks or sources of greenhouse gases, in particular, CO₂ and N₂O (Lal 2004). Hence, it is desirable to understand the interactions between climate and irrigation. Water availability is an important factor for crop production (Fig. 1) and also for the amount of organic residues in soils. In many regions of the world, water deficiency is one of the most yield-restricting factors. Due to drought and water deficiency, only 30 % of worldwide maximum attainable yields are approached (Deng et al. 2005). The use of irrigation to overcome the lack of reliable rainfall is growing worldwide. Around the world 306,247,000 ha of agricultural land are irrigated. That is 22.1 % of the arable land and 6.2 % of the worldwide agricultural area (FAO Statistical Yearbook 2010). The importance of irrigation is expected to grow further to meet the rising global demand for agricultural products and to adapt to increasing water scarcity due to climate change. Table 1 shows the current distribution of area equipped for irrigation for different continents and selected countries. Despite the huge relevance for worldwide agricultural production, the potential contribution of irrigation to net greenhouse gas emissions is little investigated compared with other agromonic activities (King et al. 2009).

The objective of this review is to compile results from 22 investigations about the impact of irrigation on soil organic carbon contents and N₂O emissions. Furthermore, the underlying processes and interactions of irrigation with other management factors such as tillage and fertilisation are discussed. Conclusions on potential effects of irrigation on net greenhouse gas emissions and necessary further research are drawn.

2 Basic effects of irrigation on CO₂ and N₂O emission processes

Irrigation may influence the CO₂ and N₂O emissions of arable land via several processes. Some processes make soil a sink for CO₂ while others may promote a release (Fig. 2).

Fig. 1 Irrigation, an essential factor in crop production

Table 1 Irrigated area (data from FAO Statistical yearbook 2010)

Region	Irrigated area, ha (2008)
World	306,247,000
Asia	195,461,000
India	62,286,000
China	64,141,000
Pakistan	19,870,000
Afghanistan	3,199,000
Europe	29,564,000
Russia	4,346,000
Italy	3,950,000
Spain	3,800,000
Romania	3,157,000
North and Central America	31,968,000
USA	23,000,000
Mexico	6,300,000
Cuba	870,000
Canada	855,000
South America	12,082,000
Brazil	4,500,000
Argentina	1,550,000
Peru	1,195,000
Colombia	900,000
Africa	13,576,000
Egypt	3,530,000
Sudan	1,863,000
South Africa	1,498,000
Morocco	1,457,000
Middle East	22,277,000
Iran	8,993,000
Turkey	5,215,000
Iraq	3,525,000
Saudi Arabia	1,731,000
Australia	2,550,000

An improved water supply leads, on the one hand, to an increased biomass generation and therefore to a higher input of carbon into the soil in the form of roots and dead plant material (Entry et al. 2008; Kochsiek et al. 2009; Roldan et al. 2005). On the other hand, the application of water and consequently higher soil moisture enhances soil microbial activity. This may result in an increased decomposition of soil organic matter and therefore in rising CO₂ emissions (Jabro et al. 2008; Kochsiek et al. 2009; Liu et al. 2008). The increased microbial decomposition of soil organic matter may lead to lower soil organic carbon contents (Dersch and Bohm 2001; Getaneh et al. 2007).

The effects of irrigation on soil organic carbon content not only depend on the decrease or increase in soil moisture.

Fig. 2 Basic effects of irrigation on soil organic carbon content and nitrous oxide (N₂O) emissions (increase, decrease)

The interaction with other factors like fertilisation, tillage or the activity of soil organisms may also influence the development of soil organic carbon.

Another important greenhouse gas emitted from arable land and influenced by irrigation is N₂O. Nitrification and denitrification are the main processes for N₂O formation (Bremner 1997; Phillips 2008), and both irrigation and precipitation can influence these microbial processes. Improved living conditions for microorganisms by increased soil moisture may cause enhanced activity of nitrifying bacteria (Jha et al. 1996). An increase in water-filled pore volume over 70 % may lead to reduced soil aeration resulting in low oxygen concentrations to anaerobic conditions which support denitrification (Amha and Bohne 2011; Ruser et al. 2006; Scheer et al. 2008). An increased soil microbial activity may lead to a decrease in the soil oxygen concentration as well (Loecke and Robertson 2009; Potthoff et al. 2005). Thus, irrigation may contribute to CO₂ mitigation and simultaneously enhance N₂O emissions. Increased N₂O emissions might reduce or offset a potential positive effect of enhanced carbon sequestration (Ball et al. 2008; Chatskikh and Olesen 2007; Li et al. 2005; Smith et al. 2000).

3 Soil carbon contents under irrigation

3.1 Overview of long-term field experiments

An overview of 14 long-term field experiments created to observe the soil carbon content under irrigation compared with non-irrigation is given in Table 2. In some cases, soil organic carbon contents were significantly higher under irrigation, while other results do not show significant differences between the treatments.

Several experiments on the development of soil organic carbon contents on irrigated and non-irrigated plots were conducted in arid or semiarid regions where irrigation is essential for crop production. Denef et al. (2008) conducted investigations in an arid region of the USA. The improved water availability under irrigation led to an increase in biomass production and thus, to an accumulation of organic matter in the soil. As a consequence, the contents of soil organic carbon were significantly higher than in the non-irrigated cultivated plots. However, on the experimental site with the higher precipitation, the highest soil organic carbon contents were found on plots with natural vegetation.

Table 2 Overview of experiments to investigate the effects of irrigation on the soil organic carbon content

Reference	Location	Climate/annual precipitation/mean annual temperature	Soil	Crops/tillage	Fertilization	Years under irrigation	C-content
Denef et al. (2008)	USA, Nebraska, Central Great Plains	Arid/570 mm/9.7 °C	sand loamy sand	corn, wheat, soybean/NT	n.r.	30–40	Native vegetation 3,809 g C m ⁻² Irrig. 3,180 g C m ⁻² a Non-irrig. 2,544 g C m ⁻² (0–20 cm soil depth)
							Native vegetation 2,155 g C m ⁻² Irrig. 2,958 g C m ⁻² a Non-irrig. 2,192 g C m ⁻² (0–20 cm soil depth)
							Native vegetation 3,974 g C m ⁻² Irrig. 3,301 g C m ⁻² a Non-irrig. 2,664 g C m ⁻² (0–20 cm soil depth)
							Native vegetation 5,910 g C m ⁻² Irrig. arable land CT 7,290 g C m ⁻² a Irrig. arable land NT 8,010 g C m ⁻² a Irrig. pasture 10,140 g C m ⁻² a (0–100 cm soil depth)
Entry et al. (2004)	USA Idaho Snake River Plain	Semiarid-arid/175–305 mm	Silty loam	Pasture, alfalfa, wheat, potato, field bean/n.r.	n.r.	Pasture, 30 Arable land, 8	Native vegetation 3,380 g C m ⁻² After 25 years irrig. 3,330 g C m ⁻² After 30 years irrig. 3,370 g C m ⁻² After 45 years irrig. 4,330 g C m ⁻² a After 55 years irrig. 4,910 g C m ⁻² a (0–100 cm soil depth)
							Native vegetation 3,380 g C m ⁻² After 25 years irrig. 3,330 g C m ⁻² After 30 years irrig. 3,370 g C m ⁻² After 45 years irrig. 4,330 g C m ⁻² a After 55 years irrig. 4,910 g C m ⁻² a (0–100 cm soil depth)
							Native vegetation 3,380 g C m ⁻² After 25 years irrig. 3,330 g C m ⁻² After 30 years irrig. 3,370 g C m ⁻² After 45 years irrig. 4,330 g C m ⁻² a After 55 years irrig. 4,910 g C m ⁻² a (0–100 cm soil depth)
							Native vegetation 3,380 g C m ⁻² After 25 years irrig. 3,330 g C m ⁻² After 30 years irrig. 3,370 g C m ⁻² After 45 years irrig. 4,330 g C m ⁻² a After 55 years irrig. 4,910 g C m ⁻² a (0–100 cm soil depth)
Wu et al. (2008)	USA California Wasco	Semiarid/n.r.	Loam	Corn/n.r.	n.r.	55	Native vegetation 3,380 g C m ⁻² After 25 years irrig. 3,330 g C m ⁻² After 30 years irrig. 3,370 g C m ⁻² After 45 years irrig. 4,330 g C m ⁻² a After 55 years irrig. 4,910 g C m ⁻² a (0–100 cm soil depth)
							Native vegetation 3,380 g C m ⁻² After 25 years irrig. 3,330 g C m ⁻² After 30 years irrig. 3,370 g C m ⁻² After 45 years irrig. 4,330 g C m ⁻² a After 55 years irrig. 4,910 g C m ⁻² a (0–100 cm soil depth)
							Native vegetation 3,380 g C m ⁻² After 25 years irrig. 3,330 g C m ⁻² After 30 years irrig. 3,370 g C m ⁻² After 45 years irrig. 4,330 g C m ⁻² a After 55 years irrig. 4,910 g C m ⁻² a (0–100 cm soil depth)
							Native vegetation 3,380 g C m ⁻² After 25 years irrig. 3,330 g C m ⁻² After 30 years irrig. 3,370 g C m ⁻² After 45 years irrig. 4,330 g C m ⁻² a After 55 years irrig. 4,910 g C m ⁻² a (0–100 cm soil depth)
Li et al. (2006)	USA California Holtville	Arid/n.r.	Silty clay	Alfalfa, wheat, corn, sugar beet/n.r.	n.r.	90	Native vegetation 3,380 g C m ⁻² After 25 years irrig. 3,330 g C m ⁻² After 30 years irrig. 3,370 g C m ⁻² After 45 years irrig. 4,330 g C m ⁻² a After 55 years irrig. 4,910 g C m ⁻² a (0–100 cm soil depth)
							Native vegetation 3,380 g C m ⁻² After 25 years irrig. 3,330 g C m ⁻² After 30 years irrig. 3,370 g C m ⁻² After 45 years irrig. 4,330 g C m ⁻² a After 55 years irrig. 4,910 g C m ⁻² a (0–100 cm soil depth)
							Native vegetation 3,380 g C m ⁻² After 25 years irrig. 3,330 g C m ⁻² After 30 years irrig. 3,370 g C m ⁻² After 45 years irrig. 4,330 g C m ⁻² a After 55 years irrig. 4,910 g C m ⁻² a (0–100 cm soil depth)
							Native vegetation 3,380 g C m ⁻² After 25 years irrig. 3,330 g C m ⁻² After 30 years irrig. 3,370 g C m ⁻² After 45 years irrig. 4,330 g C m ⁻² a After 55 years irrig. 4,910 g C m ⁻² a (0–100 cm soil depth)
Li et al. (2006)	China Gansu Province	Yongchang arid/0 mm/7.3 °C Jiuquan arid/61 mm/8.1 °C Wuwei arid/158 mm/7.6 °C Gulang arid/150 mm/6.8 °C	Aridisols	Corn, barley, sunflower, sugarbeet, wheat/n.r.	n.r.	Yongchang, 23 Jiuquan, 30 Wuwei, 50 Gulang, 113	Yongchang: After 0 years irrig. 3.5 g C kg ⁻¹ soil After 5 years irrig. 4.5 g C kg ⁻¹ soil After 9 years irrig. 4.0 g C kg ⁻¹ soil After 23 years irrig. 7.1 g C kg ⁻¹ soil Jiuquan: After 0 years irrig. 2.3 g C kg ⁻¹ soil After 6 years irrig. 4.6 g C kg ⁻¹ soil After 8 years irrig. 3.6 g C kg ⁻¹ soil After 12 years irrig. 3.6 g C kg ⁻¹ soil After 23 years irrig. 6.1 g C kg ⁻¹ soil After 30 years irrig. 6.9 g C kg ⁻¹ soil Wuwei: After 0 years irrig. 2.6 g C kg ⁻¹ soil
							Yongchang: After 0 years irrig. 3.5 g C kg ⁻¹ soil After 5 years irrig. 4.5 g C kg ⁻¹ soil After 9 years irrig. 4.0 g C kg ⁻¹ soil After 23 years irrig. 7.1 g C kg ⁻¹ soil Jiuquan: After 0 years irrig. 2.3 g C kg ⁻¹ soil After 6 years irrig. 4.6 g C kg ⁻¹ soil After 8 years irrig. 3.6 g C kg ⁻¹ soil After 12 years irrig. 3.6 g C kg ⁻¹ soil After 23 years irrig. 6.1 g C kg ⁻¹ soil After 30 years irrig. 6.9 g C kg ⁻¹ soil Wuwei: After 0 years irrig. 2.6 g C kg ⁻¹ soil
							Yongchang: After 0 years irrig. 3.5 g C kg ⁻¹ soil After 5 years irrig. 4.5 g C kg ⁻¹ soil After 9 years irrig. 4.0 g C kg ⁻¹ soil After 23 years irrig. 7.1 g C kg ⁻¹ soil Jiuquan: After 0 years irrig. 2.3 g C kg ⁻¹ soil After 6 years irrig. 4.6 g C kg ⁻¹ soil After 8 years irrig. 3.6 g C kg ⁻¹ soil After 12 years irrig. 3.6 g C kg ⁻¹ soil After 23 years irrig. 6.1 g C kg ⁻¹ soil After 30 years irrig. 6.9 g C kg ⁻¹ soil Wuwei: After 0 years irrig. 2.6 g C kg ⁻¹ soil
							Yongchang: After 0 years irrig. 3.5 g C kg ⁻¹ soil After 5 years irrig. 4.5 g C kg ⁻¹ soil After 9 years irrig. 4.0 g C kg ⁻¹ soil After 23 years irrig. 7.1 g C kg ⁻¹ soil Jiuquan: After 0 years irrig. 2.3 g C kg ⁻¹ soil After 6 years irrig. 4.6 g C kg ⁻¹ soil After 8 years irrig. 3.6 g C kg ⁻¹ soil After 12 years irrig. 3.6 g C kg ⁻¹ soil After 23 years irrig. 6.1 g C kg ⁻¹ soil After 30 years irrig. 6.9 g C kg ⁻¹ soil Wuwei: After 0 years irrig. 2.6 g C kg ⁻¹ soil

Table 2 (continued)

Reference	Location	Climate/annual precipitation/mean annual temperature	Soil	Crops/tillage	Fertilization	Years under irrigation	C-content	
Li et al. (2009)	China Gao tai region	Arid /79 mm/7.6 °C	Loamy sand	Wheat, barley, corn, alfalfa, robinia/n.r.	Corn 515 kg N ha ⁻¹ year ⁻¹ 86 kg P ha ⁻¹ year ⁻¹ Alfalfa 69 kg N ha ⁻¹ year ⁻¹ 300–400 kg N ha ⁻¹ year ⁻¹ 90–150 kg P ₂ O ₅ ha ⁻¹ year ⁻¹ 60–90 kg K ₂ O ha ⁻¹ year ⁻¹ 3,000–6,000 kg farmyard manure ha ⁻¹ year ⁻¹ n.r.	10	After 3 years irrig.	4.0 g C kg ⁻¹ soil
							After 7 years irrig.	6.9 g C kg ⁻¹ soil
							After 11 years irrig.	4.8 g C kg ⁻¹ soil
							After 50 years irrig.	5.1 g C kg ⁻¹ soil
							Gulang:	
							After 0 years irrig.	3.3 g C kg ⁻¹ soil
							After 7 years irrig.	3.5 g C kg ⁻¹ soil
							After 112 years irrig.	8.6 g C kg ⁻¹ soil
							After 113 years irrig.	7.9 g C kg ⁻¹ soil
							Uncultivated	6.7 g C kg ⁻¹ soil
Su et al. (2010)	China Gansu province	Arid/117 mm/7.6 °C	Sandy loam	Cotton, wheat, com/n.r.	300–400 kg N ha ⁻¹ year ⁻¹ 90–150 kg P ₂ O ₅ ha ⁻¹ year ⁻¹ 60–90 kg K ₂ O ha ⁻¹ year ⁻¹ 3,000–6,000 kg farmyard manure ha ⁻¹ year ⁻¹ n.r.	40	After 5 years irrig. wheat/barley + 5 years corn	10.1 g C kg ⁻¹ soil ^a
							After 5 years irrig. wheat/barley + 5 years alfalfa	7.8 g C kg ⁻¹ soil ^a
							After 6 years irrig. wheat/barley + 4 yr robinia	12.0 g C kg ⁻¹ soil ^a
							After 0 years irrig.	0.90 g C kg ⁻¹ soil
							After 3 years irrig.	1.34 g C kg ⁻¹ soil
							After 5 years irrig.	2.36 g C kg ⁻¹ soil
							After 10 years irrig.	2.18 g C kg ⁻¹ soil ^a
							After 14 years irrig.	3.74 g C kg ⁻¹ soil ^a
							After 23 years irrig.	4.47 g C kg ⁻¹ soil ^a
							After 30 years irrig.	4.29 g C kg ⁻¹ soil ^a
Fallahzade and Hajabbasi (2012)	Iran Abarkooh Plain	Arid/60 mm	Clay-loam	Wheat, alfalfa	n.r.	30	Desert wheat irrig. alfalfa irrig.	1.28 g C kg ⁻¹ soil 5.06 g C kg ⁻¹ soil ^a 7.10 g C kg ⁻¹ soil ^a
							Grain sorghum monoculture CT non-irrig.	5.5 g SOM kg ⁻¹ soil
							NT non-irrig.	6.4 g SOM kg ⁻¹ soil
							CT irrig.	6.1 g SOM kg ⁻¹ soil
							NT irrig.	7.2 g SOM kg ⁻¹ soil
							Wheat monoculture CT non-irrig.	7.6 g SOM kg ⁻¹ soil
							NT non-irrig.	6.7 g SOM kg ⁻¹ soil
							CT irrig.	7.6 g SOM kg ⁻¹ soil
							NT irrig.	8.2 g SOM kg ⁻¹ soil
Bordovsky et al. (1999)	USA Texas	Semiarid/n.r.	Loamy sand	Grain, sorghum, wheat/NT, CT	n.r.	10	Grain sorghum monoculture CT non-irrig.	5.5 g SOM kg ⁻¹ soil
							NT non-irrig.	6.4 g SOM kg ⁻¹ soil
							CT irrig.	6.1 g SOM kg ⁻¹ soil
							NT irrig.	7.2 g SOM kg ⁻¹ soil
							Wheat monoculture CT non-irrig.	7.6 g SOM kg ⁻¹ soil
							NT non-irrig.	6.7 g SOM kg ⁻¹ soil
							CT irrig.	7.6 g SOM kg ⁻¹ soil
							NT irrig.	8.2 g SOM kg ⁻¹ soil

Table 2 (continued)

Reference	Location	Climate/annual precipitation/mean annual temperature	Soil	Crops/tillage	Fertilization	Years under irrigation	C-content
Ellmer and Baumecker (2002)	Germany Brandenburg Thyrow	Humid/495 mm/8.9 °C	Low silty sand	Orchard grass, potato, winter barley, flax, winter rye/n.r.	0 kg N ha ⁻¹ year ⁻¹ 72 kg N ha ⁻¹ year ⁻¹ 144 kg N ha ⁻¹ year ⁻¹ + removal of harvest residues	32	0 kg N ha ⁻¹ irrig. 72 kg N ha ⁻¹ irrig. 144 kg N ha ⁻¹ irrig. 144 kg N ha ⁻¹ removal residues irrig. 0 kg N ha ⁻¹ non-irrig. 144 kg N ha ⁻¹ non-irrig. 72 kg N ha ⁻¹ non-irrig. 144 kg N ha ⁻¹ removal residues non-irrig. Irrig. Non-irrig. (0–21 cm soil depth) Irrig. Non-irrig. (0–21 cm soil depth)
Presley et al. (2004)	USA Kansas Richfield	Semiarid/375–500 mm	Silty loam	Corn, soybean/n.r.	n.r.	28–30	5.78 g C kg ⁻¹ soil 6.31 g C kg ⁻¹ soil 6.28 g C kg ⁻¹ soil 4.96 g C kg ⁻¹ soil 5.63 g C kg ⁻¹ soil 6.19 g C kg ⁻¹ soil 6.13 g C kg ⁻¹ soil 4.84 g C kg ⁻¹ soil 23.8 g C kg ⁻¹ soil 25.1 g C kg ⁻¹ soil
	USA Kansas Keith						19.8 g C kg ⁻¹ soil 22.1 g C kg ⁻¹ soil
Dersch and Bohm (2001)	Austria Marchfeld	Humid/540 mm/9.1 °C	Loamy sand	Cereals, sugar beet, oil crops/n.r.	N, P, K in four steps zero medium optimal excessive	27	4,930 g C m ⁻² 5,030 g C m ⁻² 5,310 g C m ⁻² 5,040 g C m ⁻² 4,640 g C m ⁻² 4,800 g C m ⁻² 4,970 g C m ⁻² 4,940 g C m ⁻² 3,224 g C m ⁻² 3,106 g C m ⁻² 3,114 g C m ⁻² 3,281 g C m ⁻² 3,177 g C m ⁻²
De Bona et al. (2008)	Brazil Rio Grande do Sul	Humid/1446 mm summer droughts possible	Sandy loam	Oat, sweet pea/n.r.	n.r.	8	Irrig. no N Irrig. medium N Irrig. optimal N Irrig. excessive N Native vegetation CT irrig. CT non-irrig. NT irrig. NT non-irrig. (0–25 cm soil depth)
Getaneh et al. (2007)	Ethiopia Oromia Wollega-district	Subhumid–humid	Sandy clayey loam clayey loam clay	n.r./n.r.	n.r.	11–60	0.8 % lower contents of SOC in the irrigated variants

CT conventional tillage, irrig. irrigated, non-irrig. non-irrigated, NT no tillage or reduced tillage, n.r. not reported, SOC soil organic carbon, SOM soil organic matter

^a Significant difference between irrigated and non-irrigated variants

Gillabel et al. (2007) also observed higher soil carbon contents in an arid region in fields irrigated for 30 years in the USA. Irrigation led to a better plant growth and thus to a high input of organic carbon by harvest residues. Also, in this case, the highest soil organic carbon contents were obtained in the plots under natural vegetation. After 30 years of investigations, Entry et al. (2004) found higher soil carbon contents in irrigated arable land compared with non-irrigated plots and natural steppe vegetation in a semi-arid to arid region in the USA. Here, the soil organic carbon contents on plots with natural vegetation were lower than on plots under cultivation.

Similar effects were detected by Wu et al. (2008) who analyzed data from 50 to 90 years of experiments in the USA comparing irrigated arable land with native vegetation in an arid and semiarid climate. They found that the carbon content was significantly higher in the irrigated arable land after 30 years. Li et al. (2006) investigated changes in soil carbon content in an arid region of China after converting desert to irrigated arable land. Some years after conversion, the soil carbon contents in all crop rotations studied were higher than under desert vegetation. The large difference in biomass production between desert and irrigated arable land was reported to be probably the main reason for the fast increase in soil organic carbon. Similar results were found by Li et al. (2009) and Su et al. (2010). A significant difference in soil organic carbon content was observed 10 years after the change from uncultivated dry land to irrigated and fertilised cropland. Irrigation and fertilisation led to a strong increase in plant growth and, thus, to a higher input of organic matter into the soil. Similar results were reported by Fallahzade and Hajabbasi (2012) whose investigations were conducted in Iran to analyse the effects of converting desert to irrigated cropland. In contrast, Bordovsky et al. (1999) found only slightly higher contents of soil organic matter 10 years after implementation of irrigation in a semiarid region in the USA.

The situation is different if precipitation is higher or soils have higher initial soil organic carbon content. In Germany (humid climate) (Fig. 3), Ellmer and Baumecker (2002) found slightly but not significantly higher soil organic carbon contents after 32 years of irrigation (Table 2). Presley et al. (2004) reported that no differences in the soil organic carbon content were observed on soils with higher initial contents after 28 to 30 years of irrigation on plots in a semiarid region the USA. After 27 years of field trials in Austria (humid climate), Dersch and Bohm (2001) did not find significant differences in the soil organic carbon content between irrigated and non-irrigated plots. The authors assume increased mineralisation under irrigation to be the reason. Mineralisation decreased during dry periods in the non-irrigated plots while it was not reduced under irrigation. The same effect seems to predominate in the investigations

Fig. 3 Long-term irrigation field trial in Germany

of De Bona et al. (2008). They also found no significant differences in soil organic carbon contents between irrigated and non-irrigated plots after 10 years of investigation in Brazil.

Getaneh et al. (2007) report a slight and not significant decrease in soil organic carbon from plots in a subhumid to humid region of Ethiopia. In this experiment, over 11 to 60 years, the high cultivation intensity and the complete removal of harvest residues are crucial. The increased soil moisture under irrigation enhanced microbial activity, and the input of organic matter was too low to compensate the increased decomposition.

In summary, among the 14 long-term field experiments shown in Table 2, there were eight cases where significantly higher soil organic carbon contents under irrigation were obtained. All of these investigations were carried out on arid or desert sites. Especially on desert or arid sites with low precipitation, scanty natural vegetation and low initial soil organic carbon contents irrigation led to increases in soil organic carbon. In semiarid and arid regions with a better natural plant growth and higher precipitation, soil organic carbon contents of irrigated and non-irrigated arable land were generally lower than those of sites with natural vegetation. However, among the plots used as arable land, irrigation led to higher soil organic carbon contents compared with non-irrigation.

No significant differences between irrigated and non-irrigated plots or slight decreases in soil organic carbon contents were found in six cases. These investigations were conducted in wetter regions like Germany, Austria and Brazil and on arable soils with higher initial soil organic carbon content. The potential for decomposition of soil organic carbon is higher in these regions. Ogle et al. (2005) investigated the loss of soil organic matter after change from natural vegetation to arable land in different climates. The largest difference was found in the humid tropics. The differences between soil organic matter contents under natural vegetation and arable land decrease with

increased aridity and decreased mean annual temperature. The analysis of the 14 long-term investigations has shown a similar result. Figure 4 shows a summary of all 14 investigations, ordered by climate and land use. The largest changes were found on irrigated desert soil with an average increase in soil organic carbon of 242.6 %. In contrast, the average increase in regions with higher precipitation was smaller, about 17 to 25 % in arid or semiarid climates. In some cases, soil organic carbon even decreased. In humid climates, irrigation showed a minimal effect. The mean increase in all investigations of this climate was about 2.0. Also, the duration of experiments plays an important role in investigations on the development of soil organic carbon contents. In order to study significant changes in soil organic carbon, long-term experiments are necessary (Li et al. 1997). Results will be more meaningful with an increasing duration of investigations on equal terms.

In the reviewed investigations, further management factors like tillage and fertilisation were also varied. Hence, the observed effects on soil organic carbon content might be the consequence of interaction of irrigation and other management factors. These factors are discussed below.

3.2 Interaction of irrigation and N fertilisation

When considering the effects of irrigation, possible interactions with other agrotechnical activities have to be regarded. One of such activity is N fertilisation.

Many investigations show that N fertilisation leads to higher biomass generation and to an increase in root and harvest residues. Thus, N fertilisation may contribute to increase the soil organic carbon content in arable land (Liu and Greaver 2009; Schlesinger 2000). On the other hand, N fertilisation may lead to a decrease in the soil carbon/nitrogen ratio and hence to a higher decomposability of soil organic matter (Li et al. 2009).

Field experiments that allow for observing the interaction between irrigation and N fertilisation are rare. Among the investigations listed in Table 2, only Ellmer and Baumecker (2002) and Dersch and Böhm (2001) include varying N fertilisation rates on irrigated and non-irrigated plots. Both report that plots with N fertilisation have higher soil organic carbon contents than plots without N fertilisation at the same intensity of irrigation. As previously mentioned, they did not observe additional increases in soil organic carbon contents by irrigation.

Fig. 4 Changes (mean, minimum maximum) in soil organic carbon content under irrigation compared with non-irrigated conditions and native vegetation in different climates, based on 14 long-term investigations (references: 1=Denef et al. (2008); 2=Gillabel et al. (2007); 3=Entry et al. (2004); 4=Wu et al. (2008); 5=Li et al. (2006); 6=Li et al.

(2009); 7=Su et al. (2010); 8=Fallahzade and Hajabbasi (2012); 9=Bordovsky et al. (1999); 10=Ellmer and Baumecker (2002); 11=Presley et al. (2004); 12=Dersch and Böhm (2001); 13=De Bona et al. (2008); 14=Getaneh et al. (2007))

3.3 Interaction of irrigation and tillage

The intensity of tillage may affect the soil organic carbon content remarkably due to its effects on soil physical and biological conditions. Bulk density, pore volume and, thus, aeration and water infiltration are strongly influenced (Amézketa 1999; Šimon et al. 2009; Stubbs et al. 2004). With regard to the combined impact of tillage and irrigation, two effects are relevant. One is the independent effect of tillage intensity on soil organic carbon. The second effect results from the influence of tillage intensity on water productivity.

Conservation tillage is well-known to be humus-preserving and leads to soil organic matter accumulation as a rule (Rusu et al. 2008). While conventional tillage causes a strong disaggregation and aeration of soil, thus supporting the decomposition of organic matter by increased oxygen availability, the impact on soil structure is strongly reduced under conservational tillage and especially under no-till. Harvest residues are shallowly incorporated into the soil or remain on the soil surface. The increasing amount of organic matter protects the soil against wind and water erosion and stimulates the edaphon (Šimon et al. 2009; Stubbs et al. 2004).

Reduced or no-tillage contribute as well to increasing water productivity, i.e., the amount of crop output generated per unit water input. Both reduced and no tillage increases the soil's water storage capacity due to humus preservation and decreased evaporation from soil. In the case of no-till, additional air moisture can be absorbed into the mulch cover. Altogether, reduced tillage and no-till may increase biomass generation per unit of water input via precipitation and irrigation and, thus, enlarge the potential positive effect of irrigation on soil carbon sequestration (Drastig et al. 2011; Rusu et al. 2008).

Consequently, the combination of irrigation and reduced tillage is assumed to have a larger potential to increase soil organic carbon contents than irrigation in combination with conventional tillage (Martens et al. 2005). This is confirmed by the results of Entry et al. (2004) and Bordovsky et al. (1999) (Table 2). Significantly higher soil organic carbon contents were found under irrigation and reduced tillage compared with plots under irrigation and conventional tillage. In contrast, De Bona et al. (2008) did not observe significant differences between plots under conventional and no-tillage with or without irrigation.

3.4 Effects of irrigation on soil aggregate stability and soil biota

The integration of soil organic matter into microaggregates (50 to 250 µm diameter) and macroaggregates (>250 µm diameter) protect it against decomposition. Carbon

compounds are difficult to disrupt when bound in the matrix of silt and clay particles in connection with solid chemical exudates (Blanco-Canqui and Lal 2004; Gillabel et al. 2007). Kong et al. (2005) consider the content of microaggregates an ideal indicator for the carbon sequestration potential of agricultural soils since additional carbon inputs are mainly fixed in these microaggregates. Other results accentuate the importance of macroaggregates for carbon sequestration. Investigations of Degens and Sparling (1995) have shown that organic carbon bound in macroaggregates was not increasingly decomposed after repeated wetting and consequently intensified microbial activity.

Soil moisture variation and hence irrigation has an important influence on soil aggregate building (Amézketa 1999). On the one hand, it is assumed that the alternation between drying and wetting has a negative impact on the stability of macroaggregates (Lehrsch et al. 1991; Mulla et al. 1992). On the other hand, investigations have shown that a continued change between wetting and drying has led to an increase in water-stable aggregates (Utomo and Dexter 1982; Dexter 1988; Barzegar et al. 1995). The effect seems to depend on soil type and soil composition. Pore volume, pore diameter, contents of clay and organic matter are important factors (Amézketa 1999; Six et al. 2004). The velocity of moisture infiltration in soil aggregates may be relevant as well (Barzegar et al. 1995). If moisture infiltrates slowly, there is little effect on soil aggregate stability in most cases. In contrast, a rapid moisture infiltration may lead to disaggregation. This depends on the water saturation of the soil in particular (Amézketa 1999).

Likewise, the intensity of irrigation may affect aggregate building and therefore the soil organic carbon content. In their investigations on deficit irrigation, Blanco-Canqui et al. (2010) found that an increase in the amount of applied irrigation water led to more soil aggregates with a diameter over 0.5 mm, while aggregates under 0.5 mm decreased. Also, the soil organic carbon content increased with higher amounts of applied water. Besides the rate and duration, the system of irrigation may also influence structural stability. Furrow and flood irrigation led to disaggregation by compression of entrapped air due to rapid soil wetting (Amézketa 1999). In addition, sprinkler irrigation may influence aggregate forming at the soil surface. Water drops may increase the aggregate breakdown on the soil surface by their impact forces. This depends on drop size and fall height (Shainberg et al. 1992).

The effects of irrigation on mesofauna are relevant as well with respect to aggregate building.

The important role of soil organisms like earth worms for building clay-humus complexes makes them essential for long-term carbon storage in soils (Pulleman et al. 2005). Amador et al. (2005) registered that earth worms (*Lumbricus terrestris* L.) decompose more organic matter

in a wet soil than under dry conditions. Also, the amounts of springtails (*Collembola*), mites (*Acari*) and nematodes (*Nematoda*) are higher under irrigation (Lindberg et al. 2002).

Soil-borne fungi are essential for aggregate building. Mycorrhiza fungi, particularly, act in multifaceted ways on the formation of soil aggregates. This occurs primarily by the growth of hyphae, which supports the complex building of microaggregates and by the excretion of secondary synthesis products. The secondary synthesis products could act directly on soil particles or influence other soil living organisms (Rilling and Mummy 2006). The effect of irrigation on mycorrhiza fungi has not been definitely ascertained. Mitra et al. (2006) did not find a significant increase of mycorrhiza infections on wheat roots under irrigation. Rillig et al. (2001) found that increased soil moisture affected hyphae length negatively.

Another mechanism to affect soil biota via irrigation might be the cooling of the soil surface. Investigations in Mongolia conducted by Mariko et al. (2007) showed that irrigation led to a fast decrease in soil surface temperature from 30 °C to 20 °C and a short-term increase in CO₂ generation, which indicates higher microbial activity. Optimal soil temperature for microorganisms depends on the specific soil microbial communities. O'Connel (1990) sees the optimum at 30 °C, while Thierron and Laudelout (1996) found the highest microbial activity between 20 °C and 25 °C. In hot climates, where the temperatures of soil surface are often high, irrigation may support microbial activity not only by providing moisture but also by cooling the soil. This may lead to an increased decomposition of soil organic matter.

3.5 Summary on irrigation and soil organic carbon

The results of long term-investigations show that the effect of irrigation strongly depends on climate and initial soil organic carbon content. Positive effects of irrigation on soil organic carbon become less pronounced at higher initial soil organic carbon contents and higher precipitation. Desert soils with very low soil moisture contents and minimal carbon contents have a lower natural activity of soil biota. Inputs of carbon by the cultivation of crops exceed the microbial decomposition. Soils with higher initial carbon contents and higher soil moisture offer better living conditions for microorganisms and usually have a higher natural activity of soil biota. In these soils, there is often a balance between carbon input and carbon decomposition. Therefore, the effect of irrigation may be twofold. Irrigation, while enhancing the carbon input through larger amounts of plant residues, could additionally improve the living conditions for microorganisms, causing higher decomposition rates. Thus, the increased input of carbon could be decomposed

completely. A further reduction in soil organic carbon content is even possible. The combination of irrigation with other agronomic management factors also influences the development of soil organic carbon content. Nitrogen fertilisation promotes plant growth but may lead to a change in the carbon/nitrogen ratio and hence to a higher decomposability of soil organic carbon. Tillage influences soil aeration and soil structure and hence the decomposability of soil organic carbon and the activity of soil living microorganisms. In addition to increased input of carbon by improved plant growth, irrigation shows direct effects on soil aggregate building and thus on the ability of soil to fix organic carbon long-term.

4 Irrigation and nitrous oxide emissions

4.1 Overview of experiments

There are few investigations directly comparing N₂O emissions from irrigated and non-irrigated land. An overview of eight field experiments carried out in different regions of the world is given in Table 3. In most cases, N₂O emissions increased under irrigation. Investigations carried out in Finland by Simojoki and Jaakkola (2000) show that N₂O emissions were significantly higher under irrigation, both with and without N fertilisation. These investigations demonstrate that the availability of reactive nitrogen compounds has an essential influence on the amount of N₂O emissions. The N₂O emissions from a soil under vegetation were clearly lower than from a fallow soil without vegetation. Plants take up nitrogen from soil and, thus, reduce the nitrification and denitrification potential. Liu et al. (2011), who carried out investigations in China, likewise observed that irrigation events led to higher N₂O emissions only in combination with adequate nitrogen availability. Comparing irrigated wheat and maize fields with and without N fertilisation, higher N₂O emissions were obtained from the fertilised fields.

Livesly et al. (2010), who investigated the effects of fertilisation, irrigation and mulching on N₂O emissions from urban lawns in Australia, report a continuous increase in N₂O emissions under weekly irrigation, whereas the increase after fertilising was short-term. They conclude that the effect of regular irrigation is higher than that of N fertilisation.

Horváth et al. (2010) report an increase in N₂O emissions from irrigated pastures in Hungary. This effect was observed in 2 of 3 years, in particular, when precipitation was low and the differences in water-filled pore volume between the irrigated and non-irrigated plots were large.

Scheer et al. (2008) investigated cotton fields in Uzbekistan. They found that a reduced application of water

Table 3 Overview of experiments to investigate the influences of irrigation on nitrous oxide emissions

Reference	Location	Annual precipitation/ mean annual temperature	Soil	Crops	Fertilization	N ₂ O emission values
Simojoki and Jaakkola (2000)	Finland Jokioinen	n.r.	Loamy clay pH-value, 5.8; total nitrogen content, 0.2 %	Fallow	0 kg N ha ⁻¹	2,030 g N ₂ O-N ha ⁻¹ Irrig.
					103 kg N ha ⁻¹	1,680 g N ₂ O-N ha ⁻¹ Non-irrig.
						3,540 g N ₂ O-N ha ⁻¹ Irrig.:
						2,030 g N ₂ O-N ha ⁻¹ Non-irrig.
					Barley undersown grass	850 g N ₂ O-N ha ⁻¹ Irrig.
Liu et al. (2011)	China Shanxi Province	562 mm 14.8 °C	Loamy clay pH-value, 8.7; total N content, 1.12 g N kg ⁻¹ soil	Wheat, maize	103 kg N ha ⁻¹	120 g N ₂ O-N ha ⁻¹ Non-irrig.
						1,150 g N ₂ O-N ha ⁻¹ Irrig.
					Wheat, 0 kg N ha ⁻¹	570 g N ₂ O-N ha ⁻¹ Non-irrig.
					Maize, 220 kg N ha ⁻¹	20.3 µg N ₂ O-N m ⁻² h ⁻¹ Irrig.
					Maize, 210 kg N ha ⁻¹	78.5 µg N ₂ O-N m ⁻² h ⁻¹ Irrig.
Livesley et al. (2010)	Australia Melbourne	681 mm n.r.	Urban soil pH-value, 6.0–6.2	Lawn	0 kg N ha ⁻¹	27 µg N ₂ O-N m ⁻² h ⁻¹ Irrig.
					0 kg N ha ⁻¹	18 µg N ₂ O-N m ⁻² h ⁻¹ Non-irrig.
					48 kg N ha ⁻¹	26 µg N ₂ O-N m ⁻² h ⁻¹ Irrig.
					48 kg N ha ⁻¹	15 µg N ₂ O-N m ⁻² h ⁻¹ Non-irrig.
					0 kg N ha ⁻¹	1,260 g N ₂ O-N ha ⁻¹ year ⁻¹ 2002 irrig.
Horváth et al. (2010)	Ungarn Gödöllő	582 mm 10.1 °C	Sandy loess soil N content, 0.2 %	Grass		1,560 g N ₂ O-N ha ⁻¹ year ⁻¹ 2003 irrig.
						780 g N ₂ O-N ha ⁻¹ year ⁻¹ 2004 irrig.
						940 g N ₂ O-N ha ⁻¹ year ⁻¹ 2002 non-irrig.
						920 g N ₂ O-N ha ⁻¹ year ⁻¹ 2003 non-irrig.
						750 g N ₂ O-N ha ⁻¹ year ⁻¹ 2004 non-irrig.
Scheer et al. (2008)	Uzbekistan	<100 mm 13.6 °C	Silty loam	Cotton	250 kg N ha ⁻¹	Irrig. (applied water: 463 mm) 102 µg N ₂ O-N m ⁻² h ⁻¹
					250 kg N ha ⁻¹	Irrig. (applied water: 373 mm) 55 µg N ₂ O-N m ⁻² h ⁻¹
					0 kg N ha ⁻¹	2004: irrig. 9,000 g N ₂ O-N ha ⁻¹
					0 kg N ha ⁻¹	2004: non-irrig. 9,800 g N ₂ O-N ha ⁻¹
					0 kg N ha ⁻¹	2005: irrig. 26,700 g N ₂ O-N ha ⁻¹
Rochette et al. (2010)	Canada	2004: 517 mm 2005: 897 mm n.r.	Limnic haplochromist organic soil	Carrots, onions, celery, lettuce	0 kg N ha ⁻¹	2005: non-irrig. 23,300 g N ₂ O-N ha ⁻¹
					50 kg N ha ⁻¹	2004 irrig. 10,200 g N ₂ O-N ha ⁻¹
					50 kg N ha ⁻¹	2004: non-irrig. 11,800 g N ₂ O-N ha ⁻¹
					50 kg N ha ⁻¹	2005: irrig. 32,200 g N ₂ O-N ha ⁻¹
					50 kg N ha ⁻¹	2005: non-irrig. 40,200 g N ₂ O-N ha ⁻¹
					100 kg N ha ⁻¹	2004 irrig. 7,900 g N ₂ O-N ha ⁻¹
					100 kg N ha ⁻¹	2004: non-irrig. 5,000 g N ₂ O-N ha ⁻¹
					100 kg N ha ⁻¹	2005: irrig. 31,300 g N ₂ O-N ha ⁻¹
					100 kg N ha ⁻¹	2005: non-irrig. 13,000 g N ₂ O-N ha ⁻¹
					150 kg N ha ⁻¹	2004 irrig. 7,500 g N ₂ O-N ha ⁻¹
					150 kg N ha ⁻¹	2004: non-irrig. 3,600 g N ₂ O-N ha ⁻¹
					150 kg N ha ⁻¹	2005: irrig. 22,600 g N ₂ O-N ha ⁻¹
					150 kg N ha ⁻¹	2005: non-irrig. 14,600 g N ₂ O-N ha ⁻¹

Table 3 (continued)

Reference	Location	Annual precipitation/ mean annual temperature	Soil	Crops	Fertilization	N ₂ O emission values
Liu et al. (2008)	Mongolia	335 mm 0.7 °C	calicic Chernozem pH value, 6.7 N-content, 2.28–3.16 mg N g ⁻¹ soil	Pasture	n.r.	2004; non-irrig.: 1.2 µg N ₂ O-N m ⁻² h ⁻¹ 2005; irrig.: 0.9 µg N ₂ O-N m ⁻² h ⁻¹ 2005; non-irrig.: 0.7 µg N ₂ O-N m ⁻² h ⁻¹
Wulf et al. (1999)	Kenya Turkana district	302 mm 30 °C	Sandy loam pH-value, 8.6 N-content, 0.4 g N kg ⁻¹ soil	Thornbush savanna for rainfed treatment <i>Acacia saligna</i> for irrigated treatment	n.r.	For 1 month: 64 g N ₂ O-N ha ⁻¹ Non-irrig.: 54 g N ₂ O-N ha ⁻¹ Irrig.

irrig, irrigated, MT minimum tillage, non-irrig, non-irrigated, n.r. not reported

by less frequent irrigation events could lead to lower N₂O emissions as a result of lower soil moisture.

Rochette et al. 2010, who investigated the effects of irrigation and N fertilisation on a drained organic soil in a 2-year study in Canada, obtained higher N₂O emissions on irrigated plots where nitrogen fertiliser was applied. Moreover, higher amounts of precipitation and irrigation water in the second year led to increased N₂O emissions.

In contrast, no significant differences between N₂O emissions from irrigated and non-irrigated land are reported from two further field experiments. Liu et al. (2008) did not observe significant increases in N₂O emissions from irrigated pastures in the Mongolian steppe. They consider the reason to be the low content of reactive nitrogen compounds in the soil. Also, Wulf et al. (1999), who conducted their investigations in Kenya, found that the amount of N₂O emissions did not differ significantly from irrigated and non-irrigated plots. After irrigation or precipitation events, increases of N₂O emissions were observed in both variants. Under irrigation, the increase was slightly smaller. This is explained by the availability of reactive nitrogen compounds, since the nitrate contents in the irrigated plots were lower. Without irrigation, the microbial activity and, hence, mineralisation, nitrification and denitrification processes were reduced in the dry period. These processes were enhanced rapidly with an increase in soil moisture after precipitation and caused a strong release and conversion of reactive nitrogen compounds. Living conditions for microorganisms are more favourable and constant under irrigation so that mineralisation and conversion proceed more steadily (Wulf et al. 1999).

4.2 Influence of soil water content and soil aeration on nitrous oxide emissions

N₂O arises as an intermediate in denitrification and nitrification. Both processes are strongly influenced by soil water content and soil aeration. Low oxygen contents and anaerobic conditions caused by high soil water content can lead to intensification of denitrification processes (Beare et al. 2009). N₂O emissions arising from denitrification processes increase strongly at a water filled pore volume over 70 % (Amha and Bohne 2011; Ruser et al. 2006). However, a nearly complete filling of the pore volume over a long time may lead to a decrease of N₂O emissions since, under strict anaerobic conditions, the intermediate in denitrification N₂O is completely deoxidised to N₂ (Huang et al. 2007).

Nitrification is the primary reason for N₂O formation under oxygen availability. A water-filled pore volume of 30 % to 60 % supports N₂O release via nitrification (Kavdir et al. 2008; Horváth et al. 2010). Also, the kind of tillage may affect soil aeration and, thus, the processes generating N₂O. There are many investigations regarding

the impact of reduced or minimum tillage on N₂O release, reporting that reduced tillage leads to higher N₂O emissions in comparison to conventional tillage (Abdalla et al. 2010; Ball et al. 2008; Rochette 2008; Smith et al. 2000). In contrast, some results have shown higher N₂O emissions from soils under conventional tillage (Chatskikh and Olesen 2007; Gregorich et al. 2007; Mutegi et al. 2010). The increased N₂O emissions under reduced or minimum tillage can be explained by a reduced gas exchange and resultant intensified denitrification processes (Abdalla et al. 2010; Ball et al. 2008; Rochette 2008). Also, increased CO₂ production from microbial respiration caused by the accumulation of organic matter in the top soil under no-till leads to a decrease in oxygen concentration (Abdalla et al. 2010). The reason for higher N₂O emissions under conventional tillage could be a strong disaggregation and enhanced soil aeration. Hence, the living conditions of aerobic nitrifying bacteria may be improved (Elmi et al. 2003; Gregorich et al. 2007).

4.3 Nitrous oxide emissions under irrigation in combination with nitrogen fertilising

In those investigations where higher N₂O emissions were observed under irrigation, the availability of nitrogen was

change in amount of N₂O emissions (%)

Fig. 5 Changes (mean, minimum maximum) in N₂O emissions under irrigation compared with non-irrigated conditions, based on six investigations (references: 1=Simojoki and Jaakkola (2000); 2=Livesley et

Fig. 6 Drip irrigation, partial wetting of soil

most often the more relevant factor (Fig. 5). An average increase of about 87 % was estimated under irrigation in combination with nitrogen fertilising. Without nitrogen, fertilising the estimated average increase was about 7 %. But exceptions with high increases in N₂O emissions under irrigation without nitrogen fertilisation are possible. According to Robertson et al. (2000), the primary reason

al. (2010); 3=Horváth et al. (2010); 4=Rochette et al. (2010); 5=Liu et al. (2008); 6=Wulf et al. (1999))

for high N_2O emissions is the existence of a high concentration of reactive nitrogen compounds in the soil. Irrigated arable land as a rule is intensively farmed and, therefore, has a high fertiliser input. Consequently, it offers a high potential for N_2O formation (Ellert and Janzen 2008). It has been affirmed in numerous investigations that an increase in nitrogen fertilising leads to higher N_2O emissions (Clayton et al. 1997; Hao et al. 2001; Lin et al. 2011; Liu and Greaver 2009; Yao et al. 2010). An increase in the water-filled pore volume can intensify this effect additionally (Abbasi and Adams 2000). Significantly higher emissions were observed in several investigations if precipitation and irrigation events proceeded after the application of nitrogen fertiliser (Dobbie and Smith 2003; Liu et al. 2010; Hutchinson and Mosier 1979; Scheer et al. 2008).

4.4 Effects of irrigation systems on nitrous oxide emissions

The irrigation technology and hence the distribution of water in the soil may affect the amount of N_2O emissions. Sanchez-Martin et al. (2010) and Kallenbach et al. (2010) found that N_2O emissions were lower under drip irrigation than under furrow irrigation. According to Kallenbach et al. (2010), the reason is the partial wetting of soil under drip irrigation (Fig. 6). The soil pore volume is filled with water only at the spots where the drippers are located. Thus, denitrification proceeds on a considerably smaller area in comparison to furrow irrigation. Nelson and Terry (1996) compared N_2O emissions under sprinkler irrigation and furrow irrigation. They observed strong changes in soil physical parameters under furrow irrigation. Increases in bulk density and crusting led to a decrease in aeration and, therefore, to an intensification of denitrification processes. At the same time, crusting may prevent N_2O from directly escaping from the soil to the atmosphere, so that N_2O may possibly deoxidise to N_2 (Mahmood et al. 2008).

4.5 Summary on irrigation and N_2O emissions

In summary, general statements on the effect of irrigation on the amount of N_2O emissions cannot yet be derived. In various cases, an increase in soil moisture intensified nitrification and denitrification processes and, thus, enhanced N_2O emissions. Direct comparisons between the amount of N_2O emissions from irrigated and non-irrigated land are rare. Those available did not always find higher N_2O emissions from the irrigated plots. In most cases, the availability of reactive nitrogen compounds seems to be more important for the release of N_2O . Different technologies and intensities of irrigation may also influence N_2O emissions.

5 Conclusions

In most cases, irrigation leads to significantly higher soil organic carbon contents in arid soils and deserts than non-irrigated conditions. For humid climates and on soils with higher initial soil organic carbon content, irrigation has no significant effects on soil organic carbon contents. N_2O emissions as a rule increase under irrigation when reactive nitrogen compounds are adequately available. Soil carbon sequestration and N_2O release are affected by environmental factors and management measures interacting with irrigation. It is difficult to estimate the effects of some of these factor combinations. The net effects of irrigation on greenhouse gas emissions have not been estimated yet. A full balance is necessary to assess the impact of irrigation on net greenhouse gas emissions.

References

- Abbasi MK, Adams WA (2000) Gaseous N emission during simultaneous nitrification–denitrification associated with mineral N fertilization to a grassland soil under field conditions. *Soil Biol Biochem* 32:1251–1259. doi:10.1016/S0038-0717(00)00042-0
- Abdalla M, Jones M, Williams M (2010) Simulation of N_2O fluxes from Irish arable soils: effect of climate change and management. *Biol Fert Soils* 46:247–260. doi:10.1007/s00374-009-0424-5
- Alexandratos N, Bruinsma J (2012) World agriculture towards 2030/2050: the 2012 revision. ESA Working paper, Rome, FAO, No. 12–03
- Amador JA, Görres JH, Savin MC (2005) Role of soil water content in carbon and nitrogen dynamics of *Lumbricus terrestris* L. burrow soil. *Appl Soil Ecol* 28:15–22. doi:10.1016/j.apsoil.2004.06.009
- Amézketa E (1999) Soil aggregate stability: a review. *J Sustain Agric* 14:83–151
- Amha Y, Böhne H (2011) Denitrification from the horticultural peats: effects of pH, nitrogen, carbon and moisture contents. *Biol Fert Soils* 47:293–302. doi:10.1007/s00374-010-0536-y
- Ball BC, Crichton I, Horgan GW (2008) Dynamics of upward and downward N_2O and CO_2 fluxes in ploughed or no-tilled soils in relation to water-filled pore space, compaction and crop presence. *Soil Till Res* 101:20–30. doi:10.1016/j.still.2008.05.012
- Barzegar AR, Rengasamy P, Oades JN (1995) Effects of clay type and rate of wetting on the mellowing of compacted soils. *Geoderma* 68:39–40. doi:10.1016/0016-7061(95)00022-g
- Beare MH, Gregorich EG, St-Georges P (2009) Compaction effects on CO_2 and N_2O production during drying and rewetting of soil. *Soil Biol Biochem* 41:611–621. doi:10.1016/j.soilbio.2008.12.024
- Blanco-Canqui H, Lal R (2004) Mechanisms of carbon sequestration in soil aggregates. *Crit Rev Plant Sci* 23:481–504. doi:10.1080/07352680490886842
- Blanco-Canqui H, Klocke NL, Schlegel AJ, Stone LR, Rice CW (2010) Impacts of deficit irrigation on carbon sequestration and soil physical properties under no-till. *Soil Sci Soc Am J* 74:1301–1309. doi:10.2136/sssaj2009.0364
- Bordovsky DG, Choudhary M, Gerard CJ (1999) Effect of tillage, cropping, and residue management on soil properties in the Texas Rolling Plains. *Soil Sci* 164:331–340. doi:10.1097/00010694-199905000-00005
- Bremner JM (1997) Sources of nitrous oxide in soils. *Nutr Cycl Agroecosyst* 49:7–16. doi:10.1023/a:1009798022569

- Bruinsma J (2009) The resource outlook to 2050: by how much do land, water and crop yields need to increase by 2050? How to feed the world in 2050. Proceedings of a technical meeting of experts. FAO (Food and Agriculture Organization of the United Nations), Rome, Italy, <http://www.fao.org/wsfs/forum2050/wsfs-background-documents/wsfs-expert-papers/en/> Accessed 10 December 2012
- Chatskikh D, Olesen JE (2007) Soil tillage enhanced CO₂ and N₂O emissions from loamy sand soil under spring barley. *Soil Till Res* 97:5–18. doi:10.1016/j.still.2007.08.004
- Clayton H, McTaggart IP, Parker J, Swan L (1997) Nitrous oxide emissions from fertilised grassland: a 2-year study of the effects of N fertiliser from environmental conditions. *Biol Fert Soils* 25:252–260. doi:10.1007/s003740050311
- De Bona FD, Bayer C, Dieckow J, Bergamaschi H (2008) Soil quality assessed by carbon management index in a subtropical Acrisol subjected to tillage systems and irrigation. *Aust J Soil Res* 46:469–475. doi:10.1071/sr08018
- Degens BP, Sparling GP (1995) Repeated wet–dry cycles do not accelerate the mineralization of organic C involved in the macro-aggregation of a sandy loam soil. *Plant Soil* 175:197–203. doi:10.1007/BF00011355
- Denef K, Stewart CE, Brenner J, Paustian K (2008) Does long-term center-pivot irrigation increase soil carbon stocks in semi-arid agro-ecosystems? *Geoderma* 145:121–129. doi:10.1016/j.geoderma.2008.03.002
- Deng X-P, Shan L, Inanaga S, Inoue M (2005) Water-saving approaches for improving wheat production. *J Sci Food Agric* 85:1379–1388. doi:10.1002/jsfa.2101
- Dersch G, Böhm K (2001) Effects of agronomic practices on the soil carbon storage potential in arable farming in Austria. *Nutr Cycl Agroecosyst* 60:49–55. doi:10.1023/a:1012607112247
- Dexter AR (1988) Advances in characterization of soil structure. *Soil Till Res* 11:199–238. doi:10.1016/0167-1987(88)90002-5
- Dobbie KE, Smith A (2003) Nitrous oxide emissions factors for agricultural soils in Great Britain: the impact of soil water-filled pore space and other controlling variables. *Glob Change Biol* 9:204–218. doi:10.1046/j.1365-2486.2003.00563.x
- Drastig K, Prochnow A, Baumecker M, Berg W, Brunsch R (2011) Agricultural water management in Brandenburg. *Erde* 142:119–140
- Elmi AA, Madramootoo C, Hamel C, Liu A (2003) Denitrification and nitrous oxide to nitrous oxide plus dinitrogen ratios in the soil profile under three tillage systems. *Biol Fert Soils* 38:340–348. doi:10.1007/s00374-003-0663-9
- Ellert BH, Janzen HH (2008) Nitrous oxide, carbon dioxide and methane emissions from irrigated cropping systems as influenced by legumes, manure and fertilizer. *Can J Soil Sci* 88:207–217. doi:10.4141/CJSS06036
- Ellmer F, Baumecker M (2002) 65 years long-term experiments at Thyrow. Results for sustainable crop production at Sandy Soils. *Arch Acker Pfl Boden* 48:521–531. doi:10.1080/03650340215639
- Entry JA, Sojka RE, Shewmaker GE (2004) Irrigation increases inorganic carbon in agricultural soils. *Environ Manage* 33:309–317. doi:10.1007/s00267-003-9140-3
- Entry JA, Mills D, Mathee K, Jayachandran K, Sojka RE, Narasimhan G (2008) Influence of irrigated agriculture on soil microbial diversity. *Appl Soil Ecol* 40:146–154. doi:10.1016/j.apsoil.2008.03.012
- Fallahzade J, Hajabbasi MA (2012) The effects of irrigation and cultivation on the quality of desert soil in central Iran. *Land Degrad Dev* 23:53–61. doi:10.1002/ldr.1049
- FAO (Food and Agriculture Organization of the United Nations) Statistical yearbook (2010) <http://www.fao.org/economic/ess/ess-publications/ess-yearbook/ess-yearbook2010/yearbook2010-resources/en/> Accessed 22 August 2011
- Getaneh F, Deressa A, Negassa W (2007) Influence of small scale irrigation on selected soil chemical properties. Tropentag, Witzenhausen, Germany, 2007 October 9–11
- Gillabel J, Denef K, Brenner J, Merckx R, Paustian K (2007) Carbon sequestration and soil aggregation in center-pivot irrigated and dryland cultivated farming systems. *Soil Sci Soc Am J* 71:1020–1028. doi:10.2136/sssaj2006.0215
- Gregorich EG, Rochette P, St-Georges P, Mckim UF, Chan C (2007) Tillage effects on N₂O emissions from soils under corn and soybeans in Eastern Canada. *Can J Soil Sci* 88:153–161. doi:10.4141/CJSS06041
- Hao X, Chang C, Carefoot JM, Janzen HH, Ellert BH (2001) Nitrous oxide emissions from an irrigated soil as affected by fertilizer and straw management. *Nutr Cycl Agroecosys* 60:1–8. doi:10.1023/A:1012603732435
- Horváth L, Grosz B, Machon A, Tuba Z, Nagy Z, Czöbel SZ, Balogh J, Péli E, Föti SZ, Weidinger T, Pintér K, Führer E (2010) Estimation of nitrous oxide emissions from Hungarian semi-arid sandy and loess grasslands; effect of soil parameters, grazing, irrigation and use of fertilizer. *Agr Ecosyst Environ* 139:255–263. doi:10.1016/j.agee.2010.08.011
- Huang S, Pant HK, Lu J (2007) Effects of water regimes on nitrous oxide emission from soils. *Ecol Eng* 31:9–15. doi:10.1016/j.ecoleng.2007.04.001
- Hutchinson GL, Mosier AR (1979) Nitrous oxide emissions from an irrigated cornfield. *Science* 205:1125–1127. doi:10.1126/science.205.4411.1125
- Jabro JD, Sainju U, Stevens WB, Evans RG (2008) Carbon dioxide flux as affected by tillage and irrigation in soil converted from perennial forages to annual crops. *J Environ Manage* 88:1478–1484. doi:10.1016/j.jenvman.2007.07.012
- Jha PB, Singh JS, Kashyap AK (1996) Dynamics of viable nitrifier community and nutrient availability in dry tropical forest habitat as affected by cultivation and soil texture. *Plant Soil* 180:277–285. doi:10.1007/bf00015311
- Kallenbach CM, Rolston DE, Horwath WR (2010) Cover cropping affects soil N₂O and CO₂ emissions differently depending on type of irrigation. *Agr Ecosyst Environ* 137:251–260. doi:10.1016/j.agee.2010.02.010
- Kavdir Y, Hellebrand HJ, Kern J (2008) Seasonal variations of nitrous oxide emissions in relation to nitrogen fertilization and energy crop types in sandy soil. *Soil Till Res* 98:175–186. doi:10.1016/j.still.2007.11.002
- King AP, Evatt KJ, Six J, Poch RM, Rolston DE, Hopmans JW (2009) Annual carbon and nitrogen loadings for a furrow-irrigated field. *Agr Water Manage* 96:925–930. doi:10.1016/j.agwat.2009.01.001
- Kochsiek AE, Knops JMH, Walters DT, Arkebauer TJ (2009) Impacts of management on decomposition balance in irrigated and rainfed no till agricultural systems. *Agr Forest Meteorol* 149:1983–1993. doi:10.1016/j.agrformet.2009.07.004
- Kong AYY, Six J, Bryant DC, Denison RF, van Kessel C (2005) The relationship between carbon input, aggregation, and soil organic carbon stabilization in sustainable cropping systems. *Soil Sci Soc Am J* 69:1078–1085. doi:10.2136/sssaj2004.0215
- Lal R (2004) Soil carbon sequestration impacts on global climate change and food security. *Science* 304:1623–1627. doi:10.1126/science.1097396
- Lehrsch GA, Sojka RE, Carter DL, Jolley PM (1991) Freezing effects on aggregate stability affected by texture, mineralogy, and organic matter. *Soil Sci Soc Am J* 55:1401–1406
- Li C, Frolking S, Butterbach-Bahl K (2005) Carbon sequestration in arable soils is likely to increase nitrous oxide emissions, offsetting reductions in climate radiative forcing. *Clim Chang* 72:321–338. doi:10.1007/s10584-005-6791-5
- Li C, Frolking S, Crocker GJ, Grace PR, Klir J, Körchens M, Poulton PR (1997) Simulating trends in soil organic carbon in long-term experiments using the DNDC model. *Geoderma* 81:45–60. doi:10.1016/S0016-7061(97)00080-3

- Li XG, Li YK, Li FM, Ma Q, Zhang PL, Yin P (2009) Changes in soil organic carbon, nutrients and aggregation after conversion of native desert soil into irrigated arable land. *Soil Till Res* 104:263–269. doi:10.1016/j.still.2009.03.002
- Li XH, Li FM, Rengel Z, Bhupinderpal S, Wang ZF (2006) Cultivation effects on temporal changes of organic carbon and aggregate stability in desert soil of Hexi Corridor in China. *Soil Till Res* 91:22–29. doi:10.1016/j.still.2005.10.004
- Lin S, Iqbal J, Hu R, Wu J, Zhao J, Ruan L, Malghani S (2011) Nitrous oxide emissions from rape field as affected by nitrogen fertilizer management: a case study in Central China. *Atmos Environ* 45:1775–1779. doi:10.1016/j.atmosenv.2011.01.003
- Lindberg N, Engtsson JB, Persson T (2002) Effect of experimental irrigation and drought on the composition and diversity of soil fauna in a coniferous stand. *J Appl Ecol* 39:924–936. doi:10.1046/j.1365-2664.2002.00769.x
- Liu L, Greaver TL (2009) A review of nitrogen enrichment effects on three biogenic GHGs: the CO₂ sink may be largely offset by stimulated N₂O and CH₄ emission. *Ecol Lett* 12:1103–1117. doi:10.1111/j.1461-0248.2009.01351.x
- Liu CY, Holst J, Brüggemann N, Butterbach-Bahl K, Yao ZS, Han SH, Han XG, Zheng XH (2008) Effects of irrigation on nitrous oxide, methane and carbon dioxide fluxes in an inner Mongolian steppe. *Adv Atmos Sci* 25:748–756. doi:10.1007/s00376-008-0748-3
- Liu CY, Wang K, Meng S, Zheng XH, Zhou ZX, Han SH, Chen D, Yang ZP (2011) Effects of irrigation, fertilization and crop straw management on nitrous oxide and nitric oxide emissions from a wheat-maize rotation field in northern China. *Agr Ecosyst Environ* 140:226–233. doi:10.1016/j.agee.2010.12.009
- Liu CY, Zheng XH, Zhou ZX, Han SH, Wang YH, Wang K, Liang WG, Li M, Chen DL, Yang ZP (2010) Nitrous oxide and nitric oxide emissions from an irrigated cotton field in Northern China. *Plant Soil* 332:123–134. doi:10.1007/s11104-009-0278-5
- Livesly SJ, Dougherty BJ, Smith AJ, Navaud D, Wylie LJ, Arndt SK (2010) Soil-atmosphere exchange of carbon dioxide, methane and nitrous oxide in urban garden systems: impact of irrigation, fertilizer and mulch. *Urban Ecosyst* 13:273–293. doi:10.1007/s11252-009-0119-6
- Loecke TD, Robertson GP (2009) Soil resource heterogeneity in terms of litter aggregation promotes nitrous oxide fluxes and slows decomposition. *Soil Biol Biochem* 41:228–235. doi:10.1016/j.soilbio.2008.10.017
- Mahmood T, Ali R, Iqbal J, Robab U (2008) Nitrous oxide emissions from an irrigated cotton field under semiarid subtropical conditions. *Biol Fertil Soils* 44:773–781. doi:10.1007/s00374-008-0276-4
- Mariko S, Urano T, Asanuma J (2007) Effects of irrigation on CO₂ and CH₄ fluxes from Mongolian steppe soil. *J Hydrol* 333:118–123. doi:10.1016/j.jhydrol.2006.07.027
- Martens DA, Emmerich W, McLain JET, Johnsen TN (2005) Atmospheric carbon mitigation potential of agricultural management in southwestern USA. *Soil Till Res* 83:95–119. doi:10.1016/j.still.2005.02.011
- Mitra S, Singh S, Bandyopahyay SK, Singh CS (2006) Effect of irrigation, N, and bio-fertilizers on yield and nutrient uptake by wheat. *Trop Agr* 83:87–94
- Mulla DJ, Huyck LM, Reganold JP (1992) Temporal variation in soil aggregate stability on conventional and alternative farms. *Soil Sci Soc Am J* 56:1620–1624
- Mutegi JK, Munkholm LJ, Petersen BM, Hansen EM, Petersen SO (2010) Nitrous oxide emissions and controls as influenced by tillage and crop residue management strategy. *Soil Biol Biochem* 42:1701–1711. doi:10.1016/j.soilbio.2010.06.004
- Nelson SD, Terry RE (1996) The effects of soil physical properties and irrigation method on denitrification. *Soil Sci* 161:242–249. doi:10.1097/00010694-199604000-00005
- O’Connel AM (1990) Microbial decomposition (respiration) of litter in eucalypt forests of south-western Australia: an empirical model based on laboratory incubations. *Soil Biol Biochem* 22:153–160. doi:10.1016/0038-0717(90)90080-J
- Ogle SM, Breidt FJ, Paustian K (2005) Agricultural management impacts on soil organic carbon storage under moist and dry climatic conditions of temperate and tropical regions. *Biogeochemistry* 72:87–121. doi:10.1007/s10533-004-0360-2
- Phillips RL (2008) Denitrification in cropping systems at sub-zero soil temperatures. A review. *Agron Sustain Dev* 28:87–93. doi:10.1051/agro:2007045
- Potthoff M, Dyckmans J, Flessa H, Muhs A, Beese F, Joergensen RG (2005) Dynamics of maize (*Zea mays* L.) leaf straw mineralization as affected by the presence of soil and the availability of nitrogen. *Soil Biol Biochem* 37:1259–1266. doi:10.1016/j.soilbio.2004.11.022
- Presley DR, Ransom MD, Kluitenberg GJ, Finnell PR (2004) Effects of thirty years of irrigation on the genesis and morphology of two semiarid soils in Kansas. *Soil Sci Soc Am J* 68:1916–1926
- Pulleman MM, Six J, Uyl A, Marinissen JCY, Jongmans AG (2005) Earthworms and management affect organic matter incorporation and microaggregate formation in agricultural soils. *Appl Soil Ecol* 29:1–15. doi:10.1016/j.apsoil.2004.10.003
- Rillig MC, Wright SF, Kimball BA, Pinter PJ, Wall GW, Ottman MJ, Leavitt SW (2001) Elevated carbon dioxide and irrigation effects on water stable aggregates in a sorghum field: a possible role for arbuscular mycorrhizal fungi. *Glob Change Biol* 7:333–337. doi:10.1046/j.1365-2486.2001.00404.x
- Rillig MC, Mummey DL (2006) Mycorrhizas and soil structure. *New Phytol* 171:41–53. doi:10.1111/j.1469-8137.2006.01750.x
- Robertson GP, Paul EA, Harwood RR (2000) Greenhouse gases in intensive agriculture: contributions of individual gases to the radiative forcing of the atmosphere. *Science* 289:1922–1925. doi:10.1126/science.289.5486.1922
- Rochette P (2008) No-till only increases N₂O emissions in poorly-aerated soils. *Soil Till Res* 101:97–100. doi:10.1016/j.still.2008.07.011
- Rochette P, Tremblay N, Fallon E, Angers DA, Chantign MH, MacDonald JD, Bertrand N, Parent L-É (2010) N₂O emissions from an irrigated and non-irrigated organic soil in eastern Canada as influenced by N fertilizer addition. *Eur J Soil Sci* 61:186–196. doi:10.1111/j.1365-2389.2009.01222.x
- Roldan A, Salinas-Garcia JR, Alguacil MM, Caravaca F (2005) Changes in soil enzyme activity, fertility, aggregation and C sequestration mediated by conservation tillage practices and water regime in a maize field. *Appl Soil Ecol* 30:11–20. doi:10.1016/j.apsoil.2005.01.004
- Ruser R, Flessa H, Russow R, Schmidt G, Buegger F, Munch JC (2006) Emissions of N₂O, N₂ and CO₂ from soil fertilized with nitrate: effect of compaction, soil moisture and rewetting. *Soil Biol Biochem* 38:263–274. doi:10.1016/j.soilbio.2005.05.005
- Rusu T et al (2008) The influence of minimum soil tillage systems on water and humus conservation in some soils of Romania. *Agricultural and biosystems engineering for a sustainable world. International Conference on Agricultural Engineering, Hersonissos, Crete, Greece, pp 177–183, 23–25 June 2008*
- Sanchez-Martin L, Meijide A, Garcia-Torres L, Vallejo A (2010) Combination of drip irrigation and organic fertilizer for mitigating emissions of nitrogen oxides in semiarid climate. *Agr Ecosyst Environ* 137:99–107. doi:10.1016/j.agee.2010.01.006
- Scheer C, Wassmann R, Kienzler K, Ibragimov N, Eschanov R (2008) Nitrous oxide emissions from fertilized cotton (*Gossypium hirsutum* L.) in the Aral Sea Basin, Uzbekistan: influence of nitrogen applications and irrigation practices. *Soil Biol Biochem* 40:290–301. doi:10.1016/j.soilbio.2007.08.007
- Schlesinger WH (2000) Some cautions amidst optimism. *Agr Ecosyst Environ* 82:121–127
- Shainberg I, Levy GJ, Rengasamy P, Frenkel H (1992) Aggregate stability and seal formation as affected by drops impact energy

- and soil amendments. *Soil Sci* 154:113–119. doi:[10.1097/00010694-199208000-00004](https://doi.org/10.1097/00010694-199208000-00004)
- Simojoki A, Jaakkola A (2000) Effect of nitrogen fertilization, cropping and irrigation on soil air composition and nitrous oxide emissions in a loamy clay. *Eur J Soil Sci* 51:413–424. doi:[10.1046/j.1365-2389.2000.00308.x](https://doi.org/10.1046/j.1365-2389.2000.00308.x)
- Šimon T, Javůrek M, Mikanová O, Vach M (2009) The influence of tillage systems on soil organic matter and soil hydrophobicity. *Soil Till Res* 105:44–48
- Six J, Bossuyt H, Degryze S, Denef K (2004) A history of research on the link between (micro)aggregates, soil biota, and soil organic matter dynamics. *Soil Till Res* 79:7–31. doi:[10.1016/j.still.2004.03.008](https://doi.org/10.1016/j.still.2004.03.008)
- Smith P, Goulding KWT, Smith KA, Powlson DS, Smith JU, Falloon P, Coleman K (2000) Including trace gas fluxes in estimates of the carbon mitigation potential of UK agricultural land. *Soil Use Manage* 16:251–259. doi:[10.1111/j.1475-2743.2000.tb00204.x](https://doi.org/10.1111/j.1475-2743.2000.tb00204.x)
- Stubbs TL, Kennedy AC, Schillinger WF (2004) Soil ecosystem changes during the transition to no-till cropping. *J Crop Imp* 11:105–135. doi:[10.1300/J411v11n01_06](https://doi.org/10.1300/J411v11n01_06)
- Su YZ, Yang R, Liu WJ, Wang XF (2010) Evolution of soil structure and fertility after conversion of native sandy desert soil to irrigated cropland in arid region, China. *Soil Sci* 175:246–254. doi:[10.1097/SS.0b013e3181e04a2d](https://doi.org/10.1097/SS.0b013e3181e04a2d)
- Thierron V, Laudelout H (1996) Contribution of root respiration to total CO₂ efflux from the soil of a deciduous forest. *Can J Forest Res* 26:1142–1148. doi:[10.1139/x26-127](https://doi.org/10.1139/x26-127)
- Utomo WH, Dexter AR (1982) Changes in soil aggregate water stability induced by wetting and drying cycles in non-saturated soil. *J Soil Sci* 33:623–637. doi:[10.1111/j.1365-2389.1982.tb01794.x](https://doi.org/10.1111/j.1365-2389.1982.tb01794.x)
- Wu L, Wood Y, Jiang P, Li L, Pan G, Lu J, Chang AC, Enloe HA (2008) Carbon sequestration and dynamics of two irrigated agricultural soil in California. *Soil Sci Soc Am J* 72:808–814. doi:[10.2136/sssaj2007.0074](https://doi.org/10.2136/sssaj2007.0074)
- Wulf S, Lehmann J, Zech W (1999) Emissions of nitrous oxide from runoff-irrigated and rainfed soils in semiarid north-west Kenya. *Agr Ecosyst Environ* 72:201–205. doi:[10.1016/S0167-8809\(98\)00172-8](https://doi.org/10.1016/S0167-8809(98)00172-8)
- Yao Z, Zhou Z, Zheng Y, Xie B, Mei B, Wang R, Butterbach-Bahl K, Zhu J (2010) Effects of organic matter incorporation on nitrous oxide emissions from rice-wheat rotation ecosystems in China. *Plant Soil* 327:315–330. doi:[10.1007/s11104-009-0056-4](https://doi.org/10.1007/s11104-009-0056-4)