Responses of labile soil organic carbon and enzyme activity in mineral soils to forest conversion in the subtropics

Qingkui Wang, Fuming Xiao, Tongxin He, Silong Wang

► To cite this version:

Qingkui Wang, Fuming Xiao, Tongxin He, Silong Wang. Responses of labile soil organic carbon and enzyme activity in mineral soils to forest conversion in the subtropics. *Annals of Forest Science*, 2013, 70 (6), pp.579-587. 10.1007/s13595-013-0294-8 . hal-01201499

HAL Id: hal-01201499

<https://hal.science/hal-01201499>

Submitted on 17 Sep 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Responses of labile soil organic carbon and enzyme activity in mineral soils to forest conversion in the subtropics

Qingkui Wang · Fuming Xiao · Tongxin He · Silong Wang

Received: 1 December 2012 / Accepted: 30 April 2013 / Published online: 16 May 2013
© INRA and Springer-Verlag France 2013

Abstract

• **Aims** Globally, extensive areas of native forest have been almost replaced by plantations to meet the demands for timber, fuel material and other forest products. This study aimed to evaluate the effects of forest conversion on labile soil organic C (SOC), soil respiration, and enzyme activity, and to quantify their relationship in subtropical forest ecosystems.

• **Methods** Surface mineral soil (0–20 cm) was collected from a *Cunninghamia lanceolata* Hook. plantation, *Pinus massoniana* Lamb. plantation, *Michelia macclurei* Dandy plantation, and an undisturbed native broadleaf forest. Soil microbial biomass C, dissolved organic C, permanganate-oxidizable C, basal respiration, and six enzyme activities were investigated.

• **Results** Soil microbial biomass C was higher by 45.9 % in native broadleaf forest than that in *M. macclurei* Dandy plantation. The ratio of soil microbial biomass C to total SOC was 27.6 % higher in the *M. macclurei* Dandy plantation than in the native broadleaf forest. The soil respiration increased by 25.2 % and 21.7 % after conversion from native broadleaf forest to *P. massoniana* Lamb. and *M. macclurei* Dandy plantations respectively. The effects of

forest conversion on the soil enzyme activities differed among the tree species. Soil microbial biomass C had higher correlation with soil respiration than with the other SOC fractions. Moreover, soil microbial biomass C was positively correlated with urease and negatively correlated with cellulase activity. Soil respiration had higher correlation with soil microbial biomass C, dissolved organic C and permanganate-oxidizable C.

• **Conclusion** Forest conversion affected the soil microbial biomass C, soil respiration, invertase, cellulase, urease, catalase, acid phosphatase, and polyphenol oxidase activities, but their response depended on tree species. Soil respiration was mainly controlled by labile SOC, not by total SOC.

Keywords Land-use change · Tree species · Labile soil organic C · Soil respiration · Soil enzyme

1 Introduction

Globally, plantations with short rotation and high productivity are widely established to meet the demand for timber, fuel material, and other forest products. In 2010, the total area of plantations was approximately 264 million ha, corresponding to 6.6 % of the forest area (FAO 2010). Most plantations were derived from native forests. The conversion of a native forest into a plantation affects the forest ecological processes and functions (Schroth et al. 2002). In past decades, most studies focused on the effects of forest conversion on soil properties such as total soil organic C (SOC) and other physicochemical properties (Schroth et al. 2002). The simultaneous examination of the changes in labile SOC and soil biological properties induced by forest conversion is important beyond the insights it gives into the effects of forest conversion on the C cycle, because both labile SOC and soil microbe contribute to the soil CO₂ emission, which results in the loss of stored C in forest soils.

Handling Editor: Gilbert Aussenac

Q. Wang (✉) · T. He · S. Wang
Huitong Experimental Station of Forest Ecology,
State Key Laboratory of Forest and Soil Ecology,
Institute of Applied Ecology, Chinese Academy of Sciences,
Shenyang 110164, People's Republic of China
e-mail: wqkui@163.com

F. Xiao
Jiangxi Academy of Forestry, Nanchang 330032,
People's Republic of China

T. He
University of Chinese Academy of Sciences, Beijing 100039,
People's Republic of China

However, few studies have focused on the simultaneous response of labile SOC and soil biological properties to forest conversion.

Total SOC is heterogeneous, and consists of different functional and biological pools with different turnover rates (von Lützow et al. 2007). Based on turnover rates and residence time, the SOC can be divided into a labile C pool, which exists for weeks to years, and a recalcitrant C pool, which exists for decades or even centuries (Rovira and Vallejo 2002). The labile C pool mainly consists of simple sugars, organic acids, microbial biomass, and metabolic compounds of incorporated plant residues, given its rapid turnover (Cochran et al. 2007). Labile SOC fractions include soil microbial biomass C (MBC), dissolved organic C (DOC), and permanganate-oxidizable C (POC). Although the labile fraction is more sensitive to short-term vegetation or land-use change (Sicardi et al. 2004; von Lützow et al. 2007), measuring labile SOC is more laborious and costly than measuring total SOC. However, whether labile SOC is still more sensitive than total SOC on a long-term scale (e.g., 30 years) after forest conversion is unknown.

Soil respiration as a key ecosystem process that releases C as CO₂ from soil is affected by forest conversion. Some studies have shown that the conversion of a natural forest into a plantation leads to a net accretion of soil respiration (Liu et al. 2011). By contrast, Mohanty and Panda (2011) noted that natural forest conversion reduces soil respiration, and Xu et al. (2008) reported that forest conversion has no effect. The inconsistency in the effect of forest conversion on soil respiration limits our understanding of the forest role in C cycle. Therefore, the effects of forest conversion on soil respiration must still be quantified. Based on small-scale experiments with similar soil temperature and moisture conditions, soil C availability is important for soil respiration. Although C availability affects soil respiration (Dube et al. 2009; Iqbal et al. 2010), the relationship between soil respiration and labile SOC fractions is still poorly understood.

Soil enzymes produced by microbes play key roles in the biochemical functions of organic matter decomposition and nutrient cycling (Waldrop et al. 2004). Soil enzyme activities, which represent microbial activity, are affected by land-use and vegetation changes (Sicardi et al. 2004; Acosta-Martínez et al. 2007; Weand et al. 2010). However, some enzyme activities do not consistently respond to forest vegetation changes (Chang et al. 2011). Therefore, a better understanding of the effects of forest conversion on soil enzyme activity, and the relationship between soil enzyme activity and labile SOC, potentially provides an opportunity to assess the response of the soil biogeochemical cycle to land-use change. However, the relationship between different SOC fractions and enzymes is still not clear.

In this study, we investigated the enzyme activities known to play key roles in the mineralization of C, N, and

P nutrients in soils (Dilly and Nannipieri 2001; Acosta-Martínez et al. 2007). For example, acid phosphatase and urease are important in soil P and N mineralization because they catalyze the hydrolysis of various organic P and N respectively (Acosta-Martínez et al. 2007). Catalase catalyzes hydrogen peroxide decomposition (Brzezińska et al. 2005). Polyphenol oxidase is involved in lignin degradation, and plays an important role in soil C stabilization by favoring humic substance formation through the catalysis of polyphenol oxidation reactions (Weand et al. 2010). Cellulase can break down cellulose to cellobiose, a sugar composed of two glucose units. Invertase catalyzes the hydrolysis of sucrose into glucose and fructose.

In this study, we compared effects of tree species after forest conversion based on a base line of undisturbed forest. Based on the analysis of different labile SOC fractions, soil respiration, and activities of six soil enzymes in four different subtropical forests, this study aimed to (1) examine the effect of forest conversion on labile SOC, soil respiration, and enzyme activity, (2) quantify the relationship among different labile SOC, soil respiration, and enzyme activity, and (3) determine whether labile SOC is a more sensitive indicator of soil quality than total SOC over the long term after forest conversion. We tested the following hypotheses: (1) that the conversion of a native broadleaved forest into a plantation decreases labile SOC, soil respiration, and enzyme activity, (2) that the soil MBC has a higher correlation with soil respiration and enzyme activity than the other SOC fractions, and (3) that labile SOC is a more sensitive indicator of soil quality than total SOC over the long term after forest conversion.

2 Materials and methods

2.1 Study area

The study was conducted at the Huitong National Research Station of Forest Ecosystem (26°50' N, 109°36' E). This region belongs to the transition zone from the Yun-Gui plateau to the lower mountains and hills along the southern bank of the Yangtze River. The altitude ranges from 300 m to 1,000 m above sea level. With a relatively deeper layer, red earth or reddish-yellow earth are generally distributed from 300 m to 600 m above sea level. This region has a humid mid-subtropical monsoon climate, with a mean annual temperature and precipitation of 16.5 °C and 1,200 mm respectively. The native vegetation is evergreen broadleaved forest typical of subtropics, with *Castanopsis fargesii* Franch. and *Quercus lotungensis* Chun as the dominant species. The region is a major production area because of the rich water and heat resources. Therefore, many highly productive commercial tree species (e.g., *Cunninghamia*

lanceolata Hook. and *Pinus massoniana* Lamb.) are planted after native forest clearcutting.

In the 1980s, most native evergreen broadleaved forests were cut and replaced by plantations. In the current study, we selected *Cunninghamia lanceolata* Hook. plantation, *P. massoniana* Lamb. plantation, and *Michelia macclurei* Dandy plantation forests established in 1984, 1987, and 1983 respectively, and an native broadleaf forest undisturbed for 50 years, representing four forest-use types. Three plots (i.e., three replications) of 20 m × 20 m in each forest type were randomly established. The distance between the plots in each site is about 80 m to 100 m. In each plot, trees with heights more than 1.3 m and diameters at breast height (DBHs) more than 1.5 cm were investigated. The dominant tree species in the native broadleaf forest were *Castanopsis fargesii* Franch., *Q. lotungensis* Chun, *Machilus pauhoi* Kaneh., *Meliosma rigida* Siebold, and *Eurya nitida* Korthals. In the NBF plots, the highest values of tree height and DBH were 30 m and 110 cm respectively. The soil was classified as an oxisol based on the US soil taxonomy (Soil Survey Staff 1999). Detailed information about the structural characteristics of the five forests and main soil physicochemical properties at investigation are shown in Table 1. The background values including soil profile characteristics, textures, and mineral composition were almost identical in the different plots (Chen et al. 2000). Consequently, any observed differences in soil properties among the forests are probably mainly due to management.

2.2 Soil sampling and analysis

To examine effects of forest conversion on labile SOC pools and microbial activity in mineral soil layer, the litter and organic horizon was removed, and six soil cores (diameter

45 mm) in each plot were collected in July 2011 from the upper 20 cm soil layer to pool into a composite soil sample. All soil samples were immediately sieved through a 2-mm mesh. Some of the soil samples were stored at 3 °C for determining soil MBC, DOC, basal respiration, and enzyme activities within 7 days. The other samples were then dried for determining soil physicochemical properties and POC.

Soil MBC was measured through the chloroform fumigation- K_2SO_4 extraction method described by Vance et al. (1987). The fumigated and unfumigated soils were extracted using $0.5 \text{ mol} \cdot \text{L}^{-1} K_2SO_4$ with a ratio of 1:4 (soil: K_2SO_4 , w:v). The extracted C was measured using an elemental analyzer (Elementar High TOC, Germany). Soil MBC was calculated using a factor of 2.2. Soil basal respiration was determined using the method of alkali absorption. Briefly, the field-moist soils (50 g, oven-dry weight) were aerobically incubated at 28 °C in a 500 ml sealed glass jar for 24 h. The CO_2 that evolved from the soil was trapped in $0.1 \text{ mol} \cdot \text{L}^{-1} NaOH$ solution. The NaOH solution was then extracted for precipitation processing using a saturated $BaCl_2$ solution. The amount of CO_2 absorbed was determined by titration using $0.05 \text{ mol} \cdot \text{L}^{-1} HCl$ solution and phenolphthalein as a visual indicator. The microbial metabolic quotient (qCO_2) was calculated by dividing the hourly basal respiration rate by the corresponding MBC.

POC encompasses all the organic components that can be readily oxidized by $KMnO_4$, including labile humic material and polysaccharides (Conteh et al. 1999). POC is defined as the organic C oxidized by $0.333 \text{ mol} \cdot \text{L}^{-1} KMnO_4$. POC was determined using the method recommended by Blair et al. (1995). Briefly, 100 mg of soil sample was added into 50-ml plastic screw cap centrifuge tubes, followed by 25 ml of $333 \text{ mmol} \cdot \text{L}^{-1} KMnO_4$. The centrifuge tubes were tightly sealed and shaken for 1 h. The tubes were then centrifuged

Table 1 Structural characteristics of the four forests studied, and main soil physicochemical properties in subtropical China

Slope aspect	NBF NE	CLP NW	PMP NW	MMP SE
Mean tree height (m)	9.00	17.9	19.1	17.4
Mean DBH (cm)	13.9	22.1	14.3	16.1
Stem density (no.·ha ⁻¹)	1500	1080	2100	1890
Mean soil water content (%)	22.9	33.7	23.8	33.4
Soil total N (g·kg ⁻¹)	1.70	1.61	1.48	1.70
Soil total P (g·kg ⁻¹)	0.23	0.26	0.24	0.27
Mineral N (mg·kg ⁻¹)	9.39	14.52	8.13	10.89
Available P (mg·kg ⁻¹)	1.07	1.17	1.14	1.35
Bulk density (g·cm ⁻³)	1.18	1.23	1.25	1.08
pH (H ₂ O)	4.43	4.32	4.30	4.38
Sand (%)	6.9	11.4	12.2	11.2
Clay (%)	43.5	47.6	43.6	38.2
Silt (%)	49.6	41.0	44.2	50.6

NBF, CLP, PMP, and MMP denote an undisturbed native broadleaf forest, *Cunninghamia lanceolata* Hook. plantation, *Pinus Massoniana* Lamb. plantation, and *Michelia macclurei* Dandy plantation, respectively.

for 5 min at 2,000 rpm, and the supernatants were diluted with deionized water. The absorbances of the diluted samples and standards were measured using an atomic absorption spectrophotometer (Varian AA240, America) at 565 nm.

DOC was extracted with distilled water using the method described by Liang et al. (1998). Briefly, 10 g of fresh soil and 25 ml of ultra-pure water were added into a centrifuge tube. For extraction, the mixture was shaken for 30 min on a reciprocal shaker (Labnet Pro 30, America), and then centrifuged at 6,000 rpm for 15 min. The extract was filtered through a 0.45- μ m glass fiber filter. The C in the extract was measured using an elemental analyzer (Elementar High TOC, Germany).

Invertase activity was assessed as described by Guan (1986). Briefly, 5 g of soil was mixed with 15 ml of 8 % sucrose, 5 ml phosphate buffer at pH 5.5, and 0.1 ml toluene. The mixture was incubated for 24 h at 37 °C. The glucose released by the invertase reacted with 3,5-dinitrosalicylic acid and 3-amino-5-nitrosalicylic acid, and was then measured on an atomic absorption spectrophotometer (Varian AA240, America) at 508 nm. The results were expressed as mg released glucose by 1 g soil per day. Soil urease activity was also determined using a standard colorimetric assay described by Guan (1986), and the results were expressed as mg of released $\text{NH}_3\text{-N}$ by 1 kg of soil per hour at 37 °C.

Cellulase activity was assessed using the modified method of Deng and Tabatabai (1994). Briefly, 5 g of soil sample was mixed with 20 ml of 50 $\text{mmol}\cdot\text{l}^{-1}$ acetate buffer with pH 5.5 and 2 % of carboxymethyl cellulose. The mixture was incubated for 4 h at 50 °C and then centrifuged. The supernatant was treated with the Somogyi–Nelson reagent. Ammonium molybdate (25 mg) was dissolved in 450 ml of distilled water. Then, 21 ml of concentrated H_2SO_4 was mixed into the solution. $\text{Na}_2\text{HASO}_4\cdot 7\text{H}_2\text{O}$ (3 mg) was dissolved in 25 ml of H_2O and mixed. The solution was centrifuged once as described above, before the color measurement of reducing sugars at 520 nm.

Acid phosphatase activity was determined using the method described by Wu et al. (2006). Briefly, 10 ml of di-sodiumphenyl phosphate solution (25 $\text{g}\cdot\text{l}^{-1}$) used as a substrate was added to 5 g of soil. Soil samples were incubated for 12 h at 37 °C in acetate buffer (pH 5.0). The buffer solution (5 ml) and filtrate (2 ml) were transferred into a 50 ml volumetric flask, and then diluted with distilled water to 25 ml. Subsequently, 0.5 ml of 2,6-dibromoquinone-chlorimide solution was added to the flask, which was incubated for 20 min at room temperature. Finally, we diluted the solution to 50 ml, and determined the released phenol at 600 nm on an atomic absorption spectrophotometer (Varian AA240, America). Acid phosphatase activity was expressed as mg hydrolyzed phenol by 1 kg soil per hour at 37 °C.

Catalase activity was determined according to the method of Guan (1986). Briefly, 2 g of soil with 40 ml distilled water and 5 ml 0.3 % H_2O_2 were added into centrifuge

tubes, and then shaken for 20 min. The filtrate was titrated with 25 $\text{mmol}\cdot\text{l}^{-1}$ KMnO_4 . The results were expressed as ml 25 $\text{mmol}\cdot\text{l}^{-1}$ $\text{KMnO}_4\cdot\text{g}^{-1}\cdot\text{h}^{-1}$.

Polyphenol oxidase activity was determined using the method described by Guan (1986). Soil samples (5 g) were mixed with 10 ml of distilled water, 6 ml of 0.1 % ascorbic acid, and 10 ml of 0.02 $\text{mol}\cdot\text{l}^{-1}$ catechol, and then incubated for 2 min in a water bath at 30 °C. We then added 3 ml of 10 % phosphoric acid. The filtrate was titrated with 0.005 $\text{mol}\cdot\text{l}^{-1}$ iodine. The results were expressed as ml 0.005 $\text{mol}\cdot\text{l}^{-1}$ $\text{I}_2\cdot\text{g}^{-1}\cdot\text{h}^{-1}$.

2.3 Statistical analyses

The effects of land use on the labile SOC fractions, soil respiration, and enzyme activities were tested via one-way analysis of variance (ANOVA) using the SPSS software version 17.0 for Windows (SPSS Inc., Chicago, IL, USA). Assumptions of normality and homogeneity of variances were checked, and dependent variables were natural log transformed where necessary. Tukey's honestly significant difference test was performed as a post-hoc test to separate the means when differences were significant. The relationships between the SOC fractions and soil respiration, as well as soil enzyme activity, were modeled via the Pearson linear procedure using the SPSS software. Pearson linear correlations were also analysed to assess the relationships between the labile SOC fractions and soil enzyme activity. All effects were considered significant at the $P < 0.05$ level.

3 Results

3.1 SOC fractions

Forest conversion from native broadleaf forest to plantation significantly affected soil MBC (Fig. 1). Significant differences in soil MBC were observed only between native broadleaf forest and *Michelia macclurei* Dandy plantation. Soil MBC in *M. macclurei* Dandy plantation was 45.9 % higher than that in native broadleaf forest. However, total SOC, DOC, and POC were not significantly affected by forest conversion. Forest conversion significantly affected the ratios of labile SOC fractions to total SOC. The ratio of soil MBC to total SOC was 27.6 % higher in *M. macclurei* Dandy plantation than in native broadleaf forest. Forest conversion from native broadleaf forest to *C. lanceolata* Hook. plantation decreased the ratio of DOC to total SOC, and the conversion from native broadleaf forest into *P. massoniana* Lamb. plantation decreased the ratio of DOC to total SOC. The labile SOC fractions were significantly correlated with total SOC, but POC had the highest correlation (Fig. 2).

Fig. 1 Changes in total soil organic C, microbial biomass C (MBC), dissolved organic C (DOC), and permanganate-oxidizable C (POC), as well as ratios of MBC, DOC, and POC to total SOC in the four subtropical forests. NBF, CLP, PMP, and MMP denote an undisturbed native broadleaved forest, *Cunninghamia lanceolata* Hook. plantation, *Pinus massoniana* Lamb. plantation, and *Michelia macclurei* Dandy plantation, respectively. Different letters on the bar denote significance at $P < 0.05$

3.2 Soil respiration and enzyme activity

Soil respiration increased by 25.2 % and 21.7 % after conversion from native broadleaf forest to *P. massoniana* Lamb. plantation and *M. macclurei* Dandy plantation respectively (Fig. 3). Forest conversion tended to reduce the $q\text{CO}_2$, but not significantly. Soil basal respiration and labile SOC fractions had comparatively stronger relationships (Fig. 4). However, the relationship between soil basal respiration and total SOC was weaker.

Forest conversion significantly affected soil enzyme activities (Fig. 5), but the response patterns of the soil enzyme activities differed. Conversion from native broadleaf forest to plantations

increased the invertase and polyphenol oxidase activities, but decreased the cellulase activity. Urease activity increased after conversion from native broadleaf forest to *M. macclurei* Dandy plantation, but the catalase activity decreased. Conversion from native broadleaf forest to *P. massoniana* Lamb. plantation decreased the acid phosphatase activity.

3.3 Relationship between soil enzyme activities and SOC fractions

The soil enzyme activities had inconsistent relationships with the labile SOC fractions (Table 2). Total SOC and

Fig. 2 Relationship of labile SOC fractions [microbial biomass C (MBC), dissolved organic C (DOC), and permanganate-oxidizable C (POC)] with total SOC in the subtropical forests. * and ** denote significance at $P < 0.05$ and 0.01 respectively

Fig. 3 Soil basal respiration and microbial metabolic quotient in the four subtropical forests. NBF, CLP, PMP, and MMP denote an undisturbed native broadleaved forest, *C. lanceolata* Hook. plantation, *P. Massoniana* Lamb. plantation, and *M. macclurei* Dandy plantation, respectively. Different letters on the bar denote significance at $P < 0.05$

POC were not correlated with any soil enzyme activity. MBC was positively correlated with the urease activity and negatively correlated with the cellulase activity. DOC was positively correlated with urease and acid phosphatase activities.

4 Discussion

In contrast to our first hypothesis, we found that forest conversion from native broadleaf forest to plantation affected only the soil MBC but not the other labile SOC fractions. Our results were supported by other previous studies (Wang and Wang 2007; Xu et al. 2008). These results indicate that the response of labile SOC fractions to forest conversion depends on forest vegetation. Litter and root are the primary source of SOC. Therefore, the different quantities and

qualities of organic matter input through litterfall and root activity (e.g., turnover and exudates) induced by the different tree species were responsible for the different responses of labile SOC to forest conversion (Wang and Wang 2007; Vesterdal et al. 2008). This finding was supported by the observations of Wang et al. (2007) in similar regions, which showed that tree species affect litter production and dynamics. The significant increase in soil MBC induced by forest conversion is important for C emission and storage in soil. Soil microbial activity utilizes SOC as C source and releases CO₂ to atmosphere, which increases the C emission from soil to atmosphere and reduces soil C storage. The higher soil respiration in plantations than in NBF also supported our hypotheses, although total SOC had no significant change.

Unexpectedly, the soil enzyme activities showed different patterns of response to forest conversion. Hou et al. (2006) also found that urease and catalase activities differed among

Fig. 4 Relationship of soil basal respiration (mg CO₂ kg⁻¹ dry soil d⁻¹) with total soil organic C (SOC; g kg⁻¹), microbial biomass C (MBC; mg kg⁻¹), dissolved organic C (DOC; mg kg⁻¹) and permanganate-oxidizable C (POC; g kg⁻¹) in the subtropical forests. * denotes significance at $P < 0.05$

Fig. 5 Activities of soil invertase (mg released glucose·kg⁻¹ soil·d⁻¹), cellulase (mg reducing sugar·kg⁻¹ soil·h⁻¹), urease (mg released NH₃-N·kg⁻¹ soil·h⁻¹), catalase (ml 25 mol⁻¹ KMnO₄·kg⁻¹ soil·h⁻¹), acid phosphatase (mg hydrolyzed phenol·kg⁻¹ soil·h⁻¹), and polyphenol oxidase (ml 0.05 mol⁻¹ I₂·kg⁻¹ soil·h⁻¹) in the four subtropical forests. NBF, CLP, PMP, and MMP denote an undisturbed native broadleaved forest, *C. lanceolata* Hook. plantation, *P. massoniana* Lamb. plantation, and *M. macclurei* Dandy plantation, respectively. Different letters on the bar denote significance at *P* < 0.05

three forest types in Yellow River Delta. The increase in invertase activity after forest conversion indicates that the microbial activity and C availability increased (Wallenstein et al. 2009), which was supported by the increase in soil MBC and respiration. Given the key roles of PPO in soil carbon stabilization, the significant increase in PPO activity suggests that forest conversion increased the soil C stability. Waldrop et al. (2004) found that PPO activity decreases with increasing mineral N availability. Therefore, the relatively lower mineral N and total N in soils from *P. massoniana* Lamb. plantation was partly responsible for the higher PPO activity. The different patterns of soil enzyme activities in the plantations suggest that soil enzymes have tree species-specific response to forest conversion.

In a subtropical montane forest with limited P, microbial P mineralization is essential for supplying trees with available P. Acid phosphatase had different responses after forest conversion based on the tree species, which is consistent with the observations of Ushio et al. (2010) in a tropical montane forest. Phosphatase activity is correlated with the available P concentration (Acosta-Martínez et al. 2007), and is suppressed by P addition (Allison and Vitousek 2005). Therefore, the differences in acid phosphatase activity under

different forests may be due to the differences in soil available P induced by different tree species.

Some of the enzyme activities were correlated with the labile SOC fractions, which confirms some previous studies (Brzezińska et al. 2005; Acosta-Martínez et al. 2007). However, Shackle et al. (2000) reported that phosphatase activity is not regulated by C availability in constructed wetlands. Generally, enzyme activities are correlated with SOC content because the latter plays a key role as a precursor for enzyme synthesis. The significant inter-correlation among cellulase, urease, catalase, and acid phosphatase activities (data not shown) suggests that these enzymes have similar origin and C source in soil. The higher correlation between the soil enzyme activities and MBC than SOC and POC also suggests that soil enzyme activity was partly controlled by soil C availability. Significant relationships between soil respiration and labile SOC fraction were also found, which is in accordance with some studies (Dube et al. 2009; Iqbal et al. 2010). In addition, soil respiration was highly related to MBC. The aforementioned results confirm our hypothesis that the soil MBC has higher correlation with soil respiration and enzyme activity than the other SOC fractions.

Table 2 Pearson product-moment correlation coefficients between soil enzymes and SOC fractions in the four subtropical forests

	Invertase	Cellulase	Urease	Catalase	Acid phosphatase	Polyphenol oxidase
SOC	0.371	-0.417	0.080	-0.035	-0.219	0.019
MBC	0.279	-0.515*	0.626*	-0.613*	0.170	0.344
DOC	-0.188	-0.085	0.697*	-0.421	0.524*	-0.176
POC	0.263	-0.305	0.022	0.025	-0.094	-0.243

SOC, MBC, DOC, and POC denote total soil organic C, microbial biomass C, dissolved organic C, and permanganate-oxidizable C respectively. * denotes significance at *P* < 0.05

Total SOC was insignificantly changed 30 years after forest conversion. However, significant correlations between SOC and labile fractions were found, which confirms the previous results of Wang et al. (2007) in a similar study region. The higher correlation between POC and total SOC suggests that POC is more closely related to the amount of total SOC, accounting for 21.5 % of total SOC. The POC fraction accounts for 5 % to 30 % of total SOC (Blair et al. 1995; Conteh et al. 1999). Therefore, the amount of labile SOC was controlled by the amount of total SOC in long-term scales after forest conversion. In contrast to our hypothesis, the aforementioned results indicate that measuring the more laborious labile forms with the exception of MBC provides little additional value. Furthermore, MBC is a sensitive indicator of soil quality over the long term after land-use changes compared with other SOC fractions, although soil quality is a complex concept, and its determination is the result of assessing several biological, chemical, and physical soil components.

In conclusion, forest conversion from native broadleaf forest to monoculture plantations increased only the MBC but not the other labile fractions and total SOC. As expected, the soil MBC had higher correlation with soil respiration and enzyme activity than the other SOC fractions. Soil C availability partly controlled the microbial and enzyme activities. MBC was a more sensitive indicator of soil quality compared with total SOC over the long term after forest conversion. Changes in soil biodiversity and its consequence on the organisms under the trees in monoculture plantations should be further studied. On the other hand, monitoring abiotic changes also should be implemented.

Acknowledgments We are grateful to Micai Zhong, Guangbiao Xu, Xiuyong Zhang, and Zhengqi Shen for their assistance during soil sampling and analysis.

Funding This research was supported by the National Basic Research Program of China (973 Program, 2012CB416905), the “Strategic Priority Research Program—Climate Change: Carbon Budget and Related Issues” of the Chinese Academy of Sciences (XDA05070305), and the National Natural Science Foundation of China (31070436 and 41030533).

References

- Acosta-Martínez V, Cruz L, Sotomayor-Ramírez D, Pérez-Alegria L (2007) Enzyme activities as affected by soil properties and land use in a tropical watershed. *Appl Soil Ecol* 35:35–45
- Allison SD, Vitousek PM (2005) Response of extracellular enzymes to simple and complex nutrient inputs. *Soil Biol Biochem* 37:937–944
- Blair GJ, Lefroy RDB, Lisle L (1995) Soil carbon fractions based on their degree of oxidation and the development of a carbon management index for a cultural systems. *Aust J Agric Res* 46:1459–1466
- Brzezińska M, Włodarczyk T, Stepniowski W, Przywara G (2005) Soil aeration status and catalase activity. *Acta Agrophysica* 5:555–565
- Chang E, Chen C, Chen T, Chiu C (2011) Soil microbial communities and activities in sand dunes of subtropical coastal forests. *Appl Soil Ecol* 49:256–262
- Chen CY, Liao LP, Wang SL (2000) Ecology of Chinese Fir plantation forest. Science Press, Beijing, pp 1–39, 84–96 (in Chinese)
- Cochran RL, Collins HP, Kennedy A, Bezdicsek DF (2007) Soil carbon pools and fluxes after land conversion in a semiarid shrub-steppe ecosystem. *Biol Fertil Soils* 43:479–489
- Conteh A, Blair GT, Lefroy RDB, Whitbread AM (1999) Labile organic carbon determined by permanganate oxidation and its relationships to other measurements of soil organic carbon. *Hum Substit Environ J* 1:3–15
- Deng SP, Tabatabai MA (1994) Cellulase activity of soils. *Soil Biol Biochem* 26:1347–1354
- Dilly O, Nannipieri P (2001) Response of ATP content, respiration rate and enzyme activities in an arable and a forest soil to nutrient addition. *Biol Fertil Soils* 34:64–72
- Dube F, Zagal E, Stolpe N, Espinosa M (2009) The influence of land-use change on the organic carbon distribution and microbial respiration in a volcanic soil of the Chilean Patagonia. *Forest Ecol Manag* 257:1695–1704
- Food and Agriculture Organization of the United Nations (2010) Global forest resources assessment 2010: main report. FAO Viale delle Terme di Caracalla, Rome
- Guan S (1986) Soil enzymes and their methodology. Agricultural Press, Beijing, pp 274–340
- Hou B, Ma F, Song Y, Hou L, Cui X (2006) A study on effect of different tree species on soil nutrients, enzyme activities and microorganisms. *Acta Agric Univ Jiangxiensis* 28:734–738
- Iqbal J, Hu R, Feng M, Lin S, Malghani S, Ali IM (2010) Microbial biomass, and dissolved organic carbon and nitrogen strongly affect soil respiration in different land uses: a case study at Three Gorges Reservoir Area, South China. *Agric Ecosyst Environ* 137:294–307
- Liang BC, MacKenzie AF, Schnitzer M, Monreath CM, Voroney PR, Beyaert RP (1998) Management-induced changes in labile soil organic matter under continuous corn in eastern Canadian soils. *Biol Fertil Soils* 26:88–94
- Liu J, Jiang P, Wang H, Zhou G, Wu J, Yang F, Xi Q (2011) Seasonal soil CO₂ efflux dynamics after land use change from a natural forest to Moso bamboo plantations in subtropical China. *Forest Ecol Manag* 262:1131–1137
- Mohanty RB, Panda T (2011) Soil respiration and microbial population in a tropical deciduous forest soil of Orissa, India. *Flora-Morphol Distrib Funct Ecol Plants* 206:1040–1044
- Rovira P, Vallejo VR (2002) Labile and recalcitrant pools of carbon and nitrogen in organic matter decomposing at different depth in soil: an acid hydrolysis approach. *Geoderma* 107:109–141
- Schroth G, D’Angelo SA, Teixeira WG, Haag D, Lieberei R (2002) Conversion of secondary forest into agroforestry and monoculture plantations in Amazonia: consequences for biomass, litter and soil carbon stocks after 7 years. *Forest Ecol Manag* 163:131–150
- Shackleton VJ, Freeman C, Reynolds B (2000) Carbon supply and the regulation of enzyme activity in constructed wetlands. *Soil Biol Biochem* 32:1935–1940
- Sicardi M, García-Prézac F, Frioni L (2004) Soil microbial indicators sensitive to land use conversion from pastures to commercial *Eucalyptus grandis* (Hill ex Maiden) plantations in Uruguay. *Appl Soil Ecol* 27:125–133
- Soil Survey Staff (1999) Soil taxonomy: a basic system of soil classification for making and interpreting soil surveys. United States Department of Agriculture. Soil Conservation Service. U.S. Government Printing Office, Washington, DC
- Ushio M, Kitayama K, Balser TC (2010) Tree species effects on soil enzyme activities through effects on soil physicochemical and

- microbial properties in a tropical montane forest on Mt. Kinabalu, Borneo. *Pedobiologia* 53:227–233
- Vance ED, Brookes PC, Jenkinson DS (1987) An extraction method for measuring soil microbial biomass C. *Soil Biol Biochem* 19:703–707
- Vesterdal L, Schmidt IK, Callesen I, Nilsson LO, Gundersen P (2008) Carbon and nitrogen in forest floor and mineral soil under six common European tree species. *Forest Ecol Manag* 255:35–48
- von Lützow M, Kögel-Knabner I, Ekschmitt K, Flessa H, Guggenberger G, Matzner E, Marschner B (2007) SOM fractionation methods: relevance to functional pools and to stabilization mechanisms. *Soil Biol Biochem* 39:2183–2207
- Waldrop MP, Zak DR, Sinsabaugh RL, Gallo M, Lauber C (2004) Nitrogen deposition modifies soil carbon storage through changes in microbial enzymatic activity. *Ecol Appl* 14:1172–1177
- Wallenstein MD, McMahon SK, Schimel JP (2009) Seasonal variation in enzyme activities and temperature sensitivities in Arctic tundra soils. *Glob Chang Biol* 15:1631–1639
- Wang Q, Wang S (2007) Soil organic matter under different forest types in southern China. *Geoderma* 142:349–356
- Wang Q, Wang S, Fan B, Yu X (2007) Litter production, leaf litter decomposition and nutrient return in *Cunninghamia lanceolata* plantations in south China: effect of planting conifers with broadleaved species. *Plant Soil* 297:201–211
- Weand MP, Arthur MA, Lovett GM, McCulley RL, Weathers KC (2010) Effects of tree species and N additions on forest floor microbial communities and extracellular enzyme activities. *Soil Biol Biochem* 42:2161–2173
- Wu J, Lin Q, Huang Q, Xiao H (2006) Soil microbial biomass—methods and application. China Meteorological Press, Beijing, pp 127–129
- Xu Z, Ward S, Chen C, Blumfield T, Prasolova N (2008) Soil carbon and nutrient pools, microbial properties and gross nitrogen transformations in adjacent natural forest and hoop pine plantations of subtropical Australia. *J Soil Sediment* 8:99–105