



Integrative responses of pig adipose tissues to high-fat high-fiber diet: towards key regulators of energy flexibility

Florence Gondret, Annie Vincent, Magalie Houee, Sandrine Lagarrigue, Anne Siegel, David Causeur, Isabelle Louveau

► To cite this version:

Florence Gondret, Annie Vincent, Magalie Houee, Sandrine Lagarrigue, Anne Siegel, et al.. Integrative responses of pig adipose tissues to high-fat high-fiber diet: towards key regulators of energy flexibility. ASAS/ADSA midwest meeting, Mar 2015, Des Moines, United States. Journal of Animal Science, 93 (Suppl. 2), 2015, Abstract book of the ASAS/ADSA midwest meeting. hal-01210925

HAL Id: hal-01210925

<https://hal.science/hal-01210925>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Meeting Information

When: March 16 - 18, 2015

Where: Des Moines, IA

Abstract N° 77

Integrative responses of pig adipose tissues to high-fat high-fiber diet: towards key regulators of energy flexibility

Wednesday, March 18, 2015: 8:45 AM

314-315 (Community Choice Credit Union Convention Center)

Florence Gondret , AgroCampus-Ouest UMR1348 PEGASE, Rennes, France

Annie Vincent , INRA UMR1348 PEGASE, Saint Gilles, France

Magalie Houee , AgroCampus-Ouest UMR6625 IRMAR, Rennes, France

Sandrine Lagarrigue , INRA UMR1348 PEGASE, Saint Gilles, France

Anne Siegel , CNRS-Université de Rennes 1-INRIA, UMR6074 IRISA, Rennes, France

David Causeur , AgroCampus-Ouest UMR6625 IRMAR, Rennes, France

Isabelle Louveau , AgroCampus-Ouest UMR1348 PEGASE, Rennes, France

Abstract Text:

The competition between food and feed challenges the use of alternative resources such as fibrous feedstuffs in diets for pigs. Adding fat to high fiber diets appears as a relevant strategy to improve dietary energy value and feed efficiency, but this changes the nutrients and energy source compared to a standard low fat diet. This study aimed to elucidate the transcriptional mechanisms involved in variations of adiposity when pigs were fed a high fat high fiber diet. Growing barrows (Large White, $n = 48$) divergently selected for feed efficiency were offered during 10 weeks either a low fat, low fiber diet (LF) or a high fat, high fiber diet (HF) where oils and wheat straw were used to partially substitute cereals ($n = 24$ per diet). At 132 days of age, HF pigs displayed lower ($P < 0.001$) proportions of perirenal (PRAT; -16%) and subcutaneous (SCAT; -28%) fat tissues than LF pigs. Analyses using a porcine microarray showed that diet had pronounced effects on adipose tissue transcriptomes. The number of differentially-expressed genes (DEG) was greater in PRAT than in SCAT, with 1,251 and 825 unique genes being up-regulated and 2,440 and 1,279 unique genes being down-regulated by the HF diet in PRAT and SCAT, respectively (cutoffs for p corrected BH < 0.10 , raw $p < 0.01$ and fold-changes between conditions > 1.1 or < 0.9). A multiple factor analysis revealed large similarities between the two adipose tissues in response to diets. Notably, different genes related to protein catabolism (33 DEG), protein transport (33 DEG), apoptosis (31 DEG), phosphate metabolic process including ATP synthesis (34 DEG), response to stress (26 DEG) and glucose metabolism (13 DEG) were commonly down-regulated in HF pigs. Conversely, IGF1R participating to the negative regulation of apoptotic process was up-regulated by the HF diet in the two adipose tissues. Correlation modules also stressed the up-regulation by the HF diet of genes related to immunity and defense response specifically in PRAT. Causality graph analysis highlighted MLXIPL, SREBF1, peroxisome proliferator-activated receptors (PPARG, PPARD) and their heterodimer partner RXRA as candidate upstream regulators of

these processes. qPCR analyses confirmed dietary-related differences in expression levels of these regulators in the two adipose tissues. Altogether, high fiber intake in growing pigs was associated with lower body fatness, which was related to lower glucose metabolism in adipose tissues; its effect on immune factors in the perirenal fat deserves further studies.

Keywords: Pig; Fiber diet; Adipose tissue

See more of: [Growth, Development, Meat Science & Muscle Biology](#)

See more of: [Growth, Development, Muscle Biology, and Meat Science](#)