On the use of intra-rectal valium in patients with dravet syndrome: family's experience

Nathalie Coqué, Nicole Chemaly, Rima Nabbout

► To cite this version:

Nathalie Coqué, Nicole Chemaly, Rima Nabbout. On the use of intra-rectal valium in patients with dravet syndrome: family's experience. 10. European Congress on Epileptology, Sep 2012, Londres, United Kingdom. 245 p., 10.1111/j.1528-1167.2012.03677.x . hal-01239518

HAL Id: hal-01239518

<https://hal.science/hal-01239518>

Submitted on 7 Dec 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives| 4.0 International License

On the use of intra-rectal valium in patients with Dravet Syndrome : family's experience

Nathalie Coqué^{1,2,3,#}, Nicole Chemaly⁴, Rima Nabbout⁴

Abstract

Intra-rectal Diazepam (DZ) is the first rescue medication for acute prolonged convulsive seizures in children in many countries. In this study, we aimed at assessing the experience of the families of patients presenting Dravet Syndrome (DS) with respect to the use of intra-rectal DZ.

Situation of the problem

Intra-rectal DZ is prepared from a glass vial and pumped with a syringe that is then inserted into the rectum with or without a rectal tube. The administrator must administer the prescribed dose according to the weight of the child. We assessed different aspects of this treatment from the family's point of view.

Methods

We sent an auto-administered on-line questionnaire to 53 families of patients with DS. Questions addressed different aspects of intra-rectal DZ use:

- usual administrators, and time required to prepare the medicine (parents or other care givers)
- facility, and risk of errors in the preparation and the administration.

Parents were questioned about their feelings concerning the efficacy and utility in emergency settings, facility of intra-rectal DZ administration, and possible refuse of care givers to use this method of DZ administration.

Result

50 families answered the questionnaire. DZ was usually administered by mothers in 96% of cases, by fathers in 38% and less frequently by other care givers (14%). Almost all parents (78%) agreed on its efficacy. However, they found the use difficult by non-parents care givers, and that preparation and administration are subject to errors when carried out of the parents presence. Many educational and non-medical institutions refused to use DZ administered intra-rectally.

Figure 5 : Estimation of the ease of DZ usage by parents and non-parents

Figure 6 : Estimation of the risk of error in the preparation and administration of intra-rectal DZ by parents and non-parents

Figure 7 : Feeling of parents concerning the usefulness and the efficacy of intra-rectal DZ in emergency settings

Figure 8 : Percentage of incident in which care givers refuse to use intra-rectal DZ

Conclusion

Parents confirm the efficacy of intra-rectal DZ. However, they mention the humiliation of children and adults who are obliged to disrobe in order to receive treatment, and also the method of administration which is often considered as a medical practice by the care givers and refused by them. As such, they emphasize the need for a medication that is easier to prepare and to administer.