Innovative bioguided fractionation of cranberry juice for *Candida* biofilm treatment and infected-wound healing

Leslie Boudesocque-Delaye, T. Puy, Marion Girardot, Arnaud Lanoue, Laurent Bazinet, Charles Bodet, Christine Imbert, Cécile Enguehard-Gueiffier

► To cite this version:

Leslie Boudesocque-Delaye, T. Puy, Marion Girardot, Arnaud Lanoue, Laurent Bazinet, et al.. Innovative bioguided fractionation of cranberry juice for *Candida* biofilm treatment and infected-wound healing. 7. International Conference on Polyphenols and Health, Oct 2015, Tours, France. , 2015. hal-01256486

HAL Id: hal-01256486

<https://hal.science/hal-01256486>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

P817**Innovative bioguided fractionation of cranberry juice for *Candida* biofilm treatment and infected-wound healing.**

L. Boudesocque-Delaye,^{1,*} T. Puy,² M. Girardot,² A. Lanoue,³ L. Bazinet,⁴ C. Bodet,⁵ C. Imbert,² C. Enguehard-Gueiffier⁶

^{1*} Université de Tours, UMR INRA 1282 ISP, 31 avenue Monge, 37200 Tours, France, leslie.boudesocque@univ-tours.fr

² Université de Poitiers, UMR-CNRS 7267 – EBI, Bât. D1, 6 rue de la Milétrie, TSA 51115 86073 Poitiers Cedex 9, France

³ Université de Tours, EA 2106 BBV, 31 avenue Monge, 37200 Tours, France

⁴ Institut sur la Nutrition et les Aliments Fonctionnels (INAF), Département de sciences des Aliments, Université Laval, 2425 Rue de l'agriculture, Québec, QC, G1V 0A6, Canada

⁵ Université de Poitiers, A 4331 LITEC, Pole Biologie Santé, 1 rue Georges Bonne TSA 51106, 86073 Poitiers Cedex 9, France

⁶ Université de Tours, INSERM U1069 N2C, 31 avenue Monge, 37200 Tours, France

Cranberry polyphenols are intensively studied for their anti-infectious and antioxidant properties. Nevertheless, in biofilm and infected-wound healing, the identity of active polyphenols still remains unclear. In order to identify active polyphenols, an innovative fractionation scheme based on Centrifugal Partition Chromatography (CPC) was developed. CPC is a support free liquid/liquid chromatographic technology used here for the first time to fractionate cranberry juice. The fractionation protocol involved 3 main steps: firstly polyphenol and sugars separation; then polyphenols fractionation based of molecular weight and polarity, and finally CPC or on LH-20 resin.

Following the second step, the 6 main fractions obtained were screened on *Candida* biofilm and toward infected wound. Fractions containing low molecular weight proanthocyanidins (Fr III and IV) and containing high molecular weight PAC (Fr V ad VI) exhibited the most interesting activities. Concerning *Candida albicans* biofilm a prophylactic effect was observed by prevention of adhesion. On infected wound, a wound-healing potential related to anti-inflammatory effect but not to antibacterial activity against *Staphylococcus aureus* was highlighted.

These four active fractions were further purified using either CPC or LH-20 resin, to lead to almost 40 sub-fractions. Again fractions containing PACs of medium and high molecular weight exhibited the highest activity against adhesion step of *C. albicans* strains and infected wound healing.

This work is a step further to understand mechanisms of action of cranberry polyphenols on *C. albicans* biofilm prevention and in infected wound healing.

Financial support: ARC Tours Poitiers 2014 grant, FERI Région Centre 2013 grant

P818**Chemical fingerprints and biochemical activities of edible congolese hibiscus leaves**

P. Mutwale^{1,2*}, N. Ngombe², C. Tsumbu¹, T. Franck³, A. Mouithys-Mickalad³, D. Mumba^{4,5}, D. Tshala-Katumbay^{4,6}, P. Kalenda², D. Sertheyn³, M. Tits¹, M. Frédéricich

¹ Laboratory of Pharmacognosy, Center for Interdisciplinary Research on Medicines (CIRM), University of Liège, CHU, B-4000 Liège 1, Belgium

² Centre d'Etudes des Substances Naturelles d'Origine Végétale (CESNOV), Faculty of Pharmaceutical Sciences, University of Kinshasa, Kinshasa, BP 212 Kin XI, Democratic Republic of Congo

³ Centre for Oxygen Research and Development (C.O.R.D.), Institute of Chemistry B6a, University of Liège, B- 4000 Liège, Belgium

⁴ Faculty of Medicine, University of Kinshasa, Kinshasa, BP 834 Kin XI, Democratic Republic of Congo

⁵ Institut National de Recherches Biomédicales (INRB), Kinshasa, BP1197 Kin-Gombe, Democratic Republic of Congo

⁶ Department of Neurology, Oregon Health & Science University, Portland, OR97239, USA

Hibiscus acetosella, *Hibiscus cannabinus* and *Hibiscus sabdariffa* leaves are widely consumed as vegetables in the form of sauces by Congolese people (Democratic Republic of Congo).

The aim of this study was to compare the chemical fingerprints and the correlations between anti-inflammatory (myeloperoxidase (MPO) activity) and antioxidant properties of polar and non polar extracts of these plants. Antioxidant properties were evaluated by using 2',7'-dichlorofluorescein-diacetate (DCFH-DA) and lucigenin on neutrophils and HL-60 cells. Chemical fingerprints were performed by Thin Layer Chromatography and High Performance Liquid Chromatography.

Chemical fingerprintings revealed that *H. acetosella* has a fingerprint different to *H. cannabinus* and *H. sabdariffa* that have the same fingerprint. Major compounds are phenolic acids represented by caffeoyhydroxycitric acid for *H. acetosella* and neochlorogenic acid for the others. All extracts displayed high ABTS and DPPH radical-scavenging activities at the concentration range of 10 – 80 µg mL⁻¹ and antioxidant activity assays at 0.05 – 10 µg mL⁻¹, according to the following order: *H. acetosella* > *H. cannabinus* > *H. sabdariffa*. Non polar extracts exhibited a strongest superoxide

CONTENTS

Welcoming message	3
Committees	4
Program at a glance	5
Conference center plan	6
Scientific program	7
Guidelines for scientific communications	18
Social program	19
General information	20
Sponsors	23
Invited speaker biographies	24
Abstracts	
Invited lectures	35
Selected communications	79
Posters	
A: Polyphenol bioavailability and exposure assessment.....	103
B: Polyphenols in obesity, metabolic syndrome and diabetes	127
C: Neuro-cognitive effects of polyphenols	159
D: Polyphenols and cancer prevention	169
E: Polyphenols in cardiovascular diseases	183
F: Innovative approaches in polyphenol research	201
G: Mechanisms of action of polyphenols	213
H: Polyphenols and gut Microbiota interactions	235
I: Polyphenols in other diseases	243
J: Polyphenols and Food Processing	261
K: Other topics	277
Authors index	303
COST Action FA1403 POSITIVE satellite symposium.....	317