



**CORRECTION OF PHOTBLEACHING FOR THE  
ASSESSMENT OF PHARMACOKINETIC  
PARAMETERS USING DYNAMIC FLUORESCENCE  
MICROSCOPY**

Marc Derieppe, Clemens Bos, M. De Greef, Chrit Moonen, Baudouin Denis de  
Senneville

► **To cite this version:**

Marc Derieppe, Clemens Bos, M. De Greef, Chrit Moonen, Baudouin Denis de Senneville. CORRECTION OF PHOTBLEACHING FOR THE ASSESSMENT OF PHARMACOKINETIC PARAMETERS USING DYNAMIC FLUORESCENCE MICROSCOPY. IEEE International Symposium on Biomedical Imaging (ISBI), May 2015, New-York, United States. 2015.

**HAL Id: hal-01259351**

**<https://hal.archives-ouvertes.fr/hal-01259351>**

Submitted on 20 Jan 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# CORRECTION OF PHOTBLEACHING FOR THE ASSESSMENT OF PHARMACOKINETIC PARAMETERS USING DYNAMIC FLUORESCENCE MICROSCOPY

*M. Derieppe, C. Bos, M. de Greef, C. Moonen, B. Denis de Senneville*

## ABSTRACT

Local drug delivery in oncology aims at depositing high doses of anticancer agents while limiting their toxic side effects. Biological barriers, such as cell plasma membranes, hinder their delivery and requires strategies to address this challenge.

Previously [1], we demonstrated the feasibility to monitor in real-time, with dynamic fluorescence microscopy, the intracellular delivery of a hydrophilic model drug mediated by ultrasound (US), and to quantify the pharmacokinetic parameters derived from a two-compartment model. We evaluate here the impact of the photobleaching (PB) effect experimentally, and compute the PB-corrected uptake kinetics.

## 1. METHODS

US-mediated uptake of Sytox Green model drug, a fluorescent intercalating dye, was monitored using Fibred Confocal Fluorescence Microscopy [1]. Cell-nuclei were first detected and tracked over all acquired images [2]. Fluorescence signal enhancement was then fitted with a 2-compartment model, representing an extracellular and an intracellular compartment separated by a plasma membrane, as follows:

$$I(t) = A[1 - e^{-k(t-T)}] \quad (1)$$

with  $A$  the asymptote,  $T$  the uptake onset,  $k$  the uptake rate.

A 3-compartment model was then performed to represent PB as an efflux rate  $k_{pb}$  of the fluorescence in the inner cell compartment, as follows:


$$I(t) = \frac{A \times k}{k - k_{pb}} \times [e^{-k_{pb}(t-T)} - e^{-k(t-T)}] \quad (2)$$

After 6 minutes, the acquisition was interrupted for 9 minutes to stop PB, while the fluorophore concentration further equilibrated (Fig. 1). Then, a second 6-minute acquisition served to evaluate the PB rate. This estimate was then used as a fixed parameter in the 3-compartment model. The kinetic parameters were compared using the Mann-Whitney test (significance:  $p < 0.05$ ).

Research supported by ERC project 268906 "Sound Pharma".

M. Derieppe, C. Bos, M. de Greef, C. Moonen and B. Denis de Senneville are with the University Medical Center Utrecht, Imaging Division, Utrecht, Netherlands (corresponding author to provide phone: +31887569665; fax: +31887555850; e-mail: m.derieppe@umcutrecht.nl).

B. Denis de Senneville is also with the Mathematical Institute of Bordeaux, UMR 5251 CNRS/University of Bordeaux.



**Fig. 1.** Representative fluorescence signal enhancement of the uptake (a) and evaluation of photobleaching (b).

## 2. RESULTS

PB rates displayed a significant increase ( $p < 0.001$ ) of  $1.0 \cdot 10^{-3} \text{ s}^{-1}$  ( $4.4 \cdot 10^{-4} \text{ s}^{-1}$ ,  $n = 85$ ),  $2.1 \cdot 10^{-3} \text{ s}^{-1}$  ( $6.4 \cdot 10^{-4} \text{ s}^{-1}$ ,  $n = 103$ ) and  $2.8 \cdot 10^{-3} \text{ s}^{-1}$  ( $1.1 \cdot 10^{-3} \text{ s}^{-1}$ ,  $n = 73$ ) at 25 %, 50 % and 75 % of laser power, respectively. In a representative experiment, from which data in Fig. 1 are taken, application of the 3-compartment model, with the PB rate as an input, modified significantly the estimated uptake rate ( $p < 0.001$ ), from  $1.2 \cdot 10^{-2} \text{ s}^{-1}$  ( $1.1 \cdot 10^{-2} \text{ s}^{-1}$ ,  $n = 62$ ), using the 2-compartment model, to  $6.2 \cdot 10^{-3} \text{ s}^{-1}$  ( $4.7 \cdot 10^{-3} \text{ s}^{-1}$ ,  $n = 62$ ) using the 3-compartment model.

## 3. CONCLUSION

This study demonstrates that processing of photobleaching is critical for the assessment of pharmacokinetic parameters in dynamic fluorescence imaging.

## 4. REFERENCES

- [1] M. Derieppe et al., "Real-time assessment of ultrasound-mediated drug delivery using fibred confocal fluorescence microscopy," *MIB*, vol. 15(1), pp. 3–11, 2013.
- [2] M. Derieppe et al., "Tracking of cell nuclei for assessment of in vitro uptake kinetics in ultrasound-mediated drug delivery using fibred confocal fluorescence microscopy," *MIB*, vol. 16(5), pp. 642–651, 2014.