



Cyclicity in the harmonic Dirichlet space

Evgueni Abakumov, Omar El-Fallah, Karim Kellay, Thomas Ransford

► **To cite this version:**

Evgueni Abakumov, Omar El-Fallah, Karim Kellay, Thomas Ransford. Cyclicity in the harmonic Dirichlet space . Conference on Harmonic and Functional, Analysis, Operator Theory and Applications., Jun 2015, Bordeaux, France. Theta Series in Advanced Mathematics. <hal-01261077>

HAL Id: hal-01261077

<https://hal.archives-ouvertes.fr/hal-01261077>

Submitted on 23 Jan 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CYCLICITY IN THE HARMONIC DIRICHLET SPACE

E. ABAKUMOV, O. EL-FALLAH, K. KELLAY, AND T. RANSFORD

ABSTRACT. The harmonic Dirichlet space is the Hilbert space of functions $f \in L^2(\mathbb{T})$ such that

$$\|f\|_{\mathcal{D}(\mathbb{T})}^2 := \sum_{n \in \mathbb{Z}} (1 + |n|) |\widehat{f}(n)|^2 < \infty.$$

We give sufficient conditions for f to be cyclic in $\mathcal{D}(\mathbb{T})$, in other words, for $\{\zeta^n f(\zeta) : n \geq 0\}$ to span a dense subspace of $\mathcal{D}(\mathbb{T})$.

1. INTRODUCTION

Let \mathcal{X} be a topological linear space of complex functions on the unit circle \mathbb{T} such that the shift operator S , given by

$$S(f)(\zeta) := \zeta f(\zeta), \quad f \in \mathcal{X},$$

is an isomorphism of \mathcal{X} onto itself.

A closed subspace \mathcal{M} of \mathcal{X} is called *invariant* if $S(\mathcal{M}) \subset \mathcal{M}$. It is said to be *1-invariant* (or *simply invariant*) for S if $S(\mathcal{M}) \subsetneq \mathcal{M}$, and it is called *2-invariant* (or *doubly invariant*) if $S(\mathcal{M}) = \mathcal{M}$. The latter condition is equivalent to the invariance of \mathcal{M} under multiplication by both ζ and $\bar{\zeta}$.

Let \mathbb{Z} denote the integers and let $\mathbb{N} := \{n \in \mathbb{Z} : n \geq 0\}$. Given $f \in \mathcal{X}$, we write

$$\begin{aligned} [f]_{\mathbb{N}} &:= \overline{\text{Span}}^{\mathcal{X}} \{z^n f : n \in \mathbb{N}\}, \\ [f]_{\mathbb{Z}} &:= \overline{\text{Span}}^{\mathcal{X}} \{z^n f : n \in \mathbb{Z}\}. \end{aligned}$$

A function f is said to be *1-invariant* if the space $[f]_{\mathbb{N}}$ is 1-invariant for S . We say that a function $f \in \mathcal{X}$ is *cyclic* (resp. *bicyclic*) for \mathcal{X} if $[f]_{\mathbb{N}} = \mathcal{X}$ (resp. $[f]_{\mathbb{Z}} = \mathcal{X}$).

2010 *Mathematics Subject Classification*. Primary 46E22; Secondary 31A05, 31A15, 31A20, 47B32.

Key words and phrases. Harmonic Dirichlet space, capacity, cyclic vectors.

Research of OEF partially supported by Hassan II Academy of Science and Technology. Research of TR supported by NSERC and the Canada Research Chairs program.

Let us begin with the classical case, namely $\mathcal{X} = L^2(\mathbb{T})$. By a well-known theorem of Wiener, the 2-invariant subspaces have the form

$$\mathcal{M} = \{f \in L^2(\mathbb{T}) : f = 0 \text{ a.e. on } \mathbb{T} \setminus \sigma\},$$

where σ is a Borel subset of \mathbb{T} (see e.g. [11, p.8, Theorem 1.2.1]). It follows from Szegő's infimum theorem that a function f is 1-invariant in $L^2(\mathbb{T})$ if and only if $\log |f| \in L^1(\mathbb{T})$ (see e.g. [10, p.12, Corollary 4]).

For $\mathcal{X} = C^\infty(\mathbb{T})$, Makarov [7] gave a complete description of the invariant subspaces of S . He also obtained the following characterization of 1-invariant functions of $C^\infty(\mathbb{T})$.

Theorem (Makarov [7, p.3]). *A function f is 1-invariant in $C^\infty(\mathbb{T})$ if and only if $\log |f| \in L^1(\mathbb{T})$.*

The case $\mathcal{X} = C^n(\mathbb{T})$ ($n \geq 1$) is more complicated, and no characterization of 1-invariant functions is known ([7, p.3], see also [8, Theorem 1.3] or [9, Theorem 5]).

We shall focus our attention on the *harmonic Dirichlet space* $\mathcal{D}(\mathbb{T})$. This is the set of functions $f \in L^2(\mathbb{T})$ whose Fourier coefficients satisfy

$$\mathcal{D}(f) := \sum_{n \in \mathbb{Z}} |\widehat{f}(n)|^2 |n| < \infty.$$

It becomes a Hilbert space if endowed with the norm $\|\cdot\|_{\mathcal{D}(\mathbb{T})}$, given by

$$\|f\|_{\mathcal{D}(\mathbb{T})}^2 := \|f\|_{L^2(\mathbb{T})}^2 + \mathcal{D}(f) = \sum_{n \in \mathbb{Z}} |\widehat{f}(n)|^2 (1 + |n|).$$

According to Douglas' formula [3], we have

$$\mathcal{D}(f) = \frac{1}{4\pi^2} \iint_{\mathbb{T}^2} \frac{|f(\zeta) - f(\zeta')|^2}{|\zeta - \zeta'|^2} |d\zeta'| |d\zeta|.$$

This can also be written

$$\mathcal{D}(f) = \frac{1}{2\pi} \int_{\mathbb{T}} \mathcal{D}_\zeta(f) |d\zeta|,$$

where $\mathcal{D}_\zeta(f)$ is the so-called local Dirichlet integral of f at ζ , given by

$$\mathcal{D}_\zeta(f) := \frac{1}{2\pi} \int_{\mathbb{T}} \frac{|f(\zeta) - f(\zeta')|^2}{|\zeta - \zeta'|^2} |d\zeta'|, \quad \zeta \in \mathbb{T}.$$

As $\mathcal{D}(\mathbb{T})$ is dense in $L^2(\mathbb{T})$, if a function f is cyclic for $\mathcal{D}(\mathbb{T})$, then it is also cyclic for $L^2(\mathbb{T})$. So by Szegő's theorem we have

$$f \text{ is cyclic for } \mathcal{D}(\mathbb{T}) \Rightarrow \int_{\mathbb{T}} \log |f(\zeta)| |d\zeta| = -\infty.$$

In [12], Ross, Richter and Sundberg gave a complete characterization of the 2-invariant subspaces \mathcal{M} of $\mathcal{D}(\mathbb{T})$ in terms of their zero

sets. In order to state their result, we need to introduce the notion of logarithmic capacity (see for instance [6, §2.4]).

The *energy* of a Borel probability measure μ on \mathbb{T} is defined by

$$I(\mu) := \iint_{\mathbb{T}^2} \log \frac{1}{|\zeta - \zeta'|} d\mu(\zeta) d\mu(\zeta') = \sum_{n=1}^{\infty} \frac{|\widehat{\mu}(n)|^2}{n}.$$

Then we define the *logarithmic capacity* of a Borel subset E of \mathbb{T} by

$$c(E) := 1 / \inf\{I(\mu) : \mu \in \mathcal{P}(E)\},$$

where $\mathcal{P}(E)$ denotes the set of probability measures supported on a compact subset of E . We say that a property holds *quasi-everywhere* (*q.e.*) if it holds everywhere outside a set of logarithmic capacity zero.

It is known that, if $f \in \mathcal{D}(\mathbb{T})$, then the radial limit of the Poisson integral of f exists q.e. and is equal to f a.e. (for more details we refer to [12] and the references therein). In the sequel, f will denote this limit, and will therefore be defined q.e. on \mathbb{T} . We shall write $\mathcal{Z}(f)$ for the zero set of f , namely

$$\mathcal{Z}(f) := \{\zeta \in \mathbb{T} : f(\zeta) = 0\}.$$

Note that this set is defined up to sets of logarithmic capacity zero.

Theorem (Richter–Ross–Sundberg [12]). *\mathcal{M} is a 2-invariant subspace of $\mathcal{D}(\mathbb{T})$ if and only if there exists a measurable set $E \subset \mathbb{T}$ such that*

$$\mathcal{M} = \mathcal{D}_E := \{f \in \mathcal{D}(\mathbb{T}) : f|_E = 0 \text{ q.e.}\}.$$

Note that the problem of characterization of 1-invariant subspaces of $\mathcal{D}(\mathbb{T})$ remains open. It was proved in [1, 13] that, for each $n \in \mathbb{N} \cup \{\infty\}$, there exists an invariant subspace \mathcal{M} of $\mathcal{D}(\mathbb{T})$ such that $\dim(\mathcal{M}/S(\mathcal{M})) = n$. This suggests that the lattice of 1-invariant subspaces has a very complicated structure.

As a direct consequence of the Richter–Ross–Sundberg theorem, we obtain the following necessary conditions for cyclicity in $\mathcal{D}(\mathbb{T})$.

Theorem 1. *If f is cyclic for $\mathcal{D}(\mathbb{T})$, then*

$$\int_{\mathbb{T}} \log |f(\zeta)| |d\zeta| = -\infty \quad \text{and} \quad c(\mathcal{Z}(f)) = 0.$$

Our goal in this paper is to give sufficient conditions for a function $f \in \mathcal{D}(\mathbb{T})$ to be cyclic.

For $\beta \in (0, 1]$, we shall denote by $\text{Lip}_\beta(\mathbb{T})$ the set of functions f continuous on \mathbb{T} such that

$$\|f\|_{\text{Lip}_\beta(\mathbb{T})} := \|f\|_{C(\mathbb{T})} + \sup_{\zeta, \zeta' \in \mathbb{T}} \frac{|f(\zeta) - f(\zeta')|}{|\zeta - \zeta'|^\beta} < \infty.$$

For $\alpha \in (0, 1)$, we set

$$\mathcal{C}^{1+\alpha}(\mathbb{T}) := \{f \in C^1(\mathbb{T}) : f' \in \text{Lip}_\alpha(\mathbb{T})\}.$$

Of course, if f belongs to $\text{Lip}_\beta(\mathbb{T})$ or $C^{1+\alpha}(\mathbb{T})$, then $\mathcal{Z}(f)$ is closed in \mathbb{T} .

We shall establish the following result.

Theorem 2. *Let $f \in \mathcal{D}(\mathbb{T})$ such that $|f| \in \mathcal{C}^{1+\alpha}(\mathbb{T})$, where $\alpha \in (0, 1)$. Suppose further that $\log |f| \notin L^1(\mathbb{T})$. Then $[f^2]_{\mathbb{N}} = \mathcal{D}_{\mathcal{Z}(f)}$.*

Combining Theorems 1 and 2, we deduce

Corollary. *Let $f \in \mathcal{D}(\mathbb{T})$ such that $|f| \in \mathcal{C}^{1+\alpha}(\mathbb{T})$, where $\alpha \in (0, 1)$. Then the following assertions are equivalent:*

- (1) f^2 is cyclic for $\mathcal{D}(\mathbb{T})$;
- (2) $\log |f| \notin L^1(\mathbb{T})$ and $c(\mathcal{Z}(f)) = 0$.

A closed set $E \subset \mathbb{T}$ is said to be a *Carleson set* (and we write $E \in (C)$) if

$$\int_{\mathbb{T}} \log \frac{1}{d(\zeta, E)} |d\zeta| < \infty.$$

For background information on Carleson sets, see e.g. [6, §4.4]. Note that, if $f \in \text{Lip}_\beta(\mathbb{T})$ and $\mathcal{Z}(f) \notin (C)$, then $\log |f| \notin L^1(\mathbb{T})$.

It is known that $\text{Lip}_\beta(\mathbb{T}) \subset \mathcal{D}(\mathbb{T})$ if and only if $\beta > 1/2$. The inclusion $\text{Lip}_\beta(\mathbb{T}) \subset \mathcal{D}(\mathbb{T})$ for $\beta > 1/2$ can easily be obtained from Douglas' formula.

We shall establish the following theorem.

Theorem 3. *Let $f \in \text{Lip}_\beta(\mathbb{T})$, where $\beta \in (\frac{1}{2}, 1]$. If $\mathcal{Z}(f) \notin (C)$, then $[f]_{\mathbb{N}} = \mathcal{D}_{\mathcal{Z}(f)}$. If furthermore $c(\mathcal{Z}(f)) = 0$, then f is cyclic for $\mathcal{D}(\mathbb{T})$.*

2. PROOF OF THEOREM 2

For the proof of Theorem 2, we shall need the following standard result.

Lemma 4. *Let $f \in \mathcal{D}(\mathbb{T})$. The following assertions are equivalent:*

- (1) $[f]_{\mathbb{N}} = [f]_{\mathbb{Z}}$;
- (2) $f \in [Sf]_{\mathbb{N}}$;
- (3) $\inf\{\|pf\|_{\mathcal{D}(\mathbb{T})} : p \in H^\infty, pf \in \mathcal{D}(\mathbb{T}) \text{ and } p(0) = 1\} = 0$.

Proof. Since S is invertible, (1) and (2) are equivalent.

If $f \in [Sf]_{\mathbb{N}}$, then there is a sequence (p_n) of polynomials such that $p_n(0) = 1$ and $\|(1 - p_n)f\|_{\mathcal{D}(\mathbb{T})} \rightarrow 0$. This proves that (2) implies (3).

Finally, suppose that (3) holds. Let $(p_n) \subset H^\infty$ be a sequence such that $p_n(0) = 1$, $p_n f \in \mathcal{D}(\mathbb{T})$ and $\|p_n f\|_{\mathcal{D}(\mathbb{T})} \rightarrow 0$. Writing $p_n = 1 - zq_n$, by [12, Proposition 3.4] we have $zq_n f \in [Sf]_{\mathbb{N}}$. Since $zq_n f$ converges to f , it follows that $f \in [Sf]_{\mathbb{N}}$, so that (2) holds. \square

We shall also need the following result, which is a special case of a theorem due to Carleson–Jacobs–Havin–Shamoyan [2, Theorem 6.1].

Lemma 5. *Let F be an outer function on \mathbb{D} that is continuous on $\overline{\mathbb{D}}$. If $|F| \in \mathcal{C}^{1+\alpha}(\mathbb{T})$, where $\alpha \in (0, 1)$, then $F \in \text{Lip}_{(1+\alpha)/2}(\overline{\mathbb{D}})$. Furthermore, the Lipschitz constant associated to F on $\overline{\mathbb{D}}$ depends only on the Lipschitz constants and bounds for the derivatives of $|F|$ on \mathbb{T} .*

Proof of Theorem 2. Let p_ϵ be the outer function such that

$$|p_\epsilon(\zeta)| = \frac{e^{-M_\epsilon}}{|f(\zeta)| + \epsilon} \quad \text{a.e. on } \mathbb{T},$$

where the constant M_ϵ is chosen so that $p_\epsilon(0) = 1$. Thus

$$M_\epsilon = \int_{\mathbb{T}} \log \left(\frac{1}{|f(\zeta)| + \epsilon} \right) \frac{|d\zeta|}{2\pi},$$

and since $\log |f| \notin L^1(\mathbb{T})$, it follows that $M_\epsilon \rightarrow \infty$ as $\epsilon \rightarrow 0^+$. We are going to prove that

$$\lim_{\epsilon \rightarrow 0^+} \|p_\epsilon f^2\|_{\mathcal{D}(\mathbb{T})} = 0.$$

If this holds, then by Lemma 4 we have $[f^2]_{\mathbb{N}} = [f^2]_{\mathbb{Z}}$, and since clearly $\mathcal{Z}(f^2) = \mathcal{Z}(f)$, we can apply the Richter–Ross–Sundberg theorem to obtain the desired result.

We have

$$\|p_\epsilon f^2\|_{\mathcal{D}(\mathbb{T})}^2 = \|p_\epsilon f^2\|_{L^2(\mathbb{T})}^2 + \mathcal{D}(p_\epsilon f^2).$$

For the first term, we have

$$\|p_\epsilon f^2\|_{L^2(\mathbb{T})}^2 = \int_{\mathbb{T}} \frac{e^{-2M_\epsilon} |f|^4 |d\zeta|}{(|f| + \epsilon)^2} \leq e^{-2M_\epsilon} \|f^2\|_{L^2(\mathbb{T})}^2 \rightarrow 0 \quad \text{as } \epsilon \rightarrow 0^+.$$

The second term we estimate using Douglas' formula, namely

$$\mathcal{D}(p_\epsilon f^2) = \frac{1}{4\pi^2} \iint_{\mathbb{T}^2} \frac{|(p_\epsilon f^2)(\zeta) - (p_\epsilon f^2)(\zeta')|^2}{|\zeta - \zeta'|^2} |d\zeta'| |d\zeta|.$$

Let

$$\Gamma := \{(\zeta, \zeta') \in \mathbb{T}^2 : |f(\zeta')| \leq |f(\zeta)|\}.$$

Then, by symmetry,

$$\mathcal{D}(p_\epsilon f) = 2 \frac{1}{4\pi^2} \iint_{\Gamma} \frac{|(p_\epsilon f)(\zeta) - (p_\epsilon f)(\zeta')|^2}{|\zeta - \zeta'|^2} |d\zeta'| |d\zeta|.$$

Now, for all $\zeta, \zeta' \in \mathbb{T}$, we have

$$\begin{aligned} & |(p_\epsilon f^2)(\zeta) - (p_\epsilon f^2)(\zeta')|^2 \\ &= |p_\epsilon(\zeta)(f^2(\zeta) - f^2(\zeta')) + f^2(\zeta')(p_\epsilon(\zeta) - p_\epsilon(\zeta'))|^2 \\ &\leq 2|p_\epsilon(\zeta)|^2 |f^2(\zeta) - f^2(\zeta')|^2 + 2|f^2(\zeta')|^2 |p_\epsilon(\zeta) - p_\epsilon(\zeta')|^2. \end{aligned}$$

Hence

$$\iint_{\Gamma} \frac{|(p_{\epsilon}f^2)(\zeta) - (p_{\epsilon}f^2)(\zeta')|^2}{|\zeta - \zeta'|^2} |d\zeta'| |d\zeta| \leq 2A_{\epsilon} + 2B_{\epsilon},$$

where

$$A_{\epsilon} := \iint_{\Gamma} |p_{\epsilon}(\zeta)|^2 \frac{|f^2(\zeta) - f^2(\zeta')|^2}{|\zeta - \zeta'|^2} |d\zeta'| |d\zeta|$$

and

$$B_{\epsilon} := \iint_{\Gamma} |f^2(\zeta')|^2 \frac{|p_{\epsilon}(\zeta) - p_{\epsilon}(\zeta')|^2}{|\zeta - \zeta'|^2} |d\zeta'| |d\zeta|.$$

We estimate A_{ϵ} directly as follows:

$$\begin{aligned} A_{\epsilon} &= e^{-2M_{\epsilon}} \iint_{\Gamma} \frac{|f(\zeta) + f(\zeta')|^2}{(|f(\zeta)| + \epsilon)^2} \frac{|f(\zeta) - f(\zeta')|^2}{|\zeta - \zeta'|^2} |d\zeta'| |d\zeta| \\ &\leq 4e^{-2M_{\epsilon}} \iint_{\mathbb{T}^2} \frac{|f(\zeta) - f(\zeta')|^2}{|\zeta - \zeta'|^2} |d\zeta'| |d\zeta| \\ &\leq 4e^{-2M_{\epsilon}} 4\pi^2 \mathcal{D}(f). \end{aligned}$$

Hence $A_{\epsilon} \rightarrow 0$ as $\epsilon \rightarrow 0^+$.

To estimate B_{ϵ} , we consider the outer function F_{ϵ} such that

$$|F_{\epsilon}(\zeta)| = |f(\zeta)| + \epsilon \quad \text{a.e. on } \mathbb{T}.$$

By Lemma 5, since $|F_{\epsilon}| \in \mathcal{C}^{1+\alpha}(\mathbb{T})$, we have $F_{\epsilon} \in \text{Lip}_{(1+\alpha)/2}(\mathbb{T}) \subset \mathcal{D}(\mathbb{T})$ and there exists a positive constant D , depending only on $|f|$, such that $\mathcal{D}(F_{\epsilon}) \leq D$ for all $\epsilon \in (0, 1)$. We then have

$$\begin{aligned} B_{\epsilon} &= \iint_{\Gamma} \frac{e^{-2M_{\epsilon}} |f^2(\zeta')|^2}{(|f(\zeta)| + \epsilon)^2 (|f(\zeta')| + \epsilon)^2} \frac{|1/p_{\epsilon}(\zeta) - 1/p_{\epsilon}(\zeta')|^2}{|\zeta - \zeta'|^2} |d\zeta'| |d\zeta| \\ &\leq e^{-2M_{\epsilon}} \iint_{\Gamma} \frac{|F_{\epsilon}(\zeta) - F_{\epsilon}(\zeta')|^2}{|\zeta - \zeta'|^2} |d\zeta'| |d\zeta| \\ &\leq e^{-2M_{\epsilon}} 4\pi^2 \mathcal{D}(F_{\epsilon}). \end{aligned}$$

Thus $B_{\epsilon} \rightarrow 0$ as $\epsilon \rightarrow 0^+$. This completes the proof of Theorem 2. \square

3. PROOF OF THEOREM 3

To prove Theorem 3, we shall need the following additional lemma.

Lemma 6. *Let $f \in \text{Lip}_{\beta}(\mathbb{T})$, where $\beta > 1/2$. Then, for $\eta \in (0, \frac{2\beta-1}{2\beta})$, we have*

$$\iint_{\mathbb{T}^2} \frac{|f(\zeta) - f(\zeta')|^{2-2\eta}}{|\zeta - \zeta'|^2} |d\zeta'| |d\zeta| < +\infty.$$

Proof. Since $\beta > 1/2(1 - \eta)$, we get

$$\iint_{\mathbb{T}^2} \frac{|f(\zeta) - f(\zeta')|^{2-2\eta}}{|\zeta - \zeta'|^2} |d\zeta'| |d\zeta| \lesssim \iint_{\mathbb{T}^2} \frac{|d\zeta'| |d\zeta|}{|\zeta - \zeta'|^{2(1-(1-\eta)\beta)}} < \infty. \quad \square$$

Note that here, and in what follows, we write $A \lesssim B$ to mean that there is an absolute constant C such that $A \leq CB$.

Proof of Theorem 3. By [12], it suffices to prove that $[f]_{\mathbb{N}}$ is 2-invariant, which is equivalent to proving that $f \in [Sf]_{\mathbb{N}}$.

Let $\epsilon, \gamma > 0$, where γ will be taken small. Let E be a closed subset of $\mathcal{Z}(f)$ such that $|E| = 0$ and $E \notin (C)$. Let p_ϵ be the outer function satisfying

$$|p_\epsilon(\zeta)| = \frac{e^{-M_\epsilon}}{(d(\zeta, E)^\gamma + \epsilon)^{1/2}} \quad \text{a.e. on } \mathbb{T},$$

where the constant M_ϵ is chosen so that $p_\epsilon(0) = 1$. Thus

$$M_\epsilon := \frac{1}{2} \int_{\mathbb{T}} \log \left(\frac{1}{d(\zeta, E)^\gamma + \epsilon} \right) \frac{|d\zeta|}{2\pi},$$

and since $E \notin (C)$, it follows that $M_\epsilon \rightarrow \infty$ as $\epsilon \rightarrow 0^+$. By Lemma 4, it suffices to prove that

$$\lim_{\epsilon \rightarrow 0^+} \|p_\epsilon f\|_{\mathcal{D}(\mathbb{T})} = 0.$$

Now

$$\|p_\epsilon f\|_{\mathcal{D}(\mathbb{T})}^2 = \|p_\epsilon f\|_{L^2(\mathbb{T})}^2 + \mathcal{D}(p_\epsilon f).$$

For the first term, we have

$$\|p_\epsilon f\|_{L^2(\mathbb{T})}^2 \lesssim e^{-2M_\epsilon} \int_{\mathbb{T}} \frac{d(\zeta, E)^{2\beta}}{d(\zeta, E)^\gamma} |d\zeta|.$$

Thus $\|p_\epsilon f\|_{L^2(\mathbb{T})} \rightarrow 0$ as $\epsilon \rightarrow 0^+$, provided that $\gamma < 2\beta$.

For the second term we again use Douglas' formula, namely

$$\mathcal{D}(p_\epsilon f) = \frac{1}{4\pi^2} \iint_{\mathbb{T}^2} \frac{|(p_\epsilon f)(\zeta) - (p_\epsilon f)(\zeta')|^2}{|\zeta - \zeta'|^2} |d\zeta'| |d\zeta|.$$

Let

$$\Gamma := \{(\zeta, \zeta') \in \mathbb{T}^2 : d(\zeta', E) \leq d(\zeta, E)\}.$$

Arguing as in the proof of Theorem 2, we have

$$\begin{aligned} \mathcal{D}(p_\epsilon f) &\lesssim \iint_{\Gamma} |p_\epsilon(\zeta)|^2 \frac{|f(\zeta) - f(\zeta')|^2}{|\zeta - \zeta'|^2} |d\zeta| |d\zeta'| \\ &\quad + \iint_{\Gamma} |f(\zeta')|^2 \frac{|p_\epsilon(\zeta) - p_\epsilon(\zeta')|^2}{|\zeta - \zeta'|^2} |d\zeta| |d\zeta'| \\ &= A_\epsilon + B_\epsilon, \text{ say.} \end{aligned}$$

To estimate A_ϵ , let $\eta \in (0, (2\beta - 1)/(2\beta))$ and let $\gamma < 2\beta\eta$. Then

$$\begin{aligned} A_\epsilon &= e^{-2M_\epsilon} \iint_{\Gamma} \left(\frac{|f(\zeta) - f(\zeta')|^{2\eta}}{d(\zeta, E)^\gamma + \epsilon} \right) \left(\frac{|f(\zeta) - f(\zeta')|^{2-2\eta}}{|\zeta - \zeta'|^2} \right) |d\zeta| |d\zeta'| \\ &\leq 2C e^{-2M_\epsilon} \iint_{\mathbb{T}^2} \frac{d(\zeta, E)^{2\beta\eta}}{d(\zeta, E)^\gamma} \left(\frac{|f(\zeta) - f(\zeta')|^{2-2\eta}}{|\zeta - \zeta'|^2} \right) |d\zeta| |d\zeta'| \\ &\leq 2C^{-2M_\epsilon} \iint_{\mathbb{T}^2} \frac{|f(\zeta) - f(\zeta')|^{2-2\eta}}{|\zeta - \zeta'|^2} |d\zeta| |d\zeta'|, \end{aligned}$$

where C depends only on f , and the double integral is finite, thanks to Lemma 6. Hence $A_\epsilon \rightarrow 0$ as $\epsilon \rightarrow 0^+$.

To estimate B_ϵ , we introduce the outer function F_ϵ satisfying

$$|F_\epsilon(\zeta)| = d(\zeta, E)^\gamma + \epsilon \quad \text{a.e. on } \mathbb{T}.$$

By the Carleson–Richter–Sundberg formula [6, Theorem 7.4.2], we have

$$\mathcal{D}_\zeta(F_\epsilon) = \int_{\mathbb{T}} \frac{|F_\epsilon(\zeta)|^2 - |F_\epsilon(\zeta')|^2 - 2|F_\epsilon(\zeta')| \log |F_\epsilon(\zeta)/F_\epsilon(\zeta')|}{|\zeta - \zeta'|^2} \frac{|d\zeta'|}{2\pi}.$$

Therefore

$$\begin{aligned} B_\epsilon &\lesssim e^{-2M_\epsilon} \iint_{\Gamma} \frac{d(\zeta', E)^{2\beta}}{(d(\zeta, E)^\gamma + \epsilon)(d(\zeta', E)^\gamma + \epsilon)} \frac{|F_\epsilon(\zeta) - F_\epsilon(\zeta')|^2}{|\zeta - \zeta'|^2} |d\zeta| |d\zeta'| \\ &\lesssim e^{-2M_\epsilon} \iint_{\mathbb{T}^2} \frac{|F_\epsilon(\zeta) - F_\epsilon(\zeta')|^2}{|\zeta - \zeta'|^2} d(\zeta', E)^{2(\beta-\gamma)} |d\zeta| |d\zeta'| \\ &\lesssim e^{-2M_\epsilon} \int_{\mathbb{T}} \mathcal{D}_\zeta(F_\epsilon) d(\zeta, E)^{2(\beta-\gamma)} |d\zeta| \\ &\lesssim e^{-2M_\epsilon} \iint_{\mathbb{T}^2} \left(\frac{(|F_\epsilon(\zeta)|^2 - |F_\epsilon(\zeta')|^2) \log |F_\epsilon(\zeta)/F_\epsilon(\zeta')|}{|\zeta - \zeta'|^2} \right) \\ &\quad \times \left(d(\zeta, E)^{2(\beta-\gamma)} + d(\zeta', E)^{2(\beta-\gamma)} \right) |d\zeta| |d\zeta'|. \end{aligned}$$

Exchanging the roles of ζ and ζ' , and taking the average, we obtain

$$\begin{aligned} B_\epsilon &\lesssim e^{-2M_\epsilon} \iint_{\mathbb{T}^2} \left(\frac{(|F_\epsilon(\zeta)|^2 - |F_\epsilon(\zeta')|^2) \log |F_\epsilon(\zeta)/F_\epsilon(\zeta')|}{|\zeta - \zeta'|^2} \right) \\ &\quad \times \left(d(\zeta, E)^{2(\beta-\gamma)} + d(\zeta', E)^{2(\beta-\gamma)} \right) |d\zeta| |d\zeta'|. \end{aligned}$$

Thus

$$B_\epsilon \lesssim e^{-2M_\epsilon} \iint_{\mathbb{T}^2} \frac{\delta^\gamma - \delta'^\gamma}{|\zeta - \zeta'|^2} \log \left(\frac{\delta^\gamma + \epsilon}{\delta'^\gamma + \epsilon} \right) (\delta^{2(\beta-\gamma)} + \delta'^{2(\beta-\gamma)}) |d\zeta| |d\zeta'|, \quad (3.1)$$

where $\delta := d(\zeta, E)$ and $\delta' := d(\zeta', E)$.

Let (I_j) be the connected components of $\mathbb{T} \setminus E$, and set

$$N_E(t) := 2 \sum_j 1_{\{|I_j| > 2t\}}, \quad 0 < t < 1.$$

Then, for every measurable function $\Omega : [0, \pi] \rightarrow \mathbb{R}^+$, we have

$$\int_{\mathbb{T}} \Omega(d(\zeta, E)) |d\zeta| = \int_0^\pi \Omega(t) N_E(t) dt.$$

Using similar ideas to those in [4, 5], we obtain

$$\begin{aligned} J &:= \iint_{\mathbb{T}^2} \frac{\delta^\gamma - \delta'^\gamma}{|\zeta - \zeta'|^2} \log\left(\frac{\delta^\gamma + \epsilon}{\delta'^\gamma + \epsilon}\right) (\delta^{2(\beta-\gamma)} + \delta'^{2(\beta-\gamma)}) |d\zeta| |d\zeta'| \\ &\lesssim \int_0^\pi \int_0^\pi \frac{((s+t)^\gamma - t^\gamma)}{s^2} \log\left(\frac{(s+t)^\gamma + \epsilon}{t^\gamma + \epsilon}\right) (t+s)^{2(\beta-\gamma)} N_E(t) ds dt \\ &\lesssim \int_0^\pi \int_0^t \frac{((s+t)^\gamma - t^\gamma) \log[(s+t)^\gamma/t^\gamma]}{s^2} (t+s)^{2(\beta-\gamma)} N_E(t) ds dt \\ &\quad + \int_0^\pi \int_t^\pi \frac{((s+t)^\gamma - t^\gamma)}{s^2} \log[1/(t^\gamma + \epsilon)] (t+s)^{2(\beta-\gamma)} ds N_E(t) dt \\ &= J_1 + J_2, \end{aligned}$$

where

$$\begin{aligned} J_1 &\lesssim \int_0^\pi t^{2\beta-\gamma-1} \int_0^1 \frac{((1+x)^\gamma - 1)}{x^2} \log(1+x) dx N_E(t) dt \\ &\lesssim \int_0^\pi t^{2\beta-\gamma-1} N_E(t) dt = O(1), \end{aligned}$$

and

$$\begin{aligned} J_2 &\lesssim \int_0^\pi t^{2\beta-\gamma-1} \log[1/(t^\gamma + \epsilon)] \int_1^{\pi/t} \frac{(1+x)^\gamma - 1}{s^2} (1+x)^{2(\beta-\gamma)} dx N_E(t) dt \\ &\lesssim \int_0^\pi \log[1/(t^\gamma + \epsilon)] N_E(t) dt \\ &\lesssim \int_{\mathbb{T}} |\log(d(\zeta, E)^\gamma + \epsilon)| |d\zeta| = 2M_\epsilon. \end{aligned}$$

Thus $J = O(M_\epsilon)$. Combining this with the estimate (3.1), we get

$$B_\epsilon \lesssim M_\epsilon e^{-2M_\epsilon}.$$

Hence $B_\epsilon \rightarrow 0$ as $\epsilon \rightarrow 0^+$. This completes the proof of Theorem 3. \square

4. CONCLUDING REMARKS

1. In order to produce cyclic functions for $\mathcal{D}(\mathbb{T})$ using Theorem 3, we need to construct closed subsets $E \subset \mathbb{T}$ such that $E \notin (C)$ and $c(E) = 0$. An easy example can be given by countable sets. Indeed, taking $E_\beta := \{e^{i/(\log n)^\beta} : n \geq 2\}$ with $\beta \leq 1$ provides such an example. Using Cantor-type sets, it is also possible to construct perfect sets E such that $E \notin (C)$ and $c(E) = 0$.

2. One can consider weighted harmonic Dirichlet spaces instead of the classical harmonic Dirichlet space. More precisely, given $\alpha \in [0, 1)$, the *weighted harmonic Dirichlet space* $\mathcal{D}_\alpha(\mathbb{T})$ is the space of functions $f \in L^2(\mathbb{T})$ such that

$$\|f\|_{\mathcal{D}_\alpha(\mathbb{T})}^2 := \sum_{n \in \mathbb{Z}} |\widehat{f}(n)|^2 (1 + |n|)^{1-\alpha} < \infty.$$

We define the α -*capacity* of a Borel subset $E \subset \mathbb{T}$ by

$$c_\alpha(E) = 1 / \inf\{I_\alpha(\mu) : \mu \in \mathcal{P}(E)\},$$

where $\mathcal{P}(E)$ is the set of all probability measures supported on a compact subset of E and $I_\alpha(\mu) := \sum_{n \geq 1} |\widehat{\mu}(n)|^2 / n^{1-\alpha}$ is the α -*energy* of μ . We say that a property holds c_α -*quasi-everywhere* if it holds everywhere outside a set of c_α -capacity zero.

It is well known that $\text{Lip}_\beta(\mathbb{T}) \subset \mathcal{D}_\alpha(\mathbb{T})$ if and only $\beta > (1 - \alpha)/2$. Theorem 3 may be extended to show that, if $f \in \text{Lip}_\beta(\mathbb{T})$, where $\beta \in ((1 - \alpha)/2, 1]$, and if $\mathcal{Z}(f) \notin (C)$, then

$$[f]_{\mathbb{N}} = \{g \in \mathcal{D}(\mathbb{T}) : g|_{\mathcal{Z}(f)} = 0 \text{ } c_\alpha\text{-quasi-everywhere}\}.$$

3. One can equally well consider the holomorphic Dirichlet space, namely $\mathcal{D} := \{f \in \mathcal{D}(\mathbb{T}) : \widehat{f}(n) = 0 \text{ } (n < 0)\}$. Here too the problem of characterizing the cyclic functions is still open. For more on this topic, see e.g. [6, Chapter 9].

ACKNOWLEDGEMENT

Part of the research for this article was carried out while the authors were visiting the University of Bordeaux to participate at the Conference on Harmonic Analysis, Function Theory, Operator Theory and Applications in honor of Jean Esterle. The authors thank the Institut de Mathématiques de Bordeaux for its support and hospitality.

REFERENCES

- [1] A. Aleman, S. Richter, W. Ross, Bergman spaces on disconnected domains, *Canad. J. Math.* 48 (1996), no. 2, 225–243.

- [2] J. E. Brennan, Approximation in the mean by polynomials on non-Carathéodory domains, *Ark. Mat.* 15 (1977), no. 1, 117–168.
- [3] J. Douglas, Solution of the problem of Plateau, *Trans. Amer. Math. Soc.* 33 (1931), no. 1, 263–321.
- [4] O. El-Fallah, K. Kellay, T. Ransford, On the Brown–Shields conjecture for cyclicity in the Dirichlet space, *Adv. Math.* 222 (2009), no. 6, 2196–2214.
- [5] O. El-Fallah, K. Kellay, T. Ransford, Cantor sets and cyclicity in weighted Dirichlet spaces, *J. Math. Anal. Appl.* 372 (2010), no. 2, 565–573.
- [6] O. El-Fallah, K. Kellay, J. Mashreghi, T. Ransford, *A Primer on the Dirichlet Space*, Cambridge University Press, Cambridge, 2014.
- [7] N. G. Makarov, Invariant subspaces of the space C^∞ , *Mat. Sb. (N.S.)* 119(161) (1982), no. 1, 1–31, 160.
- [8] N. G. Makarov, Sets of simple invariance. Investigations on linear operators and the theory of functions, X, *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)* 107 (1982), 104–135, 230.
- [9] N. G. Makarov, Sets of 1-invariance and 1-invariant subspaces (smooth functions), *Dokl. Akad. Nauk SSSR* 262 (1982), no. 5, 1072–1075.
- [10] N. K. Nikolskii, *Treatise on the Shift Operator*, Springer, Berlin, 1986.
- [11] N. K. Nikolski. *Operators, Functions, and Systems: An Easy Reading, Vol. 1: Hardy, Hankel, and Toeplitz*, Amer. Math. Soc., Providence RI, 2002.
- [12] S. Richter, W. Ross, C. Sundberg, Hyperinvariant subspaces of the harmonic Dirichlet space, *J. Reine Angew. Math.* 448 (1994), 1–26.
- [13] W. Ross, Invariant subspaces of the harmonic Dirichlet space with large co-dimension, *Proc. Amer. Math. Soc.* 124 (1996), no. 6, 1841–1846.

LAMA, UMR CNRS 8050. UNIVERSITÉ PARIS–EST, 5 BOULEVARD DESCARTES,
CHAMPS–SUR–MARNE, 77454 MARNE–LA–VALLÉE CEDEX 2, FRANCE
E-mail address: `evgueni.abakoumov@u-pem.fr`

LABORATOIRE ANALYSE ET APPLICATIONS URAC/03, MOHAMMED V UNI-
VERSITY IN RABAT, B.P. 1014 RABAT, MOROCCO
E-mail address: `elfallah@fsr.ac.ma`

IMB, UNIVERSITÉ DE BORDEAUX, 351 COURS DE LA LIBÉRATION, 33405
TALENCE, FRANCE
E-mail address: `kkellay@math.u-bordeaux1.fr`

DÉPARTEMENT DE MATHÉMATIQUES ET DE STATISTIQUE, UNIVERSITÉ LAVAL,
1045 AVENUE DE LA MÉDECINE, QUÉBEC (QC), CANADA G1V 0A6
E-mail address: `ransford@mat.ulaval.ca`