



HAL
open science

Photodissociation of protonated Leucine-Enkephalin peptide in the VUV range

Milos Lj. Ranković, Victor Z. Cerovski, Francis Canon, Laurent Nahon, Alexandre Giuliani, Aleksandar R. Milosavljević

► **To cite this version:**

Milos Lj. Ranković, Victor Z. Cerovski, Francis Canon, Laurent Nahon, Alexandre Giuliani, et al.. Photodissociation of protonated Leucine-Enkephalin peptide in the VUV range. 29. International Conference on Photonic, Electronic, and Atomic Collisions (ICPEAC), Jul 2015, Toledo, Spain. 10.1088/1742-6596/635/11/112030 . hal-01264208

HAL Id: hal-01264208

<https://hal.science/hal-01264208>

Submitted on 28 Jan 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Photodissociation of protonated Leucine-Enkephalin peptide in the VUV range

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2015 J. Phys.: Conf. Ser. 635 112030

(<http://iopscience.iop.org/1742-6596/635/11/112030>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 138.102.192.62

This content was downloaded on 12/01/2016 at 10:15

Please note that [terms and conditions apply](#).

Photodissociation of protonated Leucine-Enkephalin peptide in the VUV range

M. Lj. Ranković^{1*}, V. Cerovski¹, F. Canon², L. Nahon³, A. Giuliani^{3,4} and A. R. Milosavljević¹

¹Institute of Physics Belgrade, University of Belgrade, Pregrevica 118, 11080 Belgrade, Serbia

²INRA, UMR1324 Centre des Sciences du Goût et de l'Alimentation, F-21000 Dijon, France

³SOLEIL, l'Orme des Merisiers, St Aubin, BP48, 91192 Gif sur Yvette Cedex, France

⁴INRA, UAR1008, CEPIA, Rue de la Géraudière, BP 71627, 44316 Nantes, France

Synopsis We present results of VUV action spectroscopy of gas-phase protonated Leucine-Enkephalin peptide. The experiment has been performed by coupling a linear quadrupole ion trap with a synchrotron radiation VUV beamline.

Recent development of experimental methods that use electrospray ion source (ESI) to bring macromolecules into gas phase and collect the ions in an ion trap, which is coupled to bright synchrotron radiation sources, has allowed the investigation of large biopolymers (proteins and nucleotides) under well-defined conditions. These studies can provide more detailed insight into fundamental properties of biopolymers and their interaction with high energy photons.

We present the results of VUV photon action spectroscopy of Leucine-Enkephalin (Leu-Enk) peptide in 5-15 eV energy domain. This peptide has been studied with wide range of spectrometry techniques and has become a standard biomolecular target [1].

The experiment has been performed at the beamline DESIRS [2] of the synchrotron radiation facility SOLEIL near Paris, France. More details about the experimental setup can be found in recent publications [3, 4]. Briefly, precursor ions $[\text{Leu-Enk}+\text{H}]^+$ were produced by ESI and injected into the ion trap from the front side. Ions were irradiated with monochromatized VUV photon beam, during 500 ms from the back side of the ion trap. After irradiation all ion fragment species were ejected from the ion trap and tandem mass spectra (MS^2) as a function of the photon energies were recorded. Leu-Enk molecules were provided from Sigma Aldrich and dissolved in water/acetonitrile at 10 μM .

Photodissociation ion yields were extracted from the MS^2 and normalized to the total ion current and the photon flux. The ion yields show spectroscopic structure allowing the studying of absorption-like bands of the peptide.

Figure 1 shows the relative ion yield of one of the backbone fragments. The present experimental results are in good agreement with our time dependent density functional theory (TD DFT) calculations.




Figure 1. Relative ion yield of fragment b_3 from the photodissociation of precursor $[\text{Leu-Enk}+\text{H}]^+$.

We acknowledge the support by the ANR, France, under project ANR-08-BLAN-0065. A.R.M. and M.L.J.R. acknowledge support by the MESTD RS (Projects No. 171020). The SOLEIL synchrotron radiation facility is acknowledged for providing beamtime under project 20130388.

References

- [1] S. Bari et al 2011, *J. Chem. Phys.* **18** **134**, 024314.
- [2] L. Nahon et al 2012, *J. Sync. Rad.* **19**, 508 – 520.
- [3] A. Milosavljević et al 2011, *Phys. Chem. Chem. Phys.* **13**, 15432–15436.
- [4] A. Milosavljević et al 2012, *J. Sync. Rad.* **19**, 174.

* E-mail: mrankovic@ipb.ac.rs

