



**HAL**  
open science

# On p-adic differential equations with separation of variables

Pierre Lairez, Tristan Vaccon

► **To cite this version:**

Pierre Lairez, Tristan Vaccon. On p-adic differential equations with separation of variables. 2016. hal-01265226v1

**HAL Id: hal-01265226**

**<https://hal.science/hal-01265226v1>**

Preprint submitted on 31 Jan 2016 (v1), last revised 10 May 2016 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# On $p$ -adic differential equations with separation of variables

Pierre Lairez  
Technische Universität  
Berlin, Germany  
pierre@lairez.fr

Tristan Vaccon  
JSPS-Rikkyo University  
Tokyo, Japan  
vaccon@rikkyo.ac.jp

## ABSTRACT

Several algorithms in computer algebra involve the computation of a power series solution of a given ordinary differential equation. Over finite fields, the problem is often lifted in an approximate  $p$ -adic setting to be well-posed. This raises precision concerns: how much precision do we need on the input to compute the output accurately? In the case of ordinary differential equation with separation of variables, we apply the recent technique of differential precision to obtain optimal bounds on the stability of the Newton iteration. It applies, for example, to algorithms for manipulating algebraic numbers over finite fields, for computing isogenies between elliptic curves or for deterministically finding roots of polynomials in finite fields. The new bounds lead to significant speedups in practice.

## Categories and Subject Descriptors

I.1.2 [Computing Methodologies]: Symbolic and Algebraic Manipulations—*Algebraic Algorithms*

## Keywords

Ordinary differential equation,  $p$ -adic numbers, Newton iteration, numerical stability, differential precision

## 1. INTRODUCTION

We study in a  $p$ -adic context the loss of precision occurring during the computation of a power series solution of a certain class of differential equations. We use the method of differential precision that relies on a first-order analysis.

### 1.1 The $p$ -adic context

Let  $\mathbb{Z}_p$  be the ring of  $p$ -adic integer, for a given prime  $p$ , and  $\mathbb{Q}_p$  its field of fractions. The  $p$ -adic valuation on  $\mathbb{Q}_p$  is denoted  $v_p$ , and the  $p$ -adic norm is defined by  $|a| = p^{-v_p(a)}$ , for  $a \in \mathbb{Q}_p$ . For  $a \in \mathbb{Q}_p$  and  $\varepsilon > 0$  a positive real number, let  $a + O(\varepsilon)$  denote the set of all  $b \in \mathbb{Q}_p$  such that  $|a - b| \leq \varepsilon$ .

For  $a, b \in \mathbb{Z}_p$ , we have  $a = b + O(\varepsilon)$  if and only if  $a \equiv b \pmod{p^{-\lceil \log_p \varepsilon \rceil}}$ .

A computer can handle  $p$ -adic numbers given with bounded precision: a  $p$ -adic number  $a$  is approximately represented by a rational number  $a'$  and a radius  $\varepsilon$  such that  $a \in a' + O(\varepsilon)$ . This leads to a *ball arithmetic* over  $\mathbb{Q}_p$ . The ultrametric nature of  $\mathbb{Q}_p$  makes ball arithmetic particularly convenient since errors do not propagate when adding two numbers:

$$(a + O(\varepsilon)) + (b + O(\eta)) = a + b + O(\max(\varepsilon, \eta)),$$

or multiplying them:

$$(a + O(\varepsilon))(b + O(\eta)) = ab + O(\max(\eta|a|, \varepsilon|b|)),$$


or dividing them:

$$\frac{a + O(\varepsilon)}{b + O(\eta)} = \frac{a}{b} + O(\max(\eta|a||b|^{-2}, \varepsilon|b|^{-1})).$$

These formulas are optimal: the equalities are set equalities and not only left-to-right inclusions. When considering an algorithm performing additions, multiplications and divisions over  $p$ -adic numbers, it is possible to track the precision during all the intermediate steps. Thus, it is possible to run the algorithm on inputs given approximately as balls, and to return the result as a ball with the guarantee that whatever are the *exact* values of the input, the *exact* result lies in the ball returned. However, even if for every single operation the formulas above give the optimal precision of the result, the optimality does not compose. This is the well-known *dependency problem*. It is a major obstacle to the application of ball arithmetic over  $\mathbb{Q}_p$ , in the same way as it constricts interval arithmetic over  $\mathbb{R}$ .

For example, let us consider the computation of the determinant of a matrix with  $p$ -adic integer coefficients, given at precision  $\varepsilon$ . Since the determinant is an integral polynomial function of the coefficients, the determinant is also known at precision at least  $\varepsilon$ . However, if it is computed through a Gaussian elimination, and if at some point of the computation, one of the pivot has a positive valuation, then basic precision tracking will indicate that the result is only correct at precision less than  $\varepsilon$ . The *intrinsic* loss of precision is null, or even negative (Caruso, Roe, and Vaccon 2015), while the *algorithmic* loss, that depends on the algorithm, may be positive.

Caruso, Roe, and Vaccon (2014) have shown, in a  $p$ -adic setting, how to use first order analysis to obtain rigorous and optimal precision bounds on both the intrinsic and the algorithmic loss of precision. The method relies on the following fact:



LEMMA 1 (Caruso, Roe, and Vaccon 2014). *Let  $\varphi : \mathbb{Q}_p^n \rightarrow \mathbb{Q}_p^m$  be a differentiable map and let  $x \in \mathbb{Q}_p^n$ . If the differential  $d_x\varphi$  at  $x$  is surjective, then  $\varphi(x+B) = \varphi(x) + d_x\varphi(B)$  for any zero-centered small enough ball  $B$ .*

In other words, if the precision on the input  $x$  is  $B$ , then the precision on the output  $\varphi(x)$  is  $d_x\varphi(B)$ ; this is the inclusion of  $\varphi(x+B)$  in  $\varphi(x) + d_x\varphi(B)$ . And conversely, every perturbation of  $\varphi(x)$  up to  $d_x\varphi(B)$  comes from a perturbation from the input  $x$  up to  $B$ ; this is the converse inclusion of  $\varphi(x) + d_x\varphi(B)$  in  $\varphi(x+B)$ . What *small enough* is can be expressed in terms of the norms of the higher differentials of  $\varphi$  at  $x$ , in the case that  $\varphi$  is analytic at  $x$ .

## 1.2 Main result

We study here the computation of a power series with  $p$ -adic coefficients solution of a given first-order ordinary differential equation with separation of variables. Let  $g$  and  $h$  be power series in  $\mathbb{Z}_p[[t]]$  such that  $h(0) = 1$  and  $g(0) \neq 0$ . We consider the following differential equation:

$$y' = g \cdot h(y), \quad y(0) = 0. \quad (\text{E})$$

It has a unique solution  $y \in \mathbb{Q}_p[[t]]$ . We make the assumption that this solution has integer coefficients, that is  $y \in \mathbb{Z}_p[[t]]$ . Given the  $n$  first coefficients of  $f$  and  $g$ , at bounded precision, how far can we approximate  $y$ ? at which computational cost?

The case of a general initial condition  $y(0) = c$  reduces to the equation above by changing  $h(t)$  in  $h(t+c)$ . In particular, the linear equation  $y' = g \cdot y$ , with  $y(0) = 1$  can be written  $z' = g \cdot (1+z)$ , with  $z(0) = 0$ , thanks to the transformation  $z = y - 1$ . Nevertheless, the initial condition  $y(0) = 0$  ensures that the composition  $h(y)$  is well defined when  $h$  is a general power series.

DEFINITION. For  $\varepsilon$  a positive real number and  $n$  a positive integers, an *approximation modulo  $(p^\kappa, t^n)$*  of a power series  $f = \sum_{k \geq 0} a_k t^k \in \mathbb{Z}_p[[t]]$  is a power series  $\tilde{f} = \sum_{k \geq 0} b_k t^k \in \mathbb{Z}_p[[t]]$  such that  $a_k = b_k \pmod{p^\kappa}$  for any for all  $k < n$ .

The complexity of computing  $y$  depends on the complexity of the power series multiplication and the composition  $h(f)$  for a general power series  $f$ . Let  $M_{\mathbb{Z}}(p^\lambda, n)$  the number of bit operations required to compute the product of two polynomials of degree  $n$  with coefficients in  $\mathbb{Z}/p^\lambda\mathbb{Z}$ . Let  $C_h(p^\lambda, n)$ , the number of bit operations needed to compute an approximation modulo  $(p^\lambda, t^n)$  of  $h(f)$  given an approximation modulo  $(p^\lambda, t^n)$  of a power series  $f \in \mathbb{Z}_p[[t]]$ . We assume that  $C_h(p^\lambda, 2n) \geq 2C_h(p^\lambda, n)$ . In most applications, the composition with  $h$  is easy to compute:  $h$  is a rational function of small degree or a radical of such a rational function so that  $C_h(p^\lambda, n) = O(M_{\mathbb{Z}}(p^\lambda, n))$ . In the general case, Kedlaya and Umans (2011) proved the quasi-optimal bound  $C_h(p^\lambda, n) = O((n\lambda \log p)^{1+o(1)})$ .

Our main result is then the following:

THEOREM 2. *Let  $n > 0$ ,  $\kappa > 0$  (or  $\kappa > 1$  if  $p = 2$ ) and let  $\lambda = \kappa + \lfloor \log_p n \rfloor$ . One can compute an approximation modulo  $(p^\kappa, t^{n+1})$  of the solution  $y$  of (E) given approximations modulo  $(p^\lambda, t^n)$  of  $g$  and  $h$ , using  $O(M_{\mathbb{Z}}(p^\lambda, n) + C_h(p^\lambda, n))$  bit operations.*

The result was known in the linear case (Bostan et al. 2005; Grenet, Hoeven, and Lecerf 2015). In the non-linear case, Lecerf and Sirvent (2008) showed that an approximation modulo  $(p^\kappa, t^n)$  can be computed from approximations modulo  $(p^{\kappa+O(\log(n)^2)}, t^n)$  of  $g$  and  $h$ .

## 1.3 Applications

### 1.3.1 Newton sums

The problem studied by Bostan et al. (2005) is the recovery of a polynomial given its Newton sums. Let  $f \in \mathbb{Z}_p[t]$  be a monic polynomial of degree  $d$ , and let  $\nu_n$  be the  $n$ th Newton sum of  $f$ . If  $\alpha_1, \dots, \alpha_d$  are the roots of  $f$  in  $\overline{\mathbb{Q}_p}$ , then  $\nu_n$  is sum  $\alpha_1^n + \dots + \alpha_d^n$ , it is an element of  $\mathbb{Z}_p$ . How can we recover  $f$  given Newton sums  $\nu_0, \dots, \nu_d$ ? Let  $g$  be the polynomial  $x^d f(1/x)$ , and  $N_f$  be the generating function  $N_f(t) = \sum_{n \geq 0} \nu_{n+1} t^n$ . Then  $g' = -N_f g$ , so that  $g$  is a solution of a first-order linear differential equation (Schönhage 1993). Therefore, knowing an approximation modulo  $(p^\kappa, t^d)$  of  $N_f$  makes possible to recover each coefficients of  $g$  (and hence  $f$ ) modulo  $p^{\kappa - \lfloor \log_p d \rfloor}$ .

An interesting application is the computation over  $\mathbb{F}_p$  of composed products; composed sums can be treated similarly (Bostan et al. 2005). Let  $f$  and  $g$  be monic polynomials of  $\mathbb{F}_p[t]$  of degree  $d$  and  $e$  respectively, with associated roots  $(\alpha_i)_{1 \leq i \leq d}$  and  $(\beta_j)_{1 \leq j \leq e}$  in  $\overline{\mathbb{F}_p}$ . We define the composed product of  $f$  and  $g$  to be

$$f \oplus g = \prod_{i,j} (t - \alpha_i \beta_j) = \text{res}_y (y^p f(t/y), g(y)),$$

this is a polynomial in  $\mathbb{F}_p[t]$  of degree  $p+q$ . Then  $N_{f \oplus g}$  is the coefficientwise product of the two power series  $N_f$  and  $N_g$  (also known as the Hadamard product). This gives a strategy to compute efficiently  $f \oplus g$ . Firstly, arbitrarily lift  $f$  and  $g$  as polynomials in  $\mathbb{Z}_p[t]$ , denoted  $\tilde{f}$  and  $\tilde{g}$ . The composed product  $\tilde{f} \otimes \tilde{g}$  is a polynomial in  $\mathbb{Z}_p[t]$  and equals  $f \oplus g$  modulo  $p$ . Secondly, compute approximations modulo  $(p^\kappa, t^{d+e+1})$  of  $N_{\tilde{f}}$  and  $N_{\tilde{g}}$ , with  $\kappa = 1 + \lfloor \log_p(d+e) \rfloor$ , and, with a coefficientwise product, an approximation modulo  $(p^\kappa, t^{d+e})$  of  $N_{\tilde{f} \otimes \tilde{g}}$ . Thirdly, compute an approximation modulo  $(p, t^{d+e+1})$  of  $\tilde{f} \otimes \tilde{g}$  using Theorem 2, and deduce the value of  $f \oplus g$ .

Another application of this procedure to root finding in finite fields is developed by Grenet, Hoeven, and Lecerf (2015). Our work does not improve the complexity given by Bostan et al. (2005), in the case of a linear equation, but it gives a simpler proof that generalizes to nonlinear differential equations.

### 1.3.2 Isogeny computation

To compute normalized isogenies between elliptic curves, Bostan et al. (2008) and Lecerf and Sirvent (2008) studied the differential equation

$$y'^2 = g \cdot h(y), \quad (1)$$

where  $g$  and  $h$  are series in  $\mathbb{Z}_p[[t]]$ , which is known, in their context, to admit a solution in  $\mathbb{Z}_p[[t]]$ . As the previous example, it comes from a lift of a problem over  $\mathbb{F}_p$ . This equation rewrites equivalently as  $y' = \sqrt{g} \sqrt{h(y)}$ , and when  $p \neq 2$ , the series  $\sqrt{g}$  and  $\sqrt{h}$  are still in  $\mathbb{Z}_p[[t]]$ , so we can apply the Theorem 2. The study of this equation when  $p = 2$  is still an open problem.

In order to compute an approximation of  $y$  modulo  $(p, t^{n+1})$ , we obtain that it is enough to have approximations of  $g$  and  $h$  modulo  $(p^{1+\lfloor \log_p n \rfloor}, t^n)$ . This improves upon the result of Lecerf and Sirvent (2008) which requires approximations modulo  $(p^{O(\log(n)^2)}, t^n)$ .

## Acknowledgements

We are grateful to Alin Bostan, Xavier Caruso, Luca De Feo, Éric Schost and Kazuhiro Yokoyama for fruitful discussions. This work is partially funded by the grant BU 1371/2-2 of the Deutsche Forschungsgemeinschaft.

## 2. THE ALGORITHM

We may consider the more general setting of an unramified finite extension  $K$  of  $\mathbb{Q}_p$ , this is useful, for example, for the computation of isogenies. Let  $\mathcal{O}_K$  denote the ring of integers of  $K$ . For example, we may naturally consider  $K = \mathbb{Q}_p$  and  $\mathcal{O}_K = \mathbb{Z}_p$ .

Let  $h \in \mathcal{O}_K[[t]]$ , a power series with integer coefficients, with  $h(0) = 1$ . For  $g \in K[[t]]$ , let  $Y(g)$  be the unique  $y \in K[[t]]$  such that  $y(0) = 0$  and  $y' = g \cdot h(y)$ . Existence and uniqueness are clear because the differential equation rewrites equivalently into a well posed recurrence relation on the coefficients of  $y$ .

For  $g \in K[[t]]$ , let  $N_g$  denote the Newton operator:

$$N_g(u) = u - h(u) \int \left( \frac{u'}{h(u)} - g \right),$$

where  $\int f$ , for  $f \in K[[t]]$ , denotes the unique power series  $F \in K[[t]]$  such that  $F' = f$  and  $F(0) = 0$ .

**PROPOSITION 3.** *Let  $g, u \in K[[t]]$  be formal power series, and let  $n > 0$ . If  $u = Y(g) \pmod{t^n}$  then*

$$N_g(u) = Y(g) \pmod{t^{2n}}.$$

*Proof.* Let  $e = u'/h(u) - g$ . Since  $u = Y(g) \pmod{t^n}$ , we have  $e = 0 \pmod{t^{n-1}}$  and  $\int e = 0 \pmod{t^n}$ . With  $v = N_g(u)$ , we compute

$$v' - g h(v) = u' - u' h'(u) \int e - h(u)e - g h(v).$$

Then, the first-order expansion of  $h(v)$  at  $h(u)$  gives

$$h(v) = h(u) - h'(u)h(u) \int e \pmod{t^{2n}},$$

and using the equality  $u' = h(u)e + g h(u)$ , we obtain

$$v' - g h(v) = -e h(u) h'(u) \int e = 0 \pmod{t^{2n-1}}.$$

This implies that  $v = Y(g) \pmod{t^{2n}}$ .  $\square$

The iteration of the Newton operator leads to Algorithm 1. In an exact setting, the correctness of this procedure would be clear, thanks to Proposition 3. In a  $p$ -adic setting, where the coefficients of the power series  $g$  and  $h$  are known with finite precision only, what can be obtained with Newton iteration is not clear because the operation  $\int$  involves divisions.

Let us begin with a quick analysis of Algorithm 1. On input  $(g, h, n)$ , it performs a recursive call and computes  $u = \text{DSOL}(g, h, m)$ , with  $m = \lceil \frac{n-1}{2} \rceil$ . Let us assume that  $u$  is a approximation modulo  $(p^\kappa, t^{m+1})$  of  $Y(g)$ , for some  $\kappa > 0$ . Then  $N_g(u)$  is an approximation modulo  $(p^{\kappa - \lfloor \log_p n \rfloor}, t^{n+1})$ : Indeed, the computation of  $N_g(u)$  involves divisions by the integers from 2 to  $n$  on distinct coefficients so the loss of precision is at most the maximum valuation of these integers, which is  $\lfloor \log_p n \rfloor$ . Thus, if we define  $\mu(0) = 0$  and  $\mu(n) = \lfloor \log_p n \rfloor + \mu(\lceil \frac{n-1}{2} \rceil)$ , we obtain that  $\text{DSOL}(g, h, n)$  is an approximation modulo  $(p^{\lambda - \mu(n)}, t^{n+1})$  of  $Y(g)$ . We can check that  $\mu(n) = O(\log(n)^2)$ . Theorem 5 improves on that analysis and shows that the precision of the result is at least  $\lambda - \lfloor \log_p n \rfloor$  and matches the intrinsic loss of precision.

---

*Algorithm 1.* The Newton iteration to solve a first-order differential equation.

**Input.**  $g$  and  $h \in \mathcal{O}_K[[t]]$  given modulo  $(p^\lambda, t^n)$  — that is, given as polynomials of degree less than  $n$  with coefficients in  $\mathcal{O}_K/p^\lambda \mathcal{O}_K$ .

**Output.** A power series  $u \in \mathcal{O}_K[[t]]$  given modulo  $(p^\lambda, t^{n+1})$ .

**Specification.** If  $Y(g) \pmod{t^{n+1}}$  has integer coefficients and if  $\lambda \geq \kappa + \lfloor \log_p n \rfloor$ , then  $u$  is an approximation of  $Y(g)$  modulo  $(p^\kappa, t^{n+1})$ .

---

```

function DSOL( $g, h, n$ )
  if  $n = 0$  then
    return 0 (mod  $t$ )
  else
     $u \leftarrow \text{DSOL}(g, h, \lceil \frac{n-1}{2} \rceil)$ 
    return  $N_g(u)$  (mod  $t^{n+1}$ )
     $\triangleright$  Compute at fixed precision  $\lambda$ .

```

---

We assume the *fixed precision model* for computing with  $p$ -adic numbers: At precision  $\lambda$ , it amounts to work over the ring  $\mathcal{O}_K/p^\lambda \mathcal{O}_K$ . When a division  $a/b$  arises, with the approximation of  $a$  and  $b \in \mathcal{O}_K$  given in  $\mathcal{O}_K/p^\lambda \mathcal{O}_K$ , three cases may arise:

- $v_p(b) = 0$ , in which case  $b$  is invertible in  $\mathcal{O}_K/p^\lambda \mathcal{O}_K$  and the division is well defined.
- $v_p(a) \geq v_p(b) > 0$ , in which case  $a/b$  is in  $\mathcal{O}_K$  but its approximation in  $\mathcal{O}_K/p^\lambda \mathcal{O}_K$  is not fully determined by the approximations of  $a$  and  $b$ , so  $a/b$  is arbitrarily defined in this model to be the class in  $\mathcal{O}_K/p^\lambda \mathcal{O}_K$  of the smallest integer  $c \geq 0$  such that  $a = bc \pmod{p^\lambda}$ .
- $v_p(a) < v_p(b)$ , in which case  $a/b$  is not an integer and an error is raised.

The main argument for the correctness of Algorithm 1 is the following proposition, proved in Section 3.

**PROPOSITION 4.** *Let  $n > 0$  and  $\kappa > 0$  (or  $\kappa > 1$  if  $p = 2$ ) be integers, and let  $g \in \mathcal{O}_K[[t]]$  such that  $Y(g) \pmod{t^{n+1}}$  has integer coefficients. For any  $y \in K[[t]]$  the following are equivalent:*

1.  $y = Y(\bar{g}) \pmod{t^{n+1}}$  for some power series  $\bar{g} \in \mathcal{O}_K[[t]]$  such that  $\int(\bar{g} - g) = 0 \pmod{p^\kappa}$ ;
2.  $y = Y(g) \pmod{p^\kappa, t^{n+1}}$ .

**THEOREM 5.** *Algorithm 1 is correct: if  $\kappa > 0$  (or  $\kappa > 1$  if  $p = 2$ ) and  $\lambda \geq \lfloor \log_p n \rfloor + \kappa$ , then for all  $g \in \mathcal{O}_K[[t]]$  such that  $Y(g)$  has integer coefficients, the output of the procedure  $\text{DSOL}(g, h, n)$  equals  $Y(g) \pmod{p^\kappa, t^{n+1}}$ .*

*Moreover, it performs  $\mathcal{O}(M_{\mathcal{O}_K}(p^\lambda, n) + C_h(p^\lambda, n))$  bit operations, where  $M_{\mathcal{O}_K}(p^\lambda, n)$  is the cost of computing the product of two polynomials of degree  $n$  with coefficients in  $\mathcal{O}_K/p^\lambda \mathcal{O}_K$ .*

*Proof.* We proceed by induction on  $n$ . The case  $n = 0$  is trivial, so let us assume that  $n > 0$ . Let  $g \in \mathcal{O}_K[[t]]$  such that  $Y(g)$  has integer coefficients, let  $m = \lceil \frac{n-1}{2} \rceil$  and let  $u \in K[[t]]$  be the output of  $\text{DSOL}(g, h, m)$ . By induction hypothesis,

$u = Y(g) \pmod{p^\kappa, t^{m+1}}$ . (In particular  $u$  has integer coefficients.) By Proposition 4, this implies that  $y = Y(\bar{g}) \pmod{t^{m+1}}$  for some  $\bar{g} \in \mathcal{O}_K[[t]]$  such that  $\int(g - \bar{g}) \pmod{p^\kappa}$ . Proposition 3 gives that  $Y(\bar{g}) = N_{\bar{g}}(u) \pmod{t^{n+1}}$  and Proposition 4 gives further that  $Y(\bar{g}) = Y(g) \pmod{p^\kappa, t^{n+1}}$ . We check that

$$N_g(u) = N_{\bar{g}}(u) - h(u) \int(g - \bar{g})$$

and since  $h(u)$  has integer coefficients, this implies that  $N_g(u) = Y(g) \pmod{p^\kappa, t^{n+1}}$ .

We now relate  $N_g(u)$  to the output of the procedure DSOL( $g, h, n$ ). Let  $e = u'/h(u) - g$ . By definition, the output is  $N_g(u) = u - h(u) \int e$ , computed over  $\mathcal{O}_K/p^\lambda \mathcal{O}_K$ , in the so-called fixed precision model. Let  $E$  be the primitive  $\int e \pmod{t^{n+1}}$  computed in this model, so that the output is exactly  $u - h(u)E$ . Clearly  $E = \int e + \int \eta \pmod{t^{n+1}}$  for some  $\eta = 0 \pmod{p^\lambda}$  that reflects the indeterminacies in the divisions. Since  $\lambda \geq \lfloor \log_p n \rfloor + \kappa$ ,  $\int \eta = 0 \pmod{p^\kappa}$  and thus, the output  $N_g(u) + \int \eta$  equals  $Y(g) \pmod{p^\kappa, t^{n+1}}$ . This concludes the proof of correctness.

Concerning the complexity, the last iteration involves a composition by  $h$  with cost  $C_h(p^\lambda, n)$ , a few multiplications with cost  $M_{\mathcal{O}_K}(p^\lambda, n)$  and an inversion  $1/h(u)$  with cost  $\mathcal{O}(M_{\mathcal{O}_K}(p^\lambda, n))$  too with a Newton iteration (Kung 1974). With the assumption that the cost of an iteration is greater than twice the cost of the previous, it is well known that the cost of a Newton algorithm is dominated by the cost of the last iteration, which gives the result.  $\square$

The condition  $\lambda \geq \lfloor \log_p n \rfloor + \kappa$  cannot be improved further: it matches the *intrinsic* loss of precision. This is shown, for example, by the differential equation  $y' = at^{b-1}$ , with  $a \in K$ , whose solution is  $\frac{a}{b}t^b$ . If we take  $b = \lfloor \log_p n \rfloor$  and if  $a$  is known at precision  $\lambda$  then  $y$  is known at precision no more than  $\lambda - \lfloor \log_p n \rfloor$ .

### 3. DIFFERENTIAL PRECISION

We apply the method of Caruso, Roe, and Vaccon (2014) to study the loss in precision in the resolution of the differential equation (E) and give a proof of Proposition 4.

Let  $n > 0$  and let  $E$  and  $F$  denote respectively the two  $n$ -dimensional  $K$ -vector spaces  $K[[t]]/(t^n)$  and  $tK[[t]]/(t^{n+1})$ . Let  $\mathcal{Y}$  be the polynomial map

$$\begin{aligned} \mathcal{Y} : E &\longrightarrow F \\ [u] &\longmapsto [Y(u)], \end{aligned}$$

which is well defined because the  $n+1$  first coefficients of  $Y(u)$  depend only on the  $n$  first coefficients of  $u$ . Let  $g \in \mathcal{O}_K[[t]]$  be such that  $\mathcal{Y}(g)$  has integer coefficients in the monomial basis. Let  $d\mathcal{Y}$  denote the first differential of  $\mathcal{Y}$ : for any  $g \in E$ ,  $d_g\mathcal{Y}$  is a linear map  $E \rightarrow F$ . Let  $d^k\mathcal{Y}$  denote the higher differentials: for any  $g \in E$ ,  $d_g^k\mathcal{Y}$  is a multilinear map  $E^k \rightarrow F$ .

LEMMA 6. *For any  $w \in E$ ,  $d\mathcal{Y}(w) = h(\mathcal{Y}) \int w$ . Moreover, for any  $k \geq 1$ , there exists a polynomial  $P_k \in \mathbb{Z}[u_0, \dots, u_{k-1}]$  such that for any  $w_1, \dots, w_k \in E$ ,*

$$\begin{aligned} d^k\mathcal{Y}(w_1, \dots, w_k) = \\ P_k \left( h(\mathcal{Y}), h'(\mathcal{Y}), \dots, h^{(k-1)}(\mathcal{Y}) \right) \prod_{i=1}^k \int w_i. \quad (2) \end{aligned}$$

*Proof.* Differentiating with respect to  $g$  the defining relation  $\mathcal{Y}(g)' = g \cdot h(\mathcal{Y}(g)) \pmod{t^n}$  leads to

$$[d_g\mathcal{Y}(w)]' = w \cdot h(\mathcal{Y}(g)) + g \cdot h'(\mathcal{Y}(g)) \cdot d_g\mathcal{Y}(w),$$

which is a first-order inhomogeneous linear differential equation in  $d_g\mathcal{Y}(w)$ . The initial condition  $d_g\mathcal{Y}(w)(0) = 0$  determines a unique solution, namely  $h(\mathcal{Y}(g)) \int w$ .

The second claim follows by induction. Equation (2) holds for  $k = 1$  with  $P_1 = u_0$ ; differentiating it leads to the recurrence relation

$$P_{k+1}(u_0, \dots, u_k) = u_0 \sum_{i=0}^{k-1} \frac{\partial P}{\partial u_i} u_{i+1}. \quad \square$$

The space  $F$  is endowed with the maximum norm in the monomial basis, denoted  $\|\cdot\|_F$ . In particular, an element  $u$  of  $F$  has integer coefficients if and only if  $\|u\|_F \leq 1$ . The space  $E$  is endowed with the norm

$$\|u\|_E \stackrel{\text{def}}{=} \|\int u\|_F.$$

Let  $\|d_g^k\mathcal{Y}\|$  denote the operator norm of  $d_g^k\mathcal{Y}$ , that is

$$\|d_g^k\mathcal{Y}\| = \sup \left\{ \|d_g^k\mathcal{Y}(w_1, \dots, w_k)\|_F \mid \|\int w_i\|_F \leq 1 \right\}. \quad (3)$$

LEMMA 7.  $\|d_g^k\mathcal{Y}\| \leq 1$ , for any  $k \geq 1$ .

*Proof.* Since  $h$  and  $\mathcal{Y}(g)$  have integer coefficients, this follows easily from Lemma 6 and Equation (3).  $\square$

PROPOSITION 8. *For any  $\varepsilon \leq \frac{1}{p}$  (or  $\varepsilon \leq \frac{1}{4}$  for  $p = 2$ ),*

$$\mathcal{Y}(g + B_\varepsilon) = \mathcal{Y}(g) + d_g\mathcal{Y}(B_\varepsilon),$$

where  $B_\varepsilon = \{w \in E : \|w\|_E \leq \varepsilon\}$ .

*Proof.* We apply the result of Caruso, Roe, and Vaccon (2014, Corollary 3.16). Using their notations, we can use  $C = 1$  because the closed ball of radius 1 in  $F$  (that is the set of elements with integer coefficients) is included in  $d_g\mathcal{Y}(B_1)$ : Indeed for any  $u \in F$  we have  $d_g\mathcal{Y}((u/\mathcal{Y}(g))') = u$ , and if  $\|u\|_F \leq 1$  then  $\|(u/\mathcal{Y}(g))'\|_E = \|(u/\mathcal{Y}(g))\|_F \leq 1$ , because  $u$  and  $\mathcal{Y}(g)$  have integer coefficients.

For  $k \geq 2$ , let  $M_k$  denote  $\|\frac{1}{k!} d_g^k\mathcal{Y}\|$ . By Lemma 7, this is simply  $|\frac{1}{k!}|$ . Corollary 3.16 (ibid.), with  $\rho = 1$  in their notations, implies that  $\mathcal{Y}(g + B_\varepsilon) = \mathcal{Y}(g) + d_g\mathcal{Y}(B_\varepsilon)$  as long as  $\varepsilon$  satisfies

$$\varepsilon < \exp \left( \inf_{k \geq 2} \frac{-\log M_k}{k-1} \right) = \inf_{k \geq 2} p^{-\frac{v_p(k!)}{k-1}}.$$

Let  $A$  denote the right-hand side. Legendre's formula for the  $p$ -adic valuation of  $k!$  shows that  $v_p(k!) \leq \frac{k}{p-1}$ . Therefore  $A \geq p^{-2/(p-1)}$ . For  $p \geq 5$ , this bound gives  $A > \frac{1}{p}$ , which proves the claim. For  $p = 3$ , we have


$$A \geq \min \left( 3^{-v_3(2!)}, \inf_{k \geq 3} 3^{-\frac{v_3(k!)}{k-1}} \right) \geq \min \left( 1, 3^{-\frac{3}{4}} \right) > 3^{-1},$$

and for  $p = 2$ , we have

$$A \geq \min \left( 2^{-v_2(2!)}, \inf_{k \geq 3} 2^{-\frac{v_2(k!)}{k-1}} \right) \geq \min \left( 2^{-1}, 2^{-\frac{3}{2}} \right) > 2^{-2},$$

which concludes the proof.  $\square$

Figure 1. Timings in seconds, measured on a laptop, of Algorithm 1 run at precision  $\lambda_{\text{old}}$  (upper curve) and  $\lambda_{\text{new}}$  (lower curve) in order to compute an approximation modulo  $(5, t^{4m+1})$  of the solution of Equation (4).



*Proof of Proposition 4.* Let  $\varepsilon = p^{-\kappa}$ . The norm of an element  $v \in F$  is given by  $p^{-\lambda}$  where  $\lambda$  is the largest integer such that  $v = 0 \pmod{p^\lambda}$ . Since  $h(\mathcal{Y}(g))$  is invertible modulo  $(p^\lambda, t^{n+1})$ , for any  $\lambda > 0$ , this shows that  $\|v\|_F = \|h(\mathcal{Y}(g))v\|_F$  for any  $v \in F$ . Therefore, with Lemma 6 and the definition of the norms,

$$d_g \mathcal{Y}(B_\varepsilon) = \{v \in F \mid v = 0 \pmod{p^\kappa}\}.$$

Moreover,  $B_\varepsilon = \{u \in E \mid \int u = 0 \pmod{p^\kappa}\}$ , and Proposition 4 now appears as a rewording of Proposition 8.  $\square$

## 4. EXPERIMENTS


Let us consider the differential equation

$$y' = \sqrt{\frac{1 + \frac{1}{4}m^2y^2 + m^6y^6}{1 + \frac{1}{4}t^2 + t^6}}, \quad y(0) = 0, \quad (4)$$

inspired from algorithms for computing isogenies (Bostan et al. 2008; Lercier and Sirvent 2008). Using an implementation in Magma (Bosma, Cannon, and Playoust 1997) of Algorithm 1, we computed the power series expansion of  $y \pmod{(5, t^{4m+1})}$  for several  $m$ . We compared (Figure 1) the CPU time spent on the computation when using on the one hand the precision  $\lambda_{\text{new}} = 1 + \lfloor \log_5(4m) \rfloor$ , following Theorem 5, and using on the other hand the precision  $\lambda_{\text{old}} = 1 + \mu(4m) = \mathcal{O}(\log(m)^2)$  found by a straightforward precision analysis — see the discussion in §2 for the definition of  $\mu$ . For example, with  $m = 104281$ , we compute  $\lambda_{\text{old}} = 72$  and  $\lambda_{\text{new}} = 9$ . The number of arithmetic operations performed does not depend on the precision  $\lambda$ , only on  $m$ , but the number of bit operations does since the base ring for the computation is  $\mathbb{Z}/5^\lambda\mathbb{Z}$ . Thus, the expected speedup is  $\lambda_{\text{old}}/\lambda_{\text{new}}$ , which is close to what we observed (Figure 2). The implementation is available at

<https://gist.github.com/lairez/d648b0d7b5392d0fef74>

Figure 2. Practical speedup obtained with the new precision analysis compared with the theoretical improvement ( $m$ -axis in logarithmic scale). (■) ratio  $\lambda_{\text{old}}/\lambda_{\text{new}}$ ; (●) actual speedup.



## References

- W. Bosma, J. Cannon, and C. Playoust (1997). “The Magma algebra system. I. The user language”. In: *J. Symbolic Comput.* 24.3-4. Computational algebra and number theory (London, 1993), pp. 235–265. DOI: [10.1006/jsc.1996.0125](https://doi.org/10.1006/jsc.1996.0125).
- A. Bostan, L. González-Vega, H. Perdry, and É. Schost (2005). “From Newton sums to coefficients: complexity issues in characteristic  $p$ ”. In: *Proceedings of the 8th international symposium on effective methods in algebraic geometry*. MEGA 2005 (Porto Conte, Italy).
- A. Bostan, F. Morain, B. Salvy, and É. Schost (2008). “Fast algorithms for computing isogenies between elliptic curves”. In: *Math. Comput.* 77.263, pp. 1755–1778. DOI: [10.1090/S0025-5718-08-02066-8](https://doi.org/10.1090/S0025-5718-08-02066-8).
- X. Caruso, D. Roe, and T. Vaccon (2014). “Tracking  $p$ -adic precision”. In: *LMS J. Comput. Math.* 17.suppl. A, pp. 274–294. DOI: [10.1112/S1461157014000357](https://doi.org/10.1112/S1461157014000357).
- (2015). “ $p$ -Adic Stability In Linear Algebra”. In: *Proceedings the 40th international symposium on symbolic and algebraic computation*. ISSAC 2015, Bath, United Kingdom, pp. 101–108. DOI: [10.1145/2755996.2756655](https://doi.org/10.1145/2755996.2756655).
- B. Grenet, J. van der Hoeven, and G. Lecerf (2015). “Deterministic root finding over finite fields using Graeffe transforms”. In: *Applicable Algebra in Engineering, Communication and Computing*. To appear. DOI: [10.1007/s00200-015-0280-5](https://doi.org/10.1007/s00200-015-0280-5).
- K. S. Kedlaya and C. Umans (2011). “Fast polynomial factorization and modular composition”. In: *SIAM J. Comput.* 40.6, pp. 1767–1802. DOI: [10.1137/08073408X](https://doi.org/10.1137/08073408X).
- H. T. Kung (1974). “On computing reciprocals of power series”. In: *Numer. Math.* 22, pp. 341–348.
- R. Lercier and T. Sirvent (2008). “On Elkies subgroups of  $\ell$ -torsion points in elliptic curves defined over a finite field”. In: *J. Théorie des Nombres des Bordeaux* 20, pp. 783–797.
- A. Schönhage (1993). “Fast parallel computation of characteristic polynomials by Leverrier’s power sum method adapted to fields of finite characteristic”. In: *Automata, languages and programming (Lund, 1993)*. Vol. 700. Lecture Notes in Comput. Sci. Berlin: Springer, pp. 410–417. DOI: [10.1007/3-540-56939-1\\_90](https://doi.org/10.1007/3-540-56939-1_90).