Less carbon emissions of wheat–maize intercropping under reduced tillage in arid areas

Falong Hu, Qiang Chai, Aizhong Yu, Wen Yin, Hongyan Cui, Yantai Gan

► To cite this version:

Falong Hu, Qiang Chai, Aizhong Yu, Wen Yin, Hongyan Cui, et al.. Less carbon emissions of wheat–maize intercropping under reduced tillage in arid areas. *Agronomy for Sustainable Development*, 2015, 35 (2), pp.701-711. 10.1007/s13593-014-0257-y . hal-01284283

HAL Id: hal-01284283

<https://hal.science/hal-01284283>

Submitted on 7 Mar 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Less carbon emissions of wheat–maize intercropping under reduced tillage in arid areas

Falong Hu · Qiang Chai · Aizhong Yu · Wen Yin ·
Hongyan Cui · Yantai Gan

Accepted: 26 September 2014 / Published online: 29 October 2014
© INRA and Springer-Verlag France 2014

Abstract Intercropping is used to increase grain production in many areas of the world. However, this increasing crop yield costs large amounts of water used by intercropped plants. In addition, intercropping usually requires higher inputs that induce greenhouse gas emissions. Actually, it is unknown whether intercropping can be effective in water-limited arid areas. Here, we measured crop yield, water consumption, soil respiration, and carbon emissions of wheat–maize intercropping under different tillage and crop residue management options. A field experiment was conducted at Wuwei in northwest China in 2011 and 2012. Our results show that wheat–maize intercropping increased grain yield by 61 % in 2011 and 63 % in 2012 compared with the average yield of monoculture crops. The intercropping under reduced tillage with stubble mulching yielded 15.9 t ha^{-1} in 2011 and 15.5 t ha^{-1} in 2012, an increase of 7.8 % in 2011 and 8.1 % in 2012, compared to conventional tillage. Wheat–maize intercropping had carbon emission of $2,400 \text{ kg C ha}^{-1}$ during the growing season, about 7 % less than monoculture maize, of $2,580 \text{ kg C ha}^{-1}$. Reduced tillage decreased C emission over conventional tillage by 6.7 % for the intercropping, 5.9 % for monoculture maize, and 7.1 % for monoculture wheat. Compared to monoculture maize, wheat–maize intercropping used more water but emitted 3.4 kg C per hectare per millimeter of water used, which was 23 % lower than monoculture

maize. Overall, our findings show that maize–wheat intercropping with reduced tillage coupled with stubble mulching can be used to increase grain production while effectively lower carbon emissions in arid areas.

Keywords Intercropping · Reduced tillage · Stubble retention · Soil respiration · Carbon emission

1 Introduction

Soils are the vital pools of carbon sequestration in the terrestrial ecosystems, and it is estimated that a global total of 2,500 Gt of C is stored in soils, 3.3 times the amount of C stored in the atmosphere and 4.5 times the amount of C stored in biotic pool (Lal et al. 2004). In a global scale, a total of 75 Pg of C has been emitted annually (Schlesinger and Andrews 2000), with a large portion of the emission from agriculture (Lal et al. 2004). Soil respiration, a pathway releasing CO_2 from terrestrial ecosystems to the atmosphere (Raich and Tufekcioglu 2000), contains all CO_2 fluxes originated from rhizodeposits, crop litter, and soil organic matter decomposition. A small change of soil respiration can have a large impact on CO_2 concentration in the atmosphere (Lal et al. 2004; Grace and Rayment 2000). Hence, soil respiration in agroecosystems is crucial in the global C cycle and C budget. The development of low-carbon agriculture systems to store as much C as possible in soils is considered an urgent measure in the 21st century (Fedoroff et al. 2010). A key strategy in reducing C emission from agricultural soils is to adopt improved farming practices in crop production (Gan et al. 2011b), such as the use of conservation tillage and intercropping (Fig. 1).

Conservation tillage is increasingly used in crop production due to their environmental advantages over traditional moldboard plow systems. Studies have shown that reduced tillage

F. Hu · Q. Chai (✉) · A. Yu · W. Yin · H. Cui · Y. Gan
Gansu Provincial Key Laboratory of Aridland Crop Science,
Lanzhou 730070, China
e-mail: Chaiq@gsau.edu.cn

F. Hu · Q. Chai · A. Yu · W. Yin · H. Cui
College of Agronomy, Gansu Agricultural University,
Lanzhou 730070, China

Y. Gan
Agriculture and Agri-Food Canada, Semiarid Prairie Agricultural
Research Centre, Swift Current, SK S9H 3X2, Canada

Fig. 1 Wheat/maize intercropping system tested at Wuwei experimental station, China, at **a** early co-growth period, **b** late co-growth period, **c** after wheat harvesting with stubble standing, and **d** after wheat harvesting with stubble mulching

decreases soil disturbance (Alletto et al. 2010), inhibits soil microbial activities, and lowers CO₂ emission from the soil (Boeckx et al. 2011). Furthermore, less disturbance to the soil due to reduced tillage in combination with crop residue retention on the soil surface has the potential to lower the emission of CO₂ (Fuentes et al. 2011). However, most of the published studies are concerned on monoculture crops, i.e., sole cropping such as maize (*Zea mays*) and wheat (*Triticum aestivum*), and less attention has been paid on intercropping systems.

Intercropping has been widely adopted in many parts of the world as a means of increasing crop productivity (Chai et al. 2013), aiming at closing yield gaps between current crop yields and their potentials (Mueller et al. 2012). Also, cropping intensification is recognized as a key farming strategy in reducing global warming potential (Gan et al. 2012). The adoption of intercropping has significant advantages on carbon sequestration than conventional sole cropping (Asgedom and Kebeab 2011; Gan et al. 2011a). For example, maize–soybean (*Glycine max* L.) intercropping (Dyer et al. 2012) and maize–wheat intercropping (Beedy et al. 2010) enhance soil carbon sequestration and reduce carbon emission significantly compared to the corresponding monocultures.

In the Hexi Corridor of northwestern China, a typical oasis agricultural region, water availability is extremely low, with less than 150 mm of annual precipitation but with greater than 2,400 mm of annual evaporation (Chai et al. 2013). One of the effective strategies in conserving soil moisture in this area is to reduced tillage coupled with stubble retention. Crop residues

on the soil surface form a barrier against evaporation and help maintain soil moisture. Also, the more the crop residues returned to the soil, the more the carbon is sequestered because residues are precursors of the soil organic matter pool. However, more crop residue input in the soil may mean more CO₂ emission when soil organic matter decomposition occurs. A sustainable strategy is urgently required to increase soil water conservation with stubble retention while reducing carbon emission. Here, we determined (i) the responses of grain yield, evapotranspiration, and carbon emission of wheat–maize intercropping to various tillage and stubble retention options and (ii) carbon emission per unit of water and the emission efficiency of intercropping systems in arid irrigation areas. We tested the hypothesis that reduced tillage in combination with stubble retention under wheat/maize intercropping could increase crop yield, conserve water, and reduce carbon emissions compared to conventional monoculture cropping systems.

2 Materials and methods

2.1 Experimental site

The field experiment was conducted in 2011 and 2012 at the Gansu Agricultural University Research Station, Wuwei, China (37° 96' N, 102° 64' E). This station with altitude 1,506 m, located in the eastern part of the Hexi Corridor of northwestern China, is at the temperate arid zone in the

hinterland of the Eurasia Continent (Fan et al. 2012). The soil was an Aridisol (FAO/UNESCO 1988) with soil bulk density in the 0–1.1-m soil profile averaged 1.4 g cm^{-3} . Based on long-term (1960–2009) weather data, the solar radiation is $6,000 \text{ MJ m}^{-2}$, annual sunshine duration is higher than 2,945 h, and annual mean temperature is 7.2°C with accumulated temperature above 0°C higher than $3,513^\circ\text{C}$ and above 10°C higher than $2,985^\circ\text{C}$, and the frost-free period is 155 days. Mean annual precipitation is 156 mm, but the potential evaporation is 2,400 mm. Total nitrogen (N), phosphorous (P), and organic matter of top (0–60 cm) soil is 0.78, 1.14, and 14.3 g kg^{-1} , respectively.

2.2 Experimental design and plot management

In 2010, a pre-experiment was conducted to create proper wheat straw retention and tillage patterns that were prepared readily for the implementation of various treatments in the following years. In 2011, the four tillage and stubble retention patterns were implemented: (a) no till with 25 cm wheat stubble standing in the field (designated as RT1, stubble standing), (b) no till with 25 cm height of wheat stubble chopped and spread evenly on the soil surface (RT2, stubble mulching), (c) tillage with 25 cm height of stubble was incorporated in the soil (RT3, stubble incorporated), and (days) conventional tillage (CT, tillage without stubble retention, i.e., all stubble was removed out of the field to mimic the local farming practice where stubble was transported to households for feeding and heating use). No-till and reduced tillage were represented by RT in the experiment. The wheat–maize intercropping was planted in strips with six rows of wheat alternated with two rows of maize in the set of 80:80 cm in strip width. The different tillage and stubble retention patterns were applied to the wheat strips only, and all the maize strips were tilled. Conventional tillage without stubble retention was used as the control where a deep tillage (30 cm) in the previous fall was used for weed control and a pre-seeding rotary tillage was used for seedbed preparation. In both reduced and conventional tillage treatments, maize stalks were removed out of the fields for animal feeding. The sole maize and sole wheat were alternated among years, i.e., the plots grown with sole maize in the previous year were planted with sole wheat in the current year and vice versa. Similarly, in the wheat–maize intercropping, the maize strips in the previous year were planted with wheat in the current year and vice versa.

Due to low precipitation at the experimental areas, 150 mm annually, supplemental irrigation was applied according to the recommended irrigation schemes for the crops of study (Yang et al. 2011). All plots received an amount of 120 mm of irrigation in the previous winter just before soil freezing, and then, various irrigation quotas were applied at the different growth stages in the current year to satisfy the treatment requirements. A hydrant pipe system was used for irrigation,

and a flow meter was installed at the discharging end of the pipe to measure and record the irrigation amounts.

Spring wheat (the cultivar Yong-Liang no. 4) was planted on 28 Mar. in 2011 and 19 Mar. in 2012 and harvested on 22 July in 2011 and 18 July in 2012, and the maize (the cultivar Wu-Ke no. 2) was planted on 17 Apr. and 20 Apr. and harvested on 29 Sep. and 30 Sep. in the 2 years, respectively. Plot size was $4.8 \text{ m} \times 10 \text{ m}$. Wheat and maize were alternated in 160-cm-wide strips with wheat strip (80 cm wide) consisted of six rows with a row space of 12 cm, and maize strip (80 cm wide) had two rows of maize with 40 cm interrow. Maize strips were mulched with plastic films at seeding, an innovative technology for boosting maize productivity in arid environments (Gan et al. 2013). Planting density was $6,750,000 \text{ plants ha}^{-1}$ for wheat and $82,500 \text{ plants ha}^{-1}$ for maize. The N rates to sole wheat and sole maize were 225 and 360 kg N ha^{-1} , and P_2O_5 rates were 66 and 158 kg P ha^{-1} . For intercrops, each species received the same rate of fertilizers on a per hectare basis as the sole crops; i.e., the N and P rates were halved for each particular species because each species occupied one-half areas as in the sole cropping. All N and P were applied as base fertilizer to wheat, while maize received 30 % of N as base fertilizer at sowing, 60 % as topdressing at jointing, and the remaining 10 % as topdressing at grain filling.

2.3 Measurements and statistical analysis

2.3.1 Soil respiration

Soil respiration was measured with a CFX-2 system (Soil CO_2 Flux System, CFX-2, PP System Hitchin, UK) connected with a proprietary respiration chamber. Before measuring, all crop residues and other litters on soil surface were removed, and a hole with a diameter the same as the respiration chamber size was made on the maize strips in order to release the stored CO_2 efflux, at least 12 h before the measurement. The chamber, with a sharp edging point at the bottom, was placed on the soil surface and then pushed to the depth of 20 mm. Measurements were made at three places randomly selected in each plot, five values were recorded for each place within 180 s, and the average value was used for each plot. For intercropping, measurements were taken for each of the intercrops by placing the instrument in the wheat strip and maize strips, and the averages of the two strips were used for a plot. The diurnal soil respiration was measured at 2-h interval from 8:00 am to 8:00 pm on the selected dates, and the seasonal measurements were started on 21 Apr. in 2011 and 22 Apr. in 2012, and the rest of the measurements were taken at 20-day interval from April to September in each year.

2.3.2 Evapotranspiration

The evapotranspiration (ET) (mm) was determined using water balance equation (Yang et al. 2011):

$$ET = P + I + WS_s - WS_h \quad (1)$$

where P is the effective precipitation (mm), determined by the US Department of Agriculture Soil Conservation Services method (USDA-SCS 1972), I is the irrigation quota (mm), and WS_s and WS_h are soil water storage (mm) at 0–1.2 m depth before sowing and after harvesting, respectively.

Soil water contents in the 0–30-cm depth by 10-cm increments were measured using the oven-drying method, whereas soil water in the 30–120-cm depth by 30-cm increments was measured using neutron probe (NMM 503 DR, USA). The average value of two intercrops was used for a plot.

2.3.3 Grain yield

All plots were harvested when plants reached the full maturity, and grains were air-dried, cleaned, and weighed for the yield of the sole and intercropped crops individually.

2.3.4 Carbon emission and carbon emission per unit of water

Carbon emission (kg ha^{-1}) was estimated based on soil respiration ($\text{g CO}_2 \text{m}^{-2} \text{h}^{-1}$) using the following equation described by Zhai et al. (2011):

$$CE = \sum \left[\frac{SR(i+1) + SR_i}{2} [t(i+1) - ti] \times \frac{12}{44} \right] \times 24 \times 10 \quad (2)$$

where SR was the soil respiration ($\text{g CO}_2 \text{m}^{-2} \text{hr}^{-1}$) measured at 20-day interval during the growing season and post-harvest period, $i+1$ and i were the previous and the current sampling date, respectively, and t was the days after sowing.

Based on the conception of water use efficiency, the term “carbon emission per unit of water ($\text{kg ha}^{-1} \text{mm}^{-1}$)” was determined to describe the magnitude of carbon emission (kg ha^{-1}) associated with per unit of ET (mm) and defined as carbon emission during growing season divided by ET, as follows:

$$WUE_{CE} = \frac{CE}{ET} \quad (3)$$

where carbon emission (CE) is the carbon emission (kg ha^{-1}) and ET is the evapotranspiration (mm)

2.3.5 Carbon emission efficiency

There has been a concern that higher crop productivity may result in more carbon emission during the crop production. In order to quantify the correlation between grain yield and carbon emission, the term “carbon emission efficiency (CEE)” was created in the present study, which was expressed as follows:

$$CEE = \frac{GY}{CE} \quad (4)$$

where grain yield (GY) is the grain yield (kg ha^{-1}) and CE is the carbon emission (kg ha^{-1}).

2.3.6 Land equivalent ratio

Land equivalent ratio (LER) is defined as the land area required by sole crops to produce the same volume of grain yield produced by intercrops. A value of $LER > 1.0$ indicates a yield advantage of intercropping over sole cropping and vice versa. The ratio is calculated as follows:

$$LER = \frac{Y_{iw}}{Y_{sw}} + \frac{Y_{im}}{Y_{sm}} \quad (5)$$

where Y_{iw} and Y_{sw} are grain yields of intercropped wheat and sole wheat, respectively, and Y_{im} and Y_{sm} are grain yields of intercropped maize and sole maize, respectively.

Based on the notion of LER, the LER_{CEE} can be explicated as land equivalent ratio which counted on carbon emission efficiency.

2.3.7 Statistical analysis

Data were analyzed using Statistical Analysis Software (SPSS software, 16.0, SPSS Institute Inc., USA), and treatment effects were determined using the Duncan’s multiple range test. Due to significant year $\times$ treatment interactions for most of the variables evaluated in the study, the treatment effect was assessed for each year separately. All significances were declared at the probability level of 0.05.

3 Results and discussion

3.1 The seasonal variation of soil respiration

Soil respiration is one of the most useful tools for measuring CO_2 fluxes from soils (Raich and Tufekcioglu 2000), and the

quantification of soil respiration is linchpins in global carbon budget. Crop management practices such as tillage, cropping sequence, and intensity are the main factors affecting soil respiration (Sainju et al. 2006). In the present study, we found that the seasonal patterns of soil respiration was similar for various sole crops and wheat/maize intercrops, with the respiration rates peaked in late June, began to decline in early July, and then reached the lowest in September in both 2011 and 2012. However, the absolute value of soil respiration differed significantly between treatments (Table 1). Monoculture maize had the highest soil respiration among all cropping patterns evaluated in this study, with a mean value of $1.57 \mu\text{mol m}^{-2} \text{s}^{-1}$, which was nearly twice as high as intercrops (average $0.83 \mu\text{mol m}^{-2} \text{s}^{-1}$). Intercropping reduced soil respiration by an average of 50 % in 2011 and 46 % in 2012, compared to sole maize. The maximal rate of seasonal soil respiration of intercropping was 45 and 47 % lower than that of sole maize in 2011 and 2012, respectively. The growth of maize in intercropping was partially impaired by interspecific competition (Li et al. 2008), leading to a reduction of soil respiration.

Tillage and residue management are two major factors affecting CO_2 emission from soils. Some studies reported that conventional tillage emitted more CO_2 than reduced tillage as a result of soil disturbance from plough (Fuentes et al. 2011), whereas others found no differences between the two tillage practices (Elder and Lal 2008). Our result showed that reduced tillage decreased soil respiration and lowered CO_2 emission compared to conventional tillage (Table 1). Maize monoculture with reduced tillage (average $1.51 \mu\text{mol m}^{-2} \text{s}^{-1}$)

decreased soil respiration by 9 % compared to maize monoculture with conventional tillage ($1.65 \mu\text{mol m}^{-2} \text{s}^{-1}$). Reduced tillage with stubble mulching on the soil surface was most effective in lessening soil respiration among the three reduced tillage treatments. Compared to conventional tillage (as the check), wheat/maize intercropping with reduced tillage and stubble retention on the soil surface decreased soil respiration by 20 % in 2011 and 12 % in 2012. Among all the treatments evaluated in the study, the wheat/maize intercropping with applied stubble mulching under reduced tillage had the lowest soil respiration (Table 1). That was 21 % lower than wheat/maize intercropping under reduced tillage with stubble standing in 2011 and 11 % in 2012. Wheat/maize intercropping with reduced tillage and stubble incorporated in the soil achieved 16.6 % less soil respiration than wheat/maize intercropping with conventional tillage in 2011, but none difference was found in 2012.

In the present study, the soil respiration under reduced tillage with stubble mulching was decreased by 17 to 20 % over conventional tillage, suggesting that carbon dioxide emission can be reduced substantially by use of such reduced with residue retention in the production of field crops.

3.2 Water consumption

Averaged across 2 years, water consumption of wheat–maize intercrops ranged from 682 to 754 mm, sole wheat from 378 to 416 mm, and sole maize from 589 to 644 mm (Table 1). Generally, intercropping increased water consumption by 72 % than sole wheat and by 22 % than sole maize. However,

Table 1 Averaged soil respiration (SR), grain yield, and water consumption of wheat and maize in the sole and intercropped systems in an oasis region, in 2011 and 2012

Treatment	Averaged SR ($\mu\text{mol m}^{-2} \text{s}^{-1}$)		Grain yield (t ha^{-1})		Water consumption (mm)	
	2011	2012	2011	2012	2011	2012
Intercropping						
RT1	0.94	0.85	15.83	15.61	752	676
RT2	0.75	0.76	16.17	16.05	736	670
RT3	0.78	0.82	15.57	14.95	756	685
CT	0.94	0.87	14.70	14.36	771	695
Monoculture						
RT1	0.94	1.63	6.70	13.05	414	591
RT2	0.77	1.34	6.86	13.25	404	579
RT3	0.87	1.56	6.50	12.16	423	595
CT	0.93	1.65	6.39	11.65	424	592
<i>p</i> value ^a	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
LSD (0.05)	0.07	0.09	0.13	0.21	14	10

RT1 reduced tillage with stubble standing, RT2 reduced tillage with stubble mulching, RT3 reduced tillage with stubble incorporated in the soil, CT conventional tillage without stubble retention

^a The *p* value and the LSD (0.05) were for all the treatments in the column

intercropping achieved an equivalent to or higher water use efficiency than sole maize due to greater grain yields (data not presented). All crops, i.e., intercrops and sole crops with reduced tillage, had lower water consumption than those under conventional tillage (Table 1). Intercrops with reduced tillage have the average water consumption of 748 mm in 2011 and 667 mm in 2012, which were, respectively, 3.0 and 2.6 % lower than intercrops with conventional tillage. Moreover, wheat monoculture with reduced tillage decreased water consumption by 2.4 % compared to wheat under conventional tillage.

Stubble mulching on the soil surface combined with reduced tillage had great benefits in reducing water consumption (Table 1). For intercropping, reduced tillage with stubble mulching system decreased water consumption by 4.5 and 3.6 % in 2011 and 2012 compared to conventional tillage system. Whereas, intercropping with stubble standing in the field under reduced tillage or incorporated into the soil did not have any consistent influence on water consumption in either year.

3.3 Yield response

Mixed yields are the total yields of the two component crops, i.e., wheat and maize in the intercropping systems (Table 1). The land equivalent ratios of intercropping with the four different straw management options were all greater than 1.0. On average, intercropping increased yields by 61 % in 2011 and 63 % in 2012 compared to sole crops, and the intercropping with reduced tillage significantly ($P<0.05$) improved the land equivalent ratio over conventional intercropping. These results indicate that the wheat/maize intercropping uses less land area to achieve an equivalent grain yield as sole cropping or producing more grains than sole cropping with the same land area.

Tillage with stubble retention had a significant ($P<0.05$) impact on yields (Table 1). Intercropping with reduced tillage had an average grain yield of 15.85 t ha⁻¹ in 2011 and 15.54 t ha⁻¹ in 2012, an increase of 7.8 % in 2011 and 8.1 % in 2012, compared to intercropping with conventional tillage. Similarly, sole maize with reduced tillage (average yield 12.82 t ha⁻¹) improved yield by 10 % over sole maize with conventional tillage. Among the three conservation tillage treatments, reduced tillage with straw mulching on the soil surface achieved the highest yields (Table 1). In 2011, wheat/maize intercropping under reduced tillage with stubble mulching increased yields by 9.9 % than conventional wheat/maize intercropping; sole wheat under reduced tillage with stubble mulching increased yields by 9.3 % than conventional sole wheat. In 2012, wheat/maize intercropping with reduced tillage and stubble mulching improved yields by 12 % than conventional wheat/maize intercropping, and sole maize

with reduced tillage and stubble mulching increased yields by 9 % compared to conventional sole maize.

3.4 Carbon emission characteristics

3.4.1 Carbon emission during growing seasons

Soil C storage is usually determined by the balance between C inputs from crop residues and rhizodeposits and C loss through CO₂ emission. Less carbon emission means more C sequestration in soils, thus providing adequate C resources for crop growth and mitigating greenhouse gas effects. The present research showed that wheat/maize intercropping emitted comparatively lower carbon, averaging 2,411 kg C ha⁻¹ annually, which was 10 % less than sole maize. This directly relates to lower soil respiration rate of wheat/maize intercropping system. Overall, intercropped maize emitted 26 to 50 % more carbon than the intercropped wheat (Table 2), showing that intercropped maize greatly contributed to carbon emission in the wheat/maize intercropping system. Monoculture maize, however, emitted about 2,585 kg C ha⁻¹ annually, the highest among the planting patterns.

Reduction in soil disturbance combined with residue retention reduced the emission of CO₂ compared with conventional tillage. On average, reduced tillage decreased C emission by 6.7, 5.9, and 7.1 % for intercropping, sole maize, and sole wheat, respectively (Table 2). Wheat/maize intercropping with stubble mulching significantly ($P<0.05$) lessened carbon emission by 10 % over conventional tillage. Compared to conventional tillage, reduced tillage with stubble mulching or with stubble incorporated in the soil of maize monoculture decreased C emission by 14 and 5 %, respectively, and the two respective tillage managements on wheat monoculture decreased the emission by 6 and 8 %, respectively. In contrast to the two distinct tillage types, reduced tillage with stubble standing did not have a consistent effect on carbon emission, suggesting that stubble standing is an unsuitable type of straw management for this area. However, stubble mulching on soil surface or stubble incorporating into topsoil requires mechanical power, which may emit more carbon. More comprehensive analysis is needed to determine the effect of straw retention options on carbon emission in arid irrigation areas.

Our results indicate that the impact of intercropping with reduced tillage on carbon emission is significant. Interplant competition when two crop species are planted in strips and the “relay” growth of the two intercrops during the growing season contribute to the decreased carbon emission. Furthermore, crop residue retention on the soil surface influences soil temperature,

soil moisture, and plant growth which greatly influences soil respiration. However, it is unknown whether or not those effects can be carried on year after year, as more residues are added to the soil with the conservation technology each year. Therefore, further studies on the correlation between the influencing factors and carbon emission are needed in this arid oasis area. Also, the complex agronomic operation of intercropping often requires more labor input or mechanical power than monoculture. Accordingly, some well-designed experiments may be needed to quantify carbon emission at a system level. Also, some systematic research on carbon footprint, defined as the total greenhouse gas emission per unit of grain produced (Gan et al. 2011b), may be required for accurately assessing the benefits of intercropping systems.

3.4.2 Carbon emission per unit of water

Water is the most important factor limiting crop production in the arid northwest China (Chai et al. 2013). Intercropping used more water but emitted less carbon (Table 2). To quantify how much carbon that plants emitted per unit of water used, we created the term “carbon emission per unit of water” in this study and determined the differences between the various cropping systems. Sole maize had the greatest carbon

emission per unit of water used among all treatments, with a mean value of $4.3 \text{ kg C ha}^{-1} \text{ mm}^{-1}$ (Table 2). The adoption of wheat/maize intercropping ($3.4 \text{ kg C ha}^{-1} \text{ mm}^{-1}$) reduced carbon emission per unit of water by 23 % over sole maize albeit increased water consumption. Reduced tillage had average carbon emission per unit of water values of 3.3, 2.3, and $4.2 \text{ kg C ha}^{-1} \text{ mm}^{-1}$ for intercropping, sole wheat, and sole maize, respectively, which was 4.0, 4.7, and 5.3 % lower compared to conventional tillage.

Among the three stubble retention treatments, stubble mulching reduced carbon emission per unit of water by 8 % for wheat/maize intercropping, 10 % for sole maize, and 9 % for sole wheat, as compared with the other two stubble management options (Table 2). In comparison to conventional tillage, reduced tillage decreased carbon emission per unit of water by 10 % for sole maize, by 6 % for sole wheat, and by 9 % for wheat/maize intercropping.

3.4.3 Carbon emission efficiency

The adoption of intercropping systems is an effective approach to increase crop productivity, reduce soil respiration, and lower carbon emission in arid lands (Qin et al. 2013). However, little is known about how effective this approach is in regard to increasing grain yield while reducing carbon

Table 2 Carbon emission during the growing season, carbon emission efficiency (CEE), and carbon emission per unit of water (WUE_{CE}) for various wheat/maize intercropping systems and the corresponding sole cropping in an oasis region, in 2011 and 2012

Treatment	Carbon emission ^a						CEE		WUE _{CE}	
	2011			2012			2011	2012	2011	2012
	Maize	Wheat	Total	Maize	Wheat	Total				
	kg ha ^{−1}						kg kg ^{−1}		kg ha ^{−1} mm ^{−1}	
Intercropping										
RT1	1,715	1,097	2,812	1,302	1,054	2,357	5.63	6.62	3.74	3.49
RT2	1,324	829	2,153	1,260	1,025	2,286	7.51	7.17	2.92	3.41
RT3	1,320	963	2,282	1,288	1,031	2,319	6.82	6.44	3.02	3.38
CT	1,549	1,060	2,609	1,401	1,066	2,466	5.64	5.82	3.38	3.55
Monoculture										
RT1	—	1,119	1,119	2,610	—	2,610	5.99	5.00	2.70	4.39
RT2	—	950	950	2,348	—	2,348	7.22	5.64	2.35	3.97
RT3	—	1,018	1,018	2,501	—	2,501	6.38	4.84	2.40	4.32
CT	—	1,107	1,107	2,641	—	2,641	5.77	4.41	2.61	4.47
<i>p</i> value ^b	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
LSD (0.05)	24	12	11	23	18	12	0.08	0.09	0.03	0.06

RT1 reduced tillage with stubble standing, RT2 reduced tillage with stubble mulching, RT3 reduced tillage with stubble incorporated in the soil, CT conventional tillage without stubble retention

^a Carbon emission was estimated using the soil respiration function (Zhai et al. 2011)

^b The *p* value and the LSD(0.05) were for all the treatments in the column

Table 3 Research hotspots on carbon emission in agro-ecosystem in the recent years, mostly focusing on conservation tillage, N fertilizer application, and crop intensification

Main object	Tillage types	N fertilization	Cropping system	Most carbon emitter	Favorable carbon decrease	Source
Tillage intensity	NT RT CT	Normal application	Monoculture	CT	NT	Boeckx et al. (2011)
Intercropping	CT	Normal application	Maize–wheat, maize–rape, maize–pea, soybean–wheat intercropping Maize, wheat, rape, pea, and soybean monoculture	Maize monoculture	Maize–rape intercropping	Chai et al. (2013)
Conservation tillage, crop residue retention, crop rotation	NT CT	Normal application	Wheat monoculture Maize monoculture Maize–wheat rotation	Maize monoculture without residue retention under CT	Maize–wheat rotation with residue retention under NT	Fuentes et al. (2011)
Diversifying cropping systems	CT	Normal application	Cereal–cereal–durum Cereal–oilseed–durum Pulse–pulse–durum	Cereal–cereal–durum	With oilseed, pulse, and cereal crops were arranged in well-defined cropping sequences, e.g., pulse–pulse–durum	Gan et al. (2011b)
Irrigation, tillage, cropping system, nitrogen fertilization	NT CT	0 kg N ha ⁻¹ (ON) 78 kg N ha ⁻¹ (FN)	Monoculture	Tillage with cropping, with N fertilization and substantial rain or irrigation	No till with reduced N fertilizer in continuous crops or CRP planting	Sainju et al. (2006)
Select beneficial management practices	NT CT	Inorganic N fertilizers	Incorporation of pulses Rotation of forages Use of cover crops	Conventional tillage with abundant N fertilizers without crop intensification	Conservation tillage with proper inorganic N fertilization and intensified cropping system	Asgedom and Kebeab (2011)

Research article: Boeckx et al. (2011); Chai et al. (2013); Fuentes et al. (2011); Gan et al. (2011a, b); Sainju et al. (2006). Review paper: Asgedom and Kebeab (2011)
 NT no tillage, RT reduced tillage, CT conventional tillage, ON zero N, with 0 kg N ha⁻¹, FN fertilize N, with 67 to 134 kg N ha⁻¹, CRP conservation reserve program

emission. In the present study, we used the term “carbon emission efficiency” to determine how much grain yield that was associated with per unit of carbon emission. The land equivalent ratios based on carbon emission efficiency for the four cropping systems were all higher than 2.0, indicating that intercropping not only saves land and increases yield, but also deeply restrains carbon emission. On average, wheat/maize intercropping produced 6.5 kg grain yield per kilogram of carbon emission or 36 % greater than sole maize (4.7 kg kg^{-1}) (Table 2). Reduced tillage with stubble mulching had a great influence on carbon emission efficiency, which contributed to the improved carbon emission efficiency by 18.7 % (5.6 vs. 4.8 kg kg^{-1}) for sole maize, 19.4 % (7.2 vs. 6.1 kg kg^{-1}) for sole wheat, and 19.2 % (7.3 vs. 6.2 kg kg^{-1}) for wheat/maize intercropping, as compared to the other stubble management options.

On average, sole wheat and sole maize under reduced tillage with stubble mulching increased carbon emission efficiency by 15 and 27 % as compared to conventional sole wheat and sole maize, and 28 % was increased by the favorable wheat/maize intercropping system annually; all were statistically significant (Table 2). Compared to conventional tillage, wheat monoculture, maize monoculture, and wheat/maize intercropping under reduced tillage with stubble incorporated in the topsoil boosted carbon emission efficiency by 11, 10, and 16 %, respectively. Sole wheat under reduced tillage with stubble standing also played important role in enhancing carbon emission efficiency, which was increased

by 13 %. Overall, stubble mulching based on reduced tillage was the optimum model in tillage management and residue retention treatments in promoting carbon emission efficiency.

3.5 General discussion

Common beneficial management practices that lead to C emission reduction from farmland are use of proper inorganic N fertilizers, cropping systems, e.g., diversifying cropping systems or crop intensification, and tillage practices, e.g., conservation tillage with or without applied crop residue retention including no-till and reduced tillage (Table 3). Increasing N fertilizer application for the sake of providing sufficient quantity of grains to meet the ever-growing population's needs has given rise to heavy carbon emissions. A well consideration of applying the N fertilizer at right rate, proper selection of the N sources, and timing application is highly recommended (Asgedom and Kebeab 2011). Cropping system is the major component of farmland, by managing farm in space and time. Adopting the diversified cropping systems in crop rotation is one of the promising strategies in mitigating GHG emissions from farming (Gan et al. 2011b). Besides, highly intensified cropping system such as maize-based intercropping is most effective in reducing carbon emission and establishing a sustainable cropping system (Chai et al. 2013).

Conservation tillage with decreased soil disturbance is most advantageous to reduce carbon emission. The less tillage intensity, the less soil respiratory CO_2 -C loss

Fig. 2 The conceptual figure of experimental design with linkages between issues, hypothesis, and results of this research

(Boeckx et al. 2011). Crop residue retention on the soil surface allows a barrier formed on the surface, helping reduce CO₂ emission (Fuentes et al. 2011). The combination of crop residue retention with reduced tillage, sometimes, will enhance such an effect. In view of this, we took the two major components, i.e., cropping system and conservation tillage, into account. Then, we designed a field experiment by well considering the crop intensification, i.e., through wheat–maize intercropping and conservation tillage, through reduced tillage managements to meet the double advantages on carbon emission, which is new and novel. The design was formulated in a rational framework. When combined it with the results after the precise analysis, a conceptual figure was drawn to clearly describe the experimental design with linkages between issues, hypothesis, and results in a concise way (Fig. 2). Results of our study demonstrate that wheat/maize intercropping in combination with conservation tillage can significantly enhance carbon emission efficiency while lower carbon emission per unit of land and per unit of water. Thereby, we strongly recommend this favorable system to meet the side benefits of food security and climate change in arid areas.

4 Conclusions

Wheat/maize intercropping can effectively reduce soil respiration while increasing grain yield with improved water use efficiency significantly; this eventually leads to lower carbon emission per unit of water used and with higher carbon emission efficiency than monocultures. Reduced tillage combined with stubble mulching reduced carbon emission by 9, 12, and 10 %, respectively, for sole maize, sole wheat, and wheat/maize intercropping. Wheat/maize intercropping under reduced tillage decreased carbon emissions per unit of water by 8 % and improved carbon emission efficiency by 19 % compared to conventional tillage. We conclude that wheat/maize intercropping systems with reduced tillage and stubble mulching can be used to effectively lower carbon emissions per unit of land and per unit of water and enhance carbon emission efficiency in arid areas. In the long run, this cropping system, when managed well, can be adapted in a wide range of environments to capture the large benefit of food security while mitigating climate change.

Acknowledgments We are grateful to the research grants provided by the National Key Technology R&D Program (granted number 2012BAD14B10), the National Natural Science Fund (granted number 31360323), the Special Fund for Agro-scientific Research in the Public Interest (granted number 201103001), and the Excellent Youth Foundation of Gansu Scientific Committee (1111RJDA006).

References

- Alletto L, Coquet Y, Benoit P, Hedddadj D, Barriuso E (2010) Tillage management effects on pesticide fate in soils. A review. *Agron Sustain Dev* 30(2). doi:10.1051/agro/2009018
- Asgedom H, Kebreab E (2011) Beneficial management practices and mitigation of greenhouse gas emissions in the agriculture of the Canadian Prairie: a review. *Agron Sustain Dev* 31(3):433–451. doi:10.1007/s13593-011-0016-2
- Beedy TL, Snapp SS, Akinnifesi FK, Sileshi GW (2010) Impact of *Gliricidia sepium* intercropping on soil organic matter fractions in a maize-based cropping system. *Agric Ecosyst Environ* 138(3–4): 139–146. doi:10.1016/j.agee.2010.04.008
- Boeckx P, Van Nieuland K, Van Cleemput O (2011) Short-term effect of tillage intensity on N₂O and CO₂ emissions. *Agron Sustain Dev* 31(3):453–461. doi:10.1007/s13593-011-0001-9
- Chai Q, Qin AZ, Gan YT, Yu AZ (2013) Higher yield and lower carbon emission by intercropping maize with rape, pea, and wheat in arid irrigation areas. *Agron Sustain Dev* 34(34):535–543. doi:10.1007/s13593-013-0161-x
- Dyer L, Oelbermann M, Echarte L (2012) Soil carbon dioxide and nitrous oxide emissions during the growing season from temperate maize-soybean intercrops. *J Plant Nutr Soil Sci* 175(3):394–400
- Elder JW, Lal R (2008) Tillage effects on gaseous emissions from an intensively farmed organic soil in North Central Ohio. *Soil Tillage Res* 98(1):45–55
- Fan ZL, Chai Q, Huang GB, Yu AZ, Huang P, Yang CH, Tao ZQ, Liu HL (2012) Yield and water consumption characteristics of wheat/maize intercropping with reduced tillage in an Oasis region. *Eur J Agron* 25121:1–7. doi:10.1016/j.eja.2012.10.010
- FAO/UNESCO (1988) Soil map of the world: revised legend/prepared by the Food and Agriculture Organization of the United Nations. World soil resources report. Unesco
- Fedoroff NV, Battisti DS, Beachy RN, Cooper PJM, Fischhoff DA, Hodges CN, Knauf VC, Lobell D, Mazur BJ, Molden D, Reynolds MP, Ronald PC, Rosegrant MW, Sanchez PA, Vonshak A, Zhu JK (2010) Radically rethinking agriculture for the 21st century. *Science* 327(5967):833–834. doi:10.1126/science.1186834
- Fuentes M, Hidalgo C, Etchevers J, De León F, Guerrero A, Dendooven L, Verhulst N, Govaerts B (2011) Conservation agriculture, increased organic carbon in the top-soil macro-aggregates and reduced soil CO₂ emissions. *Plant Soil* 355(1–2):183–197. doi:10.1007/s11104-011-1092-4
- Gan YT, Liang C, Huang GB, Malhi S, Brandt S, Katepa-Mupondwa F (2011a) Carbon footprint of canola and mustard is a function of the rate of N fertilizer. *Int J Life Cycle Asses* 17(1):58–68. doi:10.1007/s11367-011-0337-z
- Gan YT, Liang C, Wang XY, McConkey B (2011b) Lowering carbon footprint of durum wheat by diversifying cropping systems. *Field Crop Res* 122(3):199–206. doi:10.1016/j.fcr.2011.03.020
- Gan YT, Liang C, Campbell CA, Zentner RP, Lemke RL, Wang H, Yang C (2012) Carbon footprint of spring wheat in response to fallow frequency and soil carbon changes over 25 years on the semiarid Canadian prairie. *Eur J Agron* 43:175–184. doi:10.1016/j.eja.2012.07.004
- Gan YT, Siddique KHM, Turner NC, Li XG, Niu JY, Yang C, Liu LP, Chai Q (2013) Chapter seven—ridge-furrow mulching systems—an innovative technique for boosting crop productivity in semiarid rain-fed environments. In: Donald LS (ed) *Advances in agronomy*, vol 118. Academic Press, pp 429–476. doi:10.1016/B978-0-12-405942-9.00007-4
- Grace J, Rayment M (2000) Respiration in the balance. *Nature* 404:819–820
- Lal R, Griffin M, Apt J, Lave L, Morgan MG (2004) Ecology. Managing soil carbon. *Science* 304(5669):393. doi:10.1126/science.1093079

- Li H, Qiu JJ, Wang LG (2008) Characterization of farm land soil respiration and modeling analysis of contribution of root respiration. *Trans Chin Soc Agr Eng* 24(4):14–20. doi:[10.3321/j.issn:1002-6819.2008.04.003](https://doi.org/10.3321/j.issn:1002-6819.2008.04.003)
- Mueller ND, Gerber JS, Johnston M, Ray DK, Ramankutty N, Foley JA (2012) Closing yield gaps through nutrient and water management. *Nature* 490(7419):254–257. doi:[10.1038/nature11420](https://doi.org/10.1038/nature11420)
- Qin AZ, Huang GB, Chai Q, Yu AZ, Huang P (2013) Grain yield and soil respiratory response to intercropping systems on arid land. *Field Crop Res* 144:1–10. doi:[10.1016/j.fcr.2012.12.005](https://doi.org/10.1016/j.fcr.2012.12.005)
- Raich JW, Tufekcioglu A (2000) Vegetation and soil respiration: correlations and controls. *Biogeochemistry* 48:71–90
- Sainju UM, Jabro JD, Stevens WB (2006) Soil carbon dioxide emission and carbon content as affected by irrigation, tillage, cropping system, and nitrogen fertilization. *J Environ Qual* 37:98–106. doi:[10.2134/jeq2006.0392](https://doi.org/10.2134/jeq2006.0392)
- Schlesinger WH, Andrews JA (2000) Soil respiration and the global carbon cycle. *Biogeochemistry* 48:7–20
- USDA-SCS (1972) National engineering handbook. Soil Conservation Service. USDA, Washington
- Yang CH, Huang GB, Chai Q, Luo ZX (2011) Water use and yield of wheat/maize intercropping under alternate irrigation in the oasis field of northwest China. *Field Crop Res* 124:426–432. doi:[10.1016/j.fcr.2011.07.013](https://doi.org/10.1016/j.fcr.2011.07.013)
- Zhai LM, Liu HB, Zhang JZ, Huang J, Wang BR (2011) Long-term application of organic manure and mineral fertilizer on N₂O and CO₂ emissions in a red soil from cultivated maize-wheat rotation in China. *J Integr Agr* 10(11):1748–1757. doi:[10.1016/s1671-2927\(11\)60174-0](https://doi.org/10.1016/s1671-2927(11)60174-0)