Imaging bacterial cells and biofilms adhering to hydrophobic organic compounds-water interfaces

Alexis Canette, Priscilla Branchu, Régis Grimaud, Murielle M. Naitali

► To cite this version:

Alexis Canette, Priscilla Branchu, Régis Grimaud, Murielle M. Naitali. Imaging bacterial cells and biofilms adhering to hydrophobic organic compounds-water interfaces. McGenity, Terry J., Timmis, Kenneth N., Nogales Fernández, Balbina. Hydrocarbon and Lipid Microbiology Protocols, Springer, 2016, Springer Protocols Handbooks, 978-3-662-49135-5. 10.1007/8623_2015_80 . hal-01294116

HAL Id: hal-01294116

<https://hal.science/hal-01294116>

Submitted on 27 Mar 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Title

2 **Imaging bacterial cells and biofilms adhering to hydrophobic organic compounds-water**
3 **interfaces**

4 Authors

5 Alexis Canette, Priscilla Branchu, Régis Grimaud*, Murielle Naïtali

6 Affiliation

7 Alexis Canette

8 INRA, UMR 1319 Micalis, Domaine de Vilvert, 78350 Jouy-en-Josas, France
9 and
10 AgroParisTech, UMR Micalis, 1 avenue des Olympiades, 91300 Massy, France
11 alexis.canette@jouy.inra.fr

12

13 Priscilla Branchu

14 INRA, UMR 1319 Micalis, Domaine de Vilvert, 78350 Jouy-en-Josas, France
15 and
16 AgroParisTech, UMR Micalis, 1 avenue des Olympiades, 91300 Massy, France
17 priscilla.branchu@ifr.ac.uk

18

19 Régis Grimaud *Corresponding author

20 Université de Pau et des Pays de l'Adour, Equipe Environnement et Microbiologie, UMRCNRS 5254
21 IPREM, IBEAS - BP1155, 64013 Pau Cedex, France.
22 regis.grimaud@univ-pau.fr

23

24 Murielle Naïtali

25 INRA, UMR 1319 Micalis, Domaine de Vilvert, 78350 Jouy-en-Josas, France
26 and
27 AgroParisTech, UMR Micalis, 1 avenue des Olympiades, 91300 Massy, France
28 murielle.naitali@agroparistech.fr

29
30

Abstract

Assimilation of Hydrophobic Organic Compound (HOC) entails frequently the formation of biofilm at the HOC-water interface. Knowledge on the behavior of cells at the oil-water interface and within the structured biofilm is therefore important to understand the degradation of the HOC in ecosystems. The adhesion and biofilm formation on oil-water interface is best documented by microscopic observations. In this chapter we thus describe two methods for observation of bacterial cells and biofilms growing at the HOC-water interface. The first method uses CLSM (confocal laser scanning microscopy) to obtain *in situ* images of biofilm developing on thin paraffin strip which offers a flat transparent surface allowing imaging directly through the bottom of the culture dish without sampling. Alternatively, the biofilm can be grown on a paraffin strip deposited on a glass microscope slide and then imaged from the top when high resolution is needed. The second method addresses the problematic of the ultrastructure of biofilm developing on HOC. It enables to obtain by TEM (transmission electron microscopy) images of cross sections of biofilms with identification of the side in contact with the HOC.

Key Words

Hydrocarbons, lipids, adhesion, biofilm, oleolytic bacteria, CLSM, TEM

1. Introduction

Hydrophobic Organic Compounds (HOC), including the chemical classes of lipids and hydrocarbons are ubiquitous components of the organic matter in ecosystems. The so called oleolytic bacteria have the ability to use members of either one or both classes as a substrate (1,2). However, they have to face the very low water solubility of these substrates. Adhesion to the HOC-water interface and the subsequent formation of biofilms (3D-architectures of bacterial cells embedded within matrixes of biopolymers) are strategies shared by many oleolytic bacteria to overcome the low solubility of their substrates (3, 4, 5). Different mechanisms have been proposed for the stimulation of the rate of mass transfer from the oily phase to cell surface. The localization of cell in the vicinity of the interface results in reduction of the length of the diffusion pathway of the hydrocarbon from the interface to the cell surface which in turn increases the mass transfer rate (6). Extracellular matrixes of biofilms provide confined environments where biosurfactants can be accumulated and contribute to hydrocarbons uptake by substrate emulsification or pseudo-solubilization. A direct contact of bacteria with hydrocarbons is also possible via the modification of the cell surface (7) offering the possibility of direct uptake from the hydrocarbon-water interface, although this remained to be demonstrated. In the case of lipids which must undergo hydrolysis into fatty acids by extracellular lipases before uptake by the cell, adhesion and biofilm formation at the water-lipid interface provide the advantage to retain together the exo-enzymes, the lipids, the hydrolysis products and the cells thus preventing them from rapid transportation to bulk water. The behavior of cells at the oil-water interface and within the structured biofilm is therefore one important aspect of the degradation of the HOCs in ecosystems.

The adhesion and biofilm formation at the oil-water interface is best studied by microscopic observations. Although a huge number of HOCs degrading strains have been isolated over the past decades, adhesion of these strains to their substrate is rarely documented. Only few authors provided images of bacteria cells adhered to HOC-water interface using light microscopy, scanning electron microscopy and confocal laser scanning microscopy (CLSM) (8, 9, 10, 11, 12, 13). Studies of the behavior of bacteria at the HOC-water interface and within biofilms often involve examination and comparison of different strains (wild type versus mutants or different species). This can be only

achieved if the HOC-water interfaces generated are compatible with the quantitative measurement of the adhesion or biofilm formation that is hardly the case with spherical or hemispherical droplets of HOC and crystals of irregular shape. To fulfill these requirements, we propose a methodology using thin films of paraffin (15 μm) to generate a substratum to grow biofilms. Ultrathin paraffin films are obtained by slicing solid paraffin with a microtome. Paraffin films pasted to the bottom of a culture dish offer a flat surface and are thin enough to allow in situ observation through the bottom of the culture dish avoiding thus disruption of the biofilm structure. A better definition of imaging can be obtained using a microscope glass slide as solid substratum to paste the paraffin; the biofilm is then observed upside-down through a coverslip.

Determination of the ultrastructure of biofilms adhered to HOC-water interface are also of great interest. In general, biofilm cells exhibit heterogeneity in their physiology which can be reflected by the presence of different morphotypes. In the case of biofilm developing on hydrocarbons, intracellular inclusions of storage lipids have been often observed (14). For instance, biofilm cells of *Marinobacter hydrocarbonoclasticus* SP17 growing on hexadecane accumulate wax ester in their cytoplasm. These storage lipids persist after detachment from the biofilm and provide cells with energy for colonization of unoccupied interfaces (15). Lipid inclusions can be investigated using CLSM using specific fluorescent dye (figure 1). However, the ultrastructure of growing cells on HOCs is best studied using transmission electron microscopy (TEM) (1, 16). The distribution and arrangement of cells containing inclusions within the biofilm can be investigated by TEM provided that the biofilm structure is preserved and the orientation of the biofilm in respect to the HOC surface is traceable. Here, we describe a method where the biofilm is grown in a petri dish whose the bottom has been coated with paraffin. TEM requires sectioning the biofilm included in epoxy resin. As paraffin is incompatible with epoxy resin, the biofilm is removed from the paraffin surface by embedding it in agarose. Then agarose pieces are processed for TEM by paying attention to keep track of the side bearing the biofilm. This procedure enables to obtain images of cross sections of biofilms with identification of the side in contact with the paraffin. The imaging methods were both applied on a marine gram negative bacterium, *Marinobacter hydrocarbonoclasticus* SP17 and on a gram positive bacterium *Rhodococcus equi* NapRu1.

104 **2. Materials**

105 **2.1 Observation of biofilms grown on paraffin by CLSM**

106 2.1.1 Strains, culture media and substratum for biofilm growth

107 1.The protocol described here has been developed with the bacterial strains *M. hydrocarbonaclasticus*
108 SP17(*I7*)and *R.equi*NapRu1 (*3*) (Note 1)

109 2.Synthetic Sea Water (SSW): NaCl 0,2 mol.L⁻¹, KCl 10 mmol.L⁻¹, Tris-HCl 50 mmol.L⁻¹ pH 7,8,
110 NH₄Cl 56 mmol.L⁻¹, K₂HPO₄ 427 µmol.L⁻¹, FeSO₄ 8 µmol.L⁻¹, MgSO₄ 65 mmol.L⁻¹ et CaCl₂
111 13 mmol.L⁻¹) (Note 2)

112 3. Sodium lactate 2 mol.L⁻¹ adjusted to pH 7

113 4. Paraffin Normal Q Path® (VWR International, France)

114 5. Petri dishes (55 and 90 mm diameter) (Note 3)

115 6. Washed glass microscopic slides(25x75 mm) (Note 4)

116 7. Ultramicrotome (UC6, Leica microsystems, Germany)

117

118 2.1.2 CLSM observation of the biofilm

119 1.Fluorescent dye: Syto®9 (L-10316, Life Technologies, France), Syto®61 (S-11343, Life
120 Technologies), BODIPY® (D-3922, Life technologies) (Note 5)

121 2. Glass coverslips N° 1.5 (Knittel Gläser, Germany)

122 3. Silicon spacer for coverslip, Press-to-Seal™, 1 mm thick (Invitrogen™, Life Technologies)

123 4.Inverted Confocal Laser Scanning Microscope, Leica TCS SP8 AOBS (Leica Microsystems,
124 Germany) (Note 6)

125 5. Image analysis software IMARIS 7.7.2 software (Bitplane, Switzerland) and Fiji (Fiji.sc/Fiji)(Note
126 7)

127 **2.2Observation of biofilms grown on paraffin by TEM**

128 2.2.1 Strains, culture media and substratum for biofilm growth

129 See subheading 2.1.1

130 2.2.2 TEM observation of the biofilm

131 1. Sodium cacodylate buffer 0.15 or 0.1 mol.L⁻¹, pH 7.4 (Sigma-Aldrich, France) (Note 8). The pH is
132 adjusted with HCl

133 2.Fixative solution (preparedexemporaneously): 2.5% glutaraldehyde (from glutaraldehyde solution
134 Grade I, 25% in H₂O, Sigma-Aldrich) in 0.15 M sodium cacodylate buffer at pH 7.4 or 0.10 M sodium
135 cacodylate buffer at pH 7.4 (Note 8)

136 3.Post-fixative solution (preparedexemporaneously): 1% osmium tetroxide (Electron Microscopy
137 Sciences, France) in 0.15 M sodium cacodylate buffer at pH 7.4 or 0.10 M sodium cacodylate buffer
138 pH 7.4 (accordingly to the fixative solution)

139 4.2.5% agarose solution in distilled water (Seakem® LE agarose, Lonza, France).Agarose solution
140 must be maintained at 50 °Cin an incubator for example(Note 9)

141 5.Ethanol 99.8% (AnalaR NORMAPUR, VWR International)

142 6.Fresh or thawed epoxy resin (R1165, Low Viscosity Resin - Medium grade, Agar Scientific, France)
143 (Note 10)

144 7.A filtered solution of 0.05% oolong tea extract (OTE) in distilled water (OTE, Eloïse, France)
145 (Note 11)

146 9.Whatman® grade n°1 cellulose filter paper (Sigma-Aldrich)

147 10.2 mL Eppendorf®microcentrifuge tubes (Eppendorf, France) (Note 12)

- 148 11. Rotator for Eppendorf microcentrifuge tubes
- 149 12. Vacuum desiccator
- 150 13. Diamond knife for ultrathin sectioning (35 ° angle, DU3520 Diatome 2 mm) and ultramicrotome
- 151 (UC6, Leica microsystems, Germany)
- 152 14. 200 and 300 mesh copper grids (G200-Cu and G300-Cu, Electron Microscopy Sciences)
- 153 15. Flat embedding molds for TEM (70900 silicone clear Electron Microscopy Sciences or 70907
- 154 Dykstra blue, Electron Microscopy Sciences) (Note 13)
- 155 16. HT7700 transmission electron microscope (Hitachi, Japan) equipped with an 8 million pixels
- 156 format CCD camera driven by the image capture engine software AMT version 6.02.
- 157

158 3. Methods

159 3.1 Observation of biofilms grown on paraffin by CLSM

160 3.1.1. Pre-culture of bacteria

161 1. Inoculate 100 μL of a glycerol stock of bacteria (strain SP17 or NapRu1) stored at $-80\text{ }^{\circ}\text{C}$ into
162 10 mL of SSW supplemented with 20 mmol.L^{-1} sodium lactate as a carbon source in a 50 mL tube and
163 incubate overnight at 180 rpm, $30\text{ }^{\circ}\text{C}$.

164 2. Transfer 100 μL of the culture in 10 mL of SSW plus 20 mmol.L^{-1} sodium lactate and incubate
165 overnight at 180 rpm, $30\text{ }^{\circ}\text{C}$.

166 3. Centrifuge the overnight culture at $20\text{ }^{\circ}\text{C}$, $10\,000\text{ g}$ for 15 min, wash the cell pellet twice in SSW and
167 resuspend the cells in SSW at an $\text{OD}_{600\text{nm}}=0.01$.

168

169 3.1.2. Biofilm culture

170 1. Prepare the substratum

171 Melt the paraffin at $65\text{ }^{\circ}\text{C}$ and pour it into cassettes for microtome. Let the paraffin cooled-down on a
172 cool top board to allow solidification. Cut the solid paraffin with a microtome in $15\text{ }\mu\text{m}$ thick strips.
173 Strips must be transported on ice (for example in a Petri dish) then stored at $4\text{ }^{\circ}\text{C}$ until use. A thin strip
174 of paraffin is placed on a microscope glass slide within a Petri dish (90 mm) or directly on the bottom
175 of a Petri dish (55 mm). Put the Petri dish in an incubator at $45\text{ }^{\circ}\text{C}$ for 10 min to soften the paraffin and
176 make it adhere to the solid support. Before use, decontaminate by exposing the Petri dish to UV light for
177 5 min.

178 2. Inoculate the solid support

179 Pour 30 mL of the bacterial suspension ($\text{OD}_{600\text{nm}}=0.01$) in the Petri dish (90 mm) containing the thin
180 paraffin strip stuck on a microscopic slide or 5 mL in a Petri dish (55 mm) containing the thin paraffin

strip stuck on the bottom. Make sure that the bacterial suspension covers the whole surface of the solid substratum. Incubate at 30° C without shaking for 5 days (Note 14).

3.1.3. CLSM observation of the biofilm

1. Biofilm staining with fluorescent markers (Note 15)

Remove the culture medium leaving about 1 mL to keep the biofilm wet. It is important to avoid dehydration to conserve the biofilm structure. Add the selected fluorescent markers to the recommended concentration (see manufacturer's instructions) and incubate at room temperature in dark for 15 min. Here, either Syto®9 (diluted 1:1000) or the mixture Syto®61 (1:1000)/BODIPY® (final concentration 0.05 mg.mL⁻¹) are used.

2. Biofilm observation

Biofilms grown on paraffin thin film deposited on the bottom of the petri dish are imaged directly, without mounting, through the bottom of the Petri dish with an inverted microscope. This approach is preferred when biofilms are loosely attached to paraffin and mounting would result in loss in the biofilm integrity. However, if a maximal resolution is desired, the biofilm is grown on paraffin film pasted on a microscope slide. Then the biofilm is mounted between the slide and a coverslip with a silicon spacer to avoid compressing it and observed upside down on the inverted microscope. Biofilms mounted or not are observed with a water immersion HCX APO L, 0.80 NA, W U-V-I, Leica 40 X objective with a working distance of 3.3 mm. Adjust the laser excitation and fluorescence signal recuperation according to the characteristic of the markers (absorption and fluorescence emission maxima Syto®9 482/500 nm, Syto®61 628/645 nm, BODIPY® 493/503 nm).

3. Acquire xy image or 3D xyz-stack and process the image series using IMARIS 7.7.2 software and Fiji (Figure 1).

205

206 **3.2 Observation of biofilms grown on paraffin by TEM**

207 The basic principles and techniques of electron microscopy for biological applications are described
208 elsewhere(18). This protocol is designed to visualize more specifically the spatial distribution in the
209 biofilm of cells containing intracytoplasmic inclusions. In order to visualize the extracellular matrix,
210 bacterial capsules or appendices, other fixative cocktails and other electron microscopy techniques
211 have to be used (19, 20).

212 3.2.1. Pre-culture of bacteria

213 See subheading 3.1.1.

214 3.2.2. Biofilm culture

215 1.Coat the bottom of petri dishes with paraffin by pouring 15 mL of liquid paraffin heated at 50 °C.

216 Let the paraffin solidify by cooling at room temperature.

217 2.Twenty milliliters of bacterial suspension prepared as described in 3.1.1 are used to inoculate the
218 paraffin coated Petri dishes. Make sure that the whole surface of the solid paraffin is covered by the
219 cell suspension. Incubate at 30 °C without shaking for 5 days (Note 14).

220

221 3.2.3. TEM observation of the biofilm

222 To prevent dehydration, the samples must be kept immersed in solution during the all procedure,
223 except for step 3. During the change of solutions, it is therefore recommended to not discard all the
224 solution from the previous step. Addition and removal of solutions must be performed with extreme
225 care to avoid biofilm alteration. Be aware of the toxicity of the compounds used (Note 16).

226 1. Fix the biofilm

227 Remove the culture medium from the petri dish and pour 20 mL of the fixative solution, incubate at
228 room temperature for 1 h. The cacodylate buffer concentration in fixative solution is 0.15 mol.L⁻¹for
229 *M. hydrocarbonoclasticus*SP17 and 0.10 mol.L⁻¹for *R. equi* NapRu1. At this stage samples can be
230 stored few months in the fixative solution, at 4°C. Wash three times during 5 min with the cacodylate
231 bufferat the corresponding concentration.

232

233 2.Post-fix the biofilm

234 Remove the fixative solution,add 20 mL of post-fixative solution and incubate at room temperature for
235 1 h. Wash twice for 10 min in distilled water. At this stage samples can be stored few days
236 maximum, at 4 °C.

237

238 3.Embed the biofilm in agarose

239 Discard the distilled water and quickly embed the fixedbiofilm in 30 mL of a 2.5% agarose solution
240 maintained above its gelling point at 50 °C (Note 9). Tilt the Petri dish and carefully swirl the melted
241 agarose to obtain a homogenous surface. Let the agarose gellify for1 h at 4 °C.

242 Detach the agarose gel from the petri dish wall by moving all around the edges of the dish a thin blade
243 inserted between the agarose and the dish wall. Unmold carefully the agarose by detachingit from the
244 paraffin. The biofilm normally separates easily from the paraffin staying in the agarose layer.

245 Checkvisually that all the biofilm has been removed from the surface of the paraffin.Transfer the
246 agarose slab, biofilm facing down, in a Petri dish containing the first solution of dehydration (30%
247 ethanol). Make sure that the biofilm bearing face is always immersed in the solution. Cut the agarose
248 slab into big pieces(1 cm³) with a scalpel and choose pieces where the biofilm seems thicker and
249 homogeneous.

250

251 4. Dehydrate the agarose pieces in ethanol solutions series: 50%, 70%, 90% and 2x100% v/v, 15 min
252 for each step at room temperature. Removal and addition of ethanol solution must be carried out with
253 care to not damage the biofilm. Dehydration procedure can be stopped at the stage of 70% ethanol.
254 Samples are stable for few days maximum, at 4°C.

255 After dehydration reduce the thickness of the agarose layer above the biofilm to 1 mm and cut small
256 pieces of 3 mm long by 1 mm wide. Transfer each piece in an Eppendorf™ microcentrifuge tube
257 containing 700 µL of a 100% ethanol (third bath).

258

259 5. Impregnate successively the dehydrated samples with epoxy resin solutions in ethanol of increasing
260 concentrations: 25% and 33% for 10 min, 50% overnight, then 66% and 75% for 10 min.

261 Impregnations are performed at room temperature under rotary shaking (Notes 17 and 18). Then
262 impregnate with pure (100%) fresh resin, for 10 min up to few hours, with the caps of the
263 microcentrifuge tubes opened to let evaporate the last traces of ethanol. Impregnate once more with
264 pure fresh resin overnight on rotary shaking. Finally, impregnate with pure fresh resin with the caps of
265 the microcentrifuge tubes opened in under vacuum to maximize resin penetration (from 10 min to few
266 hours). After dehydration and impregnation the biofilm bearing face of agarose pieces can be easily
267 recognized by their black coloration.

268

269 6. Include samples in epoxy resin.

270 Pour a thin layer of pure fresh epoxy resin into TEM molds and allow to polymerize for 2 h at 60 °C.

271 Place each impregnated sample, in a mold, on the top of the polymerized epoxy layer in such a way
272 that the biofilm black layer is below and parallel to the bottom of mold. The viscous thin layer of resin
273 in the bottom of the mold will help to stabilize the sample during polymerization (Note 19).

274 Fill completely the cavity with pure fresh epoxy resin, considering volume reduction during
275 polymerization. Put the mold under vacuum for 10 min up to few hours to maximize resin penetration

276 and eliminate air bubbles. Polymerize at 60 °C, during 16 to 18 h, depending on the resin hardness
277 preferred for the cutting. Resin blocks can be stored *ad vitam æternam* in a temperate and dry place.

278

279 7. Make ultrathin sections (60 to 80 nm) with a diamond knife and an ultramicrotome. Deposit on 200
280 or 300 mesh copper grids (Note 20). Stain each grid with one drop of 0.05% OTE solution for 30 min,
281 washed with 4 successive drops of distilled water and dried on Whatman grade n°1 cellulose filter
282 paper.

283

284 8. Observe in a TEM microscope between 80 to 120 kV (Figure 2). Contrast increases when
285 accelerating voltage is decreased and objective aperture is more closed (*21*).

286

4. Notes

Note 1. Although this method was originally developed for *M. hydrocarbonaclasticus* SP17 and *R. equi* NapRu1, it can be adapted, by using the appropriated culture medium, to any strain that is able to form biofilm on paraffin or other HOCs, hydrocarbon or lipidsthat are solid at the growth temperature. The HOC must be soft enough to be cut in very thin slices. Crystalized compounds (such as pure alkanes or aromatics) are not suitable since they are not sliceable.

Note 2. Any other medium that allows the studied strain to grow, adhere and form biofilm on HOCcan be employed.

Note 3. Classical Petri dishes are employed. Other diameters can be used. If the biofilm is observed directly through the bottom of the dish, specific Petri dishes with a bottom designed for high-end microscopic analyses have to be used (μ -Dish50 mm, low, Ibidi) for higher resolution imaging.

Note 4. Microscopic slides and coverslips are cleaned by acetone/ethanol (50/50 V/V), dried and stored in sterile conditions in Petri dishes until used.

Note 5. Syto®9 (green fluorescent dye) and Syto®61 (red fluorescent dye) are nucleic acid stains and nonpolar BODIPY® (green fluorescent dye) labels neutral and non-polar lipid components. Other fluorescent dyes can be used according to the structures to be observed as previously described (22).

Note 6. The use of an inverted CLSM is required to observe directly biofilms *in situ* without perturbing their architecture during staining and mounting.

Note 7. Various softwares can be used as previously described (22).

Note 8. Adapt and control the osmolarity of the buffer for each type of sample.

Note 9. The agarose solution is maintained at the liquid state at a temperature above its gel point but not too high to avoid the paraffin from melting or any damage to the biofilm.

Note 10. This resin is prepared accordingly to the manufacturer's instructions, aliquotedand stored at - 20°C.

311 Note 11. OTE is a non-hazardous reagent used instead of the hazardous uranyl acetate(23).

312 Note 12. Smaller microcentrifuge tubes are not suitable for dehydration and next steps.

313 Note 13. Pay attention to the compatibility between the type of resin and the mold materials.

314 Note 14. Time and temperature depend on the strain cultivated and on the conditions studied. If
315 incubation is very long, prevent the evaporation by humidifying the atmosphere.

316 Note 15. Wear a lab coat and gloves to manipulate fluorescent markers. Use specific disposal bins.
317 Read the material safety data sheet (MSDS) of the products.

318 Note 16. Work under a fume hood wearing lab coat, gloves and glasses. Use specific disposal bins.
319 Read the MSDS of the products before use.

320 Note 17. While facilitating the early stages of impregnation, propylene oxide traditionally employed is
321 not used in this protocol. It is a hazardous product (boiling point at 34°C) that reacts with many
322 plastics. Instead, ethanol is used for mixed baths with epoxy resin although they mix less easily.

323 Note 18. Number and length of the impregnation steps are adjusted to our models and our epoxy resin.
324 They can be adapted in case of troubles during the sample cutting (block too soft to an ultrathin cut,
325 dissolution of the ultrathin section in the water tank of the knife) or during TEM observations (holes
326 visible in the ultrathin section). Other standardized protocols describing impregnation steps with other
327 resins can be found (24).

328 Note 19. When placing the samples into the cavities of the molds, it is possible to reference them by
329 adding a small paper with a printed code (pen ink diffuses in the resin) on the opposite side of the
330 sample.

331 Note 20. Use the lowest possible mesh copper grids to maximize the observation area (i.e. 200 mesh
332 copper grids). Use 300 mesh copper grids for unstable sections under electron beam.

333

Acknowledgement

We thank the MIMA2 platform (www.jouy.inra.fr/mima2) for its expertise and access to microscopy equipments.

References

1. Klein B, Grossi V, Bouriat P, Goulas P, Grimaud R (2008) Cytoplasmic wax ester accumulation during biofilm-driven substrate assimilation at the alkane–water interface by *Marinobacter hydrocarbonoclasticus* SP17. Res Microbiol 159(2):137-144
2. Tanaka D, Takashima M, Mizuta A, Tanaka S, Sakatoku A, Nishikawa A, Osawa T, Noguchi M, Aizawa SI, Nakamura S 2010 *Acinetobacter* sp. Ud-4 efficiently degrades both edible and mineral oils: isolation and characterization. Curr Microbiol 60:203-209
3. Bouchez-Naïtali M, Rakatozafy H, Marchal R, Leveau JY, Vandecasteele JP (1999) Diversity of bacterial strains degrading hexadecane in relation to the mode of substrate uptake. J Appl Microbiol 86(3):421-428
4. Grimaud R (2010) Biofilm development at interfaces between hydrophobic organic compounds and water. In: Timmis KN, McGenity T, de Lorenzo V, van der Meer JR (eds) Handbook of hydrocarbons and lipid microbiology. Springer, Berlin, pp 1491-1499
5. Mounier J, Camus A, Mitteau I, Vaysse PJ, Goulas P, Grimaud R, Sivadon P (2014) The marine bacterium *Marinobacter hydrocarbonoclasticus* SP17 degrades a wide range of lipids and hydrocarbons through the formation of oleolytic biofilms with distinct gene expression profiles. FEMS Microbiol Ecol 90:816-831
6. Harms H, Smith KEC, Wick LY (2010) Introduction: problems of hydrophobicity / bioavailability. In: Timmis KN, McGenity T, de Lorenzo V, van der Meer JR (eds) Handbook of hydrocarbons and lipid microbiology. Springer, Berlin, pp 1437-1450

358 7. Heipieper HJ, Cornelissen S, Pepi M (2010) Surface properties and cellular energetics of bacteria in
359 response to the presence of hydrocarbons. In: Timmis KN, McGenity T, de Lorenzo V, van der Meer
360 JR (eds) Handbook of hydrocarbons and lipid microbiology. Springer, Berlin, pp 1615-1624

361 8. Whyte LG, Slagman SJ, Pietrantonio F, Bourbonnière L, Koval SF, Lawrence JR, Inniss WE, Greer
362 CW 1999 Physiological adaptations involved in alkane assimilation at a low temperature by
363 *Rhodococcus* sp. strain Q15. Appl Environ Microbiol 65:2961-2968

364 9. Eriksson M, Dalhammar G, Mohn WW (2002) Bacterial growth and biofilm production on pyrene.
365 FEMS Microbiol Ecol 40:21-27

366 10. Wick LY, De Munain AR, Springael D, Harms H (2002) Responses of *Mycobacterium* sp.
367 LB501T to the low bioavailability of solid anthracene. Appl Microbiol Biotechnol 58:378-385

368 11. Rodrigues AC, Brito AG, Wuertz S, Melo LF (2005) Fluorene and phenanthrene uptake by
369 *Pseudomonas putida* ATCC 17514: Kinetics and physiological aspects. Biotechnol Bioeng
370 90:281-289

371 12. Macedo AJ, Kuhlicke U, Neu TR, Timmis KN, Abraham WR (2005) Three stages of a biofilm
372 community developing at the liquid-liquid interface between polychlorinated biphenyls and water.
373 Appl Environ Microbiol 71:7301-7309

374 13. Wouters K, Maes E, Spitz JA, Roeflaers MJB, Wattiau P, Hofkens J, Springael DA (2010) A non-
375 invasive fluorescent staining procedure allows Confocal Laser Scanning Microscopy based imaging of
376 *Mycobacterium* in multispecies biofilms colonizing and degrading polycyclic aromatic hydrocarbons.
377 J Microbiol Meth 83:317-325

378 14. Manilla-Perez E, Reers C, Baumgart M, Hetzler S, Reichelt R, Malkus U, Kalscheuer R,
379 Waltermann M, Steinbuchel A (2010) Analysis of lipid export in hydrocarbonoclastic bacteria of the
380 Genus *Alcanivorax*: Identification of lipid export-negative mutants of *Alcanivorax borkumensis* SK2
381 and *Alcanivorax jadensis* T9. J Bacteriol 192:643-656

382 15.Vaysse PJ, Sivadon P, Goulas P, Grimaud R 2011 Cells dispersed from *Marinobacter*
383 *hydrocarbonoclasticus* SP17 biofilm exhibit a specific protein profile associated with a higher ability
384 to reinitiate biofilm development at the hexadecane-water interface. *Environ Microbiol* 13:737-746.

385 16.Kalscheuer R, Stöveken T, Malkus U, Reichelt R, Golyshin PN, Sabirova JS, Ferrer M, Timmis
386 KN, Steinbüchel A (2007) Analysis of storage lipid accumulation in *Alcanivorax borkumensis*:
387 Evidence for alternative triacylglycerol biosynthesis routes in bacteria. *J Bacteriol* 189(3):918-928

388 17. Gauthier MJ, Lafay B, Christen R, Fernandez L, Acquaviva M, Bonin P, Bertrand JC (1992)
389 *Marinobacter hydrocarbonoclasticus* gen. nov., sp. nov., a new, extremely halotolerant, hydrocarbon-
390 degrading marine bacterium. *Int J Syst Bacteriol* 42:568-76.

391 18.Hayat MA (2000) Principles and Techniques of Electron Microscopy - Biological Applications.
392 Cambridge University Press, Cambridge.

393 19.Erlandsen SL, Kristich CJ, Dunny GM, Wells CL (2004) High-resolution visualization of the
394 microbial glycocalyx with low-voltage scanning electron microscopy: dependence on cationic dyes. *J*
395 *Histochem Cytochem* 52:1427-1435

396 20.Hammerschmidt S, Wolff S, Hocke A, Rosseau S, Müller E, Rohde M (2005) Illustration of
397 pneumococcal polysaccharide capsule during adherence and invasion of epithelial cells. *Infect Immun*
398 73(8):4653-4667

399 21. Dykstra MJ, Reuss LE (2003) Biological electron microscopy: theory, techniques, and
400 troubleshooting, 2nd edition. Kluwer Academic/Plenum Publishers, New York

401 22.Bridier A, Dubois-Brissonnet F, Briandet R (2014) Methods for biofilms constituents and turnover,
402 Section 1. Destructive and nondestructive methods. In: Dobretsov S, Thomason JC, Williams DN
403 (eds) Biofouling Methods. John Wiley & Sons, Ltd, Oxford, UK, pp 139-152

404 23Carpentier A, Abreu S, Trichet M, Satiat-Jeunemaitre BJ (2012) Microwaves and tea: new tools to
405 process plant tissue for transmission electron microscopy. *Microsc*247(1):94-105

406 24.Graham L, Orenstein JM (2007) Processing tissue and cells for transmission electron microscopy in
407 diagnostic pathology and research. Nat Protoc2(10):2439-2450
408

Figure legends

Figure 1. Observation of paraffin-grown biofilms by CLSM

Biofilm of *M. hydrocarbonoclasticus* SP17 (A) and of *R. equi* NapRu1 (C) stained by Syto®9 observed directly through the bottom of a plastic Petri dish and paraffin (optical lens x 40, numerical zoom 1, bar = 50 µm). Biofilm of *M. hydrocarbonoclasticus* SP17 (B) and of *R. equi* NapRu1 (D) stained by the mixture BODIPY®/Syto®61 observed upside down through a microscopic cover slip. (optical lens x 40, numerical zoom 4, bar = 10 µm). Intracytoplasmic lipids are abundant during growth of *M. hydrocarbonoclasticus* SP17 on paraffin (green staining in B) but not for *R. equi* NapRu1 (D).

Images in A and C are three-dimensional projections of biofilm structures reconstructed using the Easy 3D function (in blend mode) of the IMARIS 7.7.2 software. B and D are xy-images for which the overlay of the two channels and the scale bar were performed using Fiji software.

Figure 2. TEM observation of *M. hydrocarbonoclasticus* SP17 (A) and *R. equi* NapRu1 (B).

Only the bottom of the *R. equi* NapRu1 biofilm (i.e. the side in contact with paraffin during growth) is shown as the biofilm is very thick with no stratification within the ultrastructure of cells. The *R. equi* NapRu1 biofilm (Fig. 2B) contain few cells, as compared with the *M. hydrocarbonoclasticus* SP17 biofilm (Fig. 2A), and probably a lot of extracellular matrix. Numbers of cells of *M. hydrocarbonoclasticus* SP17 contained big inclusions in contrast to *R. equi* NapRu1 cells. These observations are in agreement with CLSM imaging (Fig. 1B, D).