

A Splitting Method for Deep Water with Bathymetry

Afaf Bouharguane, Benjamin Melinand

► **To cite this version:**

Afaf Bouharguane, Benjamin Melinand. A Splitting Method for Deep Water with Bathymetry. 2017.
hal-01326794v2

HAL Id: hal-01326794

<https://hal.archives-ouvertes.fr/hal-01326794v2>

Submitted on 14 Feb 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A SPLITTING METHOD FOR DEEP WATER WITH BATHYMETRY

AFAF BOUHARGUANE AND BENJAMIN MELINAND

ABSTRACT. In this paper we derive and prove the wellposedness of a deep water model that generalizes the Saut-Xu system for nonflat bottoms. Then, we present a new numerical method based on a splitting approach for studying this system. The advantage of this method is that it does not require any low pass filter to avoid spurious oscillations. We prove a local error estimate and we show that our scheme represents a good approximation of order one in time. Then, we perform some numerical experiments which confirm our theoretical result and we study three physical phenomena : the evolution of water waves over a rough bottom; the evolution of a KdV soliton when the shallowness parameter increases; the homogenization effect of rapidly varying topographies on water waves.

1. INTRODUCTION

1.1. Presentation of the problem. Understanding the influence of the topography on water waves is an important issue in oceanography. Many physical phenomena are linked to the variation of the topography : shoaling, rip currents, diffraction, Bragg reflection. Since the direct study on the Euler equations is quite involved, several authors derived and justified asymptotic models according to different small parameters. A usual way to derive asymptotic models is to start from the Zakharov/Craig-Sulem-Sulem formulation [38, 16, 17], which is a good formulation for irrotational water waves, and to expand the Dirichlet-Neumann operator. Then, in the shallow water regime for example, several models were obtained like the Saint-Venant equations or the Green-Naghdi or Boussinesq equations, see [2, 24], [22] for instance. The present paper addresses the influence of the bathymetry in deep water, in the sense explained below.

In this paper, a denotes the typical amplitude of the water waves, L the typical length, H the typical height and a_{bott} the typical amplitude of the bathymetry. Then, we introduce three parameters : $\varepsilon = \frac{a}{H}$ the nonlinearity parameter, $\mu = \frac{H^2}{L^2}$ the shallowness parameter and $\beta = \frac{a_{bott}}{H}$ the bathymetric parameter. We recall that assuming μ small leads to shallow water models. In deep water, which is typically the case when μ is of order 1, it is quite common to assume that the steepness parameter $\varepsilon\sqrt{\mu} = \frac{a}{L}$ is small. The first asymptotic model with a small steepness assumption was derived by Matsuno in 2D for a flat and non-flat bottom and weakly transverse 3D water waves [27, 28]. Then, Choi extended this result in 3D for flat bottom [12]. Finally, Bonneton and Lannes gave a 3D version in the case of a non-flat bottom [8]. It is important to notice that these models are only formally derived. It is proven in [2] that smooth enough solutions to these models are close to the solutions of the water waves equations but, to the best of our knowledge, the wellposedness of the Matsuno equations, even in the case of a flat bottom, is still an open problem. This system could be illposed (see Ambrose, Bona and Nicholls [3]). To avoid this difficulty, Saut and Xu ([34]) developed an equivalent system to the Matsuno system which is consistent with the water waves problem and with the same accuracy. Then, they proved that this new system is wellposed. However, this model is for a flat bottom. In this paper, we derive (see Section 2), use, and prove the wellposedness of a generalization of the Saut-Xu system with a non-flat bottom which is the following system

$$(1.1) \quad \begin{cases} \partial_t \zeta - \mathcal{H}_\mu v + \varepsilon \sqrt{\mu} \left(\frac{1}{2} v \partial_x \zeta + \frac{1}{2} \mathcal{H}_\mu (v \partial_x \mathcal{H}_\mu \zeta) + \mathcal{H}_\mu (\zeta \partial_x \mathcal{H}_\mu v) + \zeta \partial_x v \right) = \beta \sqrt{\mu} \partial_x (B_\mu v) \\ \partial_t v + \partial_x \zeta + \frac{3\varepsilon \sqrt{\mu}}{2} v \partial_x v - \frac{\varepsilon \sqrt{\mu}}{2} \partial_x \zeta \mathcal{H}_\mu \partial_x \zeta - \frac{\varepsilon \sqrt{\mu}}{2} v \mathcal{H}_\mu^2 \partial_x v = 0, \end{cases}$$

where (see Subsection 1.2 for the notations)

$$\mathcal{H}_\mu = -\frac{\tanh(\sqrt{\mu}D)}{D} \partial_x \text{ and } B_\mu = \operatorname{sech}(\sqrt{\mu}D) (b \operatorname{sech}(\sqrt{\mu}D) \cdot).$$

Many authors developed numerical approaches to study the impact of the bottom on water waves, see for instance [29], [25], [36], [18], [10], [30], [20], [6], [7]). However to the best of our knowledge, when one works with deep water, there is no convergence result in the literature. After the original work of Craig and Sulem ([16]) and the paper of Craig et al. ([14]), Guyenne and Nicholls ([19]) developed a numerical method based on a pseudospectral method and a fourth-order Runge-Kutta scheme for the time integration. The linear terms are solved exactly whereas the nonlinear terms are viewed as source terms. Their approach has been developed for the whole water waves equations but we could easily adapt it to our system. However with their scheme, we observe spurious oscillations in the wave profile that lead to instabilities. These errors seem to appear when the nonlinear part is evaluated via the Fourier transform. This is the aliasing phenomenon. Guyenne and Nicholls also observe these oscillations and, to fix it, they apply at every time step a low-pass filter. The scheme that we propose in this paper avoids this low-pass filter.

We present a new numerical method based on a splitting approach for studying nonlinear water waves in the presence of a bottom. We remark that the Saut-Xu system contains a dispersive part and a nonlinear transport part. Thus, the splitting method becomes an interesting alternative to solve the system since this approach is commonly used to split different physical terms, see for instance [32]. We also motivate our decomposition by the fact that, due to the pseudodifferential operator, some terms in the dispersive part may be computed efficiently using the fast Fourier transform. The transport part is computed by a Lax-Wendroff method. Various versions of the splitting method have been developed for instance for the nonlinear Schrodinger, the viscous Burgers equation, Korteweg-de-Vries equations [9, 21, 26, 33, 37]. Thanks to this splitting, we only use a pseudospectral method for the nonlocal terms (contrary to [16, 19]), which limits the aliasing phenomenon and allows us to avoid a low-pass filter.

We denote by Φ^t the nonlinear flow associated to the Saut-Xu system (1.1), $\Phi_{\mathcal{A}}^t$ and $\Phi_{\mathcal{D}}^t$, respectively, the evolution operator associated with the transport part (see equation (3.1)) and with the dispersive part (see (3.2)). We consider the Lie formula defined by

$$(1.2) \quad \mathcal{Y}^t = \Phi_{\mathcal{A}}^t \Phi_{\mathcal{D}}^t.$$

The Saut-Xu system (1.1) is a quasilinear system. This implies derivatives losses in the proof of the convergence. In Theorem 4.6, we show that the numerical solution converges to the solution of the Saut-Xu system (1.1) in the $H^{N+\frac{1}{2}} \times H^N$ -norm for initial data in $H^{N+\frac{1}{2}-7} \times H^{N-7}(\mathbb{R})$, where $N \geq 7$.

Notice that it is not hard to generalize the present work to the Lie formula $\Phi_{\mathcal{D}}^t \Phi_{\mathcal{A}}^t$. We also make the choice to prove a convergence result for a Lie splitting but our proof can be adapted to a Strang splitting or a more complex one. Finally, notice that our scheme can be used for other equations.

The paper is organised as follows. In the next section, we extend the Saut-Xu system by adding a topography term and we prove a local wellposedness result. We also show that the flow map Φ^t is uniformly Lipschitzian. In section 3, we split the problem and we give some estimates on $\Phi_{\mathcal{A}}^t$ and $\Phi_{\mathcal{D}}^t$. In Section 4, we prove a local error estimate and we show that the Lie method represents a good approximation of order one in time (Theorem 4.6). Finally, in Section 5, we perform some numerical experiments which confirm our theoretical result and we illustrate three physical phenomena : the evolution of water waves over a rough bottom; the evolution of a KdV soliton when the shallowness parameter increases and the homogenization effect of rapidly varying topographies on water waves.

1.2. Notations and assumptions.

- x denotes the horizontal variable and z the vertical variable. In this paper, we only study the 2D case ($x \in \mathbb{R}$).
- We assume that

$$(1.3) \quad 0 \leq \varepsilon, \beta \leq 1, \exists \mu_{\max} > \mu_{\min} > 0, \mu_{\max} \geq \mu \geq \mu_{\min}.$$

- We denote $\delta = \max(\varepsilon, \beta)$.
- We denote $\Lambda = (1 - \partial_x^2)^{1/2}$ and $H^s(\mathbb{R}) = \{u \in L^2(\mathbb{R}), \|u\|_{H^s} = \|\Lambda^s u\|_{L^2} < \infty\}$ the usual Sobolev space for $s \geq 0$.
- Let $f \in \mathcal{C}^0(\mathbb{R})$ and $m \in \mathbb{N}$ such that $\frac{f}{1+|x|^m} \in L^\infty(\mathbb{R})$. We define the Fourier multiplier $f(D) : H^m(\mathbb{R}) \rightarrow L^2(\mathbb{R})$ as

$$\forall u \in H^m(\mathbb{R}), \widehat{f(D)u}(\xi) = f(\xi)\widehat{u}(\xi).$$

- D denotes the Fourier multiplier corresponding to $\frac{\partial_x}{i}$.
- We denote by $C(c_1, c_2, \dots)$ a generic positive constant, strictly positive, which depends on parameters c_1, c_2, \dots .

2. THE SAUT-XU SYSTEM

In this part, we extend the Saut-Xu system ([34]) for a non-flat bottom. Then, we give a wellposedness result that generalizes the one of Saut and Xu.

The Matsuno system, which is a full dispersion model for deep waters, is an asymptotic model of the water waves equations with an accuracy of order $\mathcal{O}(\delta^2)$. Bonneton and Lannes [8] formulated it in the following way in the presence of a non flat topography

$$(2.1) \quad \begin{cases} \partial_t \zeta - \frac{1}{\sqrt{\mu\nu}} \mathcal{H}_\mu v + \frac{\varepsilon}{\nu} (\mathcal{H}_\mu (\zeta \partial_x \mathcal{H}_\mu v) + \partial_x (\zeta v)) = \frac{\beta}{\nu} \partial_x (B_\mu v) \\ \partial_t v + \partial_x \zeta + \frac{\varepsilon}{\nu} v \partial_x v - \varepsilon \sqrt{\mu} \partial_x \zeta \mathcal{H}_\mu \partial_x \zeta = 0, \end{cases}$$

where $\zeta = \zeta(t, x)$ is the free surface, $v = v(t, x)$ is the horizontal velocity at the surface, $\nu = \frac{\tanh(\sqrt{\mu})}{\sqrt{\mu}}$, \mathcal{H}_μ and B_μ are Fourier multipliers,

$$\mathcal{H}_\mu = -\frac{\tanh(\sqrt{\mu}D)}{D} \partial_x \text{ and } B_\mu = \text{sech}(\sqrt{\mu}D) (b \text{sech}(\sqrt{\mu}D) \cdot),$$

and $-1 + \beta b$ is the topography. (We erased the fluid part)

In [2], Alvarez-Samaniego and Lannes show that this model is consistent with the Zakharov/Craig-Sulem-Sulem formulation when $\beta = 0$ and it is not painful to generalize their result to the case when $\beta \neq 0$. In [34], Saut and Xu obtained a new model with the same accuracy with the Matsuno system

thanks to a nonlinear change of variables. Notice that this change of variables is inspired by [5]. The advantage of this model is that they proved a local wellposedness on large time for this new model. We follow their approach. We define new variables

$$(2.2) \quad \tilde{v} = v + \frac{\varepsilon\sqrt{\mu}}{2}v\mathcal{H}_\mu\partial_x\zeta \text{ and } \tilde{\zeta} = \zeta - \frac{\varepsilon\sqrt{\mu}}{4}v^2.$$

Then, up to terms of order $\mathcal{O}(\delta^2)$, $\tilde{\zeta}$ and \tilde{v} satisfy (we omit the tildes for the sake of simplicity)

$$(2.3) \quad \begin{cases} \partial_t\zeta + \left(\frac{\varepsilon}{\nu} - \frac{\varepsilon\sqrt{\mu}}{2}\right)v\partial_x\zeta - \frac{1}{\sqrt{\mu\nu}}\mathcal{H}_\mu v + \frac{\varepsilon}{\nu}\left(\frac{1}{2}\mathcal{H}_\mu(v\partial_x\mathcal{H}_\mu\zeta) + \mathcal{H}_\mu(\zeta\partial_x\mathcal{H}_\mu v) + \zeta\partial_x v\right) = \frac{\beta}{\nu}\partial_x(B_\mu v) \\ \partial_t v + \left(\frac{\varepsilon}{\nu} + \frac{\varepsilon\sqrt{\mu}}{2}\right)v\partial_x v + \partial_x\zeta - \frac{\varepsilon\sqrt{\mu}}{2}\partial_x\zeta\mathcal{H}_\mu\partial_x\zeta - \frac{\varepsilon}{2\nu}v\mathcal{H}_\mu^2\partial_x v = 0. \end{cases}$$

Since our motivation is the study of water waves in deep water (μ close to 1), we assume that $\nu = \frac{1}{\mu}$. Hence we study the following system which is a variable bottom analog of the system of Saut and Xu

$$\begin{cases} \partial_t\zeta - \mathcal{H}_\mu v + \varepsilon\sqrt{\mu}\left(\frac{1}{2}v\partial_x\zeta + \frac{1}{2}\mathcal{H}_\mu(v\partial_x\mathcal{H}_\mu\zeta) + \mathcal{H}_\mu(\zeta\partial_x\mathcal{H}_\mu v) + \zeta\partial_x v\right) = \beta\sqrt{\mu}\partial_x(B_\mu v) \\ \partial_t v + \partial_x\zeta + \frac{3\varepsilon\sqrt{\mu}}{2}v\partial_x v - \frac{\varepsilon\sqrt{\mu}}{2}\partial_x\zeta\mathcal{H}_\mu\partial_x\zeta - \frac{\varepsilon\sqrt{\mu}}{2}v\mathcal{H}_\mu^2\partial_x v = 0, \end{cases}$$

In the following, we denote $\mathbf{U} = (\zeta, v)^t$ and we define the energy of the system for $N \in \mathbb{N}$ by

$$(2.4) \quad \mathcal{E}^N(\mathbf{U}) = \frac{1}{\sqrt{\mu}}|\Lambda^N\zeta|_2^2 + \left| |D|^{\frac{1}{2}}\Lambda^N\zeta \right|_2^2 + |v|_{H^N}^2,$$

where $\Lambda = \sqrt{1 + |D|^2}$ and $D = -i\nabla$. We also denote by E_μ^N the energy space related to this norm.

Remark 2.1. Notice that if μ satisfies condition (1.3), the energy \mathcal{E}^N is equivalent to the $H^{N+\frac{1}{2}} \times H^N$ -norm.

The main result of this section is the following local wellposedness result.

Theorem 2.2. Let $N \geq 2$, $\mathbf{U}_0 \in H^{N+\frac{1}{2}}(\mathbb{R}) \times H^N(\mathbb{R})$ and $b \in L^\infty(\mathbb{R})$. We assume that ε, β, μ satisfy Condition (1.3) and

$$|\mathbf{U}_0|_{H^{N+\frac{1}{2}} \times H^N} + |b|_{L^\infty} \leq M.$$

Then, there exists a time $T_0 = T_0\left(M, \frac{1}{\mu_{\min}}, \mu_{\max}\right)$ independent of ε, μ and β and a unique solution $\mathbf{U} \in \mathcal{C}\left(\left[0, \frac{T_0}{\delta}\right], E_\mu^N\right)$ of the system (1.1) with initial data \mathbf{U}_0 . Furthermore, we have the following energy estimate, for all $t \in \left[0, \frac{T_0}{\delta}\right]$,

$$\mathcal{E}^N(\mathbf{U}(t, \cdot)) \leq e^{\delta C_0 t} \mathcal{E}^N(\mathbf{U}_0),$$

where $C_0 = C\left(M, \frac{1}{\mu_{\min}}, \mu_{\max}\right)$.

Proof. We refer to Paragraph IV in [34] for a complete proof and we focus only on the bottom contribution. For $0 \leq \alpha \leq N$, we denote $\mathbf{U}^{(\alpha)} = (\partial_x^\alpha\zeta, \partial_x^\alpha v)$. Then, applying ∂_x^α to System (1.1), we get

$$\partial_t \mathbf{U}^{(\alpha)} + \mathcal{L} \mathbf{U}^{(\alpha)} + \frac{\varepsilon \sqrt{\mu}}{2} \mathbf{1}_{\{\alpha \neq 0\}} \mathcal{B}[\mathbf{U}] \mathbf{U}^{(\alpha)} = \beta \sqrt{\mu} (\partial_x \partial_x^\alpha (B_\mu v), 0)^t + \varepsilon \sqrt{\mu} \mathcal{G}^\alpha,$$

where

$$\mathcal{L} = \begin{pmatrix} 0 & -\frac{1}{\sqrt{\mu v}} \mathcal{H}_\mu \\ \partial_x & 0 \end{pmatrix},$$

$$\mathcal{B}[\mathbf{U}] = \begin{pmatrix} \mathcal{H}_\mu (v \mathcal{H}_\mu \partial_x \cdot) + v \partial_x & \mathcal{H}_\mu (\cdot \mathcal{H}_\mu \partial_x \zeta) - \partial_x \zeta \mathcal{H}_\mu^2 \\ -\partial_x \zeta \mathcal{H}_\mu \partial_x - \mathcal{H}_\mu \partial_x \zeta \partial_x & 3v \partial_x - v \mathcal{H}_\mu^2 \partial_x \end{pmatrix},$$

and $\mathcal{G}^\alpha = (\mathcal{G}_1^\alpha, \mathcal{G}_2^\alpha)^t$ with

$$\mathcal{G}_1^\alpha = \partial_x^\alpha g(\zeta, v) - \frac{1}{2} \sum_{1 \leq \gamma \leq \alpha-1} C_\alpha^\gamma (\mathcal{H}_\mu (\partial_x^\gamma v \mathcal{H}_\mu \partial_x^{1+\alpha-\gamma} \zeta) + \partial_x^\gamma v \partial_x^{1+\alpha-\gamma} \zeta) - \frac{1}{2} \partial_x \zeta (\mathcal{H}_\mu^2 + 1) \partial_x^\alpha v$$

$$\mathcal{G}_2^\alpha = \frac{1}{2} \sum_{1 \leq \gamma \leq \alpha-1} C_\alpha^\gamma \partial_x^{1+\gamma} \zeta \mathcal{H}_\mu \partial_x^{1+\alpha-\gamma} \zeta + \sum_{1 \leq \gamma \leq \alpha} C_\alpha^\gamma \left(-\frac{3}{2} \partial_x^\gamma v \partial_x^{1+\alpha-\gamma} v + \frac{1}{2} \partial_x^\gamma v \mathcal{H}_\mu^2 \partial_x^{1+\alpha-\gamma} v \right)$$

where

$$g(\zeta, v) = -[\mathcal{H}_\mu, \zeta] \mathcal{H}_\mu \partial_x v - \zeta (\mathcal{H}_\mu^2 + 1) \partial_x v.$$

Then we can show, as in Paragraph IV. B in [34] (see the paragraph called Estimate on \mathcal{G}^α) that

$$(2.5) \quad |\mathcal{G}^\alpha|_2 + \left| |D|^{\frac{1}{2}} \mathcal{G}^\alpha \right|_2 \leq C \left(\frac{1}{\mu_{\min}} \right) \mathcal{E}^N(\mathbf{U}).$$

Like Saut and Xu we define a symmetrizer for $\mathcal{L} + \mathcal{B}[\mathbf{U}]$

$$(2.6) \quad \mathcal{S} = \begin{pmatrix} \frac{D}{\tanh(\sqrt{\mu} D)} & 0 \\ 0 & 1 \end{pmatrix}.$$

Notice that $\sqrt{(\mathcal{S} \cdot, \cdot)}$ is a norm equivalent to $\sqrt{\mathcal{E}^0}$. Then, as in Paragraph IV. B in [34] (see the paragraph called Estimate on II), we get

$$(2.7) \quad \left(\left(\mathcal{L} + \frac{\varepsilon \sqrt{\mu}}{2} \mathbf{1}_{\{\alpha \neq 0\}} \mathcal{B}[\mathbf{U}] \right) \mathbf{U}^{(\alpha)}, \mathcal{S} \mathbf{U}^{(\alpha)} \right) \leq \varepsilon C \left(\frac{1}{\mu_{\min}}, \mu_{\max} \right) \sqrt{\mathcal{E}^2(\mathbf{U})} \mathcal{E}^N(\mathbf{U}).$$

Furthermore, for the bottom contribution, we easily get

$$(2.8) \quad \left| \left(\frac{D}{\tanh(\sqrt{\mu} D)} \partial_x^\alpha \zeta, \partial_x \partial_x^\alpha \operatorname{sech}(\sqrt{\mu} D) (b \operatorname{sech}(\sqrt{\mu} D) v) \right) \right| \leq C \left(\frac{1}{\mu_{\min}} \right) |b|_\infty \mathcal{E}^N(\mathbf{U}).$$

Finally, from Equations (2.5), (2.7), (2.8), we obtain,

$$\mathcal{E}^N(\mathbf{U}) \leq \mathcal{E}^N(\mathbf{U}_0) + \delta C \left(\frac{1}{\mu_{\min}}, \mu_{\max} \right) \int_0^t \left(\mathcal{E}^N(\mathbf{U})^{\frac{3}{2}} + \mathcal{E}^N(\mathbf{U}) \right) (s) ds,$$

and there exists a time $T > 0$, such that, for all $t \in [0, \frac{T}{\delta}]$,

$$\mathcal{E}^N(\mathbf{U}(t, \cdot)) \leq C \left(\frac{1}{\mu_{\min}}, \mu_{\max}, \mathcal{E}^N(\mathbf{U}_0) \right).$$

The energy estimate follows from the Gronwall Lemma. \square

In order to use a Lady Windermere's fan argument (a well-known telescopic identity used to relate global and the local error), to prove the convergence of the numerical scheme, we need a Lipschitz property for the flow of the Saut-Xu system (1.1). We first give a control of the differential of the flow with respect to the initial datum.

Proposition 2.3. *Let $N \geq 2$, $\mathbf{V}_0 \in H^{N+\frac{1}{2}}(\mathbb{R}) \times H^N(\mathbb{R})$, $\mathbf{U}_0 \in H^{N+1+\frac{1}{2}}(\mathbb{R}) \times H^{N+1}(\mathbb{R})$, and $b \in L^\infty(\mathbb{R})$. We assume that ε, β, μ satisfy Condition (1.3) and*

$$|\mathbf{V}_0|_{H^{N+\frac{1}{2}} \times H^N} + |\mathbf{U}_0|_{H^{N+1+\frac{1}{2}} \times H^{N+1}} + |b|_{L^\infty} \leq M.$$

Then, there exists a time $T = T\left(M, \frac{1}{\mu_{\min}}, \mu_{\max}\right)$ independent of the parameters ε, μ and β such that $(\Phi^t)'(\mathbf{U}_0) \cdot (\mathbf{V}_0)$ exists on $[0, \frac{T}{8}]$. Furthermore, we have, for all $0 \leq t \leq \frac{T}{8}$,

$$\left|(\Phi^t)'(\mathbf{U}_0) \cdot (\mathbf{V}_0)\right|_{H^{N+\frac{1}{2}} \times H^N} \leq C\left(\frac{1}{\mu_{\min}}, \mu_{\max}, M\right) |\mathbf{V}_0|_{H^{N+\frac{1}{2}} \times H^N}.$$

Proof. We denote by $\mathbf{U}(t) = (\zeta(t), v(t))$ the solution of the Saut-Xu system (1.1) with initial data \mathbf{U}_0 . We denote also $(\eta(t), w(t)) = (\Phi^t)'(\mathbf{U}_0) \cdot (\mathbf{V}_0)$. Then, (η, w) satisfy the following system

$$(2.9) \quad \partial_t \begin{pmatrix} \eta \\ w \end{pmatrix} + \mathcal{L} \begin{pmatrix} \eta \\ w \end{pmatrix} + \varepsilon \sqrt{\mu} \mathcal{N}[(\zeta, v)] \partial_x \begin{pmatrix} \eta \\ w \end{pmatrix} + \varepsilon \sqrt{\mu} \mathcal{N}[(\eta, w)] \partial_x \begin{pmatrix} \zeta \\ v \end{pmatrix} = \beta \sqrt{\mu} (\partial_x (B_\mu w), 0)^t,$$

where

$$\mathcal{L} = \begin{pmatrix} 0 & -\frac{1}{\sqrt{\mu v}} \mathcal{H}_\mu \\ \partial_x & 0 \end{pmatrix} \quad \text{and} \quad \mathcal{N}[(\zeta, v)] = \begin{pmatrix} \frac{1}{2} \mathcal{H}_\mu (v \mathcal{H}_\mu \cdot) + \frac{1}{2} v \mathcal{H}_\mu (\zeta \mathcal{H}_\mu \cdot) + \zeta \\ -\frac{1}{2} \partial_x \zeta \mathcal{H}_\mu & \frac{3}{2} v - \frac{1}{2} v \mathcal{H}_\mu^2 \end{pmatrix}.$$

For $0 \leq \alpha \leq N$, we denote $\mathbf{V}^{(\alpha)} = (\partial_x^\alpha \eta, \partial_x^\alpha w)$. Then, applying ∂_x^α to System (2.9), we get

$$\partial_t \mathbf{V}^{(\alpha)} + \mathcal{L} \mathbf{V}^{(\alpha)} + \frac{\varepsilon \sqrt{\mu}}{2} \mathbb{1}_{\{\alpha \neq 0\}} \left(\mathcal{B}[\mathbf{U}] \mathbf{V}^{(\alpha)} + \mathcal{B}[\mathbf{V}] \partial_x^\alpha \mathbf{U} \right) = \sqrt{\mu} \beta \begin{pmatrix} \partial_x \partial_x^\alpha (B_\mu w) \\ 0 \end{pmatrix} + \varepsilon \sqrt{\mu} \mathcal{J}^\alpha,$$

where

$$\mathcal{B}[\mathbf{U}] = \begin{pmatrix} \mathcal{H}_\mu (v \mathcal{H}_\mu \partial_x \cdot) + v \partial_x & \mathcal{H}_\mu (\cdot \mathcal{H}_\mu \partial_x \zeta) - \partial_x \zeta \mathcal{H}_\mu^2 \\ -\partial_x \zeta \mathcal{H}_\mu \partial_x - \mathcal{H}_\mu \partial_x \zeta \partial_x & 3v \partial_x - v \mathcal{H}_\mu^2 \partial_x \end{pmatrix},$$

$$\mathcal{J}^\alpha = -\partial_x^\alpha \left(\mathcal{N}[(\zeta, v)] \partial_x \begin{pmatrix} \eta \\ w \end{pmatrix} + \mathcal{N}[(\eta, w)] \partial_x \begin{pmatrix} \zeta \\ v \end{pmatrix} \right) + \frac{1}{2} \left(\mathcal{B}[\mathbf{U}] \mathbf{V}^{(\alpha)} + \mathcal{B}[\mathbf{V}] \partial_x^\alpha \mathbf{U} \right).$$

Then, we can show, as in Paragraph IV. B in [34], that

$$(2.10) \quad |\mathcal{J}^\alpha|_2 + \left| |D|^{\frac{1}{2}} \mathcal{J}^\alpha \right|_2 \leq \varepsilon \sqrt{\mu} C \left(\frac{1}{\mu_{\min}} \right) \mathcal{E}^N(\mathbf{V}).$$

We recall that we can symmetrize \mathcal{L} thanks to

$$\mathcal{S} = \begin{pmatrix} \frac{D}{\tanh(\sqrt{\mu}D)} & 0 \\ 0 & 1 \end{pmatrix}.$$

We define the energy associated to this symmetrizer

$$F^\alpha(\mathbf{V}) = \left| \sqrt{\frac{D}{\tanh(\sqrt{\mu}D)}} \partial_x^\alpha \eta \right|_2^2 + |\partial_x^\alpha w|_2^2, \quad \text{and} \quad F^N(\mathbf{V}) = \sum_{0 \leq \alpha \leq N} F^\alpha(\mathbf{V}).$$

We have, for $\alpha \neq 0$,

$$\begin{aligned} \frac{d}{dt} F^\alpha(\mathbf{V}) &= \varepsilon \sqrt{\mu} (\mathcal{J}^\alpha, \mathcal{S}\mathbf{V}^{(\alpha)}) - \frac{\varepsilon \sqrt{\mu}}{2} \left((\mathcal{B}[\mathbf{U}]\mathbf{V}^{(\alpha)}, \mathcal{S}\mathbf{V}^{(\alpha)}) + (\mathcal{B}[\mathbf{V}]\partial_x^\alpha \mathbf{U}, \mathcal{S}\mathbf{V}^{(\alpha)}) \right) \\ &\quad + \beta \sqrt{\mu} (\partial_x \partial_x^\alpha (B_\mu v), \mathcal{S}\mathbf{V}^{(\alpha)}) \\ &= I + II + III + IIII. \end{aligned}$$

We can estimate I thanks to estimate (2.10) and II as in Paragraph IV. B in [34]. For $IIII$, we can proceed as in the previous theorem. For III , we get, thanks to Proposition A.1,

$$|III| \leq \varepsilon \sqrt{\mu} |(\zeta, v)|_{H^{N+1+\frac{1}{2}} \times H^{N+1}} |(\eta, w)|_{H^{N+\frac{1}{2}} \times H^N}$$

Then, we obtain

$$\frac{d}{dt} F^N(\mathbf{V}) \leq \delta \sqrt{\mu} C(M) \left(F^N(\mathbf{V}) + \sqrt{F^N(\mathbf{V})} \right),$$

and the result follows. \square

Proposition 2.4. *Let $N \geq 2$, $\mathbf{U}_0, \mathbf{V}_0 \in H^{N+1+\frac{1}{2}} \times H^{N+1}(\mathbb{R})$ and $b \in L^\infty(\mathbb{R})$. We assume that ε, β, μ satisfy Condition (1.3) and*

$$|\mathbf{V}_0|_{H^{N+1+\frac{1}{2}} \times H^{N+1}} + |\mathbf{U}_0|_{H^{N+1+\frac{1}{2}} \times H^{N+1}} + |b|_{L^\infty} \leq M.$$

Then, there exists a time T independent of ε, μ and β and two unique solutions \mathbf{U}, \mathbf{V} of the system (1.1) on $[0, \frac{T}{\delta}]$ with initial data \mathbf{U}_0 and \mathbf{V}_0 . Furthermore, we have the following Lipschitz estimate, for all $0 \leq t \leq \frac{T}{\delta}$,

$$(2.11) \quad |\mathbf{U}(t, \cdot) - \mathbf{V}(t, \cdot)|_{H^{N+\frac{1}{2}} \times H^N} \leq K |\mathbf{U}_0 - \mathbf{V}_0|_{H^{N+\frac{1}{2}} \times H^N},$$

where $K = C\left(\frac{1}{\mu_{\min}}, \mu_{\max}, M\right)$.

Proof. The existence of \mathbf{U}, \mathbf{V} and T follow from the previous theorem. Furthermore, we have

$$\mathbf{U}(t) - \mathbf{V}(t) = \int_{s=0}^1 (\Phi^t)'(\mathbf{V}_0 + s(\mathbf{U}_0 - \mathbf{V}_0)) \cdot (\mathbf{U}_0 - \mathbf{V}_0).$$

The result follows from Proposition 2.3. \square

3. A SPLITTING SCHEME

In this section, we split the Saut-Xu system (1.1) and we give some estimates for the sub-problems. We consider, separately, the transport part

$$(3.1) \quad \begin{cases} \partial_t \zeta + \frac{\varepsilon \sqrt{\mu}}{2} ((\mathcal{H}_\mu^2 + 1) v) \partial_x \zeta = 0 \\ \partial_t v + \frac{3\varepsilon \sqrt{\mu}}{2} v \partial_x v = 0, \end{cases}$$

and the dispersive part

$$(3.2) \quad \begin{cases} \partial_t \zeta - \mathcal{H}_\mu v + \varepsilon \sqrt{\mu} \left(\frac{1}{2} \mathcal{H}_\mu (v \partial_x \mathcal{H}_\mu \zeta) + \mathcal{H}_\mu (\zeta \partial_x \mathcal{H}_\mu v) + \zeta \partial_x v - \frac{1}{2} \partial_x \zeta \mathcal{H}_\mu^2 v \right) = \beta \sqrt{\mu} \partial_x (B_\mu v) \\ \partial_t v + \partial_x \zeta - \frac{\varepsilon \sqrt{\mu}}{2} \partial_x \zeta \mathcal{H}_\mu \partial_x \zeta - \frac{\varepsilon \sqrt{\mu}}{2} v \mathcal{H}_\mu^2 \partial_x v = 0. \end{cases}$$

We denote by $\Phi_{\mathcal{A}}^t$ the flow of System (3.1) and by $\Phi_{\mathcal{D}}^t$ the flow of System (3.2).

Remark 3.1. Notice that we keep the term $\zeta \partial_x v$ in the first equation and we decompose $v \partial_x \zeta$ as $v \partial_x \zeta = \partial_x \zeta (\mathcal{H}_\mu^2 + 1) v - \partial_x \zeta \mathcal{H}_\mu^2 v$. This will be useful for the local wellposedness of the dispersive part.

In the following, we prove the local existence on large time for Systems (3.1) and (3.2).

3.1. The transport equation. The system (3.1) is a transport equation. Then, it is easy to get the following result.

Proposition 3.2. Let $s_1 \geq 0$, $s_2 > \frac{3}{2}$ and $M > 0$. We assume that ε, μ satisfies Condition (1.3). Then, there exists a time $T_1 = T_1(M, \mu_{\max}) > 0$, such that if

$$|\zeta_0|_{H^{s_1}} + |v_0|_{H^{s_2}} \leq M,$$

we have a unique solution $(\zeta, v) \in \mathcal{C} \left(\left[0, \frac{T_1}{\varepsilon}\right], H^{s_1}(\mathbb{R}) \times H^{s_2}(\mathbb{R}) \right)$, to System (3.1) with initial data (ζ_0, v_0) . Furthermore, we have, for all $t \leq \frac{T_1}{\varepsilon}$,

$$(3.3) \quad |\zeta(t, \cdot)|_{H^{s_1}} + |v(t, \cdot)|_{H^{s_2}} \leq C(M, \mu_{\max}).$$

Finally, if $s_2 \geq 4$ and $M_1 = \max_{0 \leq t \leq \frac{T_1}{\varepsilon}} |v(t, \cdot)|_{H^{s_2-2}}$, then for all $t \leq \frac{T_1}{\varepsilon}$, we have

$$(3.4) \quad |\zeta(t, \cdot)|_{H^{s_1}} + |v(t, \cdot)|_{H^{s_2}} \leq e^{\varepsilon C_1 t} |\mathbf{U}_0|_{H^{s_1} \times H^{s_2}},$$

where $C_1 > 0$ depends on M_1 and μ_{\max} .

Proof. The proof follows from the fact that the quasilinear system (3.1) is symmetric. Thanks to the Coifman-Meyer estimate (see Proposition A.3), we get

$$\frac{d}{dt} \left(|\zeta(t, \cdot)|_{H^{s_1}}^2 + |v(t, \cdot)|_{H^{s_2}}^2 \right) \leq C \varepsilon \sqrt{\mu} \left(|\zeta(t, \cdot)|_{H^{s_1}}^2 + |v(t, \cdot)|_{H^{s_2}}^2 \right)^{\frac{3}{2}}.$$

Then, we see that the energy is bounded uniformly with respect to ε and μ and we get Estimate (3.3). For the second estimate, using the same trick that in Lemma 3.1 in [21], we notice that, if $s_1 \geq 4$,

$$\frac{d}{dt} \left(|\zeta(t, \cdot)|_{H^{s_1}}^2 + |v(t, \cdot)|_{H^{s_2}}^2 \right) \leq \varepsilon \sqrt{\mu} |v(t, \cdot)|_{H^{s_2-2}} \left(|\zeta(t, \cdot)|_{H^{s_1}}^2 + |v(t, \cdot)|_{H^{s_2}}^2 \right).$$

By applying the Gronwall lemma, we get the result. \square

3.2. The dispersive equation. The system (3.2) contains all the dispersive terms of the Saut-Xu system. We have the following estimate for the flow.

Proposition 3.3. *Let $N \geq 2$, and $b \in L^\infty(\mathbb{R})$. We assume that ε, β, μ satisfy Condition (1.3). Then, there exists a time $T_2 = T_2\left(M, \frac{1}{\mu_{\min}}, \mu_{\max}\right)$ such that if*

$$|\zeta_0|_{H^{N+\frac{1}{2}}} + |v_0|_{H^N} + |b|_{L^\infty} \leq M,$$

we have a unique solution $(\zeta, v) \in \mathcal{C}\left(\left[0, \frac{T_2}{\delta}\right], H^{N+\frac{1}{2}}(\mathbb{R}) \times H^N(\mathbb{R})\right)$ to the system (3.2) with initial data (ζ_0, v_0) . Furthermore, we have, for all $t \leq \frac{T_2}{\delta}$,

$$(3.5) \quad |\zeta(t, \cdot)|_{H^{N+\frac{1}{2}}} + |v(t, \cdot)|_{H^N} \leq C \left(M, \mu_{\max}, \frac{1}{\mu_{\min}} \right).$$

Finally, if $N \geq 7$, and

$$M_1 = \max_{0 \leq t \leq \frac{T_2}{\delta}} \left(|\zeta(t, \cdot)|_{H^{N+\frac{1}{2}-2}} + |v(t, \cdot)|_{H^{N-2}} \right),$$

then for all $t \leq \frac{T_2}{\delta}$, we have

$$(3.6) \quad |\zeta(t, \cdot)|_{H^{N+\frac{1}{2}}} + |v(t, \cdot)|_{H^N} \leq e^{\delta C_2 t} |\mathbf{U}_0|_{H^{N+1/2} \times H^N},$$

where C_2 is a positive constant which depends on $\mu_{\max}, \frac{1}{\mu_{\min}}, M_1$.

Proof. The proof is an adaptation of the proof of Theorem 2.2 and part IV in [34]. We notice that, in the proof of Saut and Xu, the transport part can be treated separately and does not influence the control of the other terms. Hence, we can use the same symmetrizer \mathcal{S} that in Theorem 2.2 (see (2.6)) and we get

$$\frac{d}{dt} \mathcal{E}^N(\zeta, v) \leq C \left(\frac{1}{\mu_{\min}} \right) \left(\frac{\varepsilon}{\nu} \mathcal{E}^N(\zeta, v)^{\frac{3}{2}} + \frac{\beta}{\nu} \mathcal{E}^N(\zeta, v) \right).$$

Then, by Remark 2.1, we get Estimate (3.5). Furthermore, we notice that we use the same trick as in Lemma 3.1 in [21]. By keeping the same notations as in Theorem 2.2, we get from Equations (2.7) and (2.8) that

$$\frac{d}{dt} \mathcal{E}^N(\mathbf{U}) \leq \delta C \left(\frac{1}{\mu_{\min}}, \mu_{\max} \right) \left(\left| \tilde{\mathcal{G}}^N \right|_{H^{1/2} \times L^2} + \sqrt{\mathcal{E}^2(\mathbf{U})} + |b|_{L^\infty} \right) \mathcal{E}^N(\mathbf{U})$$

where $\tilde{\mathcal{G}}^N = (\tilde{\mathcal{G}}_1^N, \tilde{\mathcal{G}}_2^N)^t$ with

$$\begin{aligned} \tilde{\mathcal{G}}_1^N &= -\partial_x^N ([\mathcal{H}_\mu, \zeta] \mathcal{H}_\mu \partial_x v + \zeta (\mathcal{H}_\mu^2 + 1) \partial_x v) - \frac{1}{2} \sum_{1 \leq \gamma \leq N-1} C_N^\gamma \mathcal{H}_\mu (\partial_x^\gamma v \mathcal{H}_\mu \partial_x^{1+N-\gamma} \zeta) - \frac{1}{2} \partial_x \zeta (\mathcal{H}_\mu^2 + 1) \partial_x^N v \\ \tilde{\mathcal{G}}_2^N &= \frac{1}{2} \sum_{1 \leq \gamma \leq N-1} C_N^\gamma \partial_x^{1+\gamma} \zeta \mathcal{H}_\mu \partial_x^{1+N-\gamma} \zeta + \frac{1}{2} \sum_{1 \leq \gamma \leq N} C_N^\gamma \partial_x^\gamma v \mathcal{H}_\mu^2 \partial_x^{1+N-\gamma} v. \end{aligned}$$

To explain how we can adapt the trick used in Lemma 3.1 in [21], we focus our attention to one term. For $1 \leq \gamma \leq N-1$, we have to control $\left| \partial_x^{1+\gamma} \zeta \mathcal{H}_\mu \partial_x^{1+N-\gamma} \zeta \right|_{L^2}$. If $\gamma \leq \lfloor \frac{N}{2} \rfloor$, we get from Propositions A.4 and A.1 that

$$|\partial_x^{1+\gamma} \zeta \mathcal{H}_\mu \partial_x^{1+N-\gamma} \zeta|_{L^2} \leq |\partial_x^{1+\gamma} \zeta|_{H^1} |\mathcal{H}_\mu \partial_x^{1+N-\gamma} \zeta|_{L^2} \leq C(\mu_{\max}) |\zeta|_{H^{2+\lfloor \frac{N}{2} \rfloor}} |\zeta|_{H^N},$$

whereas if $\gamma > \lfloor \frac{N}{2} \rfloor$, we have

$$|\partial_x^{1+\gamma} \zeta \mathcal{H}_\mu \partial_x^{1+N-\gamma} \zeta|_{L^2} \leq |\partial_x^{1+\gamma} \zeta|_{L^2} |\mathcal{H}_\mu \partial_x^{1+N-\gamma} \zeta|_{H^1} \leq C(\mu_{\max}) |\zeta|_{H^N} |\zeta|_{H^{2+\lfloor \frac{N}{2} \rfloor}}.$$

We can mimic this method to control the other terms of $\tilde{\mathcal{G}}^N$ and, thanks to Propositions A.1, A.2 and A.4, we obtain if $N \geq 7$ that

$$\frac{d}{dt} \mathcal{E}^N(\zeta, v) \leq \delta C \left(\frac{1}{\mu_{\min}}, \mu_{\max} \right) \left(|\zeta(t, \cdot)|_{H^{N+\frac{1}{2}-2}} + |v(t, \cdot)|_{H^{N-2}} + |b|_{L^\infty} \right) \mathcal{E}^N(\zeta, v).$$

Then, Estimate (3.6) follows. \square

Remark 3.4. *Under the assumption of Proposition 3.3 and if $N \geq 7$, we get from relations (3.4) and (3.6) that, there exists a time $T_3 > 0$, such that for all $t \in [0, \frac{T_3}{\delta}]$,*

$$|\mathcal{Y}^t \mathbf{U}_0|_{H^{N+1/2} \times H^N} \leq e^{C_3 \delta t} |\mathbf{U}_0|_{H^{N+1/2} \times H^N},$$

where $C_3 = C \left(|\mathbf{U}_0|_{H^{N+1/2-2} \times H^{N-2}}, \mu_{\max}, \frac{1}{\mu_{\min}} \right)$ and $T_3 = C \left(|\mathbf{U}_0|_{H^{N+1/2} \times H^N}, \mu_{\max}, \frac{1}{\mu_{\min}} \right)$.

4. ERROR ESTIMATES

The goal of this part is to prove the main result of this paper (Theorem 4.6). Our analysis is based on energy estimates.

4.1. The local error estimate. The local error is the following quantity

$$(4.1) \quad e(t, \mathbf{U}_0) = \Phi^t \mathbf{U}_0 - \mathcal{Y}^t \mathbf{U}_0.$$

Our approach is similar to the one developed in [11]. We use the fact that $\Phi^t \mathbf{U}_0$ satisfies a symmetrizable system. Therefore, e satisfies this system up to a remainder and then, we can control e thanks to energy estimates. In the following we give different technical lemmas in order to control the local error. We recall that the transport operator is the operator \mathcal{A}

$$\mathcal{A}(\zeta, v) = -\frac{\varepsilon \sqrt{\mu}}{2} \begin{pmatrix} ((\mathcal{H}_\mu^2 + 1)v) \partial_x \zeta \\ 3v \partial_x v \end{pmatrix}.$$

The following proposition gives an estimate of the differential of the transport operator.

Lemma 4.1. *Let $s_1, s_2 \geq 0$ and ε, μ satisfying Condition (1.3). Then,*

$$|\mathcal{A}'(\zeta, v) \cdot (\eta, w)|_{H^{s_1} \times H^{s_2}} \leq \varepsilon C(\mu_{\max}) |(\zeta, v)|_{H^{s_1+1} \times H^{s_2+1}} |(\eta, w)|_{H^{s_1+1} \times H^{s_2+1}}.$$

Proof. We have

$$\mathcal{A}'(\zeta, v) \cdot (\eta, w) = -\frac{\varepsilon \sqrt{\mu}}{2} \begin{pmatrix} ((\mathcal{H}_\mu^2 + 1)v) \partial_x \eta + ((\mathcal{H}_\mu^2 + 1)w) \partial_x \zeta \\ 3w \partial_x v + 3v \partial_x w \end{pmatrix},$$

and the estimate follows from Propositions A.1 and A.4. \square

We can do the same for the dispersive part (using also Proposition A.1). We recall that the dispersive operator is the operator \mathcal{D}

$$\mathcal{D}(\zeta, v) = \begin{pmatrix} \mathcal{H}_\mu v + \varepsilon\sqrt{\mu} \left(\frac{1}{2} \mathcal{H}_\mu (v \partial_x \mathcal{H}_\mu \zeta) + \mathcal{H}_\mu (\zeta \partial_x \mathcal{H}_\mu v) + \zeta \partial_x v - \frac{1}{2} \partial_x \zeta \mathcal{H}_\mu^2 v \right) - \beta \sqrt{\mu} \partial_x (B_\mu v) \\ -\partial_x \zeta + \frac{\varepsilon\sqrt{\mu}}{2} \partial_x \zeta \mathcal{H}_\mu \partial_x \zeta + \frac{\varepsilon\sqrt{\mu}}{2} v \mathcal{H}_\mu^2 \partial_x v \end{pmatrix}$$

Lemma 4.2. *Let $s > 0$, ε, β, μ satisfying Condition (1.3) and $b \in L^\infty(\mathbb{R})$. Then,*

$$|\mathcal{D}'(\zeta, v) \cdot (\eta, w)|_{H^s \times H^s} \leq C(\mu_{\max}) (1 + \beta |b|_{L^\infty} + \varepsilon |(\zeta, v)|_{H^{s+1} \times H^{s+1}}) |(\eta, w)|_{H^{s+1} \times H^{s+1}}.$$

Furthermore, we have to control the derivative of the flow $\Phi_{\mathcal{A}}^t$ with respect to the initial data. We denote it by $(\Phi_{\mathcal{A}}^t)'$.

Lemma 4.3. *Let $s_1, s_2 \geq 0$, $M > 0$, ε, β, μ satisfying Condition (1.3) and $b \in L^\infty(\mathbb{R})$. Let $(\zeta_0, v_0) \in H^{s_1+1} \times H^{s_2+1}(\mathbb{R}^d)$ such that,*

$$|(\zeta_0, v_0)|_{H^{s_1+1} \times H^{s_2+1}} \leq M.$$

Then, there exists a time $T = T(M, \mu_{\max})$, such that $(\Phi_{\mathcal{A}}^t)'(\zeta_0, v_0) \cdot (\eta_0, w_0)$ exists for all $t \in [0, \frac{T}{\delta}]$ and if we denote

$$\begin{pmatrix} \eta \\ w \end{pmatrix} = (\Phi_{\mathcal{A}}^t)'(\zeta_0, v_0) \cdot (\eta_0, w_0),$$

for all $0 \leq t \leq \frac{T}{\delta}$,

$$|(\eta, w)(t, \cdot)|_{H^{s_1} \times H^{s_2}} \leq |(\eta_0, w_0)|_{H^{s_1} \times H^{s_2}} C(\mu_{\max}, M).$$

Proof. The quantity (η, w) satisfies the following linear system

$$\begin{cases} \partial_t \eta + \frac{\varepsilon\sqrt{\mu}}{2} (\mathcal{H}_\mu^2 + 1) v \partial_x \eta + \frac{\varepsilon\sqrt{\mu}}{2} (\mathcal{H}_\mu^2 + 1) w \partial_x \zeta = 0, \\ \partial_t w + \frac{3\varepsilon\sqrt{\mu}}{2} v \partial_x w + \frac{3\varepsilon\sqrt{\mu}}{2} w \partial_x v = 0, \end{cases}$$

where $(\zeta, v) = \Phi_{\mathcal{A}}^t(\zeta_0, v_0)$. The result follows from energy estimates, the Gronwall lemma and Proposition 3.2. \square

In the following, we use the fact $\Phi_{\mathcal{A}}^t \circ \Phi_{\mathcal{D}}^t$ satisfies the Saut-Xu system (1.1) up to a remainder. The following lemma is the key point for the control of this remainder.

Lemma 4.4. *Let $N \geq 2$, $M > 0$, ε, β, μ satisfying Condition (1.3) and $b \in L^\infty(\mathbb{R})$. Let $\mathbf{U} = (\zeta, v) \in H^{N+\frac{1}{2}} \times H^N(\mathbb{R}^d)$ such that,*

$$|b|_{L^\infty} + |\mathbf{U}|_{H^{N+\frac{1}{2}} \times H^N(\mathbb{R})} \leq M.$$

Then, there exists a time $T = T\left(M, \mu_{\max}, \frac{1}{\mu_{\min}}\right) > 0$, such that $\Phi_{\mathcal{A}}^t(\mathbf{U})$ exists for all $0 \leq t \leq \frac{T}{\delta}$, and furthermore,

$$\left| (\Phi_{\mathcal{A}}^t)'(\mathbf{U}) \cdot \mathcal{D}(\mathbf{U}) - \mathcal{D}(\Phi_{\mathcal{A}}^t(\mathbf{U})) \right|_{H^{N-2} \times H^{N-2}} \leq \varepsilon C\left(M, \mu_{\max}, \frac{1}{\mu_{\min}}\right) t.$$

Proof. The existence of T follows from Proposition 3.2. Then, we notice that

$$(\Phi_{\mathcal{A}}^t)'(\mathbf{U}) \cdot \mathcal{D}(\mathbf{U}) - \mathcal{D}(\Phi_{\mathcal{A}}^t(\mathbf{U})) = \int_0^t \mathcal{A}'(\Phi_{\mathcal{A}}^s(\mathbf{U})) \cdot ((\Phi_{\mathcal{A}}^s)'(\mathbf{U}) \cdot \mathcal{D}(\mathbf{U})) - \mathcal{D}'(\Phi_{\mathcal{A}}^s(\mathbf{U})) \cdot \mathcal{A}(\Phi_{\mathcal{A}}^s(\mathbf{U})).$$

Using Lemmas 4.1, 4.2 and Proposition 3.2, we get,

$$\left| (\Phi_{\mathcal{A}}^t)'(\mathbf{U}) \cdot \mathcal{D}(\mathbf{U}) - \mathcal{D}(\Phi_{\mathcal{A}}^t(\mathbf{U})) \right|_{H^{N-2} \times H^{N-2}} \leq C(\mu_{\max}, M) \int_0^t \varepsilon \left| (\Phi_{\mathcal{A}}^s)'(\mathbf{U}) \cdot \mathcal{D}(\mathbf{U}) \right|_{H^{N-1} \times H^{N-1}} + |\mathcal{A}(\Phi_{\mathcal{A}}^s(\mathbf{U}))|_{H^{N-1} \times H^{N-1}}.$$

Then, using Lemma 4.3, the product estimate A.4 and the expression of \mathcal{A} , we obtain

$$\left| (\Phi_{\mathcal{A}}^t)'(\mathbf{U}) \cdot \mathcal{D}(\mathbf{U}) - \mathcal{D}(\Phi_{\mathcal{A}}^t(\mathbf{U})) \right|_{H^{N-2} \times H^{N-2}} \leq \varepsilon C(\mu_{\max}, M) \int_0^t |\mathcal{D}(\mathbf{U})|_{H^{N-1} \times H^{N-1}} + |\Phi_{\mathcal{A}}^s(\mathbf{U})|_{H^N \times H^N}^2.$$

Finally, the result follows from the expression of \mathcal{D} , the product estimate A.4 and Proposition A.1. \square

We can now give the main result of this part, the local error estimate.

Proposition 4.5. *Let $N \geq 4$, $M > 0$, ε, β, μ satisfying Condition (1.3) and $b \in L^\infty(\mathbb{R})$. Let $\mathbf{U}_0 = (\zeta_0, v_0)$ such that,*

$$|b|_{L^\infty} + |\mathbf{U}_0|_{H^{N+\frac{1}{2}} \times H^N} \leq M.$$

Then, there exists a time $T_4 = T_4\left(M, \frac{1}{\mu_{\min}}, \mu_{\max}\right) > 0$, such that the local error $e(t, \mathbf{U})$ defined in (4.1) exists for all $0 \leq t \leq \frac{T_4}{\delta}$, and furthermore,

$$|e(t, \mathbf{U}_0)|_{H^{N-4+\frac{1}{2}} \times H^{N-4}} \leq \delta C_4 t^2,$$

where $C_4 = C\left(\frac{1}{\mu_{\min}}, \mu_{\max}, M\right)$.

Proof. From Propositions 3.2 and 3.3, we obtain the existence of T . We denote

$$\mathbf{U}(t) = \begin{pmatrix} \zeta(t) \\ v(t) \end{pmatrix} = \Phi^t(\mathbf{U}_0) \quad \text{and} \quad \mathbf{V}(t) = \begin{pmatrix} \eta(t) \\ w(t) \end{pmatrix} = \Phi_{\mathcal{A}}^t(\Phi_{\mathcal{D}}^t(\mathbf{U}_0)).$$

Then, from Theorem 2.2 and Propositions 3.2 and 3.3 we also have, for all $0 \leq t \leq \frac{T}{\delta}$,

$$(4.2) \quad |\mathbf{U}(t, \cdot)|_{H^{N+\frac{1}{2}} \times H^N} + |\mathbf{V}(t, \cdot)|_{H^{N+\frac{1}{2}} \times H^N} \leq C\left(\frac{1}{\mu_{\min}}, \mu_{\max}, M\right).$$

We know that (ζ, v) satisfy the Saut-Xu system (1.1). Furthermore, (η, w) also satisfy the Saut-Xu system (1.1) up to a remainder

$$\partial_t \begin{pmatrix} \eta \\ w \end{pmatrix} = \mathcal{A}(\eta, w) + \mathcal{D}(\eta, w) + \mathcal{R}(t),$$

where $\mathcal{R}(t) = (\Phi_{\mathcal{A}}^t)'(\Phi_{\mathcal{D}}^t(\mathbf{U}_0)) \cdot \mathcal{D}(\Phi_{\mathcal{D}}^t(\mathbf{U}_0)) - \mathcal{D}(\Phi_{\mathcal{A}}^t(\Phi_{\mathcal{D}}^t(\mathbf{U}_0)))$. Therefore, the local error e satisfies the following system

$$(4.3) \quad \partial_t e = \begin{pmatrix} 0 & H_\mu \\ -\partial_x & 0 \end{pmatrix} e + \begin{pmatrix} 0 & \beta\sqrt{\mu}B_\mu \\ 0 & 0 \end{pmatrix} e + \mathcal{T}_\mu((\zeta, v), (\eta, w)) - \mathcal{R}(t),$$

where the operator $\mathcal{T}_\mu(\mathbf{U}, \mathbf{V})$ is quadratic and satisfies the following estimate, for $0 \leq s \leq N-1$,

$$(4.4) \quad |\mathcal{T}_\mu((\zeta, v), (\eta, w))|_{H^s \times H^s} \leq \varepsilon C \left(\frac{1}{\mu_{\min}}, \mu_{\max}, M \right) |e|_{H^{s+1} \times H^{s+1}}.$$

Then, since $e|_{t=0} = 0$,

$$e(t, \cdot) = \int_0^t \partial_t e(s, \cdot) ds,$$

and since e satisfies (4.3), we obtain, thanks to Estimates (4.2), (4.4) and Lemma 4.4,

$$(4.5) \quad |e(t, \cdot)|_{H^{N-2} \times H^{N-2}} \leq C \left(\frac{1}{\mu_{\min}}, \mu_{\max}, M \right) t.$$

Furthermore, we recall that the Saut-Xu system (1.1) is symmetrizable thanks to the symmetrizer (see Theorem 2.2)

$$\mathcal{S} = \begin{pmatrix} \frac{D}{\tanh(\sqrt{\mu}D)} & 0 \\ 0 & 1 \end{pmatrix}.$$

Therefore, applying \mathcal{S} to the system (4.3), and using the fact that $\sqrt{(\mathcal{S}\cdot, \cdot)}$ is a norm equivalent to the $H^{\frac{1}{2}} \times L^2$ -norm, we obtain, thanks to estimates (4.2), (4.4) and (4.5) and Lemma 4.4,

$$\frac{d}{dt} \mathcal{F}(e) \leq C \left(\frac{1}{\mu_{\min}}, \mu_{\max}, M \right) \left(\beta \mathcal{F}(e) + \varepsilon t \sqrt{\mathcal{F}(e)} \right),$$

where $\mathcal{F}(e) = \sum_{|\alpha| \leq N-4} (S \partial_x^\alpha e, \partial_x^\alpha e)$. Then, we get

$$\mathcal{F}(e)(t) \leq \delta C \left(\frac{1}{\mu_{\min}}, \mu_{\max}, M \right) \int_0^t \mathcal{F}(e)(s) + s \sqrt{\mathcal{F}(e)(s)} ds.$$

Denoting $\mathcal{M}(t) = \max_{[0,t]} \sqrt{\mathcal{F}(e)(t)}$, we have

$$\mathcal{M}(t) \leq \delta C \left(\frac{1}{\mu_{\min}}, \mu_{\max}, M \right) \int_0^t \mathcal{M}(s) + s ds,$$

and the result follows from the Grönwall's lemma. \square

4.2. Global error estimate. In this part, we prove our main result. We denote by

$$\mathbf{U}_k = (\mathcal{Y}^{\Delta t})^k \mathbf{U}_0$$

the approximate solution and by $\mathbf{U}(t_k) := \Phi^{k\Delta t} \mathbf{U}_0$ the exact solution at the time $t_k = k\Delta t$.

Theorem 4.6. *Let $N \geq 7$, $M > 0$, ε, β, μ satisfying Condition (1.3) and $b \in L^\infty(\mathbb{R})$. Let $\mathbf{U}_0 = (\zeta_0, v_0)$ such that,*

$$|b|_{L^\infty} + |\mathbf{U}_0|_{H^{N+\frac{1}{2}} \times H^N} \leq M.$$

Let $\mathbf{U}_0(t, \cdot)$ the solution of the Saut-Xu equations (1.1) with initial data \mathbf{U}_0 defined on $[0, \frac{T}{\delta}]$. Then, there exist constants $\gamma, \nu, \Delta t_0, C_0 > 0$ such that for all $\Delta t \in]0, \Delta t_0]$ and for all $n \in \mathbb{N}$ such that $0 \leq n\Delta t \leq \frac{T}{\delta}$,

$$|\mathbf{U}_n|_{H^{N+\frac{1}{2}} \times H^N} \leq \nu \text{ and } \left| \Phi^{n\Delta t}(\mathbf{U}_0) - (\mathcal{Y}^{\Delta t})^n(\mathbf{U}_0) \right|_{H^{N-4+\frac{1}{2}} \times H^{N-4}} \leq \gamma\Delta t.$$

Proof. The proof is based on a Lady's Windermere's fan argument and is similar to the one in [9] (see also [21]). In order to simplify the notations, we forget the dependence on $\frac{1}{\mu_{\min}}$ and μ_{\max} in all the constants. We denote by X^N the following space

$$X^N = H^{N+\frac{1}{2}} \times H^N.$$

Thanks to Theorem 2.2, there exists ρ such that, for all $t^k = k\Delta t \in [0, \frac{T}{\delta}]$,

$$|\mathbf{U}(t_k)|_{X^N} = |\Phi^{t_k}\mathbf{U}_0|_{X^N} \leq \rho.$$

We prove by induction that there exists $\Delta t_0, \gamma, \nu$ such that if $0 < \Delta t \leq \Delta t_0$, for all $n \in \mathbb{N}$ with $n\Delta t \leq \frac{T}{\delta}$,

- (i) $|\mathbf{U}_n - \mathbf{U}(t_n)|_{X^{N-4}} \leq \gamma\Delta t$,
- (ii) $|\mathbf{U}_n|_{X^N} \leq e^{C_3(M_1)\delta n\Delta t} |\mathbf{U}_0|_{X^N} \leq M_0$,
- (iii) $|\mathbf{U}_n|_{X^{N-2}} \leq M_1$,
- (iv) $|\mathbf{U}_n|_{X^{N-4}} \leq 2\rho$,

with

$$M_1 = e^{C_3(2\rho)T} M, M_0 = e^{C_3(M_1)T} M, \gamma = T \max(K, 1) C_4 \left(e^{C_0(M_0)T} M_0 \right),$$

$$\Delta t_0 = \min \left(T, T_0(M_0), T_3(M_0), T_4(M_0), \frac{\rho}{\gamma} \right), K = K \left(M_0 e^{TC_0(M_0)} \right)$$

where $C_0, T_0, C_3, T_3, C_4, T_4, K$ are constants from Theorem 2.2, Remark 3.4, Proposition 4.5 and Inequality (2.11). The above properties are satisfied for $n = 0$. Let $n \geq 1$, and suppose that the induction assumptions are true for $0 \leq k \leq n-1$. First, we have the following telescopic series (see [21] or [9])

$$(4.6) \quad \mathbf{U}_n - \mathbf{U}(t_n) = \sum_{0 \leq k \leq n-1} \Phi^{(n-k-1)\Delta t} \mathcal{Y}^{\Delta t} \mathbf{U}_k - \Phi^{(n-k-1)\Delta t} \Phi^{\Delta t} \mathbf{U}_k.$$

For $k \leq n-2$, since $\mathcal{Y}^{\Delta t} \mathbf{U}_k = \mathbf{U}_{k+1}$, using the induction assumption (ii), we have

$$|\mathcal{Y}^{\Delta t} \mathbf{U}_k|_{X^{N-3}} \leq M_0,$$

and from Theorem 2.2, we get

$$|\Phi^{\Delta t} \mathbf{U}_k|_{X^{N-3}} \leq e^{C_0(M_0)T} M_0.$$

Therefore, from Proposition 2.4 and up to replacing $K = K \left(M_0 e^{TC_0(M_0)} \right)$ with $\max(K, 1)$, we obtain, for $k \leq n-1$ and $n\Delta t \leq \frac{T}{\delta}$,

$$\left| \Phi^{(n-k-1)\Delta t} \mathcal{Y}^{\Delta t} \mathbf{U}_k - \Phi^{(n-k-1)\Delta t} \Phi^{\Delta t} \mathbf{U}_k \right|_{X^{N-4}} \leq K \left| \mathcal{Y}^{\Delta t} \mathbf{U}_k - \Phi^{\Delta t} \mathbf{U}(t_k) \right|_{X^{N-4}}.$$

Then, using Proposition 4.5 and Inequality (ii), we infer

$$\left| \Phi^{(n-k-1)\Delta t} \mathcal{Y}^{\Delta t} \mathbf{U}_k - \Phi^{(n-k-1)\Delta t} \Phi^{\Delta t} \mathbf{U}_k \right|_{X^{N-4}} \leq \delta C_4 \left(e^{C_0(M_0)T} M_0 \right) K (\Delta t)^2.$$

Therefore, using the telescopic series (4.6), we get

$$\left| \mathbf{U}_n - \mathbf{U}(t_n) \right|_{X^{N-4}} \leq n C_4 \left(e^{C_0(M_0)T} M_0 \right) K \delta (\Delta t)^2 \leq C_4 \left(e^{C_0(M_0)T} M_0 \right) K T \Delta t.$$

For Estimate (ii), using Remark 3.4 and the induction assumptions (iii) and (ii), we have,

$$\left| \mathbf{U}_n \right|_{X^N} = \left| \mathcal{Y}^{\Delta t} (\mathbf{U}_{n-1}) \right|_{X^N} \leq e^{\delta C_3(M_1)\Delta t} \left| \mathbf{U}_{n-1} \right|_{X^N} \leq e^{C_3(M_1)\delta n \Delta t} \left| \mathbf{U}_0 \right|_{X^N} \leq M_0.$$

We get Estimate (iii) in the same way, using the induction assumptions (iv) and (iii). Finally, for Estimate (iv), using (i), we have

$$\left| \mathbf{U}_n \right|_{X^{N-4}} \leq \left| \mathbf{U}_n - \mathbf{U}(t_n) \right|_{X^{N-4}} + \left| \mathbf{U}(t_n) \right|_{X^{N-4}} \leq \gamma \Delta t + \rho \leq 2\rho.$$

□

5. NUMERICAL EXPERIMENTS

The aim of this section is to numerically verify the Lie method convergence rate in $\mathcal{O}(\Delta t)$ for the Saut-Xu system (1.1) and to illustrate some physical phenomena.

In other works and particularly on the whole water waves problem (see for example [16], [19], [31] and references therein), several authors use a discrete Fourier transform even for the transport part. They observe spurious oscillations in the wave profile that lead to instabilities. These errors seem to appear when they evaluate the nonlinear part via Fourier transform because additional terms appear in the approximation, this is the aliasing phenomenon. To fix this problem, they apply at every time step a low-pass filter. The main interest of our scheme is that we do not need one because we use a finite difference method to approximate the nonlinear part.

For the dispersive equation (3.2), we use the forward Euler discretization in time and for the spatial discretization we consider the Fast Fourier Transform (FFT) implemented in Matlab. In this scheme, the interval $[0, 1]$ is discretized by N equidistant points, with spacing $\Delta x = 1/N$. The spatial grid points are then given by $x_j = j/N$, $j = 0, \dots, N$. Therefore, if $u_j(t)$ denotes the approximate solution to $u(t, x_j)$, the discrete Fourier transform of the sequence $\{u_j\}_{j=0}^{N-1}$ is defined by

$$\hat{u}(k) = \mathcal{F}_k^d(u_j) = \sum_{j=0}^{N-1} u_j e^{-2i\pi jk/N},$$

for $k = 0, \dots, N-1$, and the inverse discrete Fourier transform is given by

$$u_j = \mathcal{F}_j^{-d}(\hat{u}_k) = \frac{1}{N} \sum_{k=0}^{N-1} \hat{u}_k e^{2i\pi kx_j},$$

for $j = 0, \dots, N-1$. Here \mathcal{F}^d denotes the discrete Fourier transform and \mathcal{F}^{-d} its inverse. Then, in what follows the numerical scheme to solve (3.2) is given by

$$(5.1) \quad \begin{pmatrix} \zeta_j^{n+1} \\ v_j^{n+1} \end{pmatrix} = \begin{pmatrix} \zeta_j^n \\ v_j^n \end{pmatrix} - \Delta t \begin{pmatrix} F_j^n + S_j^n \\ G_j^n \end{pmatrix}$$

where $S_j^n = \beta\sqrt{\mu}\mathcal{F}_j^{-d}(ik\mathcal{F}_k^d(B_\mu v_j^n))$ and $F_j^n = I_1 + I_2$ with

$$\begin{aligned} I_1 &= \mathcal{F}_j^{-d}\left(i \tanh(\sqrt{\mu}k)\left(-1 + \frac{\varepsilon\sqrt{\mu}}{2}\mathcal{F}_k^d(v_j^n\mathcal{F}_j^{-d}(k \tanh(\sqrt{\mu}k)\hat{\zeta}_k^n))\right.\right. \\ &\quad \left.\left.+ \varepsilon\sqrt{\mu}\mathcal{F}_k^d(\zeta_j^n\mathcal{F}_j^{-d}(k \tanh(\sqrt{\mu}k)\hat{v}_k^n))\right)\right) \\ I_2 &= \zeta_j^n\mathcal{F}_j^{-d}(ik\hat{v}_k^n) + \frac{1}{2}\mathcal{F}^{-d}\left(ik\hat{\zeta}_k^n\right)\mathcal{F}_j^{-d}\left(\tanh(\sqrt{\mu}k)^2\hat{v}_k^n\right). \end{aligned}$$

To approximate the equation (3.1), we use the following finite difference scheme

$$(5.2) \quad \begin{pmatrix} \zeta_j^{n+1} \\ v_j^{n+1} \end{pmatrix} = \begin{pmatrix} \zeta_j^n \\ v_j^n \end{pmatrix} - \Delta t \frac{\varepsilon\sqrt{\mu}}{2} \begin{pmatrix} G_1^n \\ 3G_2^n \end{pmatrix}$$

where

$$G_1 = w_j^n \frac{\zeta_{j+1}^n - \zeta_{j-1}^n}{2\Delta x} - \frac{\Delta t}{2\Delta x^2} (w_j^n)^2 (\zeta_{j-1}^n - 2\zeta_j^n + \zeta_{j+1}^n)$$

with $w_j^n = -\mathcal{F}_j^{-d}(\tanh(\sqrt{\mu}k)^2\hat{v}_k^n) + v_j^n$ and

$$G_2^m = \frac{(v_{j+1}^n)^2 - (v_{j-1}^n)^2}{2\Delta x} - \frac{\Delta t}{2\Delta x^2} \left(v_{j+1/2}^n ((v_{j+1}^n)^2 - (v_j^n)^2) - v_{j-1/2}^n ((v_j^n)^2 - (v_{j-1}^n)^2) \right)$$

with $v_{j\pm 1/2}^n = \frac{v_j^n + v_{j\pm 1}^n}{2}$.

We remarked that for our numerical simulations, it is not necessary to decompose the term $v\partial_x\zeta$ (see Remark 3.1) to get the numerical convergence. Indeed, it seems that since the time step is chosen very small, we obtain a solution of the dispersive equation for each iteration. In this case, we do not need to evaluate the term $\partial_x\zeta\mathcal{H}_\mu^2 v$.

To ensure the validity of our numerical simulations, we have to be careful of the numerical instability, that why the time and the space steps are chosen in a way that the following CFL condition is satisfied:

$$(5.3) \quad |v| \frac{\Delta t}{\Delta x} < 1.$$

5.1. Example 1: Convergence curve. In this example, we consider the following initial data:

$$\zeta_0(x) = \operatorname{sech}\left(\frac{\sqrt{3}}{2}x\right), \quad v_0 = \zeta_0.$$

with two different bathymetries: a bump ($b(x) = \cos(x)$) and a ripple bottom

$$b(x) = \begin{cases} 0.5 - \frac{1}{18}(x-8)^2 & \text{if } 5 \leq x \leq 11 \\ 0 & \text{otherwise} \end{cases}.$$

Note that in order to avoid numerical reflections due to the boundaries and justify of the use of the Fast Fourier Transform, we decide to take rapidly decreasing initial data. Figures 1 and 2 display the evolution for different times of the free surface ζ for these two test cases. We decided to take $\varepsilon = 0.1, \mu = 1, \beta = \frac{1}{2}, N = 2^8, \Delta x = 2L/N, T = 10$ where N is the mesh modes number, $L = 30$ the length of the domain and T the final time. Note that the time step Δt is chosen iteratively in a way that the CFL condition (5.3) is satisfied.




FIGURE 1. Upper: Evolution of the free surface for different times $t=2.5, 5, 7.5$ and 10 . Lower: bottom topography and initial condition.




FIGURE 2. Upper: Evolution of the free surface for different times $t=2.5, 5, 7.5$ and 10 . Lower: bottom topography and initial condition.

Figure 3 displays the convergence curve for this example. We plot the logarithm of the error (in norm $H^1 \times L^2$) in function of the logarithm of the time step Δt . The convergence numerical order is then given by the slope of this curve. For reference, a small line (the dashed line) of slope one is added in this figure. We see that the numerical rate of convergence is greater than 1.

5.2. Example 2: Non smooth topographies. In this example we study the evolution of water waves over a rough bottom. This problem is still a mathematical issue. Many models derived from the Euler equations suppose that the bathymetry is smooth. Even worse, a non smooth bathymetry introduces singular terms in these models. This issue is particularly easy to see for shallow waters models. To handle this, Hamilton ([20]) and Nachbin ([30]) used a conformal mapping to derive long waves models. Notice also the work of Cathala ([10]) who derived alternatives Saint-Venant equations and Boussinesq systems with non smooth topographies which do not involve any singular terms. We notice that the Saut-Xu equations (1.1) can handle a non smooth topography (see Theorem (2.2)) and our numerical scheme too (see Theorem (4.6)).




FIGURE 3. Convergence curve (for the $H^1 \times L^2$ -norm) for the Lie method for two bottoms : bump (black line) and ripple bottom (blue line) for $T = 10$.




FIGURE 4. Evolution of the free surface (blue lines) for different times $t=0,3,6,9$ and 12 over a rough bottom (black line).

In the following, we give an example with a non smooth bathymetry. We consider the following initial conditions and bathymetry

$$\zeta_0(x) = v_0(x) = e^{-x^2} \text{ and } b(x) = \frac{\beta}{4}(1 + \tanh(100(x - 2)))(1 - \tanh(100(x - 8))).$$

We decided to take $\mu = 1$, $\beta = 0.5$, $\varepsilon = 0.1$, $N = 2^8$, $\Delta x = 2L/N$ where N is the mesh modes number, $L = 20$ the length of the domain. The time step Δt is chosen iteratively in a way that the CFL condition (5.3) is satisfied. Figure 4 displays the evolution of the surface ζ for different times over the bottom.

5.3. Example 3: Boussinesq regime. In Section 3, we crucially use the fact that μ is bounded from below. In this example, we test our scheme for small values of μ (also called the shallow water regime). We show that our scheme is still valid even if we do not have a proof of the convergence of our scheme in this regime. In the shallow water regime, there is a huge literature for asymptotic models (see for instance [24]). Among all these asymptotic models, we have the KdV equation. It is a model obtained under the Boussinesq regime, i.e. when $\varepsilon = \mu$, $\beta = 0$ and μ small. In the following, we formally derive a KdV equation from the Saut-Xu equations and we give numerical simulations in this setting.

We recall that, without the assumption $\nu = \frac{1}{\mu}$, the Saut-Xu equations are given by the system (2.3). Notice also that

$$(5.4) \quad \mathcal{H}_\mu = -\sqrt{\mu}\partial_x - \frac{1}{3}\mu^{\frac{3}{2}}\partial_x^3 + \mathcal{O}(\mu^2).$$

Then if we assume that $\mu = \varepsilon$, $\nu = 1$ (since $\nu \sim 1$ if μ is small) and we drop all the terms of order $\mathcal{O}(\mu^2)$ in System (2.3), we obtain the following equations

$$(5.5) \quad \begin{cases} \partial_t \zeta + \partial_x v + \mu v \partial_x \zeta - \frac{1}{2}\mu^{\frac{3}{2}}v \partial_x \zeta + \frac{1}{3}\mu \partial_x^3 v + \mu \zeta \partial_x v = 0, \\ \partial_t v + \partial_x \zeta + \mu v \partial_x v + \mu^{\frac{3}{2}}\frac{1}{2}v \partial_x v = 0. \end{cases}$$

Formally, the solutions of this system are close to the solutions of (2.3) with an accuracy of order $\mathcal{O}(\mu^2)$. Notice that this system is not a standard Boussinesq system (in the sense of [4] or [24]) because of our nonlinear change of variables (2.2). Using the approach developed in [35], [5], [2] (see also Part 7.1.1 in [24]) we can check that, formally, the following KdV equation is an asymptotic model of the system (5.5)

$$(5.6) \quad \partial_t f + \frac{3}{2}f \partial_x f + \frac{1}{6}\partial_x^3 f = 0.$$

This means that if we solve (5.5) with the initial data (f_0, f_0) and (5.6) with the initial datum f_0 , the solution $(\zeta, v)(t, x)$ of (5.5) is close to $(f, f)(\mu t, x - t)$. Furthermore, if we take $f_0(x) = \alpha \operatorname{sech}^2\left(\sqrt{\frac{3}{4}}\alpha x\right)$, the solution f of the KdV equation with this initial datum is the soliton $f(t, x) = f_0(x - ct)$ with $c = \frac{\alpha}{2}$. Hence, in this case, the solution of (5.5) and (2.3) are close to a soliton.

In the following we check that the solution to (1.1) is indeed close to the KdV solution when μ is small. We simulate one soliton. We took $v_0(x) = \zeta_0(x) = \operatorname{sech}^2\left(\frac{\sqrt{3}}{2}x\right)$, $\varepsilon = \mu = 0.01$, $\alpha = 1$ and the final time is $T = 10$. We decided to take $N = 2^9$, $\Delta x = 2L/N$ where N is the mesh modes number, $L = 30$ the length of the domain. The time step Δt is chosen iteratively in a way that the CFL condition (5.3) is satisfied. Figure 5 represents the evolution of this soliton at different times. Hence, our scheme is still valid when μ is small.

In deep water (μ not small), the KdV approximation ceases to be a good approximation. In order to get some insight on the range of validity of the KdV approximation, we compare in Figure 6 the solution of (1.1) to the exact soliton after a time $T = 10$ for various values of μ . We took the same numerical parameters that before. We notice that even for $\varepsilon = \mu = 0.1$ and a final time $T = \frac{1}{\mu}$, the KdV approximation remains a good approximation of the Saut-Xu system.




FIGURE 5. Evolution of the soliton at different times $t = 0, 3, 6, 9$ ($\varepsilon = 0.01$).




FIGURE 6. Difference after a time $T = 10$ between a real soliton and a soliton generated by our scheme with the same initial data and for different values of $\varepsilon = \mu$. Abscissa : value of ε ; Ordinate : quotient of the difference after a final time $T = 10$ by the maximum of the soliton.

5.4. Example 4: Rapidly varying topographies. In this example we study the evolution of water waves over a rapidly varying periodic bottom. We assume that $\mu = 1$. This problem is linked to the Bragg reflection phenomenon (see for instance [29], [25], [19]). We take

$$(5.7) \quad \zeta_0 = v_0 = \operatorname{sech}^2\left(\frac{\sqrt{3}}{2}x\right) \text{ and } b(x) = \cos(\alpha x).$$

We decided to take $N = 2^9$, $\Delta x = 2L/N$ where N is the mesh modes number, $L = 30$ the length of the domain. The time step Δt is chosen iteratively in a way that the CFL condition (5.3) is satisfied. Figure 7 compares the evolution of water waves when we take $\alpha = 10$ (blue line) and when we take $b(x) = 0$ (blue line). Figure 8 displays the difference between the case of a flat bottom and the case of a bottom of the form $b(x) = \cos(\alpha x)$ for different values of α . We observe an homogenization effect when α is large. It seems that when α goes to infinity, a solution of the Saut-Xu equations converges




FIGURE 7. Comparison between the evolution of a water wave (blue line) over a bottom of the form $b(x) = \cos(10x)$ (dashed line) and the evolution of a water wave over a flat bottom (black line). $\varepsilon = 0.05$, $\beta = 0.5$.




FIGURE 8. Difference between a water wave over a rapidly varying topography $b(x) = \cos(\alpha x)$ and a water wave over a flat bottom. Abscissa : value of α ; Ordinate : quotient of the difference after a final time $T = 10$ by the maximum of ζ_0 .

to a solution of the Saut-Xu equations with a flat bottom (corresponding to the mean of b). Notice that this result is different from what we could see in the literature (for instance [13] or [15]), since we take a bottom of the form $b(x) = \cos(\alpha x)$ and not of the form $b(x) = \frac{1}{\alpha} \cos(\alpha x)$. Our numerical simulations suggest therefore a homogenization effect for large amplitude bottom variations that has not been investigated so far.

APPENDIX A

In this part, we give some estimate for the operator \mathcal{H}_μ and some standard product and commutator estimates. For the estimates for \mathcal{H}_μ , we refer to part III in [34]. For the other estimates we refer to [1] and [23]. We recall that \mathcal{H}_μ is defined by

$$\mathcal{H}_\mu = -\frac{\tanh(\sqrt{\mu}D)}{D}\partial_x.$$

First, we show that \mathcal{H}_μ is a zero-order operator.

Proposition A.1. *Let $s \geq 0$ and μ satisfying Condition (1.3). Then,*

$$|\mathcal{H}_\mu u|_{H^s} \leq C(\mu_{\max}) |u|_{H^s}.$$

Furthermore, for all $s \geq r \geq 0$,

$$|(\mathcal{H}_\mu^2 + 1)u|_{H^s} \leq C_r \left(\frac{1}{\mu_{\min}}\right) |u|_{H^r}.$$

Then, we give a commutator estimate for \mathcal{H}_μ .

Proposition A.2. *Let $s \geq 0$, $t_0 > \frac{1}{2}$, $r \geq 0$, and μ satisfying Condition (1.3). Then,*

$$|[\mathcal{H}_\mu, a]u|_2 \leq C |a|_{H^{t_0}} |f|_2,$$

$$\left| |\xi|^s [\widehat{\mathcal{H}_\mu, a}] u \right|_2 \leq C \left(\frac{1}{\mu_{\min}}\right) |a|_{H^{r+s}} \left| \frac{(1+|\xi|)^{t_0}}{|\xi|^r} \widehat{u} \right|_2,$$

and

$$\left| |\xi|^s [\widehat{\mathcal{H}_\mu, a}] u \right|_2 \leq C \left(\frac{1}{\mu_{\min}}\right) |a|_{H^{r+s+t_0}} \left| \frac{1}{|\xi|^r} \widehat{u} \right|_2.$$

We recall the well-known Coifman-Meyer estimate. We recall also that Λ is the Fourier multiplier $\Lambda = \sqrt{1+D^2}$.

Proposition A.3. *Let $s > \frac{3}{2}$, $u \in H^s(\mathbb{R})$ and $v \in H^{s-1}(\mathbb{R})$. Then we have the following commutator estimate*

$$|[\Lambda^s, u]v|_2 \leq C |u|_{H^s} |v|_{H^{s-1}}.$$

We recall also the following product estimate.

Proposition A.4. *Let s_1, s_2, s such that $s_1 + s_2 \geq 0$, $s \leq \min(s_1, s_2)$ and $s < s_1 + s_2 - \frac{1}{2}$. Let $u \in H^{s_1}(\mathbb{R})$ and $v \in H^{s_2}(\mathbb{R})$. Then,*

$$|uv|_{H^s} \leq C |u|_{H^{s_1}} |v|_{H^{s_2}}.$$

ACKNOWLEDGMENTS

The second author has been partially funded by the ANR project Dyficolti ANR-13-BS01-0003.

REFERENCES

- [1] S. Alinhac and P. Gérard. *Opérateurs pseudo-différentiels et théorème de Nash-Moser*. Savoirs Actuels. [Current Scholarship]. InterEditions, Paris; Éditions du Centre National de la Recherche Scientifique (CNRS), Meudon, 1991.
- [2] B. Alvarez-Samaniego and D. Lannes. Large time existence for 3D water-waves and asymptotics. *Invent. Math.*, 171(3):485–541, 2008.
- [3] D. M. Ambrose, J. L. Bona, and D. P. Nicholls. On ill-posedness of truncated series models for water waves. *Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.*, 470(2166):20130849, 16, 2014.
- [4] J. L. Bona, M. Chen, and J.-C. Saut. Boussinesq equations and other systems for small-amplitude long waves in nonlinear dispersive media. I. Derivation and linear theory. *J. Nonlinear Sci.*, 12(4):283–318, 2002.
- [5] J. L. Bona, T. Colin, and D. Lannes. Long wave approximations for water waves. *Arch. Ration. Mech. Anal.*, 178(3):373–410, 2005.
- [6] P. Bonneton, E. Barthélemy, F. Chazel, R. Cienfuegos, D. Lannes, F. Marche, and M. Tissier. Recent advances in serre-green naghdi modelling for wave transformation, breaking and runup processes. *European Journal of Mechanics - B Fluids*, 30(6):589 – 597, 2011. Special Issue: Nearshore Hydrodynamics.
- [7] P. Bonneton, F. Chazel, D. Lannes, F. Marche, and M. Tissier. A splitting approach for the fully nonlinear and weakly dispersive green-naghdi model. *Journal of Computational Physics*, 230(4):1479 – 1498, 2011.
- [8] P. Bonneton and D. Lannes. Derivation of asymptotic two-dimensional time-dependent equations for surface water wave propagation. *Phys. Fluids*, 21, 2009.
- [9] R. Carles. On Fourier time-splitting methods for nonlinear Schrödinger equations in the semiclassical limit. *SIAM J. Numer. Anal.*, 51(6):3232–3258, 2013.
- [10] M. Cathala. Asymptotic shallow water models with non smooth topographies. *Monatshefte für Mathematik*, pages 1–29, 2014.
- [11] P. Chartier, L. Le treust, and F. Méhats. Uniformly accurate time-splitting methods for the semiclassical schrödinger equation part 2 : numerical analysis of the linear case. *arXiv : 1601.04825v1*, 2016.
- [12] W. Choi. Nonlinear evolution equations for two-dimensional surface waves in a fluid of finite depth. *J. Fluid Mech.*, 295:381–394, 1995.
- [13] L. Chupin. Roughness effect on Neumann boundary condition. *Asymptot. Anal.*, 78(1-2):85–121, 2012.
- [14] W. Craig, P. Guyenne, D. P. Nicholls, and C. Sulem. Hamiltonian long-wave expansions for water waves over a rough bottom. *Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.*, 461:839–873, 2005.
- [15] W. Craig, D. Lannes, and C. Sulem. Water waves over a rough bottom in the shallow water regime. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 29(2):233–259, 2012.
- [16] W. Craig and C. Sulem. Numerical simulation of gravity waves. *J. Comput. Phys.*, 108(1):73–83, 1993.
- [17] W. Craig, C. Sulem, and P.-L. Sulem. Nonlinear modulation of gravity waves: a rigorous approach. *Nonlinearity*, 5(2):497–522, 1992.
- [18] P. Guyenne and D. P. Nicholls. Numerical simulation of solitary waves on plane slopes. *Math. Comput. Simul.*, 69:269–281, 2005.
- [19] P. Guyenne and D. P. Nicholls. A high-order spectral method for nonlinear water waves over moving bottom topography. *SIAM J. Sci. Computer*, no. 1:81–101, 2007-08.
- [20] James Hamilton. Differential equations for long-period gravity waves on fluid of rapidly varying depth. *J. Fluid Mech.*, 83(2):289–310, 1977.
- [21] H. Holden, C. Lubich, and N. H. Risebro. Operator splitting for partial differential equations with Burgers nonlinearity. *Math. Comput.*, 82(281), 2013.
- [22] T. Iguchi. A shallow water approximation for water waves. *J. Math. Kyoto Univ.*, 49(1):13–55, 2009.
- [23] D. Lannes. Sharp estimates for pseudo-differential operators with symbols of limited smoothness and commutators. *J. Funct. Anal.*, 232(2):495–539, 2006.
- [24] D. Lannes. *The water waves problem*, volume 188 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2013. Mathematical analysis and asymptotics.
- [25] Y. Liu and D.K.P. Yue. On generalized Bragg scattering of surface waves by bottom ripples. *Journal of Fluid Mechanics*, 356:297–326, 1998.
- [26] C. Lubich. On splitting methods for Schrödinger-Poisson and cubic nonlinear Schrödinger equations. *Math. Comp.*, 77(264):2141–2153, 2008.
- [27] Y. Matsuno. Nonlinear evolutions of surface gravity waves on fluid of finite depth. *Phys. Rev. Lett.*, 69(4):609–611, 1992.
- [28] Y. Matsuno. Nonlinear evolution of surface gravity waves over an uneven bottom. *J. Fluid Mech.*, 249:121–133, 1993.

- [29] C. C. Mei. Resonant reflection of surface water waves by periodic sandbars. *Journal of Fluid Mechanics*, 152:315–335, 3 1985.
- [30] André Nachbin. A terrain-following Boussinesq system. *SIAM J. Appl. Math.*, 63(3):905–922, 2003.
- [31] D. P. Nicholls and F. Reitich. Stability of high-order perturbative methods for the computation of Dirichlet–Neumann operators. *Journal of Computational Physics*, 170(1):276–298, 2001.
- [32] D. L. Ropp and J. N. Shadid. Stability of operator splitting methods for systems with indefinite operators: advection-diffusion-reaction systems. *J. Comput. Phys.*, 228(9):3508–3516, 2009.
- [33] A. Sacchetti. Spectral splitting method for nonlinear Schrödinger equations with singular potential. *J. Comput. Phys.*, 227(2):1483–1499, 2007.
- [34] J. C. Saut and L. Xu. Well-posedness on large time for a modified full dispersion system of surface waves. *J. Math. Phys.*, 53(11):115606, 23, 2012.
- [35] G. Schneider and C. E. Wayne. Corrigendum: The long-wave limit for the water wave problem I. The case of zero surface tension [mr1780702]. *Comm. Pure Appl. Math.*, 65(5):587–591, 2012.
- [36] R. A. Smith. An operator expansion formalism for nonlinear surface waves over variable depth. *J. Fluid Mech.*, 363:333–347, 1998.
- [37] T. R. Taha and M. J. Ablowitz. Analytical and numerical aspects of certain nonlinear evolution equations. II. Numerical, nonlinear Schrödinger equation. *J. Comput. Phys.*, 55(2):203–230, 1984.
- [38] V.E Zakharov. Stability of periodic waves of finite amplitude on the surface of a deep fluid. *J. Applied Mech. Tech. Phys.*, 9:190–194, 1968.

UNIV. BORDEAUX, IMB, INRIA MEMPHIS, UMR 5251, F-33400 TALENCE, FRANCE.

E-mail address: afaf.bouharguane@math.u-bordeaux1.fr

UNIV. BORDEAUX, IMB, UMR 5251, F-33400 TALENCE, FRANCE.

E-mail address: benjamin.melinand@math.u-bordeaux1.fr