Nitrogen isotope variations in the solar system

Evelyn Füri, Bernard Marty

► To cite this version:

Evelyn Füri, Bernard Marty. Nitrogen isotope variations in the solar system. Nature Geoscience, 2015, 10.1038/NGEO2451 . hal-01345978

HAL Id: hal-01345978

<https://hal.science/hal-01345978>

Submitted on 20 Jul 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Nitrogen isotope variations in the solar system

Evelyn Füri¹ and Bernard Marty^{1,*}

¹ Centre de Recherches Pétrographiques et Géochimiques, CNRS-Université de Lorraine, 15 rue Notre Dame des Pauvres, BP20, 54501 Vandoeuvre-lès-Nancy, France

* Corresponding author e-mail address: bmarty@crpg.cnrs-nancy.fr

The relative proportion of the two isotopes of nitrogen (^{14}N and ^{15}N) shows dramatic variations among the different solar system objects and reservoirs. NASA's Genesis mission, which provided the first direct sample of the solar wind, confirmed that the Sun, and by inference the protosolar nebula, is highly depleted in the heavy ^{15}N isotope. The inner planets, asteroids, and comets are enriched in ^{15}N by tens to hundreds of percent, with organic matter in primitive meteorites recording the most extreme $^{15}\text{N}/^{14}\text{N}$ ratios. Several lines of evidence suggest that these ^{15}N enrichments were not inherited from presolar material but are, instead, the result of N isotope fractionation processes that occurred early in solar system history. Together, these observations indicate that N isotopes are a powerful tool to investigate early material processing and large-scale disk dynamics as well as planetary formation processes. In addition, N isotopes are the tracer of choice to investigate the origin and evolution of planetary atmospheres.

The solar system formed when a fraction of a dense molecular cloud collapsed and a central star, the proto-Sun, started burning its nuclear fuel¹. The surrounding disk made of gas and dust, the protosolar nebula (PSN), was thoroughly stirred and homogenized due to large-scale heating and mixing driven by loss of the angular momentum, the energy delivered by the proto-Sun, and magneto-rotational turbulence. The efficiency of these processes is evident in primitive ("carbonaceous") meteorites, which show a remarkable homogeneity in the isotopic compositions of their constituents down to the part per million level for most elements of the

periodic table². Relics of the initial heterogeneous mixture of stellar debris can only be found in nano-to-micron-sized presolar grains that were thermally resistant enough to survive high enthalpy processing³. However, the light elements hydrogen, carbon, nitrogen, and oxygen, show significant, sometimes extreme, isotope variations among solar system objects and reservoirs, from a few percents for C and O, to tens or even hundreds of percents for H and N (ref. 4). These light elements, by far the most abundant ones in the PSN, share the property to have been predominantly in the gaseous state (H₂, CO, N₂, etc., and their ionized derivatives) in the presolar cloud and in the disk. Consequently, they were prone to efficient isotope exchange and interactions with stellar photons and cosmic rays, either in the interstellar medium (ISM)⁵, or in the presolar cloud or the PSN (e.g., refs. 4,6). Thus, these isotope compositions convey a unique record of solar system forming processes.

The largest isotope variations are observed for hydrogen and nitrogen. The deuterium/hydrogen (D/H) ratio varies by a factor of ~ 35 , from the PSN value of $21 \pm 0.5 \times 10^{-6}$ (ref. 7) to D-rich "hotspots" in meteorites with values up to 720×10^{-6} (ref. 8). Inner solar system objects ($\sim 150 \times 10^{-6}$; ref. 7) and comets ($150\text{--}500 \times 10^{-6}$; refs. 9–11) show intermediate values, and possibly define an increase of the D/H ratio with heliocentric distance. A consistent scheme emerges in which nebular H₂ poor in deuterium exchanged isotopically with H₂O at low temperature, resulting in a preferential D-enrichment of the latter. Deuterium-rich water then froze out onto icy grains and exchanged isotopically with organics and silicates as a result of turbulent transport and aqueous alteration on forming planetesimals¹². Although this scenario is not without weaknesses and is still a matter of debate, the D/H isotopic tracer offers the possibility to investigate the relationships between the different solar system reservoirs. In particular, it is central in the debate on the origin of water (cometary or asteroidal) in the inner solar system including the oceans¹⁰.

The relative proportion of the two stable isotopes of nitrogen, ^{14}N and ^{15}N , also shows outstanding variability in the solar system. For expressing the N isotope composition, geochemists and cosmochemists use the stable isotope delta notation:

$$\delta^{15}\text{N} = [({}^{15}\text{N}/{}^{14}\text{N})_{\text{sample}}/({}^{15}\text{N}/{}^{14}\text{N})_{\text{standard}} - 1] \times 1,000$$

where $\delta^{15}\text{N}$ expresses the deviation of the sample ratio relative to a standard in parts per mil (‰). The nitrogen standard is the isotope composition of atmospheric N_2 ($^{15}\text{N}/^{14}\text{N} = 3.676 \times 10^{-3}$; ref. 13). On Earth, most variations are of the order of a few to tens of permil (e.g., ref. 14). Because the range of extraterrestrial N isotope variations can be much larger than the permil level, cosmochemists use instead the absolute value of the $^{15}\text{N}/^{14}\text{N}$ ratio, following the stable isotope convention that the rare, heavy isotope is the numerator. To complicate matters further, astronomers and astrophysicists use instead the $^{14}\text{N}/^{15}\text{N}$ notation (272 for air) (despite using the D/H notation for hydrogen as do cosmochemists). Both notations are given here for the sake of better understanding by these communities.

On Earth, the N isotope composition varies by no more than 2 %, but variations can reach 500 % on a solar system scale (Figs. 1 and 2). Until recently, the causes of this variability were not understood, for two main reasons. Firstly, the solar system initial $^{14}\text{N}/^{15}\text{N}$ ratio was unknown. Secondly, nitrogen isotopes are more difficult to quantify than hydrogen isotopes because they are generally less abundant in cosmochemical material, and because they are difficult to measure at distance by spectroscopic methods. The analysis of solar wind ions returned to Earth by the Genesis mission – together with advances in high-spatial-resolution, high-sensitivity isotope analysis in the laboratory as well as in high-resolution UV spectroscopy (Box 1) – have permitted major leaps of understanding in the cosmochemistry of this element. Here, we review recent advances in the domain that are particularly relevant in the context of the measurements currently carried out on Comet 67P/Churyumov-Gerasimenko by the Rosetta spacecraft instruments.

72 Nucleosynthesis of N isotopes and galactic evolution

73 The production paths and rates of nitrogen isotopes are not fully understood¹⁵⁻¹⁷. ^{14}N is
74 produced during the cold CNO cycle in low to intermediate mass stars ($M_{\text{solar}} < M < 7 M_{\text{solar}}$) and
75 is released to the interstellar medium by dredge-up events during asymptotic giant branch
76 (AGB) phases that terminate the lives of most stars ("secondary" production, i.e., not produced
77 directly from H and He nuclei). This isotope is also produced in the so-called hot CNO cycle that
78 takes place in massive stars turning into AGBs during dredge-up from the carbon layer
79 ("primary" production from 3 alpha particles and two protons). ^{15}N is destroyed in the cold
80 CNO cycle, requiring other mechanisms such as ^{15}N production during the hot CNO cycle in
81 novae and, possibly, neutrino spallation on ^{16}O in type II supernovae, both productions being
82 "secondary". Consequently, a nucleosynthetic isotope composition cannot be predicted from
83 theory, and estimates for the N isotope production in the galaxy rely on (limited) observations.

84 In molecular clouds, the N isotope composition is mainly measured in CN, HCN, HNC, and,
85 more recently, in NHD^+ molecules¹⁷⁻²². For molecular N_2 , the signal-to-noise ratio is too low for
86 a spectroscopic measurement. The data seem to define an isotope gradient with a decrease of
87 ^{15}N relative to ^{14}N with increasing distance from the galaxy center (Fig. 1; refs. 17,18). Because
88 stars in the galaxy center tend to be more metal-rich (metallicity being here the proportion of
89 elements heavier than helium), secondary production of ^{15}N that presumably takes place in
90 novae and supernovae should decrease more rapidly with galactocentric distance than the
91 primary component of ^{14}N production, in qualitative agreement with the observation (Fig. 1).
92 Strikingly, the solar system, represented by the PSN value (see next subsection), does not fit
93 the correlation. The observed offset could result from a localized N isotope evolution since the
94 solar system isolated nitrogen from the local ISM 4.56 Ga ago. However, a $^{14}\text{N}/^{15}\text{N}$ value of $\sim$
95 ≥ 400 has been proposed for the local ISM²³ where the Sun was born, consistent with the solar
96 ratio and implying little N isotope evolution with time. Alternatively, the apparent anomaly
97 may be explained by the uncertainty in the data defining the correlation depicted in Fig. 1. CN

or HCN, the molecules for which the nitrogen isotopic compositions are measured in dense cores, could have been isotopically fractionated with respect to the source composition, and may not be representative of the local ISM values. Indeed, measurements of the N isotope compositions in three starless dense cores of the local ISM, either in $\text{NH}_3\text{-NH}^+$ (refs. 20–22) or in HCN, suggest that the nitrile formation path (leading to HCN and CNH) drastically enriches ^{15}N compared to the amine path (NH compounds) (Fig. 2; ref. 19).

Remarkably, the order of magnitude variation of the nitrogen isotope composition in the solar system, defined by the most ^{15}N -rich portion of the Isheyevo meteorite²⁴ and the protosolar nebula composition (Figs. 1 and 2; see subsection below), is comparable to the range of values measured in galactic molecular clouds all over the galaxy (Fig. 1). A straightforward interpretation would be that the solar system contains isotopically heterogeneous, nucleosynthetic nitrogen components inherited from ISM. However, other isotope systems do not show correlated isotope variations. For example, nitrogen's "brother" element carbon, whose two stable isotopes ^{12}C and ^{13}C are produced in different nucleosynthetic pathways as well and also show a diversity of isotopic compositions among molecular clouds, displays a near-constant isotopic composition in the solar system within $\sim 10\%$ (ref. 25). The isotope variations of nitrogen in the solar system are therefore unlikely to represent remnants of nucleosynthetic heterogeneities, and are instead attributed to isotope fractionation. The nature of the fractionating processes (e.g., interactions with solar/stellar photons vs. low temperature isotope exchange) are still debated (Boxes 2 and 3).

The protosolar nebula

The N isotope analysis of Jupiter's atmosphere by spectroscopy ($^{14}\text{N}/^{15}\text{N} = 526^{+585}_{-169}$, $\delta^{15}\text{N} = -483^{+245}_{-272}\text{‰}$; ref. 26) and by in-situ mass spectrometry ($^{14}\text{N}/^{15}\text{N} = 435 \pm 60$, $\delta^{15}\text{N} = -375 \pm 80\text{‰}$; ref. 27) suggested a ^{15}N -poor solar nitrogen composition. Independently, a $^{14}\text{N}/^{15}\text{N}$ ratio of 428 ± 8 ($\delta^{15}\text{N} = -364 \pm 12\text{‰}$) has been obtained for a rare TiN (osbornite) phase within a

calcium-aluminum-rich inclusion (CAI) (ref. 28). Because osbornite was presumably the first solid N-bearing phase to condense from the PSN, this value was concluded to correspond to the PSN signature at the time of solid condensation.

At present, the isotopic composition of the PSN is best represented by that of the Sun, which concentrates more than 99 % of the mass of the solar system. For the isotope analysis of the Sun, only the solar wind (SW; the corpuscular emission of the Sun which consists of matter from the solar convective zone, ionized in the photosphere and accelerated along open lines of the Sun's magnetic field) is available for sampling. The nitrogen isotope composition of the modern solar wind was recently measured directly thanks to NASA's Genesis mission (Discovery Program)²⁹. The Genesis spacecraft sampled the SW during 27 months at the Lagrangian point L2 and returned SW-irradiated material for laboratory analysis in 2004. Despite a hard landing (the parachute failed to open), the SW-implanted target material could be analyzed by two different extraction techniques for N isotopes (laser ablation-static mass spectrometry³⁰ and ion probe³¹). Both methods gave consistent results that permitted to define the PSN nitrogen isotope composition of $^{14}\text{N}/^{15}\text{N} = 441 \pm 5$ ($\delta^{15}\text{N} = -383 \pm 8 \text{‰}$; ref. 31), after moderate (24 ‰) correction for isotope fractionation in solar processing. Because the modern solar wind has a N isotope composition very close to that of Jupiter and osbornite, this comparison confirms that the Sun did neither synthesize N nor significantly fractionate N isotopes by more than about 3 % from its birth to present-day. The PSN value constitutes the anchor value to which to compare other nitrogen isotope signatures of solar system objects (Fig. 2).

The surface of the Moon

The surface of the Moon, which lacks atmospheric shielding and full magnetic field protection, has accumulated SW ions for hundreds of millions of years. The analysis of SW, either implanted into aluminum foils during the visits by the Apollo astronauts or in the lunar

150 soils that were recovered, was a science priority of the Apollo and Luna programs. One of the
151 most intriguing results of the lunar exploration was the discovery of a $\sim 30\%$ N isotope
152 variation in lunar soils and rocks^{32,33}, one order of magnitude larger than that on Earth.
153 Spallation by cosmic rays that produces ^{15}N from ^{16}O can account for some of the low $^{14}\text{N}/^{15}\text{N}$
154 values but fails to explain the occurrence of light (^{15}N -poor) nitrogen on the Moon. Noble gases
155 in lunar soils are mostly derived from SW³⁴, so that lunar nitrogen was assumed to be solar as
156 well. Thus, prior to the Genesis mission, the N isotope variations, which seemed to relate to the
157 epoch of soil exposure, were attributed to secular changes in the isotopic composition of the
158 SW^{33,35}. This possibility, however, faced serious difficulties: no nucleosynthetic process within
159 the Sun capable of changing the N isotope composition could be identified³⁶, and comparison
160 with carbon (whose isotope composition in lunar samples varies much less than that of
161 nitrogen; ref. 25) and helium (whose isotope composition in lunar soils appears constant over
162 time; ref. 37) implied a negligible secular evolution of the SW. Finally, the development of
163 coupled noble gas-nitrogen isotope analysis – together with the Genesis findings –
164 demonstrated that, contrary to noble gases, an additional, non-SW nitrogen component is
165 present in lunar soils^{38–40}. The conjoint measurement of H and N isotope variations through the
166 outer skin of lunar soil grains with an ion probe (Box 1; ref. 41) revealed that ^{15}N -depleted SW-
167 N ($^{14}\text{N}/^{15}\text{N} > 340$, $\delta^{15}\text{N} < -200\text{‰}$) is mixed to a variable extent with "heavy" (^{15}N -rich) nitrogen.
168 The N isotope signature of the non-solar end-member ($^{14}\text{N}/^{15}\text{N} = 265 \pm 5$; $\delta^{15}\text{N} = 50\text{--}100\text{‰}$) is
169 consistent with delivery of asteroidal, rather than cometary, material to the lunar surface, with
170 a flux comparable to that of interplanetary dust and micrometeorites on Earth, after scaling to
171 different gravitational focusing and surface areas of the two planetary bodies^{39,40}. An
172 alternative interpretation is that the ^{15}N -rich component is supplied by an "Earth wind" flux of
173 terrestrial atmospheric nitrogen⁴². Early in the history of the Earth-Moon system, before the
174 development of the global geomagnetic field, interaction of the SW with Earth's upper
175 atmosphere may have resulted in a significant N^+ escape flux. However, since the Moon is

tidally locked to Earth, the Earth wind contribution is expected to be absent on the lunar farside. Future sampling may permit to distinguish between an asteroidal or terrestrial origin of the ^{15}N enrichment. Hence, the lunar surface constitutes a superb opportunity to investigate the nature and flux of matter and ions delivered to planetary surfaces through time, and will certainly deserve further attention in future lunar missions aimed at investigating the origin of volatile elements (e.g., water) in the inner solar system.

Planets, asteroids and comets

The atmospheres of Jupiter^{26,27} and Saturn⁴³ are as ^{15}N -depleted as the PSN. Thus, they may represent remnants of the PSN gas, either trapped gravitationally during planetary growth or inherited from accretion of icy bodies⁴⁴. The second possibility implies that these icy planetesimals had a solar-like N isotope composition and were not as ^{15}N -rich as are present-day comets (see below).

The inner planets – Earth and Moon⁴⁵, as well as the interior of Mars⁴⁶ and the atmosphere of Venus⁴⁷ – are richer in ^{15}N than solar N by approximately 60 %, and have comparable N isotope compositions on the scale of solar system variations (Fig. 2). In primitive meteorites, nitrogen is mostly hosted in organics, particularly in insoluble organic matter (IOM) and, under reducing conditions, in nitrides. Carbonaceous chondrites, which are rich in volatile elements as their name implies, contain of the order of 1,000 ppm N, with bulk N isotope compositions within 5 % of the terrestrial value⁴⁸⁻⁵⁰. An exception is the CR carbonaceous chondrite clan, whose members are richer in ^{15}N by up to 25 % (e.g., ref. 51), and a few meteorites defining the CB-CH group that present bulk ^{15}N enrichments up to 100-150 % (refs. 52,53). The causes of these enrichments are unclear and could be related to impacts between asteroidal bodies and volatile- and/or organic-rich objects rich in ^{15}N (refs. 51,54). One member of this ^{15}N -rich group, the Isheyevo meteorite, presents so-called " ^{15}N hotspots" (ref. 55), measured on a micron scale by ion probe^{24,56}, characterized by the most extreme ^{15}N enrichments found in the

202 solar system, up to about 450 ‰ ($^{14}\text{N}/^{15}\text{N} \sim 50$; $\delta^{15}\text{N} \sim 4,500$ ‰; ref. 24). ^{15}N -rich hotspots are
203 typically found in IOM of primitive meteorites⁵⁷ and interplanetary dust particles⁵⁸ (which are
204 small [$\leq 50\mu\text{m}$] volatile-rich grains snowing onto Earth's surface, and some of which are
205 probably cometary), as well as in nano-sized globules of the Tagish Lake meteorite⁵⁹. Since IOM
206 shows systematically higher ^{15}N contents than bulk meteorites and inner planets, ^{15}N -rich
207 nitrogen hosted by organic molecules within dust grains must have been mixed with ^{15}N -poor
208 PSN N_2 to match bulk values^{30,31}.

209 Cometary matter was analyzed on Earth thanks to NASA's Stardust mission (Discovery
210 program), which succeeded in collecting grains from a Jupiter family comet (JFC) named
211 81P/Wild2 in 2004, and in returning them to Earth two years later for laboratory analyses^{60,61}.
212 As a result of the high velocity collection, ices were lost during recovery, and only
213 silicate/metal grains survived. Analyses showed that cometary dust is derived from precursors
214 that share mineralogical and isotopic similarities with carbonaceous chondrites, consistent
215 with models advocating large-scale radial mixing in the nascent solar system^{60–62}. $^{15}\text{N}/^{14}\text{N}$
216 ratios are also comparable to the values found in meteorites, and excesses of ^{15}N typical of
217 cometary values (see below) are not observed in bulk analyses⁶³, probably because such
218 excesses are hosted by cometary ice (CN, HCN, NH_3) that was lost during Stardust collection.
219 On a smaller scale, ^{15}N -rich hotspots with a maximum $\delta^{15}\text{N}$ value of 1300 ± 400 ‰ are
220 observed, similar to the highest values found in IOM and IDPs⁶². Several presolar grains have
221 been identified showing much larger C and N isotope variations (from +60 to +964 ‰ for $\delta^{13}\text{C}$,
222 and from -518 to -350 ‰ for $\delta^{15}\text{N}$; ref. 62).

223 Additional cometary N isotope data are from radio and optical spectroscopic observations
224 of CN, HCN, and NH_2 (the latter presumably derived from the photodissociation of NH_3)^{64–68}.
225 Measurements are challenging due to spectroscopic interferences and low abundances of the
226 analyzed molecules, so that data have significant uncertainties compared to laboratory
227 measurements (Box 1). All data collected to date indicate no systematic difference in the

$^{14}\text{N}/^{15}\text{N}$ ratio between these different molecules within uncertainties: The mean $^{14}\text{N}/^{15}\text{N}$ ratios of CN (18 comets), HCN (1 comet), and NH_2 (3 comets) are 148 ± 6 ($\delta^{15}\text{N} = 840 \pm 75 \text{‰}$), 139 ± 26 ($\delta^{15}\text{N} = 650 \pm 400 \text{‰}$), and $\sim 140 \pm 40$ ($\delta^{15}\text{N} = 940 \pm 500 \text{‰}$), respectively (Fig. 2). Furthermore, short- and long-period comets seem to have comparable N isotope compositions. Long-period comets are thought to come from the Oort Cloud (OC comets), whose population is attributed to scattering of icy bodies originally located within the Uranus-Neptune region^{69,70}. Short-period comets are proposed to originate from the Kuiper belt reservoir, which is beyond the present-day orbit of Neptune, and/or from the external edge of the disk. Due to gravitational instabilities, some of these icy objects would then have been scattered and injected into the Jupiter region, becoming isotropic (Jupiter Family comets - JFCs). Thus, the available cometary data is inconsistent with a systematic N isotope variation with heliocentric distance beyond Jupiter. The observation that NH_2 on one hand, and CN-HCN on the other hand, show similar N isotope signatures is at odds with the prediction of significant nitrogen isotope fractionation between amine and nitrile radicals¹⁹, unless isotope exchange subsequently erased this effect. It should be emphasized, however, that our knowledge of nitrogen isotopes in cometary material is limited. Depending on the thermal history of these bodies, it is possible that ^{15}N -poor dinitrogen from the PSN is trapped in ices, and that the bulk N isotopic composition of comets is de facto unknown. The ROSETTA space mission, which has just started to study comet 67P/Churyumov-Gerasimenko, will hopefully provide for the first time an in-situ inventory of the $^{14}\text{N}/^{15}\text{N}$ ratios of different species degassed by the comet.

Conclusions and prospects

The picture that emerges is that there are at least three isotopic reservoirs in the solar system (Fig. 2): i) the PSN, poor in ^{15}N ($^{14}\text{N}/^{15}\text{N} = 441 \pm 5$; $\delta^{15}\text{N} = -387 \pm 8 \text{‰}$), ii) the inner solar system, where planets and bulk meteorites are enriched by a factor of 1.6 relative to the PSN ($^{14}\text{N}/^{15}\text{N} = 272 \pm 15$; $\delta^{15}\text{N} = 0 \pm 50 \text{‰}$), and iii) cometary ices, enriched in ^{15}N by a factor of 3

relative to the PSN ($^{14}\text{N}/^{15}\text{N} = 147 \pm 11$; $\delta^{15}\text{N} = 850 \pm 150$ ‰). Thus, the distribution of N isotopes in the solar system is roughly consistent with an increase of ^{15}N enrichments with radial distance from the Sun, in qualitative agreement with the D/H ratio (Fig. B3). However, several lines of evidence demonstrate that a homogeneous, ^{15}N -poor isotope signature initially characterized the solar system, thus requiring a mechanism for the progressive temporal and spatial isotopic evolution. The recent detection of ^{15}N -enriched nitrogen in a CAI, which last interacted with the nebular gas ≤ 0.15 Ma ago as shown by the ^{26}Al - ^{26}Mg chronometer, suggests that at least some regions of the disk were driven toward isotopically heavy values very early in solar system history, within the first few tens to hundreds thousands of years after solar system formation⁷¹. The nature of the fractionating processes is still a matter of debate and several processes might have played a role (Boxes 2 and 3). While some of the D/H vs. $^{15}\text{N}/^{14}\text{N}$ co-variations are consistent with either kinetic isotope fractionation or ion-molecule reactions, many ^{15}N excesses are not correlated with enrichments in D, and thus require nitrogen-specific isotope fractionation (Fig. B3). Furthermore, the increase of ^{15}N enrichments with radial distance from the Sun could have resulted from different types of processes. The interaction rate between the PSN gas and UV photons could have varied with heliocentric distance, with enhanced ^{15}N fractionation occurring in the outer solar system, where the lower gas density would have allowed deeper photon penetration and/or more efficient freezing out of synthesized molecules than in the denser, hotter inner solar system. It is also possible that several processes were involved: photodissociation by solar photons in the inner solar system, and contribution of ^{15}N -rich ices formed early in the outer solar system by illumination of ice/dust grains from nearby stars. All these possibilities have profound astrophysical implications and can be tested using nitrogen isotopes, from observation and/or dedicated modeling. Irrespective of the origin(s) of the ^{15}N -enrichment, large-scale turbulence and mixing eventually regionally homogenized material available for planetary accretion, resulting in the isotopic similarity seen for the Earth and most chondritic groups. The same requirement

of efficient isotope fractionation and mixing stands for the outer solar system since all cometary N analyzed so far is enriched in ^{15}N by a factor of ~ 3 relative to inner planets (Fig. 2).

A particularly important question that remains to be addressed with nitrogen isotopes concerns the origin and evolution of the terrestrial atmosphere and oceans. The isotopes of hydrogen do not seem to provide an unambiguous tool for distinguishing between a cometary versus asteroidal origin ever since the discovery of an ocean-like D/H ratio in a comet¹⁰. So far, all cometary measurements indicate a 3-fold ^{15}N enrichment in comets compared to terrestrial N, suggesting that nitrogen isotopes represent a more powerful tracer than H isotopes for investigating the origin(s) of planetary atmospheres. Thus, N isotopes are the tracer of choice to investigate the evolution of planetary atmospheres through time upon thermal and non-thermal escape processes.

Figure Legends

Figure 1: Nitrogen isotope variation in molecular clouds from our galaxy as a function of distance from the galactic center. Adapted from refs. 17,18. Isotope compositions are given in both absolute $^{14}\text{N}/^{15}\text{N}$ ratios and $\delta^{15}\text{N}$ values (deviation in permil from the atmospheric $^{15}\text{N}/^{14}\text{N}$ ratio). The observed correlation (dashed line; ref. 17) appears consistent with secondary ^{15}N production in massive stars. The total N isotope range within the solar system, defined by the most ^{15}N -rich "hotspot" in the Isheyevo meteorite²⁴ and the protosolar nebula (PSN) composition³¹, is comparable to that in our galaxy. Error bars indicate 1σ uncertainties¹⁷.

Figure 2: Nitrogen isotope variations in solar system objects and reservoirs. The value of the protosolar nebula (PSN) is defined by the measurement of solar wind nitrogen collected by NASA's Genesis mission^{30,31}. Except for the Sun, osbornite (TiN)²⁸, and the atmospheres of the

giant planets^{26,27,43}, all solar system objects and reservoirs are significantly enriched in ^{15}N compared to the PSN. Values determined for various N species in local and galactic molecular clouds¹⁷⁻²² are shown for comparison. Uncertainties of spectroscopic measurements are shown at the 1σ level.

Figure B1: Nitrogen and hydrogen isotope variations within single lunar grains. a) Silicate grain (breccia 79035) showing $\delta^{15}\text{N}$ values (circles) as low as -250‰ together with very low δD values (blue squares) at depths ≤ 120 nm. The D-poor hydrogen component (i.e, $\delta\text{D} \approx -900$ ‰) represents implanted SW, highly depleted in deuterium due to D-burning in the Sun. Thus, the low $\delta^{15}\text{N}$ values give a first-order estimate of the isotopic signature of SW-N. b) Ilmenite grain (soil 71501) containing non-solar N – associated with D-rich (planetary) hydrogen –, which is enriched in ^{15}N to a similar extent as meteoritic nitrogen. Data from ref. 41.

Figure B2: Processes leading to a progressive ^{15}N -enrichment in the solar system. Schematic diagram of the protoplanetary disk made of gas and dust with the proto-Sun at the centre. ^{15}N -enriched $^{15}\text{NH}_x$ (and C^{15}N) is produced by ion-molecule reactions in the cold regions of the disk^{72,73} or as a result of photodissociation of N_2 by UV light from the proto-Sun or from nearby stars⁷⁴, and is removed from the gas by freezing out and/or trapping in organic matter. (Image of Beta Pictoris; courtesy of NASA).

Figure B3: Hydrogen and nitrogen isotope ratios of solar system objects and isotope fractionation processes. Adapted from refs. 75,76. Alpha H and alpha N are defined as the D/H and $^{15}\text{N}/^{14}\text{N}$ ratios normalized to the protosolar nebula values ($[2.1 \pm 0.5] \times 10^{-5}$ (ref. 7) and $[2.27 \times 0.03] \times 10^{-3}$ (ref. 31), respectively). Data and corresponding uncertainties for Moon^{45,77}; Venus^{47,78}; Mars atmosphere^{79,80}; chondrites, bulk^{6,8,50}; chondrites, IOM^{75,81};

Isheyevo hotspots^{24,56}; Saturn^{43,82}; Jupiter^{27,83}; Titan atmosphere⁸⁴; comets ^{66,67,85,86}. The observed H-N isotope compositions require various isotope fractionation processes.

Box 1: Analysis of nitrogen isotopes

For geochemical applications, nitrogen isotopes have historically been measured using isotope ratio mass spectrometers operated in dynamic pumping mode (e.g., ref. 87). However, since this technique is not suitable for the analysis of low nitrogen abundances, higher-sensitivity noble gas mass spectrometers operated under static vacuum conditions are now the standard method to determine the nitrogen isotopic composition of N-poor and/or small-sized (≤ 5 mg) terrestrial and extraterrestrial samples (e.g., ref. 88,89). In this method, nitrogen – extracted from the samples by various destructive techniques such as crushing, laser ablation, laser-, or furnace-heating, and purified from interfering carbon compounds – is analyzed in the form of N_2 at masses 28 ($^{14}N^{14}N$), 29 ($^{14}N^{15}N$) and 30 ($^{15}N^{15}N$) in mono- or multi-collection mode. Measured peak heights and isotope ratios must be corrected for isobaric mass interferences from CO, N_2H , and hydrocarbons (C_2H_x), which are ubiquitously present in the mass spectrometer system, and the instrumental precision and reproducibility is monitored by repeat analysis of atmospheric N_2 .

Secondary ionization mass spectrometry (SIMS) analysis represents a non-destructive means to determine N isotope variations with a high lateral resolution (i.e., at a scale as small as ≤ 10 μm) or a high depth resolution (i.e., ≤ 10 nm; Fig. B1; ref. 39). While the isotope analysis of nitrogen by SIMS is challenging due to its low ionization efficiency, an intense CN^- signal is formed in the presence of carbon by bombardment of the sample surface with a Cs^+ ion beam, and nitrogen isotope ratios have successfully been determined in the form of $^{12}C^{15}N^- / ^{12}C^{14}N^-$ in various extraterrestrial samples^{24,28,41,56,57,75} as well as in terrestrial diamonds⁹⁰. Furthermore, the nitrogen isotope composition of solar wind collected during NASA's Genesis mission was determined with high precision by SIMS³¹. Notably, since the ionization rate depends on the sample matrix, using a standard made of the same material as the sample and with a known N abundance and isotope composition is critical for an accurate measurement by this technique.

High-resolution radio and optical spectroscopy represents the only analytical means to remotely determine the isotopic composition of cometary (atmospheric) nitrogen. Cometary emissions are produced by absorption of solar light by the various molecules followed by re-emission of lines of specific wavelengths, a process called resonance-fluorescence. Isotope ratios ($^{14}N / ^{15}N$) are determined by comparing the observed molecular spectra with synthetic spectra of the same species. Such measurements require high signal-to-noise spectra together with a high spectral resolution, and have so far only been feasible for nitrogen in the form of CN, HCN, and NH_3 for a limited number of Oort cloud and Jupiter family comets. Notably, the Rosetta Orbiter Spectrometer for Ion and Neutral Analysis (ROSINA) on board the Rosetta spacecraft has been able to determine, for the first time, the abundance of N_2 *in-situ* in the coma of comet 67P/Churyumov-Gerasimenko⁹¹.

Box 2: Possible origin(s) of the ^{15}N -enrichment in solar system material

Two different processes are advocated to explain the ^{15}N enrichment in solar system material. In the first model, ^{15}N enrichment takes place during ion-molecule reactions in the interstellar medium with sufficient density, for example during the collapse of dense pre-stellar cores forming dark molecular clouds, and/or in the cold regions of the disk surrounding the protostar. The possibility of ^{15}N fractionation has been established experimentally⁹². Elemental ^{15}N reacts with $^{14}\text{N}_2\text{H}^+$, which recombines to give ^{15}N -rich $^{14}\text{N}^{15}\text{N}$. He^+ releases additional $^{15}\text{NH}^+$, which is incorporated into NH_3 and NH_2 . These molecules freeze out onto dust grains to yield ammonia ice rich in ^{15}N (ref. 73). However, because nitrogen is continuously cycled between N and N_2 , backward reactions tend to limit ^{15}N enrichments to 30 % or less⁷². To circumvent this problem, it has been proposed (ref. 73) that coupling between N and N_2 is removed when N_2 is deficient in the gas, e.g., frozen out with CO at very low temperature in dense cores, so that the extent of ^{15}N fractionation is no longer limited and can account for the extreme ^{15}N enrichments seen in the solar system⁵. Different reaction paths could also account for the large fractionation between amines (e.g., NH_3) and nitriles (e.g., NH_2)¹⁹. These possibilities are subject to discussion due to uncertainties in relevant chemical rates and branching ratios^{93,94}.

Other types of models advocate photodissociation of N_2 by UV light from the proto-Sun or from nearby stars. In one model, the photodissociation of $^{14}\text{N}^{14}\text{N}$ saturates with respect to that of $^{15}\text{N}^{14}\text{N}$, because the latter is much less abundant, and, therefore, photons with wavelengths able to dissociate $^{14}\text{N}^{14}\text{N}$ get consumed at a greater rate than those photodissociating $^{15}\text{N}^{14}\text{N}$. Dissociated N^+ recombines with surrounding atoms and ions (e.g., H and C) to form amines and nitriles, which can then be removed from the gas by freezing out. This process, which is called self-shielding and is known to occur in dense cores, is postulated to take place in the solar nebula, and not only accounts for ^{15}N fractionation but also for mass-independent oxygen isotope signatures⁹⁵. Another model based on quantum mechanics⁹⁶ is substantiated by photodissociation experiments with UV light from a synchrotron radiation source illuminating a $\text{N}_2\text{-H}_2$ mixture⁷⁴. The produced NH_3 , which is frozen out onto a cold finger and analyzed for N isotopes, shows dramatic ^{15}N excesses up to 12,000 ‰. Although it is not clear if these excesses are the result of a peculiar quantum effect or are due to self-shielding in the experimental reactor, this straightforward and illuminating measurement provides a quantitative case of extreme ^{15}N enrichments in ice formed during interactions between dinitrogen and UV light in a manner reminiscent of the conditions that prevailed during early solar system evolution. In both cases, the key ingredient of the recipe is photons illuminating the PSN gas, indicating that specific environments are required in which the PSN is transparent enough to allow photons to react with gaseous molecules. Possibilities include the surfaces of the disk illuminated by the proto-Sun (Fig. B2) or by other young stars, or the outer regions of the disk illuminated by nearby stars. Once formed, amines and nitriles would then be decoupled from the gas and eventually be transported to regions where they can react with further radicals to yield ^{15}N -rich ices or organics.

Box 3. H-N isotope co-variations in the solar system and constraints on the origin of terrestrial volatiles

Various isotope fractionation processes have been invoked to explain the hydrogen and nitrogen isotope co-variations recorded by the different solar system objects and reservoirs. The kinetic isotope fractionation line shown in Fig. B3 is an example of mass-dependent fractionation, proportional to the square root of mass; however, different mass dependencies would lead to distinct slopes. The ion-molecule isotope fractionation trend (from ref. 75) illustrates the effect of low temperature isotope exchange between organics and the protosolar gas. The D-enrichment of ordinary chondrite IOM with only slight or no concomitant ^{14}N enrichment (yellow triangles) has been proposed to reflect isotope exchange of the molecular host with a D-rich component similar to gas phase molecules observed in dense clouds, pre-stellar cores and class 0 protostars⁷⁵. In contrast, the ^{15}N enrichments above the kinetic isotope fractionation line require preferential N isotope fractionation compared to D/H. This could result from photodissociation of protosolar N_2 by UV light⁷⁴, or from self-shielding⁹⁵ (Box 2), and appears to have occurred within the first few tens to hundreds thousands of years after solar system formation⁷¹. Similarly, the enrichment in the heavy isotopes of oxygen (^{17}O and ^{18}O) relative to the major, light isotope (^{16}O) recorded by solar system solids compared to the protosolar oxygen isotope composition may be the result of illumination of the PSN gas by solar/stellar photons (e.g., ref. 95). This process, which has been advocated to explain the oxygen isotope composition of refractory phases in meteorites, might also have been rapid ($\leq 2\text{Ma}$), according to the chronology recorded by these refractory phases (e.g., ref. 97).

Earth shares H and N isotope signatures with bulk chondrites (Fig. B3), whereas most comets are richer in D and ^{15}N . These observations suggest an asteroidal, rather than cometary, origin for terrestrial volatiles^{76,98}. While coupled H-N systematics provide a powerful means to trace the origin of volatiles in the inner solar system, secondary fractionation processes that lead to a preferential loss of light isotopes modified the H and N isotope signature of several solar system objects (Figs. 2 and B3). The Venusian atmosphere is extremely depleted in water and rich in D as a consequence of photodissociation of H_2O and subsequent H loss⁷⁸. The enrichment in both D and ^{15}N of the Martian atmosphere^{79,80} is attributed to atmospheric escape processes. In contrast, Titan is rich in ^{15}N but not in D (ref. 84), suggesting a source effect for nitrogen rather than isotope fractionation.

References

1. Boss, A. P. & Goswami, J. N. in *Meteorites early Sol. Syst. II* (Lauretta, D. S. & McSween, H. Y.) 171–186 (University of Arizona Press, 2006).

- 441 2. Warren, P. H. Stable-isotopic anomalies and the accretionary assemblage of the Earth
442 and Mars: A subordinate role for carbonaceous chondrites. *Earth Planet. Sci. Lett.* **311**,
443 93–100 (2011).
- 444 3. Zinner, E. *et al.* NanoSIMS isotopic analysis of small presolar grains: Search for Si₃N₄
445 grains from AGB stars and Al and Ti isotopic compositions of rare presolar SiC grains.
446 *Geochim. Cosmochim. Acta* **71**, 4786–4813 (2007).
- 447 4. Clayton, R. N. Isotopes: from Earth to the solar system. *Annu. Rev. Earth Planet. Sci.* **35**, 1–
448 19 (2007).
- 449 5. Rodgers, S. D. & Charnley, S. B. Nitrogen superfractionation in dense cloud cores. *Mon.*
450 *Not. R. Astron. Soc.* **385**, L48–L52 (2008).
- 451 6. Robert, F. The D/H ratio in chondrites. *Space Sci. Rev.* **106**, 87–101 (2003).
- 452 7. Robert, F., Gautier, D. & Dubrulle, B. The solar system D/H ratio: Observations and
453 theories. *Sp. Sci. Rev.* **92**, 201–224 (2000).
- 454 8. Deloule, E., Robert, F. & Doukhan, J. C. Interstellar hydroxyl in meteoritic chondrules:
455 Implications for the origin of water in the inner solar system. *Geochim. Cosmochim. Acta*
456 **62**, 3367–3378 (1998).
- 457 9. Bockelée-Morvan, D., Crovisier, J., Mumma, M. J. & Weaver, H. A. in *Comets II* (Festou, M.
458 C., Keller, H. U. & Weaver, H. A.) 391–423 (Univ. Arizona Press, 2004).
- 459 10. Hartogh, P. *et al.* Ocean-like water in the Jupiter-family comet 103P/Hartley 2. *Nature*
460 **478**, 218–220 (2011).
- 461 11. Altwegg, K. *et al.* 67P/Churyumov-Gerasimenko, a Jupiter family comet with a high D/H
462 ratio. *Science* **347**, 1261952 (2015).
- 463 12. Jacquet, E. & Robert, F. Water transport in protoplanetary disks and the hydrogen
464 isotopic composition of chondrites. *Icarus* **223**, 722–732 (2013).
- 465 13. Nier, A. A redetermination of the relative abundances of the isotopes of carbon, nitrogen,
466 oxygen, argon, and potassium. *Phys. Rev.* **77**, 789–793 (1950).

- 467 14. Cartigny, P. & Marty, B. Nitrogen isotopes and mantle geodynamics: the emergence of life
468 and the atmosphere-crust-mantle connection. *Elements* **9**, 359–366 (2013).
- 469 15. Audouze, J., Lequeux, J. & Vigroux, L. Isotopes of carbon, nitrogen and oxygen as probes
470 of nucleosynthesis, stellar mass losses and galactic evolution. *Astron. Astrophys.* **43**, 71–
471 83 (1975).
- 472 16. Chin, Y., Henkel, C., Langer, N. & Mauersberger, R. The detection of extragalactic ^{15}N :
473 consequences for nitrogen nucleosynthesis and chemical evolution. *Astrophys. J.* **512**,
474 L143–L146 (1999).
- 475 17. Adande, G. R. & Ziurys, L. M. Millimeter-wave observations of CN and HNC and their ^{15}N
476 isotopologues: a new evaluation of the $^{14}\text{N}/^{15}\text{N}$ ratio across the galaxy. *Astrophys. J.* **744**,
477 194 (2012).
- 478 18. Dahmen, G., Wilson, T. L. & Matteucci, F. The nitrogen isotope abundance in the Galaxy, I.
479 The Galactic disk gradient. *Astron. Astrophys.* **295**, 194–198 (1995).
- 480 19. Hily-Blant, P., Bonal, L., Faure, a. & Quirico, E. The ^{15}N -enrichment in dark clouds and
481 Solar System objects. *Icarus* **223**, 582–590 (2013).
- 482 20. Bizzocchi, L., Caselli, P. & Dore, L. Detection of N^{15}NH^+ in L1544. *Astron. Astrophys.* **510**,
483 L5 (2010).
- 484 21. Gerin, M. *et al.* Detection of $^{15}\text{NH}_2\text{D}$ in dense cores: A new tool for measuring the $^{14}\text{N}/^{15}\text{N}$
485 ratio in the cold ISM. *Astron. Astrophys.* **498**, L9–L12 (2009).
- 486 22. Lis, D. C., Wootten, A., Gerin, M. & Roueff, E. Nitrogen isotopic fractionation in interstellar
487 ammonia. *Astrophys. J. Lett.* **710**, L49–L52 (2010).
- 488 23. Wielen, R. & Wilson, T. L. The evolution of the C, N, and O isotope ratios from an
489 improved comparison of the interstellar medium with the Sun. *Astron. Astrophys.* **142**,
490 139–142 (1997).
- 491 24. Briani, G. *et al.* Pristine extraterrestrial material with unprecedented nitrogen isotopic
492 variation. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 10522–10527 (2009).

- 493 25. Hashizume, K., Chaussidon, M., Marty, B. & Terada, K. Protosolar carbon isotopic
494 composition: implications for the origin of meteoritic organics. *Astrophys. J.* **600**, 480–
495 484 (2004).
- 496 26. Fouchet, T. *et al.* ISO-SWS observations of Jupiter: measurement of the ammonia
497 tropospheric profile and of the $^{15}\text{N}/^{14}\text{N}$ isotopic ratio. *Icarus* **143**, 223–243 (2000).
- 498 27. Owen, T., Mahaffy, P. R., Niemann, H. B., Atreya, S. & Wong, M. Protosolar nitrogen.
499 *Astrophys. J.* **553**, L77–L79 (2001).
- 500 28. Meibom, A. *et al.* Nitrogen and carbon isotopic composition of the Sun inferred from a
501 high-temperature solar nebular condensate. *Astrophys. J.* **656**, L33–L36 (2007).
- 502 29. Burnett, D. S. & Genesis Sci, T. Solar composition from the Genesis Discovery Mission.
503 *Proc. Natl. Acad. Sci. U. S. A.* **108**, 19147–19151 (2011).
- 504 30. Marty, B. *et al.* Nitrogen isotopes in the recent solar wind from the analysis of Genesis
505 targets: Evidence for large scale isotope heterogeneity in the early solar system.
506 *Geochim. Cosmochim. Acta* **74**, 340–355 (2010).
- 507 31. Marty, B., Chaussidon, M., Wiens, R. C., Jurewicz, A. J. G. & Burnett, D. S. A ^{15}N -poor
508 isotopic composition for the solar system as shown by Genesis solar wind samples.
509 *Science* **332**, 1533–1536 (2011).
- 510 32. Becker, R. H. & Clayton, R. N. Nitrogen abundances and isotopic compositions in lunar
511 samples. *Proc. Lunar Sci. Conf. 6th* 2131–2149 (1975).
- 512 33. Kerridge, J. F. Solar nitrogen: Evidence for a secular change in the ratio of nitrogen-15 to
513 nitrogen-14. *Science* **188**, 162–164 (1975).
- 514 34. Bogard, D. D., Nyquist, L., Hirsch, W. C. & Moore, D. Trapped solar and cosmogenic noble
515 gas abundances in Apollo 15 and 16 deep drill samples. *Earth Planet. Sci. Lett.* **21**, 52–69
516 (1973).
- 517 35. Clayton, R. N. & Thiemens, M. H. in *Anc. Sun Foss. Rec. Earth, Moon, Meteorites* (Pepin, R.
518 O., Eddy, J. A. & Merrill, R. B.) 463–473 (Pergamon Press, 1980).

- 519 36. Geiss, J. & Bochsler, P. Nitrogen isotopes in the solar system. *Geochim. Cosmochim. Acta*
520 **46**, 529–548 (1982).
- 521 37. Wieler, R. The solar noble gas record in lunar samples and meteorites. *Space Sci. Rev.* **85**,
522 303–314 (1998).
- 523 38. Wieler, R., Humbert, F. & Marty, B. Evidence for a predominantly non-solar origin of
524 nitrogen in the lunar regolith revealed by single grain analyses. *Earth Planet. Sci. Lett.*
525 **167**, 47–60 (1999).
- 526 39. Hashizume, K., Marty, B. & Wieler, R. Analyses of nitrogen and argon in single lunar
527 grains: towards a quantification of the asteroidal contribution to planetary surfaces.
528 *Earth Planet. Sci. Lett.* **202**, 201–216 (2002).
- 529 40. Füri, E., Marty, B. & Assonov, S. S. Constraints on the flux of meteoritic and cometary
530 water on the Moon from volatile element (N-Ar) analyses of single lunar soil grains, Luna
531 24 core. *Icarus* **218**, 220–229 (2012).
- 532 41. Hashizume, K., Chaussidon, M., Marty, B. & Robert, F. Solar wind record on the Moon:
533 deciphering presolar from planetary nitrogen. *Science* **290**, 1142–1145 (2000).
- 534 42. Ozima, M. *et al.* Terrestrial nitrogen and noble gases in lunar soils. *Nature* **436**, 655–659
535 (2005).
- 536 43. Fletcher, L. N. *et al.* The origin of nitrogen on Jupiter and Saturn from the $^{15}\text{N}/^{14}\text{N}$ ratio.
537 *Icarus* **238**, 170–190 (2014).
- 538 44. Owen, T. *et al.* A low-temperature origin for the planetesimals that formed Jupiter.
539 *Nature* **402**, 269–270 (1999).
- 540 45. Kerridge, J. F., Eugster, O., Kim, J. S. & Marti, K. Nitrogen isotopes in the 74001/74002
541 double-drive tube from Shorty Crater, Apollo 17. *Proc. 21st Lunar Planet. Sci. Conf* **54**,
542 291–299 (1991).
- 543 46. Mathew, K. J. & Marti, K. Early evolution of Martian volatiles: Nitrogen and noble gas
544 components in ALH84001 and Chassigny. *J. Geophys. Res.* **106**, 1401–1422 (2001).

- 545 47. Hoffman, J. H., Oyama, V. I. & von Zahn, U. Measurements of the Venus lower atmosphere
546 composition: A comparison of results. *J. Geophys. Res.* **85**, 7871 (1980).
- 547 48. Alexander, C. M. O. *et al.* The origin of chondritic macromolecular organic matter: A
548 carbon and nitrogen study. *Meteorit. Planet. Sci.* **33**, 603–622 (1998).
- 549 49. Robert, F. & Epstein, S. The concentration and isotopic composition of hydrogen, carbon
550 and nitrogen in carbonaceous meteorites. *Geochim. Cosmochim. Acta* **46**, 81–95 (1982).
- 551 50. Kerridge, J. F. Carbon, hydrogen and nitrogen in carbonaceous chondrites: abundances
552 and isotopic compositions in bulk samples. *Geochim. Cosmochim. Acta* **49**, 1707–1714
553 (1985).
- 554 51. Weisberg, M. *et al.* The CR chondrite clan. *Proc. NIPR Symp. Antart. Meteorites* **8**, 11–32
555 (1995).
- 556 52. Grady, M. M. & Pillinger, C. T. ALH 85085: nitrogen isotope analysis of a highly unusual
557 primitive chondrite. *Earth Planet. Sci. Lett.* **97**, 29–40 (1990).
- 558 53. Prombo, C. A. & Clayton, R. N. A striking nitrogen isotope anomaly in the Bencubbin and
559 Weatherford meteorites. *Science* **230**, 935–937 (1985).
- 560 54. Marty, B., Kelley, S. & Turner, G. Chronology and shock history of the Bencubbin
561 meteorite: A nitrogen, noble gas, and Ar–Ar investigation of silicates, metal and fluid
562 inclusions. *Geochim. Cosmochim. Acta* **74**, 6636–6653 (2010).
- 563 55. Ivanova, M. A. *et al.* The Isheyevo meteorite: Mineralogy, petrology, bulk chemistry,
564 oxygen, nitrogen, carbon isotopic compositions, and ^{40}Ar – ^{39}Ar ages. *Meteorit. Planet. Sci.*
565 **43**, 915–940 (2008).
- 566 56. Bonal, L. *et al.* Highly ^{15}N -enriched chondritic clasts in the CB/CH-like meteorite
567 Isheyevo. *Geochim. Cosmochim. Acta* **74**, 6590–6609 (2010).
- 568 57. Busemann, H. *et al.* Interstellar chemistry recorded in organic matter from primitive
569 meteorites. *Science* **312**, 727–730 (2006).

- 570 58. Aléon, J., Robert, F., Chaussidon, M. & Marty, B. Nitrogen isotopic composition of
571 macromolecular organic matter in interplanetary dust particles. *Geochim. Cosmochim.*
572 *Acta* **67**, 3773–3783 (2003).
- 573 59. Nakamura-Messenger, K., Messenger, S., Keller, L. P., Clemett, S. J. & Zolensky, M. E.
574 Organic globules in the Tagish Lake meteorite: remnants of the protosolar disk. *Science*
575 **314**, 1439–1442 (2006).
- 576 60. Brownlee, D. E. The Stardust mission: Analyzing samples from the edge of the solar
577 system. *Annu. Rev. Earth Planet. Sci.* **42**, 179–205 (2014).
- 578 61. Brownlee, D. Comet 81P/Wild 2 under a microscope. *Science* **314**, 1711–1716 (2006).
- 579 62. McKeegan, K. D. *et al.* Isotopic compositions of cometary matter returned by Stardust.
580 *Science* **314**, 1724–8 (2006).
- 581 63. Stadermann, F. J. *et al.* Stardust in Stardust - The C, N, and O isotopic compositions of
582 Wild 2 cometary matter in Al foil impacts. *Meteorit. Planet. Sci.* **313**, 299–313 (2008).
- 583 64. Arpigny, C. *et al.* Anomalous nitrogen isotope ratio in comets. *Science* **301**, 1522–1524
584 (2003).
- 585 65. Jehin, E., Manfroid, J., Hutsemékers, D., Arpigny, C. & Zucconi, J.-M. Isotopic ratios in
586 comets: status and perspectives. *Earth. Moon. Planets* **105**, 167–180 (2009).
- 587 66. Bockelée-Morvan, D. *et al.* Large excess of heavy nitrogen in both hydrogen cyanide and
588 cyanogen from comet 17P/Holmes. *Astrophys. J.* **679**, L49–L52 (2008).
- 589 67. Rousselot, P. *et al.* Toward a unique nitrogen isotopic ratio in cometary ices. *Astrophys. J.*
590 **780**, L17 (2014).
- 591 68. Shinnaka, Y., Kawakita, H., Kobayashi, H., Nagashima, M. & Boice, D. C. $^{14}\text{NH}_2/^{15}\text{NH}_2$ ratio
592 in comet C/2012 S1 (Ison) observed during its outburst in 2013 November. *Astrophys. J.*
593 **782**, L16 (2014).
- 594 69. Duncan, M. J. & Levison, H. F. A disk of scattered icy objects and the origin of Jupiter-
595 family comets. *Science* **276**, 1670–1672 (1997).

- 596 70. Carusi, A., Kresák, L., Perozzi, E. & Valsecchi, G. B. High-order librations of Halley-type
597 comets. *Astron. Astrophys.* **187**, 899–905 (1987).
- 598 71. Füri, E., Chaussidon, M. & Marty, B. Evidence for an early nitrogen isotopic evolution in
599 the solar nebula from volatile analyses of a CAI from the CV3 chondrite NWA 8616.
600 *Geochim. Cosmochim. Acta* **153**, 183–201 (2015).
- 601 72. Terzieva, R. & Herbst, E. The possibility of nitrogen isotopic fractionation in interstellar
602 clouds. *Mon. Not. R. Astron. Soc.* **317**, 563–568 (2000).
- 603 73. Charnley, S. B. & Rodgers, S. D. The end of interstellar chemistry as the origin of nitrogen
604 in comets and meteorites. *Astrophys. J. Lett.* **569**, L133–L137 (2002).
- 605 74. Chakraborty, S. *et al.* Massive isotopic effect in vacuum UV photodissociation of N₂ and
606 implications for meteorite data. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 14704–14709 (2014).
- 607 75. Aléon, J. Multiple origins of nitrogen isotopic anomalies in meteorites and comets.
608 *Astrophys. J.* **722**, 1342–1351 (2010).
- 609 76. Marty, B. The origins and concentrations of water, carbon, nitrogen and noble gases on
610 Earth. *Earth Planet. Sci. Lett.* **313–314**, 56–66 (2012).
- 611 77. Füri, E., Deloule, E., Gurenko, A. & Marty, B. New evidence for chondritic lunar water
612 from combined D/H and noble gas analyses of single Apollo 17 volcanic glasses. *Icarus*
613 **229**, 109–120 (2014).
- 614 78. Grinspoon, D. H. Implications of the high D/H ratio for the sources of water in Venus’
615 atmosphere. *Nature* **363**, 428–431 (1993).
- 616 79. Leshin, L. A. Insights into martian water reservoirs from analyses of martian meteorite
617 QUE94201. *Geophys. Res. Lett.* **27**, 2017–2020 (2000).
- 618 80. Leshin, L. A. *et al.* Volatile, isotope, and organic analysis of martian fines with the Mars
619 Curiosity rover. *Science* **341**, 1238937 (2013).

- 620 81. Alexander, C. M. O., Fogel, M., Yabuta, H. & Cody, G. D. The origin and evolution of
621 chondrites recorded in the elemental and isotopic compositions of their macromolecular
622 organic matter. *Geochim. Cosmochim. Acta* **71**, 4380–4403 (2007).
- 623 82. Macy, W. & Smith, W. H. Detection of HD on Saturn and Uranus, and the D/H ratio.
624 *Astrophys. J.* **222**, L73 (1978).
- 625 83. Mahaffy, P. R., Donahue, T. M., Atreya, S. K., Owen, T. C. & Niemann, H. B. Galileo probe
626 measurements of D/H and $^3\text{He}/^4\text{He}$ in Jupiter’s atmosphere. *Space Sci. Rev.* **84**, 251–263
627 (1998).
- 628 84. Niemann, H. B. *et al.* Composition of Titan’s lower atmosphere and simple surface
629 volatiles as measured by the Cassini-Huygens probe gas chromatograph mass
630 spectrometer experiment. *J. Geophys. Res.* **115**, E12006 (2010).
- 631 85. Ceccarelli, C. *et al.* in *Protostars Planets VI* (Beuther, H., Klessen, R., Dullemond, C. &
632 Henning, T.) (University of Arizona Press, 2014).
- 633 86. Manfroid, J. *et al.* The CN isotopic ratios in comets. *Astron. Astrophys.* **503**, 613–624
634 (2009).
- 635 87. Javoy, M. & Pineau, F. The volatiles record of a “popping” rock from the Mid-Atlantic
636 Ridge at 14°N: chemical and isotopic composition of gas trapped in the vesicles. *Earth*
637 *Planet. Sci. Lett.* **107**, 598–611 (1991).
- 638 88. Hashizume, K. & Marty, B. in *Handb. stable Isot. Anal. Tech.* (de Groot, P. A.) 361–375
639 (Elsevier Science, 2004).
- 640 89. Barry, P. H., Hilton, D. R., Halldórsson, S. A., Hahm, D. & Marti, K. High precision nitrogen
641 isotope measurements in oceanic basalts using a static triple collection noble gas mass
642 spectrometer. *Geochemistry, Geophys. Geosystems* **13**, (2012).
- 643 90. Hauri, E. H., Wang, J., Pearson, D. G. & Bulanova, G. P. Microanalysis of $\delta^{13}\text{C}$, $\delta^{15}\text{N}$, and N
644 abundances in diamonds by secondary ion mass spectrometry. *Chem. Geol.* **185**, 149–
645 163 (2002).

- 646 91. Rubin, M. *et al.* Molecular nitrogen in comet 67P/Churyumov - Gerasimenko indicates a
647 low formation temperature. *Science* (2015).
- 648 92. Adams, N. G. & Smith, D. $^{14}\text{N}/^{15}\text{N}$ isotope fractionation in the reaction $\text{N}_2\text{H}^+ + \text{N}_2$:
649 Interstellar significance. *Astrophys. J.* **247**, L123–L125 (1981).
- 650 93. Geppert, W. D. *et al.* Dissociative recombination of N_2H^+ : evidence for fracture of the N-N
651 Bond. *Astrophys. J.* **609**, 459–464 (2004).
- 652 94. Molek, C. D., McLain, J. L., Poterya, V. & Adams, N. G. A remeasurement of the products for
653 electron recombination of N_2H^+ using a new technique: no significant $\text{NH} + \text{N}$ production.
654 *J. Phys. Chem. A* **111**, 6760–6765 (2007).
- 655 95. Clayton, R. N. Self-shielding in the solar nebula. *Nature* **415**, 860–861 (2002).
- 656 96. Muskatel, B. H., Remale, F., Thiemens, M. H. & Levine, R. D. On the strong and selective
657 isotope effect in the UV excitation of N_2 with implications toward the nebula and Martian
658 atmosphere. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 6020–6025 (2011).
- 659 97. Davis, A. M. *et al.* in *Protostars Planets VI* (Beuther, H., Klessen, R. S., Dullemond, C. P. &
660 Henning, T.) 809–831 (University of Arizona Press, 2015).
- 661 98. Alexander, C. M. O. *et al.* The provenances of asteroids, and their contributions to the
662 volatile inventories of the terrestrial planets. *Science* **337**, 721–723 (2012).

663

664 **Additional information**

665 Correspondence and requests for materials should be addressed to B.M.

666

667 **Acknowledgements**

668 Discussions with Jérôme Aléon, Kathrin Altwegg, Dominique Bockelée-Morvan, Pierre
669 Cartigny, Marc Chaussidon, Kathie Mandt, Olivier Mousis, and François Robert were much
670 appreciated during the preparation of this work. This work was supported by the European
671 Research Council under the European Community's Seventh Framework Programme

672 (FP7/2007-2013 grant agreement no. 267255 to B.M.), the Deep Carbon Observatory, and the
673 CNRS-INSU Programme de Planétologie (PNP). This is CRPG-CNRS contribution 2372.