Abundant molecular oxygen in the coma of comet 67P/Churyumov–Gerasimenko

Andre Bieler, Kathrin Altwegg, Hans Balsiger, Akiva Bar-Nun, Jean-Jacques Berthelier, P. Bochslers, Christelle Briais, Ursina Calmonte, M. Combi, J. de Keyser, et al.

► To cite this version:

Andre Bieler, Kathrin Altwegg, Hans Balsiger, Akiva Bar-Nun, Jean-Jacques Berthelier, et al.. Abundant molecular oxygen in the coma of comet 67P/Churyumov–Gerasimenko. *Nature*, 2015, 526 (7575), pp.678-681. 10.1038/nature15707 . hal-01346075

HAL Id: hal-01346075

<https://hal.science/hal-01346075>

Submitted on 20 Jul 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Measurement of molecular oxygen in the coma of 67P/Churyumov-Gerasimenko

Authors: A. Bieler^{1, 2*}, K. Altwegg^{2, 3}, H. Balsiger², A. Bar-Nun⁴, J.-J. Berthelier⁵, P. Bochsler², C. Briois⁶, U. Calmonte², M. Combi¹, J. De Keyser⁷, E. F. van Dishoeck¹⁶, B. Fiethe⁸, S. A. Fuselier⁹, S. Gasc², T. I. Gombosi¹, K. C. Hansen¹, M. Hässig^{2, 9}, A. Jäckel², E. Kopp², A. Korth¹⁰, L. Le Roy³, U. Mall¹⁰, B. Marty¹¹, O. Mousis¹², T. Owen¹³, H. Rème^{14, 15}, M. Rubin¹, T. Sémon², C.-Y. Tzou², J. H. Waite⁹, C. Walsh¹⁶, P. Wurz^{2, 3}

Affiliations:

¹Department of Atmospheric, Oceanic and Space Sciences, University of Michigan, 2455 Hayward Street, Ann Arbor, MI 48109, USA

²Physikalisches Institut, University of Bern, Sidlerstr. 5, CH-3012 Bern, Switzerland.

³Center for Space and Habitability, University of Bern, Sidlerstr. 5, CH-3012 Bern, Switzerland.

⁴Department of Geoscience, Tel-Aviv University, Ramat-Aviv, Tel-Aviv, Israel

⁵LATMOS/IPSL-CNRS-UPMC-UVSQ, 4 Avenue de Neptune F-94100, Saint-Maur, France.

⁶Laboratoire de Physique et Chimie de l'Environnement et de l'Espace (LPC2E), UMR 6115 CNRS – Université d'Orléans, France.

⁷Belgian Institute for Space Aeronomy, BIRA-IASB, Ringlaan 3, B-1180 Brussels, Belgium.

⁸Institute of Computer and Network Engineering (IDA), TU Braunschweig, Hans-Sommer-Straße 66, D-38106 Braunschweig, Germany.

⁹Department of Space Science, Southwest Research Institute, 6220 Culebra Rd., San Antonio, TX 78228, USA.

¹⁰Max-Planck-Institut für Sonnensystemforschung, Justus-von-Liebig-Weg 3, 37077 Göttingen, Germany.

¹¹Centre de Recherches Pétrographiques et Géochimiques, CRPG-CNRS, Université de Lorraine, 15 rue Notre Dame des Pauvres, BP 20, 54501 Vandoeuvre lès Nancy, France.

¹²Aix Marseille Université, CNRS, LAM (Laboratoire d'Astrophysique de Marseille) UMR 7326, 13388, Marseille, France.

¹³Institute for Astronomy, University of Hawaii, Honolulu, HI 96822, USA

¹⁴Université de Toulouse; UPS-OMP; IRAP, Toulouse, France.

¹⁵CNRS; IRAP; 9 Avenue du Colonel Roche, BP 44346, F-31028 Toulouse Cedex 4, France.

¹⁶Leiden Observatory, Leiden University, P.O. Box 9513, 2300 RA Leiden, Netherlands.

Abstract

The composition of the neutral gas coma of a comet is dominated by H₂O, CO and CO₂, typically comprising as much as 95% of the total gas density¹. In addition to these common species, the cometary coma has been measured to contain a rich array of additional molecules including noble gases, sulfuric compounds and complex hydrocarbons. Molecular oxygen (O₂), despite its detection on other icy bodies, such as the moons of Jupiter and Saturn^{2, 3}, had however remained undetected in the cometary volatile inventory. Here we report the direct in situ measurement of molecular oxygen in the cometary coma of 67P/Churyumov-Gerasimenko with a local abundance ranging from 1% to 10% relative to H₂O and a mean value of 3.7 ± 1.5 %. Our observations indicate that the O₂/H₂O ratio is isotropic in the coma and does not systematically change over a period of several months. This suggests that O₂ was incorporated into the cometary nucleus primordially during the comet's formation. Current Solar System formation models do not predict conditions that would allow this to occur, suggesting that our detection may play a significant role in advancing our understanding of comet formation and the prevailing conditions and processes in the early stages of our Solar System.

Measurements of the coma of 67P/Churyumov-Gerasimenko (hereafter 67P) were made between September 2014 and March 2015 with the ROSINA-DFMS mass spectrometer⁴ onboard the Rosetta spacecraft. For this study we analyzed 2808 mass spectra taken in this time period. Due to the high resolving power and sensitivity of ROSINA-DFMS it is possible to unambiguously differentiate between the three main species present in the narrow mass range centered on 32 Da/e; molecular oxygen (O₂), sulfur (S) and methanol (CH₃OH), something which has not been achieved by previous in situ or remote measurements at comets. Fig. 1 shows several measurements centered at the O₂ peak. The green and orange lines show data taken before the close encounter with 67P. Only minor signatures from the tenuous neutral gas atmosphere of the Rosetta spacecraft can be identified and even after long thruster firing maneuvers, which use N₂O₄ as an oxidizer, the contamination of the O₂ signal remains small (green line in Fig. 1). Measurements while orbiting 67P, shown as the light blue, dark blue and purple line in Fig. 1, show a clear increase of the O₂ signal, indicating the presence of cometary O₂. These three measurements were taken at decreasing distances (r) from the comet nucleus and follow the predicted $1/r^2$ signal dependence that is expected for a conserved cometary species, further gaining confidence in our detection.

As previously reported, the local number densities in the coma vary spatially and temporally^{5, 6}, for different compounds. The bottom panel of Fig. 2 shows an O₂ and H₂O measurement sequence taken between the 5th and the 7th of November 2014 and the peaks and valleys occur with the rotation frequency of 12.4 hours of 67P⁷. During this time range the radial distance from the comet is nearly constant at roughly 30 km, the phase angle decreases from 115 to 102 degrees, the latitude linearly (in time) increases from -22 to 22 degrees and all longitudes are covered multiple times (due to the short

rotation of 67P). Uncertainties of the individual measurements for both species are on the order of 30% and are indicated by the shaded areas in Fig. 2. We observe a strong correlation between the absolute abundances of H₂O and O₂, not only for the time span shown in Fig. 2, but for the entire dataset. This strong correlation, with a Pearson correlation coefficient of 0.91, indicates that H₂O and O₂ are of similar origin on the nucleus and their release mechanisms are linked. Despite the overall strong correlation the O₂ ratio decreases for high H₂O abundances as can be seen in the top panel of Fig. 2. Because of the radial dependence of the number density there is however not an absolute value above which this saturation effect is observed. This effect is the main cause for the variability in the measured O₂ ratios as shown in the time series in Fig 3.

A plausible mechanism for the strong O₂/H₂O correlation is the possibility to produce O₂ by radiolysis or photolysis of water ice. Here we follow the nomenclature that photolysis refers to UV photons that break bonds, whereas radiolysis refers to more energetic photons or fast electrons and ions depositing energy into the ice and ionizing molecules. Creation of sputtered O₂ by radiolysis was demonstrated by Hart et al.⁸ and is observed for the icy moons of Jupiter; Europa, Ganymede and Callisto⁹⁻¹¹, as well as for the moons of Saturn; Dione and Rhea³. At comets, radiolysis does happen on different time scales: during billions of years while they are residing in the Kuiper belt, over the period of several years once they enter the inner Solar System and on very short time scales. In the first case, the skin depth for producing O₂ is in the range of a meter, while in the latter it is only a few micrometers. Once a comet begins its residence in the inner Solar System, it loses several meters of its surface material during each orbit around the Sun, therefore we can safely assume that all radiolysis products created in the Kuiper Belt phase are gone from 67P. Furthermore, radiolysis and photolysis by solar wind and UV radiation in the inner Solar System only affect the top few micrometers of the cometary surface. Due to 67P's continuous mass loss through outgassing, we estimate the surface loss to be in the range of several cm for the time from August 2014 to March 2015. If recent production were the source of the measured O₂, our data would show a continuous decrease of the O₂ ratio over the examined time period. Apart from the variations related to the H₂O abundance, Fig. 3 shows that we do not observe a systematic change in the O₂ ratio over several months. On-line creation of the measured O₂ by radiolysis or photolysis at 67P seems overall unlikely and would lead to anisotropic O₂ ratios. Given that radiolysis and photolysis, on any of the discussed time scales, do not seem plausible production mechanisms for O₂, our preferred explanation is the incorporation of primordial O₂ into the cometary nucleus.

Time dependent models of surface grain chemistry in molecular clouds predict an abundance of a few percent of O₂ relative to H₂O on icy grains at time scales of >10⁶ years¹². Thereby O₂ is formed together with H₂O by grain surface reactions, which would explain their correlation and be in line with the very low abundances of HO₂ and H₂O₂ that we measured in the coma of 67P (see Fig. 4). A further consequence would be that these icy grains have been incorporated into the comet mostly unaltered, a fact very much under debate, but which has recently been proposed again by Cleaves et al.¹³ and would also be in accordance with the measured D/H ratio in 67P¹⁴.

Constraints on the O₂ abundance are of great benefit for future theoretical studies and modeling efforts as the current understanding of the grain-gas interaction is still evolving. In contrast, laboratory experiments of photolysis or particle bombardment of H₂O ice find significantly higher abundances of H₂O₂ than we measured¹⁵.

Interestingly, there has not been a direct detection of O₂ in interstellar ices, but upper limits for the O₂/H₂O ratios of <0.1 have been published based on analysis of solid ¹³CO lines^{16, 17} on icy grains. These

limits then lead to a O_2/CO_2 ratio of approximately 0.75. Both of these ratios, albeit relatively uncertain, agree with our findings and previous observations of 67P². So far, this non-detection has preferably been explained by the high volatility and chemical reactivity of O_2 . This potential O_2 abundance on icy grains on the order of a few percent relative to H_2O would have significant implications on our current understanding of ices in interstellar molecular clouds and their importance for the formation of comets. The global distribution of elemental O in the interstellar medium is probably not affected by our findings, as O_2 on ice grains with an abundance of a few percent relative to H_2O accounts only for a small fraction of the total O inventory.

An alternative explanation for the O_2 is the incorporation of gaseous O_2 into water ice. Models of protoplanetary disks do show that O_2 can be abundant in the comet forming zone. A rapid cooling scenario from >100 K to less than 30 K is then needed to form water ice with trapped O_2 . However, despite great efforts by remote sensing campaigns, O_2 has only been detected in two regions in the interstellar medium so far¹⁸⁻²⁰. This lack of O_2 is not understood but as a consequence, molecular oxygen is generally considered to be present only at very low abundances. However, the abundance ratios of HO_2/O_2 and H_2O_2/O_2 determined by DFMS for the coma of 67P are very close to those observed in the rho Oph A interstellar core, one of the two regions where O_2 was actually detected in the ISM. We find ratios of $HO_2/O_2 = (1.9 \pm 0.3) \times 10^{-3}$ and $H_2O_2/O_2 = (0.6 \pm 0.07) \times 10^{-3}$, which are similar to $HO_2/O_2 \sim H_2O_2/O_2 \sim 0.6 \times 10^{-3}$ as reported for rho Oph A^{21,22}. To what extent these gas-phase abundance ratios reflect those in the ice is however still unclear.

The unique case of rho Oph A has been interpreted by having experienced slightly higher temperatures of around 20-30 K over its lifetime, compared to ~10 K for most other interstellar clouds^{22, 23}. Applied to our own Solar System this might indicate it was formed from an unusual warm molecular cloud, challenging our current understanding of the chemistry occurring during these early stages. Admittedly, the observed ratios are in contrast to simulations of interstellar ice chemistry, which predict H_2O_2 and O_3 ice to be more abundant than O_2 ice by one to two orders of magnitude²⁴. We found no evidence for the presence of ozone (O_3) (see Fig. 4) for which we give an upper limit of 1×10^{-6} relative to water.

References

- [1] Bockelée-Morvan, D., Mumma, M. J., Weaver, H. A. 2004, In *Comets II*, University of Arizona Press, Tucson, Arizona, p. 391-423.
- [2] Lanzerotti, L. J., Brown, W. L., Poate, J. M. & Augustyniak, W. M. Production of O_2 on icy satellites by electronic excitation of low-temperature water ice. *Geophys. Res. Lett.* 5, 155–158 (1978).
- [3] Noll, K. S., Roush, T. L., Cruikshank, D. P., Johnson, R. E. & Pendleton, Y. L. Detection of ozone on Saturn's satellites Rhea and Dione. *Nature* 388, 45–47 (1997).
- [4] Balsiger, H. et al. ROSINA – ROSETTA orbiter spectrometer for ion and neutral analysis. *Sp. Sci. Rev.* 128, 745-801, (2007)

- [5] Hässig, M. et al. Time variability and heterogeneity in the coma of 67P/Churyumov-Gerasimenko. Science 347 (2015). URL <http://www.sciencemag.org/content/347/6220/aaa0276.abstract>. <http://www.sciencemag.org/content/347/6220/aaa0276.full.pdf>.
- [6] Adrienne
- [7] ref on rotation period
- [8] Hart, E. G. & Platzmann, R. L. Mechanisms in Radiobiology (Academic Press, New York, 1961).
- [9] Carlson, R. W. et al. Hydrogen peroxide on the surface of Europa. Science 283, 2062–2064 (1999). URL <http://www.sciencemag.org/content/283/5410/2062.abstract>. <http://www.sciencemag.org/content/283/5410/2062.full.pdf>.
- [10] Spencer, J. R., Calvin, W. M. & Person, M. J. Charge-coupled device spectra of the Galilean satellites: Molecular oxygen on Ganymede. J. Geophys. Res. (1995).
- [11] Spencer, J. R. & Calvin, W. M. Condensed O₂ on Europa and Callisto. The Astronomical Journal 124, 3400–3403 (2002).
- [12] D’Hendecourt, L. B., Allamandola, L. J., Greenberg, J. M. Time dependent chemistry in dense molecular clouds. I - Grain surface reactions, gas/grain interactions and infrared spectroscopy. Astron. Astrophys., 152, 130-150, (1985)
- [13] Cleaves, L. I., Bergin, E. A., Alexander, C. M., Du, F., Graninger, D., Öberg, K. I., Harries, T. J. The ancient heritage of water ice in the solar system. Science, 345, (2014)
- [14] Altwegg et al. 67P/Churyumov-Gerasimenko, a Jupiter family comet with a high D/H ratio. Science, 347, (2015)
- [15] Zheng, W., Jewitt, D. & Kaiser, R. I. Formation of hydrogen, oxygen and hydrogen peroxide in electron-irradiated crystalline water ice. The Astrophysical Journal 639, 534–548 (2006).
- [16] Vandenbussche, B. et al. Constraints on the abundance of solid O₂ in dense clouds from ISO-SWS and ground-based observations. Astron. Astrophys., 346, L57-L60 (1999)
- [17] Pontoppidan, K. et al. A 3-5 μ m VLT spectroscopic survey of embedded young low mass stars I. Structure of the CO ice. Astron. Astrophys., 408, 981-1007 (2003)
- [18] Liseau, R. et al. Multi-line detection of O₂ toward rho Ophiuchi A. Astron. Astrophys., 541, A73 (2012)

- [19] Larsson, B., Liesau, R., Pagani, L. et al. Molecular oxygen in the Ophiuchi cloud. *Astron. & Astrophys.*, 466, 999 (2007)
- [20] Goldsmith, P. F., Liesau, R., Bell, T. A. et al. Herschel Measurements of molecular oxygen in Orion. *ApJ*, 737, 96, (2011)
- [21] Bergman, P. et al. Detection of interstellar hydrogen peroxide, *Astron. Astrophys.*, 531, L8 (2011)
- [22] Parise, B., Bergman, P., Du, F. Detection of the hydroperoxyl radical HO₂ toward rho Ophiuchi A. Additional constraints on the water chemical network. *Astron. Astrophys.*, 541, L11 (2012)
- [23] Du, F., Parise, B., Bergman, P. Production of interstellar hydrogen peroxide (H₂O₂) on the surface of dust grains. *Astron. Astrophys.*, 538, A91 (2012)
- [24] Taquet, V., Ceccarelli, C., Kahane, C. Multilayer modeling of porous grain surface chemistry. I. The GRAINOBLE model. *Astron. Astrophys.*, 538, A42 (2012)

Figure 1: Normalized DFMS mass spectrum at 32 Da/e. The black labels indicate the three major species found in the coma of 67P at 32 Da/e. The green labels and graph identify contamination peaks from thruster firings, the contribution to the O₂ peak is very low. The light blue, dark blue and purple lines represent measurements taken at different distances from the comet nucleus.

Figure 2: Correlation between O_2 ratio and the H_2O abundance for 5th to 7th of November 2014. Shaded areas indicate 1 sigma error limits. The O_2 signal in the bottom panel is multiplied by a factor of 15 for visual clarity. The O_2 ratio systematically drops for high H_2O abundances.

Figure 3: O_2 ratio over time. There seems to be no systematic increase or decrease of the O_2 ratio, as the variances can be explained by Figure 2.

Figure 4: DFMS spectra for some of the common products of radiolysis of water ice; O_2 , O_2H , O_2H_2 and O_3 . This data was recorded on October 20th at around 01:00 UTC. With the exception of O_3 (see panel C), all the previously mentioned species are measured and can clearly be identified in the mass spectra of DFMS. Shaded areas mark 1 sigma error boundaries.

Methods

Spacecraft outgassing background

To clearly identify the measured O₂ as cometary in origin, all non-cometary sources of O₂ must be considered and excluded. The Rosetta spacecraft produces a neutral gas cloud of its own, mainly due to diffusion of volatiles out of spacecraft material and desorption of re-deposited volatiles from the spacecraft. For example, by changing the spacecraft attitude, different components are illuminated by the Sun, which then warm up and release gas. The orange line in Fig. 1 shows the low level signals from this spacecraft contamination for O₂, S, and CH₃OH, referred to as “background”, in measurements taken several days before the encounter with 67P. It is not possible to distinguish this background signal from any potential cometary signature with DFMS, but it has been well characterized prior to the arrival at 67P and is usually orders of magnitude lower than the measured O₂ signals⁶. To keep the background influence as low as possible, we only considered mass spectra where both the O₂ and H₂O abundances are at least 10 times larger than the corresponding spacecraft contamination. Another potential source of O₂ is the oxidizer, N₂O₄, used by the Rosetta spacecraft during thruster firings. Measurements taken shortly after a large thruster firing maneuver from June 2014 (still before arrival at the comet) show minor contaminations around 32 Da/e, but not directly affecting the O₂ peak (see green curve in Fig. 1). Although contamination from thruster firings is small, DFMS measurements are usually performed hours after thruster firings, in order to minimize influence thereof. Finally we can exclude the production of O₂ inside the instrument through a careful review of all oxygen-bearing molecules up to 150 Da/e, which could potentially fragment into O₂ in the DFMS electron impact ion source. Many minor species contain O₂ but these are too low in abundance to account for the large amount of O₂ detected. The remaining possibility is CO₂, which is very abundant in the coma of 67P². However, due to its chemical structure it only fragments into CO and O, not O₂^[7]. Finally, we exclude the production of O₂ from H₂O in the instrument. For 81 mass spectra taken from May to the end of June 2014 we determine an O₂ abundance of $(0.18 \pm 0.07) \%$ relative to H₂O, which is a factor of 20 lower than the cometary values.

Correlation with H₂O

The measured O₂ signal shows a very strong dependence on radial distance (*r*) from the comet. It increases by roughly one order of magnitude when the radial distance from the comet decreases from 30 km to 10 km. This is in agreement with a predicted $1/r^2$ dependence of the number density profile of a non-reactive species. Examining the data further, we observe a strong correlation between H₂O and

O₂(see Fig. 2, Pearson correlation coefficient $R = 0.91$) for data from September 2014 to March 2015. This correlation indicates that O₂ and H₂O are both of a similar cometary origin. In contrast, there is no correlation between O₂ and H₂O for measurements taken before the arrival at the comet ($R = -0.01$). The observed temporal variations in the O₂/H₂O ratio are largely due to a non-linear correlation between H₂O and O₂ for high water densities. The O₂ ratio drops with increasing H₂O abundance as is indicated in Fig. 2. A possible explanation for this is a modification of the ice-dust matrix close to the surface, e.g. sintering or re-deposition of ice grains as surface frost which then is depleted in O₂. The correlation similarly supports the ruling out of CO₂ as a source of the O₂ because Hässig et al.² and Rubin et al.⁸ show a lack of correlation between the abundance of H₂O and other species like CO, CO₂ and N₂.

Radiolysis

The production of O₂ from water ice by radiolysis is the result of several reactions, where initially H, O, and OH are produced, followed by subsequent rearrangement to form H₂, HO₂ and H₂O₂ and ultimately O₂^[10, 11]. These radiolysis products can either be trapped¹² inside voids in the water ice that are also created by radiolysis, or can be scattered or desorbed directly from the surface

Sputtering

O₂ production through surface sputtering: Wurz et al.²¹ show that sputtering of refractory materials from the cometary surface due to the solar wind is occurring now at 67P. This work demonstrates a clear difference between the southern and northern latitudes in the measured abundances of the sputtered species. This apparent spatial difference is explained via the asymmetry of the neutral coma, with higher number densities in the northern hemisphere that is preferentially exposed to the Sun for the time span under study. The solar wind, which is responsible for the sputtering, is therefore attenuated more efficiently by these denser parts of the coma and thus has limited or no access to the surface. We find that the O₂/H₂O ratio is independent of latitude and relatively constant over a period of 7 months. Furthermore, the major part of the top surface of 67P accessible by the solar wind does not contain any water ice²². This suggests that sputtering cannot be the main source of the detected molecular oxygen. Moreover, the sputter yields by solar wind ions are orders of magnitude too low to explain the observed amount of O₂.

Recent radiolysis

O₂ production after the comet formed: Zheng et al.²⁵ argue that Kuiper belt objects, such as 67P, subject to radiolysis, reach an equilibrium abundance of O₂ and H₂O in the surface ice layer after some 10⁵ years, leading to a relative abundance of O₂/H₂O of about 0.6%. Once these objects enter the inner Solar System they lose (due to their activity and depending on their perihelion distance) material from the surface in the range of several meters per orbit around the Sun. Therefore, as most products built up during the stay in the Kuiper belt reside in the outer few meters, these should be released quickly on the first solar passages. A study by Maquet²⁶ shows that 67P's perihelion distance was within the orbit of Jupiter for the last 250 years, possibly even for more than 5000 years. Since a close encounter with Jupiter in 1959, the perihelion distance of 67P has been about 1.3 AU with an orbital period of 6.4 years. Accumulated over the last perihelion passages we can assume that 67P has lost hundreds of meters from its surface. Although Galactic Cosmic Rays with energies well above 1 GeV can penetrate down to depths of hundreds of meters, the amount of O₂ that can be produced by GCR-induced radiolysis appears to be at least several orders of magnitude too small if one assumes that the present-day high-energy GCR flux represents the average flux over the history of the Solar System. There are also uncertainties on the cross-sections, the relative proportion of high-Z cosmic rays, and the role of porosity and defects in the cometary ice^{21, 27}. Overall, it appears unlikely that most of the O₂ in 67P is of evolutionary origin.