



HAL
open science

Snapmelon (*Cucumis melo* L. *Momordica* group), an indigenous cucurbit from India with immense value for melon breeding

Narinder P.S. Dhillon, H. Singh, Michel Pitrat, Antonio J. Monforte, James D. McCreight

► **To cite this version:**

Narinder P.S. Dhillon, H. Singh, Michel Pitrat, Antonio J. Monforte, James D. McCreight. Snapmelon (*Cucumis melo* L. *Momordica* group), an indigenous cucurbit from India with immense value for melon breeding. *Acta Horticulturae*, 2015, 1102, pp.99-108. hal-01351898

HAL Id: hal-01351898

<https://hal.science/hal-01351898>

Submitted on 28 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Snampmelon (*Cucumis melo* L. *Momordica* group), an indigenous cucurbit from India with immense value for melon breeding

N.P.S. Dhillon^{1,a}, H. Singh², M. Pitrat³, A.J. Monforte⁴ and J.D. McCreight⁵

¹AVRDC – The World Vegetable Center, East and Southeast Asia, Kasetsart University, Kamphaeng Saen, Nakhon Pathom 73140, Thailand; ²Department of Vegetable Science, Punjab Agricultural University, Ludhiana 141 004, India; ³Institut National de la Recherche Agronomique (INRA), UR1052, Génétique et Amélioration des Fruits et Légumes, CS 60094, F-84143, Montfavet cedex, France; ⁴Instituto de Biología Molecular y Celular de Plantas (IBMCP) UPV-CSIC, Ciudad Politécnica de la Innovación, Edificio 8E, IngenieroFaustoElio s/n, 46022 Valencia, Spain; ⁵US Department of Agriculture, Agricultural Research Service, US Agricultural Research Station, 1636 E. Alisal St., Salinas, CA 93905, USA.

Abstract

Snampmelon (*Cucumis melo* L. *Momordica* Group; $2n=2x=24$) is native to India, where it is widely cultivated and is commonly called 'phut', which means to split. Immature fruits are cooked or eaten raw. In this paper we review the wealth of genetic resources in Indian snampmelon landraces for resistance to fungal and viral diseases, nematodes, and insects, and tolerance to drought, soil salinity, and high temperature. Global melon breeding programs have transferred many of these qualities into open-pollinated and hybrid cultivars of sweet melons cultivated in Africa, Asia, Australia, Europe, and the Americas. Snampmelon is a source of high fruit acidity, a trait that has been utilized to breed uniquely flavored melon cultivars. Resistance genes to combat pathogens and pests, and to strengthen crop resilience against climate change have been identified in snampmelon collections from various parts of India. More effort is needed to collect, characterize, evaluate and preserve snampmelon diversity in genebanks.

Keywords: disease resistance, pest resistance, molecular markers, resistance to abiotic stress, fruit quality

INTRODUCTION

Snampmelon is native to India, which is considered the center of domestication of melon by some researchers with the earliest melon remains at the Indus Valley site of Harappa dated between 2300 and 1600 BC (Vishnu-Mittre, 1974). It is widely cultivated in various Indian states such as Rajasthan, Gujarat, Punjab, Haryana, Uttar Pradesh, West Bengal and some other northeastern states. It is also cultivated in other countries of Southeast Asia, for instance Myanmar (Yi et al., 2009) and Vietnam. It is commonly called 'phut,' which means 'to split.' Fruit cracking is either longitudinal or starting in the middle of fruit, though in some instances only skin peeling (longitudinal or random) occurs (Dhillon et al., 2007). It is also known by other names such as 'phootkakari' or 'kakadia.' Several types of fruit shape are found in snampmelon: round, acorn, oblate, ovate, elongated, elliptical and pyriform (Dhillon et al., 2007). Fruit flesh color varies from cream and yellow to orange. Vines are monoecious. Immature fruits may be eaten raw or cooked, or pickled or dehydrated for off-season use. After removing the coat, seeds are used in bakery products and the traditional drink 'thandai'. Fruits are sources of vitamin C, iron and calcium (Goyal and Sharma, 2009). We review the wealth of genetic resources in Indian snampmelon landraces for resistance to fungal and viral diseases, nematodes and insects, tolerance to drought and salinity, genes for unique flavors, and the present status of genetic diversity of snampmelons in different parts of India.

^aE-mail: Narinder.Dhillon@worldveg.org



FUNGAL DISEASE RESISTANCE

Powdery mildew is a serious foliar disease of melon worldwide (Jahn et al., 2002). It affects the plant canopy and subsequently the fruit yield and quality. Cucurbit powdery mildew is mainly caused by two fungal species: *Podosphaera xanthii* (Castagne) Braun & Shishkoff (formerly *Sphaerotheca fuliginea*) and *Golovinomyces cichoracearum* (DC) V.P. Heluta (formerly *Erysiphe cichoracearum*); the first of the pathogen species is the most frequent on melon. More than 30 races of each pathogen have been identified on melon (Lebeda and Sedláková, 2004; McCreight et al., 2012). Powdery mildew was first noted on an epidemic scale in 1925 in the melon production area in the Imperial Valley of California, where it remained a serious problem for several successive years (Jagger, 1926; Jagger and Scott, 1937). A source of genetic resistance was identified in accession Calif. 525, which was a self-pollinated increase of 'Big Round' that was brought to the United States by D.N. Mehta (Second Economic Botanist, Nagpur Provinces, India) an Indian student of J.T. Rosa (Swarup, 2000; Kathleen R. Reitsma, pers. commun.; I.C. Jagger, unpublished pedigree note). This germplasm was collected from the Kathiawar region of Gujarat state in India. Powdery mildew-susceptible, orange flesh 'Hale's Best' melon (*Cucumis melo Reticulatus* Group) was crossed with Calif. 525 and the F₁ was resistant to the local strain of powdery mildew. A resistant F₂ selection that produced large elongated fruits was backcrossed to 'Hale's Best' to recombine powdery mildew resistance with superior horticultural qualities of 'Hale's Best'. Seven generations of inbreeding and selection led to the development and release of 'Powdery Mildew Resistant Cantaloupe No. 45' ('PMR 45') to the western United States melon industry in 1935 (Jagger and Scott, 1937).

Race 2 of *P. xanthii* overcoming the resistance of 'PMR 45' appeared in 'Imperial Valley' in 1938 (Jagger et al., 1938). Resistance to this new race was identified in 'Piria', donated by D. Mehta of Nagpur, Madhya Pradesh, India to the United States in 1929 and designated PI 79376 (Pryor et al., 1946; USDA ARS, 2014). Resistance to race 2 was combined with resistance to race 1 in melon cultivars 'PMR 5' and 'PMR 6', released in 1942, and 'Campo' and 'Jacumba' released in 1964 in the United States (Bohn et al., 1965). PI 124111 (MR-1) from Bihar state, India, was resistant to 26 races of *P. xanthii* (McCreight et al., 2012). This accession was also resistant to several races of *G. cichoracearum* (Pitrat et al., 1998; Lebeda et al., 2012).

Powdery mildew resistance genes identified in Calif. 525 and PI 79376 were introgressed to many melon breeding lines and cultivars (Harwood and Markarian, 1968) and are still prevalent in modern melon releases in the United States: 'Georgia 47' (Anon. 1954), 'Home Garden' (Ivanoff, 1957), 'Gulfstream' and 'Planter Jumbo' (Nugent, 1994), 'Mainstream' and 'Edisto 47' (Nugent et al., 1979), and recently released 'Chujuc' and 'Pacal' (Crosby et al., 2007, 2008). The moderately powdery mildew-resistant melon cultivar 'Punjab Sunehri', released in 1975 in Punjab state, India (Nandpuri et al., 1975; Waraitch et al., 1977), has 'Edisto' in its pedigree. This orange-fleshed, powdery mildew-resistant cultivar remained popular with Indian growers and consumers for two decades. Race 1 and race 2 powdery mildew-resistant *Reticulatus* Group cultivars 'PMR 5', 'Dulce', 'Gulfstream' and 'Jacumba' were resistant to a race-unknown population of powdery mildew in India in a controlled inoculation experiment (Waraitch et al., 1977). PI 414723 was also resistant to many races of *P. xanthii* (McCreight et al., 2012). Several race-specific genes have been described in these accessions (Dogimont, 2010-2011). Fergany et al. (2011) reported two additional snapmelon accessions resistant to powdery mildew: AM 22 to races 1 (strain Sm3) and 3 (strain 00Sm39) and AM 86 resistant to races 1 and 5 (strain 98Sm65). A large number of powdery mildew resistance genes have been described (Dogimont, 2010-2011); most of them are likely allelic. Several QTLs conferring resistance to different races have been mapped in melon Linkage Groups (LG) II, V and XII (Perchepped et al., 2005; Zhang et al., 2013).

Downy mildew caused by *Pseudoperonospora cubensis* (Berk. & Curtis) Rostov. is a common foliar disease of melons in humid production areas of the world. Six pathotypes have been identified: 1 and 2 in Japan, 3 and 6 in Israel, and 4 and 5 in the United States (Cohen et al., 2003). These pathotypes do not colonize *Luffa* ssp. (Thomas et al., 1987),

whereas the Indian and Chinese isolates of cucurbit downy mildew are able to colonize *Luffa* ssp. and are considered distinct races. Shetty et al. (2002) confirmed that the downy mildew races in the US are distinct from the race in Asia, whereas the race in Poland is similar to the races in the US. Four partially dominant resistance genes were identified in three accessions of snapmelon: PI 124111 (*Pc-1*, *Pc-2*), PI 414723 (*Pc-3*), and 5-4-2-1 (*Pc-5*) (Dogimont, 2010-2011). Resistance to *P. cubensis* races 3 and 6 in Israel has been found in PI 124111F controlled by two R genes, *At1* and *At2* (Taler et al., 2004). Interestingly, PI 124111F, reported resistant to six pathotypes of *P. cubensis*, was susceptible to an Indian isolate of *P. cubensis* (More, 2002), but IC 267353, IC 274029, KP7, and B-159 were resistant to this Indian isolate (Dhillon et al., 2007; Pandey et al., 2008). It will be interesting to test the reaction of these genotypes to the six pathotypes of *P. cubensis* available in the other parts of the world.

Melon Fusarium wilt is caused by the soil-borne fungal pathogen *Fusarium oxysporum* Schlechtend:Fr. f. sp. *melonis* (H.N. Hansen) W.C. Snyder & H.N. Hans (*Fom*). The pathogen survives in the soil as chlamydozoospores and is able to colonize crop residues and roots of most crops cultivated in rotation with melon (Gordon et al., 1989), thus rendering crop rotation as a limited tool to manage this disease. Soil solarization can reduce soil inoculum but is limited by local climate factors, i.e., temperature and relative humidity (Tamietti and Valentino, 2006) and it is not suitable for intensive vegetable farming systems where there is insufficient time for effective soil solarization. Grafting of susceptible melon scions onto Fusarium wilt-resistant rootstocks is an effective control strategy for melon Fusarium wilt, but the additional cost limits this approach to very high value melon cultivars. Use of resistant cultivars is regarded as the most effective strategy to control this disease. Melon Fusarium wilt isolates have been designated into four physiological races: 0, 1, 2, and 1.2. Two dominant resistance genes, *Fom-1* and *Fom-2*, control resistance to races 0 and 2, and 0 and 1, respectively and were identified in PI 124111F and its derivative MR-1 (Cohen and Eyal, 1987; Zink and Thomas, 1990). *Fom-2* also has been reported in PI 414723. Using MR-1 and PI 414723, these two genes have been cloned using chromosome walking strategies and belong to the NB-LRR family (TIR subfamily for *Fom-1* and non-TIR for *Fom-2*) (Joobeur et al., 2004; Brotman et al., 2013). These *Fom* genes are routinely deployed along with cucurbit powdery mildew resistance genes in modern melon commercial hybrids. Accession AM 27 exhibited uniform resistance to race 2 and segregated for resistance to race 1 (Fergany et al., 2011). Resistance to race 1.2 seems to have a complex genetic control that is hampering the development of reliable molecular markers and subsequent cloning (Oumouloud et al., 2013).

Melon Fusarium wilt and leafminer (*Liriomyza* spp.) are the most devastating disease and insect pests of melon in India. The prevailing melon cultivars grown by Indian farmers ('NS 7475', 'Punjab Sunehri', 'Punjab Hybrid 1', 'Durgapur Madhu', 'Kashi Madhu', 'Pusa Madhurus', 'Arka Jeet', 'Arka Rajhans') are susceptible to melon Fusarium wilt. Existing global melon genetic resources, including recent releases from seed companies, have been found susceptible in Indian field conditions (N.P.S. Dhillon, unpublished data; Arvind Kapur, pers. commun.).

There was a much lower incidence of melon Fusarium wilt or *Monosporascus* sudden wilt, which is incited by *Monosporascus cannonballus* (Pollack & Uecker), exhibited by snapmelon germplasm compared with 100% loss of muskmelon (*Reticulatus* Group) landraces and cultivars during melon germplasm collection expeditions in farmers' fields in the arid and semi-arid areas of Rajasthan and southern Punjab in India (N.P.S. Dhillon, unpublished data). Snapmelon accessions may have additional genes for resistance to melon Fusarium wilt and *M. cannonballus*.

Alternaria leaf blight of melons caused by *Alternaria cucumerina* (Ellis & Everh.) is widespread in wet and warm conditions (20 to 30°C) in areas with sandy soil, such as southern India and southeastern United States (Thomas, 1996). Resistance to this fungal pathogen is controlled by the single dominant gene *Ac* in MR-1, which was derived from PI 124111 (Thomas et al., 1990).



VIRAL DISEASE RESISTANCE

Numerous viruses affect melons worldwide. Three kinds of virus symptoms generally appear on the vines: 1) mosaic on leaves associated with leaf and fruit discolorations and deformation, 2) yellowing of leaves coupled with leaf thickening, and 3) necrotic spots or progressive necrosis resulting in vine death (Lecoq et al., 1998). Melon fields may be infected with more than one virus.

Cucumber mosaic virus (CMV) causes economic losses in melon worldwide. Resistance to CMV was first reported in accessions belonging to the *Conomon* Group from East Asia and is controlled by recessive oligogenes (Karchi et al., 1975; Dogimont et al., 2000; Essafi et al., 2009; Guiu-Aragonés et al., 2014). This resistance is not effective against all CMV strains and thus is not easy to use for the development of commercial F₁ hybrids. Resistance to a broad spectrum of CMV strains will likely need the combination of genes from different CMV resistance sources.

Snampmelon accessions AM 25, AM 82, IC 274014, SM 67, SM 72, SM 73, SM 82, MM 3974, MM 3982 and MM 3994 were highly resistant to CMV (Dhillon et al., 2007, 2009; Fergany et al., 2011; Malik et al., 2014). These accessions may contribute to a broad-based resistance against different strains of CMV prevailing in different parts of the world.

Zucchini yellow mosaic virus (ZYMV) is a serious virus of cucurbits worldwide (Desbiez and Lecoq, 1997). Three complementary, dominant genes in PI 414723 (*Zym-1*, *Zym-2* and *Zym-3*) impart resistance to ZYMV (Pitrat and Lecoq, 1984; Danin-Poleg et al., 1997). Accessions IC 274007, IC 274014, and PI 179905 are potentially useful sources of resistance to ZYMV (Dhillon et al., 2007).

Papaya ringspot virus watermelon strain (PRSV-W), formerly *Watermelon mosaic virus 1*, is a very common potyvirus in the tropics (Lecoq et al., 1980). Two alleles, *Prv¹* and *Prv²*, found in the weedy type melons PI 180280 and PI 180283, respectively, condition resistance to PRSV-W (Kaan, 1973; Webb, 1979; Pitrat and Lecoq, 1983). PI 414723 has the *Prv²* allele (M. Pitrat, unpublished data) and this gene has been recently isolated from the Indian accession PI 414723, encoding for a NBS-LRR type protein (Brotman et al., 2013). Nine accessions from northern India were heterogenous for resistance to PRSV-W: IC 267360, IC 267363, IC 267374, IC 267384, IC 274006, IC 274007, IC 274010, IC 274011 and IC 274013 (Dhillon et al., 2007). The genetic relationships between PRSV-W resistance genes in these accessions and *Prv* have not been established. Twenty-nine landraces from Kerala and Tamil Nadu states in southern India, exhibited necrotic symptoms in response to artificial inoculation with the potyvirus *Moroccan watermelon mosaic virus* (MWMV) (Fergany et al., 2011).

Watermelon mosaic virus (WMV), formerly *Watermelon mosaic virus 2*, is another widespread *Potyvirus* of melons. Genetic resistance to WMV was reported in PI 414723 (Munger, 1991) and is controlled by a single dominant gene, *Wmr* (Gilbert et al., 1994).

Cucurbit aphid-borne yellow virus (CABYV) is an aphid-transmitted *Polerovirus* of worldwide importance to melons. 'Faizabadiphoot' and PI 414723 were reported resistant to CABYV (Dogimont et al., 1997).

Watermelon chlorotic stunt virus (WmCSV) is an economically important *Geminivirus* in Yemen, Sudan, and Iran (Yousif et al., 2007). PI 414723 provided resistance during graft inoculation experiments and multiple field trials in Sudan (Yousif et al., 2007).

Cucurbit leaf crumple virus (CuLCrV) is a sweet potato whitefly-transmitted *Begomovirus* of melon that has appeared in commercial melon fields in the southwestern US, western Mexico, and Central America since 1977. PI124111, PI 179901, and PI 414723 exhibited partial resistance to CuLCrV in naturally infected field and controlled inoculation greenhouse tests. Resistance in PI 313970 (*Acidulus* Group) was conditioned by a single recessive gene and appeared allelic to that in the snampmelon accessions (McCreight et al., 2008).

Cucumber green mottle mosaic virus (CGMMV) is an economically significant *Tobamovirus* in greenhouse production (Hollings et al., 1975) that has been reported in Europe and Asia. A biological vector of this melon virus is unknown, but CGMMV is transmitted mechanically and through growing media (Lecoq et al., 1998). In the early

1980s, CGMMV affected 70 to 80% of plants in peri-urban cucurbit fields of Delhi, India (Raychaudhuri and Varma, 1978). Identification of resistance to CGMMV in 'Phoot' led to the development of five Indian lines (VRM 5-10, VRM 29-1, VRM 31-1-2, VRM 42-4, and VRM 43-6) that had high-level resistance to CGMMV along with improved yield and sweetness (More et al., 1993).

Kyuri green mottle mosaic virus (KGMMV) is an economically significant *Tobamovirus* in Japan, Korea, and Indonesia (Daryono et al., 2005). It is mechanically transmitted and seed-borne. PI 414723 is resistant to KGMMV (Daryono et al., 2005).

Spring (dry season) melons in the trans-Gangetic plains of India are threatened by CMV, CGMMV, SqMV, PRSV, and ZYMV, whereas whitefly transmitted begomoviruses predominate during the rainy season in this region (Sharma et al., 2007). Landrace IC 274014 is an asymptomatic host of CMV that also exhibited field resistance to an unidentified *Begomovirus* (Sharma and Kang, 2009).

ROOT-KNOT NEMATODE AND INSECT RESISTANCE

Root-knot nematode, *Meloidogyne* spp., is found in melon fields worldwide, particularly in sandy soils. Its impact on melon yield depends upon the nematode population density in the field. Current melon cultivars are susceptible to root-knot nematode. High-level resistance to *M. incognita* has been identified in landrace IC 274023 (Dhillon et al., 2007) which should be exploited to develop the first root-knot nematode-resistant melon cultivar.

Melon aphid, also called cotton melon aphid, *Aphis gossypii* Glover, is found throughout most of the temperate, subtropical and tropical regions of the world. Younger plants are more susceptible to feeding. The melon aphid is also an efficient vector of viruses including CMV and potyviruses. Strong resistance to cotton melon aphid biotype D (McCreight et al., 1992) available in PI 414723 was used in conventional breeding to develop orange-fleshed melon breeding lines 'AR Topmark', 'AR-5' and 'AR Hale's Best Jumbo' (McCreight et al., 1984). These breeding lines exhibited different levels of resistance to virus transmission by CMA (Kishaba et al., 1992). PI 414723 has three components of resistance to cotton melon aphid: antibiosis, antixenosis, and tolerance (Bohn et al., 1972). Tolerance to cotton melon aphid in PI 414723 is expressed as freedom from curling of leaves and is governed by the single dominant gene *Ag* (Bohn et al., 1973). Snapmelon landraces IC 267353, IC 267384, and IC 274010 have resistance to virus transmission by cotton melon aphid (Dhillon et al., 2007).

Cucumber beetles infest melon seedlings and fruit (Kishaba et al., 1998). Seedling and fruit resistance to western striped cucumber beetle [*Acalymma trivittata* (Mannerheim)] and spotted cucumber beetle [*Diabrotica undecimpunctata undecimpunctata* (Mannerheim)] has been identified in PI 414723 but the genetic basis of resistance was not determined.

Sweet potato whitefly, *Bemisia tabaci* Gennadius, is another economically important insect pest in the desert southwestern United States. It has several biotypes: biotype A is a vector of *Lettuce infectious yellow virus* (LIYV) whereas biotype B is a vector of *Cucurbit yellow stunting disorder virus* (CYSDV) and *Cucurbit leaf crumple virus* (CuLCrV). Resistance to *B. tabaci* biotype B was reported in PI 414723 (Boissot et al., 2003).

MELON FLAVOR ENRICHMENT

Accumulation of sugar and acid in melon fruit imparts unique taste and flavor. Fruit acidity of Indian commercial melons ranges from 0.12 to 0.2% (N.P.S. Dhillon, unpublished data). High acidity sources have been reported in snapmelon landraces IC 274021 (0.61%) and IC 267360 (0.57%) (Dhillon et al., 2007). The low-pH gene derived from Indian snapmelon accession IND-35 was exploited using marker-assisted selection by Syngenta in 2008 to develop the pleasant-tasting F₁ hybrid melon 'GWANIPA' that was commercialized in UK, Germany and Holland by Kernel Export (Jordi Garcia-Mas, pers. commun.). This melon has a lemon flavor and contains 700-800 mg citric acid per 100 g fresh weight (FW) with a pH level of 4.5 (patent no. EP 1587933 B1) (Casanueva et al., 2010).



VITAMIN AND MINERAL CONTENT

Higher concentrations (up to 34.1 mg 100 g⁻¹ FW) of vitamin C were detected in the snapmelon landraces of northern India compared to the germplasm from eastern India (up to 19.4 mg 100 g⁻¹ FW) and southern India (up to 9.0 mg 100 g⁻¹ FW) (Dhillon et al., 2007; Fergany et al., 2011; Malik et al., 2014).

Iron and zinc deficiencies are recognized as a nutritional problem worldwide (Uauy et al., 2006). Wide variation for phosphorus (2.6 to 21.4 mg 100 g⁻¹ FW), potassium (19.7 to 232.4 mg 100 g⁻¹ FW), iron (0.5 to 0.89 mg 100 g⁻¹ FW) and zinc (0.12 to 0.68 mg 100 g⁻¹ FW) have been identified in landraces (Fergany et al., 2011). This genetic variation for vitamins and minerals is important for breeding of new mineral- and vitamin-rich snapmelon cultivars. Snapmelons are consumed by poor and middle-class consumers, and the fruit is available in the market for nearly five months of the spring and rainy season. Snapmelon fruits were used as food in the two Japanese islands (Hachijo and Fukue) during the two World Wars (Fujishita, 2004).

SNAPMELON GENETIC DIVERSITY

Based on the variability at nine simple sequence repeat loci (160 alleles, polymorphism information content value 0.81), clear genetic differentiation was observed among gene pools of snapmelon germplasm from northern, southern and eastern regions of India (Dhillon et al., 2013). Global melon reference populations were distinct from this germplasm; clearly snapmelon landraces possessed unique alleles when compared with international reference accessions. Snapmelon germplasm offers opportunities to widen the genetic base of melons in the secondary centers of diversity (eastern Asia, western Mediterranean area) and proximal parts (e.g., Turkey) of the primary center of diversity.

SNAPMELONS FOR DEVELOPING CLIMATE-SMART MELONS

To maintain and increase crop productivity in increasingly hostile environments, novel sources of genetic variation must be sought for adapting crops to unstable climates. Snapmelons have little-explored gene pools that are readily available to the sweet melon gene pools through conventional hybridization. Snapmelons are drought-hardy, and are cultivated by small-scale farmers in arid and semi-arid regions of India during the rainy season (Pareek and Samadia, 2002). Two highly drought- and heat-tolerant snapmelon selections, AHS 10 and AHS 82, were bred from local landraces from the arid region of Rajasthan, India (Pareek and Samadia, 2002). Snapmelon accession RSM 50 was highly drought tolerant compared to *Reticulatus* Group cultivars in a controlled (water deficit) irrigation field experiment (Dhillon et al., 2013). Accession Calif. 525 was acknowledged for contributing high levels of tolerance to salt and high temperature along with resistance to powdery mildew in 'PMR 45', the first modern western US shipping-type melon (Jagger and Scott, 1937). Grafted melons are more tolerant of salinity than non-grafted controls (Orsini et al., 2013). Snapmelon landraces from the coastal, arid areas of India (Gujarat, Karnataka, Kerala, Andhra Pradesh, Tamil Nadu, Odisha) are potential sources of new salt-, drought- and heat-tolerant rootstocks for melon grafting.

CONCLUSION

Snapmelons originated in India and have provided yeoman service to sweet melon breeding programs worldwide. For example, PI 414723 was used in breeding as source of resistance to eight fungal and viral diseases and two insect pests. PI 124111 (MR-1) is resistant to powdery and downy mildew, Alternaria and Fusarium wilt. More than 1000 snapmelon accessions are maintained in four national genebanks in India (National Bureau of Plant Genetic Resources, New Delhi; Central Institute for Arid Horticulture, Bikaner; Central Horticultural Experiment Station, Bhubaneswar; Punjab Agricultural University, Ludhiana), but they have not been comprehensively evaluated against various biotic and abiotic stresses. Erosion of snapmelon genetic diversity in India is a real threat because of rapid urbanization, and swift adoption and spread of commercial F₁ hybrid sweet melons across India. International collaborative research efforts should be launched to collect,

characterize and evaluate the snapmelon germplasm available in India's diverse agro-ecological regions. This will result in identification of unique and useful genes to broaden the narrow gene pool of sweet melons, provide new sources for resistance to various diseases and pests of melons, and help meet the challenges of climate change to the sustainable production of melons worldwide.

Literature cited

- Anon. (1954). New vegetable varieties. Proc. Am. Soc. Hort. Sci. 63, 503–525.
- Bohn, G.W., Davis, G.N., Foster, R.E., and Whitaker, T.W. (1965). Campo and Jacumba. New cantaloupe varieties for the southwest. Calif. Agric. 19, 8–10.
- Bohn, G.W., Kishaba, A.N., and Toba, H.H. (1972). Mechanisms of resistance to melon aphid in a muskmelon line. HortScience 7, 281–282.
- Bohn, G.W., Kishaba, A.N., Principe, J.A., and Toba, H.H. (1973). Tolerance to melon aphid in *Cucumis melo* L. J. Am. Soc. Hortic. Sci. 98, 37–40.
- Boissot, N., Lafortune, D., Pavis, C., and Sauvion, N. (2003). Field resistance to *Bemisia tabaci* in *Cucumis melo*. HortScience 38, 77–80.
- Brotman, Y., Normantovich, M., Goldenberg, Z., Zvirin, Z., Kovalski, I., Stovbun, N., Doniger, T., Bolger, A.M., Troadec, C., Bendahmane, A., et al. (2013). Dual resistance of melon to *Fusarium oxysporum* races 0 and 2 and to *Papaya ring-spot virus* is controlled by a pair of head-to-head-oriented NB-LRR genes of unusual architecture. Mol. Plant 6, 235–238 <http://dx.doi.org/10.1093/mp/sss121>.
- Casanueva, A., Foncelle, B., Nicolet, J.E., Van Doon, J.E., and Oliver, S.M. (2010). Novel melon plants. Patent no. 20100034952. <http://www.faqs.org/patents/app/20100034952>.
- Cohen, Y., and Eyal, H. (1987). Downy mildew, powdery mildew and *Fusarium* wilt-resistant muskmelon breeding line PI 124111F. Phytoparasitica 15, 187–195 <http://dx.doi.org/10.1007/BF02979582>.
- Cohen, Y., Meron, I., Mor, N., and Zuriel, S. (2003). A new pathotype of *Pseudoperonospora cubensis* causing downy mildew in cucurbits in Israel. Phytoparasitica 31, 458–466 <http://dx.doi.org/10.1007/BF02979739>.
- Crosby, K., Jifon, J., and Leskovar, D. (2007). 'Pacal' orange-casaba, and 'Chujuc' western-shipper cantaloupe: two new melon cultivars from the Texas Agricultural Experiment Station. HortScience 42, 1013.
- Crosby, K.M., Jifon, J.L., and Leskovar, D.I. (2008). 'Chujuc', a new powdery mildew-resistant U.S. western-shipper melon with high sugar and β -carotene content. HortScience 43, 1904–1906.
- Danin-Poleg, Y., Paris, H.S., Cohen, S., Rabinowitch, H.D., and Karchi, Z. (1997). Oligogenic inheritance of resistance to *Zucchini yellow mosaic virus* in melons. Euphytica 93, 331–337 <http://dx.doi.org/10.1023/A:1002944432083>.
- Daryono, B.S., Somowiyarjo, S., and Natsuaki, K.T. (2005). Screening for resistance to *Kyuri green mottle mosaic virus* in various melons. Plant Breed. 124, 487–490 <http://dx.doi.org/10.1111/j.1439-0523.2005.01121.x>.
- Desbiez, C., and Lecoq, H. (1997). *Zucchini yellow mosaic virus*. Plant Pathol. 46, 809–829 <http://dx.doi.org/10.1046/j.1365-3059.1997.d01-87.x>.
- Dhillon, N.P.S., Ranjana, R., Singh, K., Eduardo, I., Monforte, A.J., Pitrat, M., Dhillon, N.K., and Singh, P.P. (2007). Diversity among landraces of Indian snapmelon (*Cucumis melo* var. *momordica*). Genet. Resour. Crop Evol. 54, 1267–1283 <http://dx.doi.org/10.1007/s10722-006-9108-2>.
- Dhillon, N.P.S., Singh, J., Fergany, M., Monforte, A.J., and Sureja, A.K. (2009). Phenotypic and molecular diversity among landraces of snapmelon (*Cucumis melo* var. *momordica*) adopted to the hot and humid tropics of eastern India. Plant Genet. Resour. 7, 291–300 <http://dx.doi.org/10.1017/S1479262109990050>.
- Dhillon, N.P.S., Monforte, A.J., Pitrat, M., Pandey, S., Singh, P.K., Reitsma, K.R., Garcia-Mas, J., Sharma, A., and McCreight, J.D. (2013). Melon landraces of India: contributions and importance. Plant Breed. Rev. 35, 85–150.
- Dogimont, C. (2010–2011). Gene list 2011 for melon. Cucurbit Genet. Coop. Rep. 33–34, 104–133.
- Dogimont, C., Bussemakers, A., Martin, J., Slama, S., Lecoq, H., and Pitrat, M. (1997). Two complementary recessive genes conferring resistance to *Cucurbit aphid borne yellows luteovirus* in an Indian melon line (*Cucumis melo* L.). Euphytica 96, 391–395 <http://dx.doi.org/10.1023/A:1003069703294>.
- Dogimont, C., Lecomte, L., Périn, C., Thabuis, A., Lecoq, H., and Pitrat, M. (2000). Identification of QTLs contributing to resistance to different strains of *Cucumber mosaic cucumovirus* in melons. Acta Hort. 510, 391–398 <http://dx.doi.org/10.17660/ActaHortic.2000.510.62>.

- Essafi, A., Díaz-Pendón, J.A., Moriones, E., Monforte, A.J., Garcia-Mas, J., and Martín-Hernández, A.M. (2009). Dissection of the oligogenic resistance to *Cucumber mosaic virus* in the melon accession PI 161375. *Theor. Appl. Genet.* *118*, 275–284 <http://dx.doi.org/10.1007/s00122-008-0897-x>.
- Fergany, M., Kaur, B., Monforte, A.J., Pitrat, M., Rys, C., Lecoq, H., Dhillon, N.P.S., and Dhaliwal, S.S. (2011). Variation in melon (*Cucumis melo* L.) landraces adapted to the humid tropics of southern India. *Genet. Resour. Crop Evol.* *58*, 225–243 <http://dx.doi.org/10.1007/s10722-010-9564-6>.
- Fujishita, N. (2004). Pedigree of melon (*Cucumis melo* L.) in ancient Japan. Paper presented at: International Symposium on Asian Plant Diversity and Systematics (Sakura, Japan).
- Fukino, N., Ohara, T., Monforte, A.J., Sugiyama, M., Sakata, Y., Kunihisa, M., and Matsumoto, S. (2008). Identification of QTLs for resistance to powdery mildew and SSR markers diagnostic for powdery mildew resistance genes in melon (*Cucumis melo* L.). *Theor. Appl. Genet.* *118*, 165–175 <http://dx.doi.org/10.1007/s00122-008-0885-1>.
- Gilbert, R.Z., Kyle, M.M., Munger, H.M., and Gay, S.M. (1994). Inheritance of resistance to *Watermelon mosaic virus* in *Cucumis melo* L. *HortScience* *29*, 107–110.
- Gordon, T.R., Okamoto, D., and Jacobson, D.J. (1989). Colonization of muskmelon and non-host crops by *Fusarium oxysporum* f.sp. *melonis* and other species of *Fusarium*. *Phytopathology* *79*, 1095–1100 <http://dx.doi.org/10.1094/Phyto-79-1095>.
- Goyal, M., and Sharma, S.K. (2009). Traditional wisdom and value addition prospects of arid foods of desert regions of North West India. *Indian J. Traditional Knowledge* *8*, 581–585.
- Guiu-Aragonés, C., Monforte, A.J., Saladié, M., Corrêa, R.X., Garcia-Mas, J., and Martín-Hernández, A.M. (2014). The complex resistance to *Cucumber mosaic cucumovirus* (CMV) in the melon accession PI 161375 is governed by one gene and at least two quantitative trait loci. *Mol. Breed.* *34*, 351–362 <http://dx.doi.org/10.1007/s11032-014-0038-y>.
- Harwood, R.R., and Markarian, D. (1968). A genetic survey of resistance to powdery mildew in muskmelon. *J. Hered.* *59*, 213–217.
- Hollings, M., Komuro, Y., and Tochigara, H. (1975). *Cucumber green mottle mosaic virus*. Descriptions of plant viruses No. 154. (Kew, Surrey, UK: Commonwealth Mycol. Inst./Assn. Appl. Biol.) www.dpvweb.net/dpv/showdpv.php?dpvno=154.
- Ivanoff, S.S. (1957). The Home garden cantaloupe, a variety with combined resistance to downy mildew, powdery mildew, and aphids. *Phytopathology* *47*, 552–556.
- Jagger, I. (1926). Powdery mildew of muskmelon in the Imperial valley of California in 1925. *Phytopathology* *16*, 1009–1010.
- Jagger, I.C., and Scott, G.W. (1937). Development of powdery mildew resistant cantaloupe No 45. *USDA Circ.* *441*, 6.
- Jagger, I.C., Whitaker, T.W., and Porter, D.R. (1938). Inheritance in *Cucumis melo* of resistance to powdery mildew (*Erysiphe cichoracearum*). *Phytopathology* *28*, 761.
- Jahn, M., Munger, H.M., and McCreight, J.D. (2002). Breeding cucurbit crops for powdery mildew resistance. In *The Powdery Mildews: a Comprehensive Treatise*, R.R. Bélanger, W.R. Bushnell, A.J. Dik, and T.L.W. Carver, eds. (St. Paul, USA: APS Press), p.239–248.
- Joobeur, T., King, J.J., Nolin, S.J., Thomas, C.E., and Dean, R.A. (2004). The *Fusarium* wilt resistance locus *Fom-2* of melon contains a single resistance gene with complex features. *Plant J.* *39*, 283–297 <http://dx.doi.org/10.1111/j.1365-313X.2004.02134.x>.
- Kaan, J.F. (1973). Recherches sur la résistance du melon aux maladies, notamment à la mosaïque de la pastèque et au *Pseudoperonospora*, appliquées au type variétal 'Cantaloup Charentais'. Paper presented at: EUCARPIA Meeting on Melon (Avignon, France).
- Karchi, Z., Cohen, S., and Govers, A. (1975). Inheritance of resistance to *Cucumber mosaic virus* in melons. *Phytopathology* *65*, 479–481 <http://dx.doi.org/10.1094/Phyto-65-479>.
- Kishaba, A.N., Castle, S., Coudriet, D.L., McCreight, J.D., and Bohn, G.W. (1992). Virus transmission by *Aphis gossypii* Glover to aphid-resistant and susceptible muskmelons. *J. Am. Soc. Hortic. Sci.* *117*, 248–254.
- Kishaba, A.N., Castle, S.I., Coudriet, D.L., McCreight, J.D., and Bohn, G.W. (1998). Resistance to western spotted and striped cucumber beetle in melon. Paper presented at: *Cucurbitaceae' 98: Evaluation and Enhancement of Cucurbit Germplasm* (Pacific Grove, California, USA: ASHS Press).
- Lebeda, A., and Sedláková, B. (2004). Disease impact and pathogenicity variation in Czech populations of cucurbit powdery mildews. Paper presented at: *Cucurbitaceae 2004*, Proceedings of the VIIIth EUCARPIA meeting on

Genetics and Breeding of *Cucurbitaceae* (Palacký University, Olomouc, Czech Republic).

Lebeda, A., Sedláková, B., Křístková, E., Vajdová, M., and McCreight, J.D. (2012). Application of a new approach for characterization and denomination of races of cucurbit powdery mildews – a case study of Czech pathogen populations. Paper presented at: *Cucurbitaceae* 2012, Proceedings of the Xth EUCARPIA meeting on Genetics and Breeding of *Cucurbitaceae* (Antalya, Turkey).

Lecoq, H., Labonne, G., and Pitrat, M. (1980). Specificity of resistance to virus transmission by aphids in *Cucumis melo*. *Ann. Phytopathol.* 12, 139–144.

Lecoq, H., Wisler, G., and Pitrat, M. (1998). Cucurbit viruses: the classic and the emerging. Paper presented at: *Cucurbitaceae* '98: Evaluation and Enhancement of Cucurbit Germplasm (Pacific Grove, California, USA: ASHS Press).

Malik, A.A., Vashisht, V.K., Singh, K., Sharma, A., Singh, D.K., Singh, H., Monforte, A.J., McCreight, J.D., and Dhillon, N.P.S. (2014). Diversity among melon (*Cucumis melo* L.) landraces from the Indo-Gangetic plains of India and their genetic relationship with USA melon cultivars. *Genet. Resour. Crop Evol.* 61, 1189–1208 <http://dx.doi.org/10.1007/s10722-014-0101-x>.

McCreight, J.D., Kishaba, A.N., and Bohn, G.W. (1984). AR Hale's Best Jumbo, AR 5, and AR Topmark, melon aphid-resistant muskmelon breeding lines. *HortScience* 19, 309–310.

McCreight, J.D., Bohn, G.W., and Kishaba, A.N. (1992). "Pedigree" of PI 414723 melon. *Cucurbit Genet. Coop. Rpt.* 15, 51–52.

McCreight, J.D., Liu, H.-Y., and Turini, T.A. (2008). Genetic resistance to *Cucurbit leaf crumple virus* in melon. *HortScience* 43, 122–126.

McCreight, J.D., Coffey, M.D., Sedláková, B., and Lebeda, A. (2012). Cucurbit powdery mildew of melon incited by *Podosphaera xanthii*: Global and Western US perspectives. Paper presented at: *Cucurbitaceae* 2012, Xth EUCARPIA meeting on Genetics and Breeding of *Cucurbitaceae* (Antalya, Turkey).

More, T.A. (2002). Enhancement of muskmelon resistance to disease via breeding and transformation. *Acta Hort.* 588, 205–211 <http://dx.doi.org/10.17660/ActaHortic.2002.588.31>.

More, T.A., Varma, A., Seshadri, V.S., Somkuwar, R.G., and Rajamony, L. (1993). Breeding and development of *Cucumber green mottle mosaic virus* (CGMMV) resistant lines in melon (*Cucumis melo* L.). *Cucurbit Genet. Coop. Rpt.* 16, 44–46.

Munger, H.M. (1991). Progress in breeding melons for watermelon mosaic resistance. *Cucurbit Genet. Coop. Rept.* 14, 43–44.

Nandpuri, K.S., Singh, S., and Lal, T. (1975). Punjab Sunehri: a new variety of muskmelon. *Progressive Farming* November issue, p.21–23.

Nhi, P.T.P., Akashi, Y., Hang, T.T.M., Tanaka, K., Aierken, Y., Yamamoto, T., Nishida, H., Long, C., and Kato, K. (2010). Genetic diversity in Vietnamese melon landraces revealed by the analyses of morphological traits and nuclear and cytoplasmic molecular markers. *Breed. Sci.* 60, 255–266 <http://dx.doi.org/10.1270/jsbbs.60.255>.

Nugent, P.E. (1994). Tetraploid 'Planters Jumbo' melon lines C883-M6-4X and 67-M6-1004X. *HortScience* 29, 48–49.

Nugent, P.E., Hoffman, J.C., and Andrus, C.F. (1979). 'Mainstream' muskmelon. *HortScience* 14, 192.

Orsini, F., Sanoubar, R., Oztekin, G.B., Kappel, N., Tepecik, M., Quacquarelli, C., Tuzel, Y., Bona, S., and Gianquinto, G. (2013). Improved stomatal regulation and ion partitioning boosts salt tolerance in grafted melons. *Funct. Plant Biol.* 40, 628–636 <http://dx.doi.org/10.1071/FP12350>.

Oumouloud, A., El-Otmani, M., Chikh-Rouhou, H., Claver, A.G., Torres, R.G., Perl-Treves, R., and Alvarez, J.M. (2013). Breeding melon for resistance to *Fusarium* wilt: recent developments. *Euphytica* 192, 155–169 <http://dx.doi.org/10.1007/s10681-013-0904-4>.

Pandey, S., Rai, M., Prassanna, H.C., and Kallou, G. (2008). Kashi Madhu: a new muskmelon cultivar with high total soluble solids. *HortScience* 43, 245–246.

Pareek, O.P., and Samadia, O.K. (2002). Promising indigenous cucurbit varieties. *Indian Hort.* July-Sept. ,p.15–18.

Perchepied, L., Bardin, M., Dogimont, C., and Pitrat, M. (2005). Relationship between Loci conferring downy mildew and powdery mildew resistance in melon assessed by quantitative trait Loci mapping. *Phytopathology* 95, 556–565 <http://dx.doi.org/10.1094/PHYTO-95-0556>.

Pitrat, M., and Lecoq, H. (1983). Two alleles for *Watermelon mosaic virus 1* resistance in melon. *Cucurbit Genet. Coop. Rept.* 6, 52–53.

Pitrat, M., and Lecoq, H. (1984). Inheritance of *Zucchini yellow mosaic virus* resistance in *Cucumis melo* L.



- Euphytica 33, 57–61 <http://dx.doi.org/10.1007/BF00022750>.
- Pitrat, M., Dogimont, C., and Bardin, M. (1998). Resistance to fungal diseases of foliage in melon. Paper presented at: *Cucurbitaceae '98: Evaluation and Enhancement of Cucurbit Germplasm* (Pacific Grove, California, USA: ASHS Press).
- Pryor, D.E., Whitaker, T.W., and Davis, G.N. (1946). The development of powdery mildew resistant cantaloupes. *Proc. Am. Soc. Hort. Sci.* 47, 347–356.
- Raychaudhuri, M., and Varma, A. (1978). Mosaic diseases of muskmelon, caused by a minor variant of *Cucumber green mottle mosaic virus*. *Phytopathol. Z.* 93, 120–125 <http://dx.doi.org/10.1111/j.1439-0434.1978.tb03645.x>.
- Sharma, A., and Kang, S.S. (2009). Cucurbits: a new host for *begomovirus* in Punjab. *Plant Dis.* 24, 51–53.
- Sharma, A., Kang, S.S., Thiara, S.K., and Kaur, S.I. (2007). Occurrence of cucurbit viruses in Punjab. Paper presented at: 10th International Plant Virus Epidemiology Symposium on Controlling Epidemics of Emerging and established Plant Virus Diseases - The Way Forward (Hyderabad, India: ICRISAT).
- Shetty, N.V., Wehner, T.C., Thomas, C.E., Doruchowski, R.W., and Shetty, K.P.V. (2002). Evidence for downy mildew races in cucumber tested in Asia, Europe, and North America. *Sci. Hortic. (Amsterdam)* 94, 231–239 [http://dx.doi.org/10.1016/S0304-4238\(02\)00013-4](http://dx.doi.org/10.1016/S0304-4238(02)00013-4).
- Swarup, V. (2000). Genetic resources in vegetable crops in India. In *Emerging Scenario in Vegetable Research and Development*, G. Kallo, and K. Singh, eds. (New Delhi, India: Research Periodicals and Book Publishing Home), p.346–355.
- Taler, D., Galperin, M., Benjamin, I., Cohen, Y., and Kenigsbuch, D. (2004). Plant eR genes that encode photorespiratory enzymes confer resistance against disease. *Plant Cell* 16, 172–184 <http://dx.doi.org/10.1105/tpc.016352>.
- Tamietti, G., and Valentino, D. (2006). Soil solarization as an ecological method for the control of *Fusarium* wilt of melon in Italy. *Crop Prot.* 25, 389–397 <http://dx.doi.org/10.1016/j.cropro.2005.07.002>.
- Thomas, C.E. (1996). *Alternaria* leaf blight. In *Compendium of Cucurbit Diseases*, T.A. Zitter, D.L. Hopkins, and C.E. Thomas, eds. (St. Paul, USA: APS Press), p.23–24.
- Thomas, C.E., Inaba, T., and Cohen, Y. (1987). Physiological specialization in *Pseudoperonospora cubensis*. *Phytopathology* 77, 1621–1624 <http://dx.doi.org/10.1094/Phyto-77-1621>.
- Thomas, C.E., McCreight, J.D., and Jourdain, E.L. (1990). Inheritance of resistance to *Alternaria cucumerina* in *Cucumis melo* line MR-1. *Plant Dis.* 74, 868–870 <http://dx.doi.org/10.1094/PD-74-0868>.
- Uauy, C., Distelfeld, A., Fahima, T., Blechl, A., and Dubcovsky, J. (2006). A NAC Gene regulating senescence improves grain protein, zinc, and iron content in wheat. *Science* 314, 1298–1301 <http://dx.doi.org/10.1126/science.1133649>.
- USDA ARS. (2014). National Genetic Resources Program. Germplasm Resources Information Network (GRIN). (Online database). National Germplasm Resources Laboratory, Beltsville, MD. <http://www.ars-grin.gov/cgi-bin/npgs/acc/display.pl?1743283>.
- Vishnu-Mittre. (1974). Palaeobotanical evidence in India. In *Evolutionary Studies in World Crops, Diversity and Changes in Indian Subcontinent*, J. Hutchinson, ed. (Cambridge, UK: Cambridge Univ. Press), p.3–30.
- Waraitch, K.S., Munshi, G.D., Nandpuri, K.S., and Lal, T. (1977). Screening of muskmelon, wild melon and snapmelon for resistance to powdery mildew (*Sphaerotheca fuliginea*). *Phytopathol. Mediterr.* 16, 37–39.
- Webb, R.E. (1979). Inheritance of resistance to watermelon mosaic virus in *Cucumis melo* L. *HortScience* 14, 265–266.
- Yi, S.S., Akashi, Y., Tanaka, K., Cho, T.T., Khaing, M.T., Yoshino, H., Nishida, H., Yamamoto, T., Win, K., and Kato, K. (2009). Molecular analysis of genetic diversity in melon landraces (*Cucumis melo* L.) from Myanmar and their relationship with melon germplasm from East and South Asia. *Genet. Resour. Crop Evol.* 56, 1149–1161 <http://dx.doi.org/10.1007/s10722-009-9438-y>.
- Yousif, M.T., Kheyr-Pour, A., Gronenborn, B., Pitrat, M., and Dogimont, C. (2007). Sources of resistance to *Watermelon chlorotic stunt virus*. *Plant Breed.* 126, 422–427 <http://dx.doi.org/10.1111/j.1439-0523.2007.01366.x>.
- Zhang, C., Ren, Y., Guo, S., Zhang, H., Gong, G., Du, Y., and Xu, Y. (2013). Application of comparative genomics in developing markers tightly linked to the *Pm-2F* gene for powdery mildew resistance in melon (*Cucumis melo* L.). *Euphytica* 190, 157–168 <http://dx.doi.org/10.1007/s10681-012-0828-4>.
- Zink, F.W., and Thomas, C.E. (1990). Genetics of resistance to *Fusarium oxysporum* f.sp. *melonis* races 0, 1, and 2 in muskmelon line MR-1. *Phytopathology* 80, 1230–1232 <http://dx.doi.org/10.1094/Phyto-80-1230>.