

# Dynamics of particles with anisotropic mass depending on time and position

T. S. Ratiu, O. G. Smolyanov

► **To cite this version:**

T. S. Ratiu, O. G. Smolyanov. Dynamics of particles with anisotropic mass depending on time and position. / Doklady Mathematics, MAIK Nauka/Interperiodica, 2015, 92 (3), pp.723-726. 10.1134/S1064562415060241 . hal-01398414

**HAL Id: hal-01398414**

**<https://hal.archives-ouvertes.fr/hal-01398414>**

Submitted on 17 Nov 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Dynamics of Particles with Anisotropic Mass Depending on Time and Position

T. S. Ratiu<sup>a</sup> and O. G. Smolyanov<sup>b</sup>

Presented by Academician V.V. Kozlov January 22, 2015

**Abstract**—Representations of solutions of equations describing the diffusion and quantum dynamics of particles in a Riemannian manifold are discussed under the assumption that the mass of particles is anisotropic and depends on both time and position. These equations are evolution differential equations with second-order elliptic operators, in which the coefficients depend on time and position. The Riemannian manifold is assumed to be isometrically embedded into Euclidean space, and the solutions are represented by Feynman formulas; the representation of a solution depends on the embedding.

We consider the following problem. Suppose that  $E = \mathbb{R}^n$ ,  $S$  is a compact Riemannian submanifold in  $E$  and  $A$  is a Borel function on  $E$  taking values in the space  $\mathcal{L}(E)$  (given a Banach space  $X$ ,  $\mathcal{L}(X)$  denotes the Banach space of all continuous linear operators on  $X$ ). By  $\Delta_A$  we denote the closed linear operator on the (real or complex) space  $\mathfrak{L}_p(E)$ ,  $1 \leq p < \infty$ , defined by  $(\Delta_A \varphi)(x) := \operatorname{tr} A(x) \varphi''(x)$  for functions  $\varphi$  (assumed to be infinitely differentiable) which form a dense subspace of  $\mathfrak{L}_p(E)$ . In what follows, we also define operators  $\Delta_A^S$  on  $\mathfrak{L}_p(S)$  associated with  $\Delta_A$ .

Let  $G$  be a function on  $[0, a)$ ,  $a > 0$ , taking values in the space of  $\mathcal{L}(E)$ -valued Borel functions on  $E$ . We shall obtain Feynman formulas for the evolution families of operators (a definition is given below) on  $\mathfrak{L}_p(S)$  generated by the families  $\Delta_{G(t)}^S$  and  $i\Delta_{G(t)}^S$  of operators associated with the families  $\Delta_{G(t)}$  and  $i\Delta_{G(t)}$ .

The method which we use can be described as follows. If  $A(x) = \mathcal{I}$  for each  $x$ , where  $\mathcal{I}$  is the identity self-mapping of  $E$ , then  $\Delta_A$  is the usual Laplacian acting on functions defined on  $E$ , and the associated operator  $\Delta_A^S$ , which acts on functions on  $S$ , is the Laplace–Beltrami operator. In the situation under consideration, it can be obtained as follows. For each  $\tau > 0$ , let  $F(\tau)$  be the integral operator on  $\mathfrak{L}_p(S)$  defined

$$\text{by } (F(\tau)\varphi)(x) := c(\tau, x) \int_S e^{-\frac{(z-x)^2}{2\tau}} \varphi(z) dz, \quad \text{where}$$

$$(c(\tau, x))^{-1} = \int_S e^{-\frac{(z-x)^2}{2\tau}} dz \text{ and } F(0) = \mathcal{I}. \text{ Then the func-}$$

tion  $F$  is continuous in the topology of pointwise convergence on  $\mathcal{L}(\mathfrak{L}_p(S))$ , and  $F'(0)$  is the Laplace–Beltrami operator  $\Delta_{\mathcal{I}}^S$  on  $\mathfrak{L}_p(S)$ . It follows that if a function  $F(\cdot)$  satisfies the estimate of Chernoff’s theorem, then this function can be used to obtain Feynman

approximations for the semigroup  $e^{t\Delta_{\mathcal{I}}^S}$ . Our method is based on the fact that a similar operator-valued function  $F(\cdot)$  can be defined for any function  $A: E \rightarrow \mathcal{L}(E)$  in such a way that  $F'(0)\varphi = \Delta_A^S \varphi$  for  $\varphi \in \operatorname{dom} \Delta_A^S$ .

The paper is organized as follows. In the first section, we give definitions used in what follows and state a theorem similar to Chernoff’s theorem for evolution families of operators (Chernoff’s classical theorem deals with one-parameter semigroups of operators). In the next section, we derive Feynman formulas for evolution families of operators on  $\mathfrak{L}_p(S)$  generated (a definition is given below) by the families of operators  $\Delta_{G(t)}^S$  and  $i\Delta_{G(t)}^S$ . The last, third, section contains two conjectures on a relationship between stochastic processes in  $E (= \mathbb{R}^n)$  and in  $S$  that are generated by  $\Delta_{G(t)}$  and  $\Delta_{G(t)}^S$ .

Main attention is focused on the algebraic structure of the problems, and assumptions of analytical character are not considered.

<sup>a</sup> *Section de Mathématiques, École Polytechnique Fédérale de Lausanne, Lausanne, CH-1015 Switzerland*

<sup>b</sup> *Mechanics and Mathematics Faculty, Moscow State University, Moscow, 119991 Russia  
e-mail: smolyanov@yandex.ru*

## 1. DEFINITIONS AND PRELIMINARY RESULTS

Let  $a > 0$ , and let  $\mathcal{A}(t)$  be a linear (unbounded in the general case) operator on a Banach space  $X$  for each  $t \in [0, a)$ ; we assume that the domains of the operators  $\mathcal{A}(t)$  coincide and denote the common domain by  $\text{dom}\mathcal{A}$ .

**Definition 1.** The evolution family on  $[0, a)$  of continuous linear operators on a Banach space  $X$  generated by an operator-valued function  $\mathcal{A}$  is a mapping  $U^{\mathcal{A}}$  of the set  $V_a$  defined by  $V_a = \{(\tau, t) \in \mathbb{R}^2 \mid 0 \leq \tau < t < a\}$  to  $\mathcal{L}(X)$  with the following properties:

(i)  $U^{\mathcal{A}}(\tau, \tau) = \mathcal{I}$  for  $\tau \in [0, a)$  (here and in what follows,  $\mathcal{I}$  denotes the identity map of the corresponding space);

(ii) for each  $\tau \in [0, a)$ , the function  $[\tau, a) \ni t \rightarrow f_\tau(t) := U^{\mathcal{A}}(\tau, t)$  is continuous with respect to the topology of pointwise convergence (the strong operator topology) on  $\mathcal{L}(X)$ , and  $\frac{1}{\delta}(f_\tau(\tau + \delta)\varphi - \varphi) \rightarrow \mathcal{A}(\tau)\varphi$  in  $X$  as  $\delta \rightarrow 0$  for all  $\varphi \in \text{dom}\mathcal{A}$ ;

(iii) for any  $0 \leq t_1 < t_2 < t_3 < a$ , the relation  $U^{\mathcal{A}}(t_2, t_3)U^{\mathcal{A}}(t_1, t_2) = U^{\mathcal{A}}(t_1, t_3)$  holds.

**Remark 1.** If  $U^{\mathcal{A}}$  is an evolution family in the sense of the above definition, then, for any  $\varphi \in X$  and any  $\tau \in [0, a)$ , the function  $[\tau, a) \ni t \rightarrow f_\tau(t)\varphi \in X$  is a solution of the Cauchy problem for the equation  $\psi'(t) = \mathcal{A}(t)\psi(t)$  with initial data  $(\tau, \varphi)$ .

**Theorem 1** (generalization of Chernoff's theorem [2, 4, 5]). *Let  $X$  be a Banach space with norm  $\|\cdot\|$ , and let  $\mathcal{L}_s(X)$  be the space  $\mathcal{L}(X)$  with the topology of pointwise convergence; suppose that, for  $a > 0$ ,  $F: V_a \rightarrow \mathcal{L}_s(X)$  is a continuous mapping with the following properties:*

(i)  $F(\tau, \tau) = \mathcal{I}$ ;

(ii) *there exists an  $\alpha > 0$  such that if  $\tau \in [0, a)$ ,  $\delta > 0$ , and  $(\tau, \tau + \delta) \in V_a$ , then  $\|F(\tau, \tau + \delta)\| \leq \exp(\alpha\delta)$ ;*

(iii) *there exists a vector subspace  $D$  of  $X$  such that*

(a) *for any  $x \in D$  and any  $\tau \in [0, a)$ , the limit*

$$F'_2(\tau, \tau) := \lim_{\delta \rightarrow 0} \frac{F(\tau, \tau + \delta)x - x}{\delta}$$

*exists in  $X$ ; this means that  $D$  is contained in the domain of the derived function  $[\tau, a) \ni t \mapsto F(\tau, \tau + t) \in \mathcal{L}_s(X)$  at zero;*

(b) *the operator  $F'_2(\tau, \tau)$  with domain  $D$  defined by  $D \ni x \mapsto F'_2(\tau, \tau)x \in X$  has a closure, which coincides with  $\mathcal{A}(\tau)$ .*

*Then, for any  $\tau \in [0, a)$  and  $t > 0$  such that  $(\tau, t) \in V_a$ ,*

*the sequence of operators  $\prod_{k=n}^{k=1} F\left(\tau + \frac{k-1}{b}t, \tau + \frac{k}{b}t\right) \in \mathcal{L}_s(X)$  converges to  $U^{\mathcal{A}}(\tau, \tau + t)$  uniformly*

*with respect to admissible  $t \in [0, T]$  for any  $T > 0$ .*

The proof of this theorem is similar to that of Chernoff's classical theorem [2, 4, 5].

## 2. FEYNMAN FORMULAS FOR EVOLUTION FAMILIES OF OPERATORS IN FUNCTION SPACES ON RIEMANNIAN MANIFOLDS

If  $X$  is a function space on a measure space and the values of the mapping  $F$  specified in Theorem 1 are integral operators on  $X$ , then the formula

$$\prod_{k=n}^{k=1} F\left(\tau + \frac{(k-1)t}{n}, \tau + \frac{kn}{n}\right) \rightarrow U^{\mathcal{A}}(\tau, \tau + t)$$

as  $n \rightarrow \infty$

is called the Feynman formula in  $X$  for the evolution family  $U^{\mathcal{A}}$  on  $[0, a)$  and the mapping  $F$ . Moreover, if the assumptions of Theorem 1 hold, then the Feynman formula in  $X$  for the evolution family  $U^{\mathcal{A}}$  on  $[0, a)$  and the mapping  $F$  is valid.

Let  $S$  be a compact Riemannian submanifold of the space  $E = \mathbb{R}^n$ ; for  $x \in S$ , by  $T_x S$  we denote the tangent space to  $S$  at  $x$  and by  $\text{pr}_{T_x S}: \mathbb{R}^n \rightarrow T_x S$ , the orthogonal projection. We use the symbol  $\nabla$  to denote the covariant derivative generated by the Riemannian (Levi-Civita) connection. Note that, given any function  $\psi \in C^\infty(S)$ , its second covariant derivative  $\nabla^2 \psi: S \rightarrow \mathcal{L}(T_x S)$  defines a linear map on each fiber  $T_x S$  of  $TS$  (we denote it by the same symbol).

**Definition 2.** The operator  $\Delta_A^S$  on  $\mathcal{L}_p(S)$ ,  $1 \leq l < \infty$ , associated with  $\Delta_A$  is defined by

$$(\Delta_A^S \psi)(x) := \text{tr}(\text{pr}_{T_x S} \mathcal{A}(x) |_{T_x S} \nabla^2 \psi(x)).$$

**Remark 2.** In the space  $\mathcal{L}(T_x S)$   $\text{pr}_{T_x S} \mathcal{A}(x) |_{T_x S} \nabla^2 \psi(x) = (\text{pr}_{T_x S} \mathcal{A}(x))(\text{pr}_{T_x S} \nabla^2 \psi(x))$ .

Let  $a > 0$ , and let  $F^j$ ,  $j = 1, 2$ , be mappings of  $V_a$  to  $\mathcal{L}(\mathcal{L}_p(S))$ ,  $1 \leq p < \infty$ , defined by the following relations (throughout the paper, we assume that all operators  $G(t)(x)$  are invertible):

$$(F^1(\tau, t)\varphi)(x) := c_1(\tau, t, x) \int_S e^{\frac{((G(\tau)(x))^{-1}(x-z), x-z)}{2(t-\tau)}} \varphi(z) dz,$$

$$F^1(0) = \mathcal{I},$$

$$(F^2(\tau, t)\varphi)(x) := c_2(\tau, t, x) \int_S e^{\frac{((G(\tau)(z))^{-1}(x-z), x-z)}{2(t-\tau)}} \varphi(z) dz,$$

$$F^2(0) = \mathcal{I},$$

where  $\varphi \in \mathcal{L}_p(S)$ , and

$$\frac{1}{c_1(\tau, t, x)} := \int_S e^{\frac{((G(\tau)(x))^{-1}(x-z), x-z)}{2(t-\tau)}} dz,$$

$$\frac{1}{c_2(\tau, t, x)} := \int_S e^{\frac{((G(\tau)(z))^{-1}(x-z), x-z)}{2(t-\tau)}} dz.$$

**Theorem 2.** If  $\varphi \in \text{dom} \Delta_{G(\tau)}^S$ , then

$$((F^j)'(\tau, \tau)_2(\varphi))(x) = \frac{1}{2}(\Delta_{G(\tau)}^S \varphi)(x), \quad j = 1, 2.$$

Let  $X = \mathfrak{L}_p(S)$ ,  $1 \leq p < \infty$ ; consider the operators  $\mathcal{A}(t)$ ,  $t \in [0, a)$ , on  $X$  defined by  $\mathcal{A}(t) := \Delta_{G(t)}^S$ .

**Theorem 3.** The Feynman formula is valid in  $\mathfrak{L}_p(S)$ ,  $1 \leq p < \infty$ , for the evolution family  $U^{\mathcal{A}}$  on  $[0, a)$  and each of the functions  $F^1$  and  $F^2$ .

The proof of Theorem 3 uses Theorems 1 and 2.

Let  $a > 0$ , and let  $F^{Qj}$ ,  $j = 1, 2$ , be mappings of  $V_a$  to  $\mathcal{L}(\mathfrak{L}_2(S))$  defined by

$$\begin{aligned} (F^{Q1}(\tau, t)\varphi)(x) &:= c_1^Q(\tau, t, x) \\ &\times \int_S e^{\frac{(i(G(\tau)(x))^{-1}(x-z), x-z)}{2(t-\tau)}} \varphi(z) dz, \quad F^{Q1}(0) = \mathcal{F}, \\ (F^{Q2}(\tau, t)\varphi)(x) &:= c_2^Q(\tau, t, x) \\ &\times \int_S e^{\frac{(i(G(\tau)(z))^{-1}(x-z), x-z)}{2(t-\tau)}} \varphi(z) dz, \quad F^{Q2}(0) = \mathcal{F}, \end{aligned}$$

where  $\varphi \in \mathfrak{L}_2(S)$ , and

$$\begin{aligned} \frac{1}{c_1^Q(\tau, t, x)} &:= \int_S e^{\frac{(i(G(\tau)(x))^{-1}(x-z), x-z)}{2(t-\tau)}} dz, \\ \frac{1}{c_2^Q(\tau, t, x)} &:= \int_S e^{\frac{(i(G(\tau)(z))^{-1}(x-z), x-z)}{2(t-\tau)}} dz. \end{aligned}$$

Here, we use the natural regularizations of integrals of complex exponentials.

Let  $X = \mathfrak{L}_2(S)$ , and let  $\mathcal{A}^{Qj}(t)$ ,  $t \in [0, a)$  be the operators on  $X$  defined by  $\mathcal{A}^{Qj}(t) := i\Delta_{G(t)}^S$ .

**Theorem 4.** The Feynman formula is valid in  $\mathfrak{L}_2(S)$  for the evolution family  $U^{\mathcal{A}^{Qj}}$  on  $[0, a)$  and each of the functions  $F_1^Q$  and  $F_2^Q$ .

**Remark 3.** Theorem 4 provides a solution of the Cauchy problem for a Schrödinger-type equation in which the Hamiltonian may be even asymmetric. Similar formulas can also be obtained for symmetric Hamiltonians.

**Remark 4.** The Feynman formulas obtained above are approximations of integrals of so-called chronological exponentials. In fact, according to the usual definition of such integrals, these integrals are equal to the limits of integrals of Feynman formulas. We also mention that, in the case under consideration, the corresponding chronological exponential can be

regarded as the generalized density of a probability measure or a Feynman pseudomeasure on a space of functions taking values in  $S$ .

**Remark 5.** Feynman formulas can also be obtained for equations of the form

$$\dot{f}(t) = \frac{1}{2}\Delta_{G(t)}^S f(t) + (\nabla(f(t)), h(t)) + V(t)f(t),$$

where  $h(t)$  is a vector field on  $S$  and  $V(t)$  is a function on  $S$  for each  $t$ . The multiple integrals in these formulas approximate integrals over spaces of functions taking values in  $S$ ; the integrands contain exponentials of stochastic integrals determined by the vector field  $h$  as multipliers.

### 3. CONJECTURES

In this section, we formulate two conjectures related to the results obtained above. The Green function of the Cauchy problem for the equation  $\dot{f}(t) = \frac{1}{2}\Delta_{G(t)}^S f(t)$  (with respect to the unknown function  $f$  taking values in  $\mathfrak{L}_p(S)$ ) and the Green function of the Cauchy problem for the equation  $\dot{\varphi}(t) = \frac{1}{2}\Delta_{G(t)}\varphi(t)$

(with respect to the unknown function  $\varphi$  taking values in  $\mathfrak{L}_p(E)$ ) generate probability measures  $\mu_S$  and  $\mu_E$  on the space of functions on  $[0, a)$  taking values in  $S$  and  $E$ , respectively (or, which is the same thing, random processes in  $S$  and  $E$ ). For each  $\varepsilon > 0$ , let  $S^\varepsilon$  be the  $\varepsilon$ -neighborhood of the manifold  $S$  in  $E$ , and let  $C_b([0, a], S)$ ,  $C_b([0, a], S^\varepsilon)$ , and  $C_b([0, a], E)$  be the spaces of continuous functions on  $[0, a]$  taking values in  $S$ ,  $S^\varepsilon$ , and  $E$ , respectively, and vanishing at  $b \in S$ . We assume that  $\mu_S$  and  $\mu_E$  are countably additive on

$C_b([0, a], S)$  and on  $C_b([0, a], E)$ . Let  $\bar{v}^\varepsilon$  be the restriction of the measure  $\mu_E$  to  $C_b([0, a], S^\varepsilon)$ ; we set  $v^\varepsilon := \bar{v}^\varepsilon / \mu_E(C_b([0, a], S^\varepsilon))$  and define  $\mu^\varepsilon$  to be the probability measure on  $C_b([0, a], E)$  being the image of  $v^\varepsilon$  under the canonical embedding of  $C_b([0, a], S)$  into  $C_b([0, a], E)$ .

**Conjecture 1.** For suitable  $G$ , there exists a probability measure  $v_S$  on  $C_b([0, a], E)$  concentrated on  $C_b([0, a], S)$  and such that  $\mu^\varepsilon \rightarrow v_S$  in the weak topology on the space  $\mathcal{M}(C_b([0, a], E))$  of measures on  $C_b([0, a], E)$  determined by the duality between the space  $\mathcal{M}(C_b([0, a], E))$  and an appropriate space consisting of bounded infinitely differentiable cylindrical functions on  $C_b([0, a], E)$  all of whose derivatives are bounded. Moreover, the measure  $v_S$  is equivalent to  $\mu_S$  (it would be useful to find the corresponding Radon–Nikodym density). In the case where  $G(t)(x)$  is the identity mapping for each  $t \in [0, a]$  and each  $x \in E$ , this conjecture was proved and the Radon–Nikodym density was found in [7].

Let  $\eta^\varepsilon$  be the probability measure generated on  $C_b([0, a], S^\varepsilon)$  by the Green function of the Cauchy–Neumann problem for the equation  $\dot{\psi} = \frac{1}{2} \Delta_{G(t)} \psi(t)$

with respect to the unknown function  $\psi$  taking values in  $\mathcal{L}_p(S^\varepsilon)$ , and let  $\gamma^\varepsilon$  be the measure on  $C_b([0, a], E)$  being the image of  $\eta^\varepsilon$  under the canonical embedding of  $C_b([0, a], S^\varepsilon)$  into  $C_b([0, a], E)$ .

**Conjecture 2.** In the space  $\mathcal{M}(C_b([0, a], E))$  endowed with the topology defined above,  $\gamma^\varepsilon \rightarrow \mu_S$ . This was proved in [6] in the case where  $G(t)(x) = \mathcal{J}$  for all  $t \in [0, a]$  and all  $x \in E$ .

**Remark 6.** The formulas given above are usually referred to as Lagrangian formulas, because the functional integrals approximated by these formulas contain action in Lagrangian form. However, the corresponding Hamiltonian formulas can be obtained as well [3]. Such formulas are useful if the classical Hamiltonian function is not the sum of the kinetic and the potential energy.

#### ACKNOWLEDGMENTS

T.S. Ratiu acknowledges the support of the Swiss National Science Foundation, grant NCCR SwissMAP.

O.G. Smolyanov acknowledges the support of the Federal Institute of Technology in Lausanne (EPFL) and of the Russian Foundation for Basic Research, project no. 14-01-00516.

#### REFERENCES

1. M. Gadella and O. G. Smolyanov, Dokl. Math. **77**, 120–123 (2008).
2. O. O. Obrezkov, O. G. Smolyanov, and A. Truman, Dokl. Math. **71**, 105–110 (2005).
3. T. Ratiu and O. G. Smolyanov, Dokl. Math. **87**, 289–292 (2013).
4. A. S. Plyashechnik, Russ. J. Math. Phys. **19** (3), 340–359 (2012).
5. A. S. Plyashechnik, Russ. J. Math. Phys. **20** (3), 377–379 (2013).
6. N. Sidorova, Infin. Dimens. Anal. Quantum Probab. Rel. Top. **7** (3), 461–471 (2004).
7. O. G. Smolyanov, H. von Weizsacker, and O. Wittich, Potential Anal. **26** (1), 1–29 (2007).

*Translated by O. Sipacheva*