Rapid and Convenient Synthesis of Original 5-Substituted Quinoli- no[3,4-b]quinoxalin-6(5H)-ones under Eco-Friendly Conditions

Omar Khoumeri, François-Xavier Vanelle, Maxime D Crozet, Thierry Terme,
Patrice Vanelle

► To cite this version:

Omar Khoumeri, François-Xavier Vanelle, Maxime D Crozet, Thierry Terme, Patrice Vanelle. Rapid and Convenient Synthesis of Original 5-Substituted Quinoli- no[3,4-b]quinoxalin-6(5H)-ones under Eco-Friendly Conditions. SYNLETT, 2016, 27 (10), pp.1547-1550. 10.1055/s-0035-1561577 . hal-01406037

HAL Id: hal-01406037

<https://hal.science/hal-01406037>

Submitted on 5 Dec 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Rapid and Convenient Synthesis of Original 5-(substituted)quinolino[3,4-*b*]quinoxalin-6(5*H*)-ones under Eco-Friendly Conditions

OmarKhoumeri^a
 François-XavierVanelle^a
 Maxime D. Crozet^a
 ThierryTerme^a
 PatriceVanelle^{a*}

^aAix-Marseille Université, Institut de Chimie Radicale ICR,
 UMR CNRS 7273, Laboratoire de Pharmaco-Chimie Radicale,
 Faculté de Pharmacie, 27 Boulevard Jean Moulin – CS 30064,
 13385 Marseille Cedex 05, France

patrice.vanelle@univ-amu.fr

[Click here to insert a dedication.](#)

12 examples

Received:
 Accepted:
 Published online:
 DOI:

Abstract We report herein the synthesis of 5-(substituted)quinolino[3,4-*b*]quinoxalin-6(5*H*)-one derivatives from ethyl 3-(2-bromophenyl)quinoxaline-2-carboxylate under one-pot Buchwald-Hartwig coupling/lactamization reaction.

Quinoline, Aromatic anilines, Buchwald-Hartwig reaction

Key words quinoxaline, aromatic anilines, Buchwald-Hartwig reaction, lactamization reaction

Introduction

Quinoxalines are common structural motifs in many natural and synthetic compounds with interesting biological activities. Many quinoxaline derivatives show a wide range of biological activities, making them potential antitumor,¹ antibiotic,² antiviral,³ anti-HIV⁴ and anti-inflammatory agents⁵. The list of their uses in medicinal chemistry is long.⁶ The olaquinox (OLA), 2-[(2-hydroxyethyl)carbamoyl]-3-methylquinoxaline 1,4-dioxide, presents strong antimicrobial activities and is used in subtherapeutics (animal feed) to promote animal growth.⁷ Quinacillin is significantly active (0.15–0.62 µg/mL, 0.32–1.35 µM) against *Staphylococcus aureus*.⁸

The phenanthridin group is present in many bioactive natural and pharmaceutical products.⁹ They show particularly interesting biological activity as anticancer agents.¹⁰ In order to synthesize structures with high biological activity potential, we combined the quinoxaline moiety with the phenanthridin moiety. Furthermore, to the best of our knowledge, few syntheses of quinolino[3,4-*b*]quinoxalin-6(5*H*)-one derivatives have been reported¹¹ and none used coupling reactions.

Palladium-catalyzed coupling reactions have been extensively used in medicinal chemistry. The synthesis process is not expensive and possesses a wide compatibility with many functional groups. This approach is an effective way of functionalizing the quinoxaline derivatives as Suzuki-Miyaura,¹² Sonogashira¹³ or Buchwald-Hartwig¹⁴ reactions, and the 2-bromophenylboronic acid (2) proved promising as a substrate for coupling/lactamization reaction.¹⁵

As part of our research program to develop new bioactive compounds,¹⁶ we report herein the first one-pot Buchwald-Hartwig coupling/lactamization reaction in a quinoxaline series for the synthesis of 5-(substituted)quinolino[3,4-*b*]quinoxalin-6(5*H*)-ones.

The starting product, for one-pot coupling/lactamization reaction, ethyl 3-(2-bromophenyl)quinoxaline-2-carboxylate **3**, was synthesized by Suzuki-Miyaura reaction between ethyl 3-chloroquinoxaline-2-carboxylate (**1**) and 2-bromophenylboronic acid (**2**) using 0.02 equiv of Pd(PPh₃)₄ as a palladium source and 5 equiv of Na₂CO₃ as a base in an acetonitrile–water mixture (3/1) at 100 °C for 5 h (Scheme 1). The key product **3** was obtained in good yields (73%).¹⁷

Scheme 1 Preparation of ethyl 3-(2-bromophenyl)quinoxaline-2-carboxylate (**3**)

Our initial strategy consisted in the introduction of the amide group from ester, first replacing the ethoxy group, which was followed by an intramolecular Buchwald-Hartwig reaction/cyclization. The amidation reaction was tested with starting product **3** and aniline in presence of *t*-BuOK at room temperature in toluene for 30 min. The formation of intermediate 3-(2-bromophenyl)-*N*-phenylquinoxaline-2-carboxamide was confirmed by TLC and LCMS. Then, we applied the coupling reaction conditions with different catalysts (Pd(OAc)₂, Pd(PPh₃)₄), in presence of bases (K₂CO₃, Cs₂CO₃, *t*-BuOK) at 110 °C. The best yield (44%) of 5-phenylquinolino[3,4-*b*]quinoxalin-6(5*H*)-one (**5a**) was obtained with palladium acetate, BINAP as supporting ligand and Cs₂CO₃ at 110 °C for 12 h.

In view of these last poor yields, a simpler strategy was developed, only involving coupling reaction conditions with ethyl 3-

(2-bromophenyl)quinoxaline-2-carboxylate (**3**). Preliminary tests were conducted with ethyl 3-(2-bromophenyl)quinoxaline-2-carboxylate (**3**) and aniline with different catalysts [Pd(OAc)₂, Pd(PPh₃)₄], in the presence of base (K₂CO₃, Cs₂CO₃, *t*-BuOK) at 110°C in toluene (Table 1).

A better yield was obtained with palladium acetate as catalyst and BINAP as ligand in presence of Cs₂CO₃ in toluene at 110°C for 5h (97% yield).

Scheme 2 Reaction of ethyl 3-(2-bromophenyl)quinoxaline-2-carboxylate (**3**) and aniline (**4a**)

Table 1 Optimization reaction between ethyl 3-(2-bromophenyl)quinoxaline-2-carboxylate and aniline^a

Entry	Catalyst ^a	Ligand ^a	Base ^a	Time (h)	Yield ^b (%)
1	Pd(OAc) ₂	BINAP	K ₂ CO ₃	15	78
2	Pd(OAc) ₂	BINAP	<i>t</i> -BuOk	15	Complex mixture
3	Pd(OAc) ₂	BINAP	Cs ₂ CO ₃	5	97
4	Pd(PPh ₃) ₄	-	K ₂ CO ₃	10	Starting material

^a All reactions are performed using 1.5 equiv of aniline, 20 mol% of Pd, 15 mol% of ligand, and 2.2 equiv of base. ^b All yields refer to chromatographically pure products based on substrate **3**.

Aromatic amine	Product ^a	Yield ^b (%)
Aniline	5a	97
2-Chloroaniline	5b	63
3-Chloroaniline	5c	80
3-Fluoroaniline	5d	72
3-(Trifluoromethyl)aniline	5e	65
3-Aminobenzonitrile	5f	82
4-Chloroaniline	5g	82
4-Fluoroaniline	5h	88
4-Methoxyaniline	5i	17 (50) ^c
3,4-Chloroaniline	5j	50 (60) ^c
2,3-Chloroaniline	5k	76
<i>p</i> -Toluidine	5l	90

Scheme 3 Synthesis of 5-(substituted)quinolino[3,4-*b*]quinoxalin-6(5*H*)-ones **5a–l**

Table 2 Generalization of one-pot coupling/lactamization reactions

^a All the reactions are performed using 1.5 equiv of aniline, 20 mol% of Pd(OAc)₂, 15 mol% of BINAP, 2.2 equiv of Cs₂CO₃ in toluene for 5h in 110°C. ^b All yields refer to chromatographically isolated pure products based

on substrate **3**.

^c The reactions are performed using 20 mol% of Pd(OAc)₂, 15 mol% of BINAP, 3 equiv of K₂CO₃ in toluene for 24h in 110°C

After optimization of the conditions (Table 1, entry 3) we generalized the coupling/lactamization reaction with different aniline derivatives and the corresponding 5-(substituted)quinolino[3,4-*b*]quinoxalin-6(5*H*)-one derivatives were obtained in good yields (50–97% Scheme 3, Table 2). Very good yields were obtained with no substituted aniline (**5a** 97%) and *p*-toluidine (**5l** 90%). Good yields were obtained with anilines substituted by halogen atom in *ortho*, *meta* or *para* position (63–88%). A yield of 82% was obtained with anilines substituted by CN or 65% yield with CF₃ group in *meta* position. With dichlorinated anilines, moderate to good yields (**5j** 60% and **5k** 76%) were obtained. No reaction was observed with anilines substituted by nitro or cyano group in *para* position. Finally, a low yield was obtained with 4-methoxyaniline (**5i** 17% with Cs₂CO₃ for 5h and 50% with K₂CO₃ for 24h).

In conclusion, we developed a very efficient one-pot reaction combining Buchwald-Hartwig and lactamization reactions for the synthesis of quinolino[3,4-*b*]quinoxalin-6(5*H*)-one derivatives. Starting material ethyl 3-(2-bromophenyl)quinoxaline-2-carboxylate (**3**) was prepared by Suzuki–Miyaura reaction from ethyl 3-chloroquinoxaline-2-carboxylate and 2-bromophenylboronic acid. The pharmacological evaluation of these highly potential bioactive compounds is under active investigation.

Acknowledgment

This work was supported by the Centre National de la Recherche Scientifique, Aix-Marseille. We express our thanks to V. Remusat for recording ¹H NMR and ¹³C NMR spectra and T. Schembri for recording mass spectra.

References and Notes

- (1) (a) Monge, A.; Martinez-Crespo, F. J.; Lopez de Cerain, A.; Palop, J. A.; Narro, S.; Senador, V.; Marin, A.; Sainz, Y.; Gonzalez, M.; Hamilton, E.; Barker, A. J. *J. Med. Chem.* **1995**, *38*, 4488. (b) Lindsley, C. W.; Zhao, Z.; Leister, W. H.; Robinson, R. G.; Barnett, S. F.; Defeo Jones, D.; Jones, R. E.; Hartman, G. D.; Huff, J. R.; Huber, H. E.; Duggan, M. E. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 761. (c) Marcus, L.; Kingi, N.; Bergman, J. *J. Med. Chem.* **2008**, *51*, 7744. (d) Lee, S.-H.; Kim, N.; Kim, S.-J.; Song, J.; Gong, Y.-D.; Kim, S.-Y. *J. Cancer Res. Clin. Oncol.* **2013**, *139*, 1279.
- (2) (a) Shoji, J.; Katakiri, K. *J. Antibiotics Ser. A* **1961**, *14*, 335. (b) Kim, Y. B.; Kim, Y. H.; Park, J. Y.; Kim, S. K. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 541.
- (3) (a) Sehlsedt, U.; Aich, P.; Bergman, J.; Vallberg, H.; Norden, B.; Graslund, A. *J. Mol. Biol.* **1998**, *278*, 31. (b) Harmenber, J.; Akesson-Johansson, A.; Graslund, A.; Malmfors, T.; Bergman, J.; Wahren, B.; Akerfeldt, S.; Lundblad, L.; Cox, S. *Antiviral Res.* **1991**, *15*, 193. (c) Wilhelmsson, L. M.; Kingi, N.; Bergman, J. *J. Med. Chem.* **2008**, *51*, 7744.
- (4) (a) Munos, M.-H.; Mayrargue, J.; Fournet, A.; Gantier, J.-C.; Hocquemiller, R.; Moskowitz, H. *Chem. Pharm. Bull.* **1994**, *42*, 1914. (b) Maga, G.; Gemma, S.; Fattorusso, C.; Locatelli, G. A.; Butini, S.; Persico, M.; Kukreja, G.; Romano, M. P.; Chiasserini, L.; Savini, L.; Novellino, E.; Nacci, V.; Spadari, S.; Campiani, G. *Biochemistry* **2005**, *44*, 9637.
- (5) (a) Badran, M.M.; Abouzid, K.A.M.; Hussein, M.H.M. *Arch. Pharm. Res.* **2003**, *26*, 107. (b) Kim, Y. B.; Kim, Y. H.; Park, J. Y.; Kim, S. K. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 541. (c) He, W.; Meyers, M. R.; Hanney, B.; Sapada, A.; Blider, G.; Galzeinski, H.; Amin, D.; Needle, S.; Page, K.; Jayyosi, Z.; Perrone, H. *Bioorg. Med. Chem. Lett.* **2003**, *13*, 3097. (d) Sakata, G.; Makino, K.; Kurasawa, Y. *Heterocycles* **1988**, *27*, 2481. (e) Khan, S. A.; Mullick, P.; Pandit, S.; Kaushik, D. *Acta Pol. Pharm.* **2009**, *66*, 169. (f) Abu-Hashem, A.-A.; Gouda, M. A.; Badria, F. A. *Eur. J. Med. Chem.* **2010**, *45*, 1976.
- (6) Cheeseman, G.W.H.; Cookson, R.F. *The Chemistry of Heterocyclic Compounds*; Weissberger, A., Taylor, E.C., Eds.; John Wiley and Sons: New York, NY, USA, **1979**, 35, 1.
- (7) (a) Carta, A.; Corona, P.; Loriga, M. *Curr. Med. Chem.* **2005**, *12*, 2259. (b) Tianci, X.; Xingjia, G.; Lei, Z.; Fang, P.; Junna, L.; Yunyu Z.; Hongjing, J. *Food. Chem. Toxicol.* **2012**, *50*, 2540.
- (8) Hugo, W. B.; Stretton, R. G. *Nature* **1964**, *202*, 1217.
- (9) (a) Michael, J. P. *Nat. Prod. Rep.* **1995**, *12*, 465. (b) Li, J.-H.; Serdyuk, L.; Ferraris, D. V.; Xiao, G.; Tays, K. L.; Kletzly, P. W.; Li, W.; Lautar, S.; Zhang, J.; Kalish, V. J. *Bioorg. Med. Chem. Lett.* **2001**, *11*, 1687. (c) Ishida, J.; Yamamoto, H.; Kido, Y.; Kamijo, K.; Murano, K.; Miyake, H.; Ohkubo, M.; Kinoshita, T.; Warizaya, M.; Iwashita, A.; Mihara, K.; Matsuokac, N.; Hattori, K. *Bioorg. Med. Chem.* **2006**, *14*, 1378. (d) Patil, S. Kamath, S. Sanchez, T. Neamati, N. Schinazi R. F. and Buolamwini, J. K. *Bioorg. Med. Chem.* **2007**, *15*, 1212. (e) Jin, Z. *Nat. Prod. Rep.* **2009**, *26*, 363. (f) Lehtio, L.; Jemth, A.-S.; Collins, R.; Loseva, O.; Johansson, A.; Markova, N.; Hammarstro, M.; Flores, A.; HolmbergSchiaivone, L.; Weigelt, J.; Helleday, T.; Schuler, H.; Karlberg, T. *J. Med. Chem.* **2009**, *52*, 3108. (g) Gui, Q.; Yang, Z.; Chen, X.; Liu, J.; Tan, Z.; Guo, R.; Yu, W. *Synlett* **2013**, *24*, 1016. (h) Pimparkar, S.; Jegannmohan, M. *ChemComm.* **2014**, *50*, 12116. (i) Yang, Y.; Huang, H.; Wu, L.; Yun, L. *Org. Biomol. Chem.* **2014**, *12*, 5351.
- (10) (a) Poschenrieder, H.; Stachel, H.-D.; Wiesend, B.; Polborn, K. *J. Heterocyclic. Chem.* **2003**, *40*, 61. (b) El-Gogary, S. R.; Waly, M. A.; Ibrahim, I. T.; El-Sepelgy, O. Z. *Monatsh. Chem.* **2010**, *141*, 1253.
- (11) Mohamed, E. A.; Gabr, Y.; Ismail, M. M.; Abass, M. *An. Quim.* **1995**, *91*, 579.
- (12) (a) Mao, L.; Sakurai, H.; Hirao, T. *Synthesis*, **2004**, *15*, 2535. (b) Ali, I.; Siyo, B.; Al-Soud, Y.; Villinger, A.; Langer, P. *Synthesis* **2012**, *44*, 1637.
- (13) (a) Armengol, M.; Joule, J. A. *J. Chem. Soc., Perkin Trans.* **2001**, *1*, 978. (b) Kang, F.-A.; Lanter, J. C.; Cai, C.; Sui, Z.; Murray, W. V. *Chem. Commun.* **2010**, *46*, 1347. (c) Shi, C.; Aldrich, C. C. *Org. Lett.* **2010**, *12*, 2286. (c) Keivanloo, A.; Bakherad, M.; Rahimi, A.; Taheri, S. A. N. *Tetrahedron Lett.* **2010**, *51*, 2409. (d) Gulevskaya, A. V.; Lazarevich, R. Y.; Pozharskii, A. F. *Tetrahedron* **2013**, *69*, 910. (e) Nakhi, A.; Rahman, M. S.; Seerapu, G. P. K.; Banote, R. K.; Kumar, K. L.; Kulkarni, P.; Haldar, D.; Pal, M. *Org. Biomol. Chem.*, **2013**, *11*, 4930. (f) Kolli, S. K.; Nakhi, A.; Medishetti, R.; Yellanki, S.; Kulkarni, P.; Ramesh Raju, R.; Pal, M. *Bioorg. Med. Chem. Lett.* **2014**, *24*, 4460.
- (14) Hung, T. Q.; Hoang, D. H.; Thang, N. N.; Dang, T. T.; Ayub, K.; Villinger, A.; Friedrich, A.; Lochbrunner, S.; Flechsig, G.U.; Langer, P. *Org. Biomol. Chem.* **2014**, *12*, 6151.
- (15) (a) Bolibrukh, K.; Khoumeri, O.; Polovkovych, V.; Terme, T.; Vanelle, P. *Synlett* **2014**, *25*, 2765. (b) Do, H.H.; Tran, H.Q.; Ohlendorf, L.; Ngo, T.N.; Dang, T.T.; Ehlers, P.; Villinger, A.; Langer, P. *Synlett* **2015**, *26*, 2429.
- (16) (a) Crozet, M. P.; Gellis, A.; Pasquier, C.; Vanelle, P.; Aune, J.-P. *Tetrahedron Lett.* **1995**, *36*, 525. (b) Montana, M.; Terme, T.; Vanelle, P. *Tetrahedron Lett.* **2006**, *47*, 6573. (c) Dunn, L.A.; Burgess, A. G.; Krauer, K. G.; Eckmann, L.; Vanelle, P.; Crozet, M. D. Gillin, F. D.; Upcroft, P.; Upcroft, J. A. *Int. J. Antimicrob. Agents* **2010**, *36*, 37. (d) Kieffer, C.; Cohen, A.; Verhaeghe, P.; Paloque, L.; Hutter, S.; Castera-Ducros, C.; Laget, M.; Rault, S.; Valentin, A.; Rathelot, P.; Azas, N.; Vanelle, P. *Bioorg. Med. Chem.* **2015**, *23*, 2377.
- (17) Ethyl 3-(2-bromophenyl)quinoxaline-2-carboxylate (**3**)
To a mixture of ethyl 3-chloroquinoxaline-2-carboxylate (**1g**, 4.22mmol) and 2-bromophenylboronic acid (0.93 g, 4.64mmol) in acetonitrile (30 mL), was added Pd(PPh₃)₄ (0.097 g, 0.084mmol) and 2M aqueous sodium carbonate solution (10 mL). The resulting mixture was stirred at 100°C for 5h. After cooling the mixture was diluted with dichloromethane and washed 3 times with water and dried over MgSO₄. After evaporation, the crude product was purified by silica gel chromatography with CH₂Cl₂ as eluant furnished the product **3** as yellow solid; yield: 1.1 g (73%); mp 100°C; ¹H NMR (250MHz, CDCl₃) δ = 1.19 (t, J = 7.1 Hz, 3H, CH₃), 4.33 (q, J = 7.1 Hz, 2H, CH₂), 7.33-7.40 (m, 1H, H_{Ar}), 7.45-7.85 (m, 2H, H_{Ar}), 7.67 (d, J = 8.0 Hz, 1H, H_{Ar}), 7.89-7.93 (m, 2H, H_{Ar}), 8.19-8.23 (m, 1H, H_{Ar}), 8.31-8.35 (m, 1H, H_{Ar}). ¹³C NMR (62.5MHz, CDCl₃) δ = 13.7, 62.3, 122.1, 127.6, 129.3, 130.0, 130.3, 130.5, 131.0, 132.1, 132.3, 139.8, 140.4, 142.0, 144.4, 153.2, 164.7. Anal. Calcd for C₁₇H₁₃BrN₂O₂: C, 57.16; H, 3.67; N, 7.84. Found: C, 57.61; H, 3.61. N, 7.86.
- (18) General procedure of one-pot Buchwald-Hartwig/lactamization reactions (**5a-l**)
A mixture of ethyl 3-(2-bromophenyl)quinoxaline-2-carboxylate (**3**) (50 mg, 0.14mmol), aniline (0.21mmol, 1.5equiv) in toluene (3mL), Cs₂CO₃ (100mg, 0.3mmol), Pd(OAc)₂ (6.2 mg, 0.028mmol), BINAP (14 mg, 0.021mmol) was added to a 4mL sealed vial. The resulting mixture was stirred at 110°C for 5h. After cooling the mixture was diluted with dichloromethane, washed with H₂O (3x40 mL) and dried over MgSO₄. After evaporation, the crude product was purified by silica gel chromatography (CH₂Cl₂/ diethylether: 97/3).
5-phenylquinolino[3,4-b]quinoxalin-6(5H)-one (**5a**) yellow solid; mp 240°C; ¹H NMR (250MHz, CDCl₃) δ = 6.73 (d, J = 7.9 Hz, 1H, H_{Ar}); 7.38-7.46 (m, 4H, H_{Ar}); 7.59-7.70 (m, 3H, H_{Ar}); 7.87-8.01 (m, 2H, H_{Ar}); 8.33 (d, J = 8.4 Hz, 1H, H_{Ar}); 8.48 (d, J = 8.5 Hz, 1H, H_{Ar}); 9.05-9.08 (m, 1H, H_{Ar}). ¹³C NMR (62.5MHz, CDCl₃) δ = 116.9, 119.6, 123.5, 125.8, 129.0, 129.2, 130.4, 130.4, 130.7, 131.0, 131.9, 132.9, 137.6, 140.5, 143.1, 144.5, 145.7, 160.3. Anal. Calcd for C₂₁H₁₃N₃O: C, 78.00; H, 4.05; N, 13.00. Found: C, 78.12; H, 4.00. N, 12.91.