Discovery of new hit-molecules targeting *Plasmodium falciparum* through a global SAR study of the 4-substituted-2-trichloromethylquinazoline antiplasmodial scaffold

Justine Desroches, Charline Kieffer, Nicolas Primas, Sébastien Hutter, Armand Gellis, Hussein El-Kashef, Pascal Rathelot, Pierre Verhaeghe, Nadine Azas, Patrice Vanelle

► To cite this version:

Justine Desroches, Charline Kieffer, Nicolas Primas, Sébastien Hutter, Armand Gellis, et al.. Discovery of new hit-molecules targeting *Plasmodium falciparum* through a global SAR study of the 4-substituted-2-trichloromethylquinazoline antiplasmodial scaffold. *European Journal of Medicinal Chemistry*, 2017, 125 (128), pp.68-86. 10.1016/j.ejmech.2016.09.029 . hal-01416990

HAL Id: hal-01416990

<https://hal.science/hal-01416990>

Submitted on 15 Dec 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Discovery of new hit-molecules targeting *Plasmodium falciparum* through a global SAR study of the 4-substituted-2-trichloromethylquinazoline antiplasmodial scaffold

Justine Desroches, Charline Kieffer, Nicolas Primas, Sébastien Hutter, Armand Gellis, Hussein El-Kashef, Pascal Rathelot, Pierre Verhaeghe, Nadine Azas, Patrice Vanelle

PII: S0223-5234(16)30759-0

DOI: [10.1016/j.ejmech.2016.09.029](https://doi.org/10.1016/j.ejmech.2016.09.029)

Reference: EJMECH 8895

To appear in: *European Journal of Medicinal Chemistry*

Received Date: 22 July 2016

Revised Date: 8 September 2016

Accepted Date: 9 September 2016

Please cite this article as: J. Desroches, C. Kieffer, N. Primas, S. Hutter, A. Gellis, H. El-Kashef, P. Rathelot, P. Verhaeghe, N. Azas, P. Vanelle, Discovery of new hit-molecules targeting *Plasmodium falciparum* through a global SAR study of the 4-substituted-2-trichloromethylquinazoline antiplasmodial scaffold, *European Journal of Medicinal Chemistry* (2016), doi: 10.1016/j.ejmech.2016.09.029.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Graphical Abstract:**Thienopyrimidine Analogues**

SAR
7 molecules

Initial Antiplasmodial Quinazolines

X = NH, O, S

SAR study
11 molecules

SAR
8 molecules

Quinoline and Quinoxaline Analogues

Discovery of new hit-molecules targeting *Plasmodium falciparum* through a global SAR study of the 4-substituted-2-trichloromethylquinazoline antiplasmodial scaffold

Justine Desroches^{1#}, Charline Kieffer^{1#}, Nicolas Primas¹, Sébastien Hutter², Armand Gellis¹, Hussein El-Kashef³, Pascal Rathelot¹, Pierre Verhaeghe^{4*}, Nadine Azas² and Patrice Vanelle^{1*}

¹Aix-Marseille Université, CNRS, ICR UMR 7273, Equipe Pharmaco-Chimie Radicalaire, Faculté de Pharmacie, 27 Boulevard Jean Moulin – CS30064, 13385 Marseille cedex 05, France.

²Aix-Marseille Université, UMR MD3, Infections Parasitaires, Transmission et Thérapeutique, Faculté de Pharmacie, 27 Boulevard Jean Moulin – CS30064, 13385 Marseille cedex 05, France.

³Department of Chemistry, Faculty of Science, Assiut University, 71516 Assiut, Egypt.

⁴Université Paul Sabatier, Faculté des Sciences Pharmaceutiques – CNRS UPR 8241, Laboratoire de Chimie de Coordination, 205 Route de Narbonne, 31077 Toulouse cedex 04, France.

#Co-first authors. *Corresponding authors: E-mail addresses: pierre.verhaeghe@univ-tlse3.fr (P. Verhaeghe), patrice.vanelle@univ-amu.fr (P. Vanelle).

Abstract:

From 4 antiplasmodial hit-molecules identified in 2-trichloromethylquinazoline series, we conducted a global Structure-Activity relationship (SAR) study involving 26 compounds and covering 5 molecular regions (I – V), aiming at defining the corresponding pharmacophore and identifying new bioactive derivatives. Thus, after studying the aniline moiety in detail, thienopyrimidine, quinoline and quinoxaline bio-isosters were synthesized and tested on the K1 multi-resistant *P. falciparum* strain, along with a cytotoxicity evaluation on the human HepG2 cell line, to define selectivity indices. SARs first showed that thienopyrimidines and quinolines were globally more cytotoxic, while quinoxaline analogs appeared as active as- and less cytotoxic than their quinazoline counterparts. Such pharmacomodulation in quinoxaline series not only provided a new antiplasmodial reference hit-molecule ($IC_{50} = 0.4 \mu M$, selectivity index = 100), but also highlighted an active ($IC_{50} = 0.4 \mu M$) and quite selective (SI = 265) synthesis intermediate.

Highlights:

► Antiplasmodial pharmacomodulation of CCl_3 -substituted-nitrogen containing heterocycles was made. ► Thienopyrimidine derivatives appeared more cytotoxic. ► Original 3-substituted-2-trichloromethylquinoxaline analogs were prepared. ► Two quinoxaline derivatives displayed *in vitro* IC_{50} values of 0.4 and 0.5 μM on the K1 multi-resistant *P. falciparum* strain. ► Cytotoxicity was assessed on the human HepG2 cell line showing low cytotoxicity ($CC_{50} \sim 40 \mu M$) and improved selectivity indices (77-100).

Keywords: Quinazoline; Quinoline; Quinoxaline; Thienopyrimidine; Trichloromethyl group; *Plasmodium falciparum*; *In vitro* antiparasitic activity; *In vitro* HepG2 cytotoxicity, Structure-Activity Relationships.

1. Introduction

Cerebral malaria, caused by *Plasmodium falciparum*, is the leading cause of death among parasitic infections worldwide. According to the 2015 World Malaria Report [1], 214 million people were infected by *Plasmodium* in 2014, leading to 438.000 deaths out of which about 88% occurred in Africa, mainly in children under five. It is to note that a significant improvement of the situation has been noted since the beginning of the 21st century, thanks to the action of the WHO and the financial involvement of several non-governmental organizations.

However, the control of the infection is facing worldwide the emergence of drug-resistant strains of the parasite which turns into a major concern for the medical and scientific community. Indeed, the WHO recommended the treatment of *P. falciparum* malaria is based on an artemisinin-combination therapy (ACT), but resistances to artemisinin derivatives are emerging in Asia [2], and it was demonstrated that they were responsible for therapeutic failures in several infected patients [3,4]. Moreover, it was also highlighted that the African *Anopheles gambiae* mosquito could transmit such Asian resistant parasites [5], indicating a major worldwide spreading risk. Thus, research efforts have to be developed, in order to discover new chemical entities presenting novel mechanisms of action, to complete and guaranty the durable efficiency of ACTs.

Several research teams working in the field of antimalarial agents previously reported novel quinazoline derivatives displaying significant antiparasitic activities, in particular when bearing an amino- [6], alkylamino- [7,8] or an aniline- substituent [9] at position 4 of the quinazoline ring.

The research activity of our group is focused on the synthesis and anti-infective evaluation of new nitrogen-containing heterocycles [10]. We have intensively studied a large series of antiparasitic derivatives based on the original 2-trichloromethylquinazoline scaffold. Thus, the synthesis and *in vitro* biological evaluation of a large quinazoline chemical library bearing an arylamino- [11,12], aryl- [13], phenoxy- [14], phenylthio- [15], sulfonamido- [16], alkynyl- [17], heteroarylamino- [18], benzyloxy- or alkoxy- [19] substituent at position 4 of the quinazoline ring, revealed several hit-molecules (A-D) which are presented in Figure 1.

Figure 1. Antiplasmodial hit-molecules identified in 2-trichloromethylquinazoline series.

In the light of these hit-molecules, within the aim of defining the antiplasmodial Structure Activity Relationships (SARs) of the corresponding series, 5 key-regions (I-V) were chosen and modulated, as summarized in **Figure 2**. Thus, for region I, we first investigated the role of the $-CCl_3$ group by replacing it by various substituents including closely related halogenated groups. For region II, we studied the influence of the combination of the halogen atoms on the aniline moiety of hit **A**. The effect of the methylation of the secondary amine nitrogen of hit **A** was then investigated in region III. Further modulations in region IV were made by replacing the quinazoline ring of hits **A-D** by a bio-isosteric thienopyrimidine. Indeed, starting from the natural quinazoline-based antimalarial molecule Febrifugine, such ring replacement was successfully carried out [20]. Moreover, some recent thienopyrimidine derivatives demonstrated promising antiplasmodial activities [21,22]. Finally, in region V, we looked for novel analogs, centered either on the quinoline nucleus, as this heterocycle is encountered in numerous antimalarial drugs such as quinine, chloroquine, mefloquine, amodiaquine, piperazine and primaquine, or on the quinoxaline nucleus, taking advantage of previously reported results obtained in our team in the pyrrolo[1,2-*a*]quinoxaline series [23]. The synthesis of all new derivatives, their *in vitro* biological evaluation and the resulting SARs is presented and discussed herein.

Figure 2. Antiplasmodial SAR study involving 5 key-regions (I-V) from the quinazoline hits

2. Results and discussion

2.1. Synthesis of hit A analogues

2.1.1. Region I: Variation of the nature of the substituent at position 2 of the quinazoline ring

The nature of the substituent at position 2 of hit **A** was first studied. Thus, the synthesis of the 2-methyl- **1**, 2-chloromethyl- **2**, 2-dichloromethyl- **3**, 2-chloro-2-difluoro-methyl- **7**, 2-trifluoromethyl- **8** and the 2-unsubstituted- **9** quinazoline analogs were operated, starting from appropriate 4-chloroquinazolines (**Scheme 1**). Compound **1** [11] was already synthesized by our team *via* an Aromatic Nucleophilic Substitution (S_NAr) reaction, starting from 4-chloro-2-trichloromethylquinazoline **E** [24]. Derivatives **2** and **3** were prepared by S_NAr reactions starting from 4-chloro-2-chloromethylquinazoline **F** [25] and 4-chloro-2-dichloromethylquinazoline **G** [26]. The synthesis of the parent 2-unsubstituted compound **9** was already reported in the literature, but to the best of our knowledge, it was never studied for its antiplasmodial potential [27]. The biological evaluation of the above prepared quinazolines is summed up in **Table 1**.

Scheme 1. SAR modulation of region I: modification of the substituent at position 2 of the quinazoline ring.

Reagents and conditions : (i) 2,4-dichloroaniline 0.8 equiv, conc. HCl cat., *i*PrOH, 70 °C, 2 h.

For the synthesis of derivative **7**, the corresponding 4-chloroquinazoline intermediate was not reported in the literature. Thus, starting from 2-aminobenzonitrile, acylation of the amine was made by reacting with 2-chloro-2-difluoroacetic acid in the presence of the coupling agent *N*-dimethylaminopropyl-*N*'-ethylcarbodiimide (EDCI) and dimethylaminopyridine

(DMAP), affording **4** in low yield (15%). Then, a smooth oxidative hydration of nitrile **4** was operated with H_2O_2 , followed by a cyclization in alkaline medium [17,28], leading to quinazolinone **5**. This last was further chlorodehydroxylated using POCl_3 , affording the expected intermediate **6** (Scheme 2).

Scheme 2. Variation of the region I: Synthesis of intermediate **6**.

Reagents and conditions : (i) 2-chloro-2-difluoroacetic acid 1 equiv, EDCI.HCl 1 equiv, DMAP 0.8 equiv, CH_2Cl_2 , 0 °C then rt, 24 h; (ii) H_2O_2 30%, NaOH, EtOH/ H_2O , 0 °C then rt, 90 min; (iii) POCl_3 , 140 °C, MW, 10 min, 800 W.

2.1.2. Region II: Variation of the halogen atoms at position 2' and 4' of the 4-anilino-substituent

The synthesis of the anilino derivatives **10-13** was easily performed under classical $\text{S}_{\text{N}}\text{Ar}$ reaction conditions by reacting the appropriate 2,4-dihaloaniline with 4-chloro-2-trichloromethylquinazoline **E** in refluxing isopropanol, in 55-70 % range yield (Scheme 3). The results of their *in vitro* biological evaluation are presented in Table 1.

Scheme 3. Chemical modulations at regions II and III of the pharmacophore

Reagents and conditions: (i) 2,4-dihaloaniline (2 equiv) or 2,4-dichloro-*N*-methylaniline (1.1 equiv), *i*PrOH, reflux, 2 h.

2.1.3. Region III: *N*-methylation of the 4-(2',4'-dichlorophenyl)amine substituent

The influence of the hydrogen atom on the secondary amine, in region III, was also studied. For this purpose, a S_NAr reaction was performed between **E** and *N*-methyl-2,4-dichloroaniline, affording the expected methylated compound **14** in low yield (**Scheme 3**). The biological evaluation of compound **14** is summarized in **Table 1**.

2.2. Biological evaluation of the series belonging to regions I, II and III

These new derivatives were evaluated *in vitro* toward both their antiplasmodial activity against the K1 multi-resistant *P. falciparum* strain (determination of the IC_{50} = inhibitory concentration 50%) and their cytotoxicity (determination of the CC_{50} = cytotoxic concentration 50%) on the HepG2 human cell line. The results were compared with three commercial antimalarial reference-drugs (atovaquone, chloroquine and doxycycline) and a cytotoxic reference drug (doxorubicine). For all tested compounds, the corresponding selectivity indices (SI) were calculated ($SI = CC_{50} / IC_{50}$). The results are presented in **Table 1**.

The aim of the modulation of region I was to confirm that the trichloromethyl group, at position 2 of the quinazoline ring, was the only substituent providing the antiplasmodial activity. Indeed, the antiplasmodial activity was totally lost when replacing it with other groups, including closely related ones such as CF_3 or $CHCl_2$. Looking at the combination of the halogen substituents on the aniline ring at position 4 (region II), the antiplasmodial activity was maintained for brominated compounds **10-12**, in comparison with hit **A** (respectively 0.3, 0.5 and 0.6 vs 0.4 μM), but the cytotoxicity was significantly increased, leading to less selective molecules. Contrary to the results obtained with brominated analogues, fluorinated analogue **13** presented the same cytotoxicity profile as hit **A** but was less active ($IC_{50} = 3.1 \mu M$). Finally in region III, the methylation of the aniline moiety (compound **14**) significantly impaired both antiplasmodial activity and cytotoxicity.

Table 1. *In vitro* antiplasmodial and cytotoxicity evaluation of compounds **1-3** and **7-14**: Modulations at regions I, II and III.

Molecule	-R ₂	-R ₄	HepG2 CC ₅₀ (μM)	K1 <i>P. falciparum</i> IC ₅₀ (μM)	Selectivity Index ^c
1	-CH ₃		15	85	0.18
2	-CH ₂ Cl		0.5	>12.5 ^d	< 0.04
3	-CHCl ₂		4.7	>50	< 0.09
7	-CF ₂ Cl		7	48.4	0.14
8	-CF ₃		>50	>50	-
9	-H		35	>50	<0.7
10	-CCl ₃		2	0.3	6.7
11	-CCl ₃		5	0.5	10
12	-CCl ₃		2.9	0.6	4.8
13	-CCl ₃		19.6	3.1	6.3
14	-CCl ₃		10	3.4	2.9
Hit A	-CCl ₃		16	0.4	40
Doxorubicine^a			0.2	-	-
Atovaquone^b			> 15.6 ^d	0.001	15600
Chloroquine^b			30	0.6	50
Doxycycline^b			20	6.0	3.3

^aDoxorubicine was used as a cytotoxic reference-drug; ^bAtovaquone, Chloroquine and doxycycline were used as antimalarial reference-drugs, ^cSelectivity indices were calculated according to the formula : SI = HepG2 CC₅₀ / K1 IC₅₀. ^dHighest concentration tested due to a lack of solubility.

2.3. Region IV: Synthesis of thieno[3,2-*d*]pyrimidine series.

The replacement of the phenyl ring of the quinazoline nucleus, by a thiophene nucleus as a bio-isostere, was next studied. We first tried to synthesize 4-substituted-2-(trichloromethyl)thieno[3,2-*d*]pyrimidine derivatives by a S_NAr reaction between the key chlorinated intermediate **20** and various nucleophilic species, corresponding to the substituents borne by hits **A-D**, respectively 2,4-dichloroaniline **a**, 3-trifluoromethylaniline **b**, 4-chlorophenol **c** and 4-chlorothiophenol **d** (Figure 3).

Figure 3. Thieno[3,2-*d*]pyrimidines initial retrosynthesis pathway.

Starting from the commercially available 3-aminothiophene-2-carboxylate, we first acetylated the amino group in presence of Ac_2O , leading to **15** in 78% yield [29]. Then, the cyclization was made by heating **15** in presence of 25% ammonia in a sealed vial, affording thienopyrimidinone **16** [30] in 63% yield. Unfortunately, in our hands, the chlorination reaction of **16**, to access **20**, did not succeed. Thus, we decided to realize the chlorination reaction in 2 steps. We first synthesized 4-chloro-2-methylthieno[3,2-*d*]pyrimidine **17** [30]. Then, **17** was engaged in a second chlorination reaction with $PCl_5/POCl_3$, under classical heating or microwave activation to convert the 2-methyl group into a 2-trichloromethyl substituent. Unfortunately, here again, no trace of the key intermediate **20** was formed, and only inseparable mixtures of polychlorinated compounds were observed by LC-MS analysis (Scheme 4).

Scheme 4. Synthesis of key intermediate **20**.

Reagents and conditions: (i) Ac_2O , AcOH cat., $120\text{ }^\circ\text{C}$, MW, 10 min; (ii) NH_4OH 25%, $105\text{ }^\circ\text{C}$, 3 h, sealed vial; (iii) PCl_5 6 equiv, POCl_3 as solvent, $100\text{ }^\circ\text{C}$, MW, 20 min, 800 W; (iv) POCl_3 , pyridine cat., reflux 24 h (molecule **17**) or 1 h (molecule **20**); (v) Et_3N , trichloroacetyl chloride, $5\text{ }^\circ\text{C}$, 30 min; (vi) AcOH saturated with gaseous HCl , trichloroacetonitrile, $100\text{ }^\circ\text{C}$, 18 h, evaporated then refluxed in *i*PrOH, 5 min.

To introduce the $-\text{CCl}_3$ group, we chose to start again from 3-aminothiophene-2-carboxylate and carried out the acylation reaction, using trichloroacetyl chloride in presence of Et_3N . The corresponding product **18** was obtained in quantitative yield. Unfortunately, the following cyclization step to lactam **19** was unsuccessful under the reaction conditions used (either NH_4OH at $105\text{ }^\circ\text{C}$ in sealed vial or NH_3 in MeOH at $120\text{ }^\circ\text{C}$ under MW). Finally, we succeeded in preparing lactam **19** by reacting 3-aminothiophene-2-carboxylate with trichloroacetonitrile in AcOH , saturated with gaseous HCl [31]. This was a similar approach to the one of Ried [32] who used a more expensive reagent: methyl 2,2,2-trichloroacetimidate. Then, the chlorodehydroxylation of lactam **19** in refluxing POCl_3 in the presence of a catalytic amount of pyridine, furnished the key intermediate **20** in almost quantitative yield. $\text{S}_\text{N}\text{Ar}$ reactions between **20** and some N, O or S centered nucleophiles were then performed in a similar way to those operated in quinazoline series. The target 4-substituted-2-(trichloromethyl)thieno[3,2-*d*]pyrimidine derivatives **21a-d** were obtained, however, in low to moderate yields (7-55%) (Scheme 5). The results of the biological evaluation of this series are presented in Table 2.

Scheme 5. Synthesis of 4-substituted-2-(trichloromethyl)thieno[3,2-*d*]pyrimidine.

Reagents and conditions: (i) aniline derivative **a** or **b** 1 equiv, conc. HCl cat., EtOH, 100 °C, 18 h, sealed vial; (ii) NaH 2 equiv, 4-chlorophenol **c** or 4-chlorothiophenol **d** 1 equiv, DMSO, rt or 50 °C, 24 h.

2.4. Region IV: Synthesis of the thieno[2,3-*d*]pyrimidine series.

To complete this SAR study with more thiophene-containing bio-isosteres, we decided to synthesize the position isomer analogues in thieno[2,3-*d*]pyrimidine series. This could be performed by a S_NAr reaction between the desired nucleophiles **a-d** and a key chlorinated intermediate **25**, obtained from 2-aminothiophene-3-carbonitrile (**Figure 4**).

Figure 4. Thieno[2,3-*d*]pyrimidines initial retrosynthesis pathway.

Thus, 2-aminothiophene-3-carbonitrile was acetylated with Ac_2O to give **22** [33]. Then, the cyclization was made in the presence of H_2O_2 30% in alkaline medium (NaOH) [17,28] leading to thienopyrimidinone **23** [34] which was then chlorodehydroxylated into **24**. Unfortunately, as previously observed in the thieno[3,2-*d*]pyrimidine series, we did not succeed in synthesizing **25**, neither by the *gem*-trichloromethylation of **24** [35], nor by the tetrachlorination of **23** with in a $PCl_5/POCl_3$ mixture. (**Scheme 6**). Acylation of 2-aminothiophene-3-carbonitrile was then realized with trichloroacetyl chloride, to give **26**. Cyclization into lactam **27** was next conducted with a mixture of polyphosphoric acid (PPA)

and phosphoric acid. Nevertheless, once again, it remained impossible to isolate key intermediate **25** via a chlorodehydroxylation reaction. It seemed that **25** was too unstable to be isolated, indeed some traces were detected by LC-MS, but direct engagement of crude **25** in S_NAr reaction did not afford target compounds.

Scheme 6. Various synthesis routes investigated for the synthesis of key intermediate **25**.

Reagents and conditions: (i) Ac_2O , $AcOH$ cat., 120 °C, MW, 10 min; (ii) H_2O_2 35%, $NaOH$, $EtOH/H_2O$, 0 to 55 °C, 15 min; (iii) PCl_5 6 equiv, $POCl_3$ as solvent, 100 °C, MW, 20 min, 800 W; (iv) $POCl_3$, pyridine cat., reflux 24 h; (v) Et_3N , trichloroacetyl chloride, rt, 36 h; (vi) H_3PO_4 , polyphosphoric acid, 70 °C, 3 h; (vii) $POCl_3$ or PCl_5 or oxalyl chloride or P_2O_5 /tetrabutylammonium chloride (TBACl) or $SOCl_2$.

Facing the difficulty to prepare the intermediate **25** and in order to access to the target thieno[2,3-*d*]pyrimidine derivatives, we tried another synthesis pathway based on the cyclization of thiophene derivatives which could be cyclized into thieno[2,3-*d*]pyrimidine rings bearing already both the CCl_3 group and an aniline (or phenol or thiophenol) moiety (**Scheme 7**). Thus the synthesis of amidines **28a** and **28b** was conducted by reacting nitrile **26** with appropriate anilines in presence of $AlCl_3$ at room temperature in CH_2Cl_2 [36]. The crudes residues obtained were a mixture of unreacted starting material **26** (even after 72 h at rt), the expected amidines **28a-b** and cyclized target derivatives **29a** and **29b**. When the temperature was increased to improve the conversion, undesirable side products appeared. Refluxing the obtained amidines **28a-b** in ethanol led to the target compounds **29a** and **29b**, respectively in 36 and 15% yields [37]. For phenol **c** and thiophenol **d**, the same conditions using $AlCl_3$ did not afford the expected derivatives. We then considered the synthesis work of Baati and al.,

describing the addition of thiols onto nitrile groups in presence of gaseous HBr in ether at 0 °C [38]. Indeed, to our delight, by using these reaction conditions, thioimide **28d** was produced, and further cyclization in refluxing ethanol afforded compound **29d** in low yield (**Scheme 7**). Unfortunately, the same reaction conditions were inefficient to obtain compounds **28c** and **29c**. Other conditions were attempted in order to produce the imide **28c** *via* the addition of 4-chlorophenol into the nitrile group, in the presence of NaH [39], Na [40], K₂CO₃ [41] or gaseous HCl [42], but all of them conducted to the same disappointing result. The results of the biological evaluation are presented in **Table 2**.

Scheme 7. Synthesis of 4-substituted-2-(trichloromethyl)thieno[2,3-*d*]pyrimidine **29a-d**.

Reagents and conditions: (i) anilines **a** or **b**, AlCl₃, CH₂Cl₂, rt, 72 h; (ii) 4-chlorothiophenol, gaseous HBr, Et₂O, - 20 °C, 30 min then 18 h at rt; (iii) *i*PrOH, reflux, 5 min.

2.5. Modulation of region IV: Biological evaluation

In region IV, the replacement of the quinazoline ring by thienopyrimidine scaffolds led to molecules displaying similar antiparasmodial activity as hit-molecules **A-D**, with IC₅₀ values in the submicromolar range (0.4 to 0.9 μM) (**Table 2**). However, all the thienopyrimidine derivatives, belonging either to the [3,2-*d*] or [2,3-*d*] series, appeared more cytotoxic (CC₅₀ = 0.7 to 13.3 μM) than the corresponding hit-molecules, leading to lower selectivity indices. Globally, SARs indicate that there is no significant difference between the two thienopyrimidine series, regarding both activity and cytotoxicity.

Table 2. *In vitro* antiplasmodial and cytotoxicity evaluation of compounds **21a-d** and **29a-d**: modulations at region IV

Molecule	Structure	HepG2 CC ₅₀ (μ M)	K1 <i>P.falciparum</i> IC ₅₀ (μ M)	Selectivity Index ^c
21a		3.2	0.6	5.3
21b		0.7	0.9	0.8
21c		4.3	0.6	7.2
21d		6.9	0.4	17.2
29a		6.2	0.5	12.4
29b		4.0	0.6	6.7
29d		13.3	0.8	16.6
Hit A		16	0.4	40
Hit B		150	1.8	83
Hit C		50	1.1	45
Hit D		>25 ^d	0.9	>28
Doxorubicine^a		0.2	-	-
Atovaquone^b		> 15.6 ^d	0.001	15600
Chloroquine^b		30	0.6	50
Doxycycline^b		20	6.0	3.3

^aDoxorubicine was used as a cytotoxic reference-drug; ^bAtovaquone, Chloroquine and doxycycline were used as antimalarial reference-drugs, ^cSelectivity indices were calculated according to the formula : SI = HepG2 CC₅₀ / K1 IC₅₀. ^dHighest concentration tested due to a lack of solubility.

2.6. Region V: Synthesis of quinoline series.

In order to evaluate the influence of the pyrimidine moiety of the quinazoline ring onto the biological profile, we chose to synthesize some analogs in 4-substituted-2-trichloromethylquinoline series bearing, at position 4, the same substituents **a-d** as the hit-molecules. Thus, starting from 4-chloro-2-trichloromethylquinoline **H** obtained by chlorination of 2-methylquinolin-4(1*H*)-one with PCl_5 and POCl_3 [42], we first tried solvent-free $\text{S}_\text{N}\text{Ar}$ reaction conditions [11] with anilines but without any conversion. Compounds **30a** and **30b** were finally obtained by using a catalytic amount of conc. HCl in refluxing EtOH to obtain the desired compounds in 45 and 55% yields respectively. For the phenol **c** and thiophenol **d**, we generated the corresponding anions using NaH before reacting **H**. Thus, phenol derivative **30c** and thiophenol derivative **30d** were obtained, respectively, in 72 and 32% yields (**Scheme 8**). The biological evaluation of these molecules is summarized in **Table 3**.

Scheme 8. Synthesis of 4-substituted-2-trichloromethylquinoline series.

Reagents and conditions : (i) aniline derivative 1 equiv, conc. HCl cat., EtOH , $100\text{ }^\circ\text{C}$, 18 h, sealed vial; (ii) NaH 2 equiv, 4-chlorophenol or 4-chlorothiophenol reagent 1 equiv, DMSO , rt or $50\text{ }^\circ\text{C}$, 24 h.

2.7. Region V: Synthesis of quinoxaline derivatives.

Previously, we reported that 2-trichloromethylquinoxaline **I** displayed an *in vitro* antiparasitic activity ($\text{IC}_{50} = 1.5\text{ }\mu\text{M}$), hindered by a significant *in vitro* cytotoxicity ($\text{CC}_{50} = 3.1\text{ }\mu\text{M}$) [23]. Moreover, the introduction of a phenyl substituent at position 3 (compound **J**) gave a more selective derivative **J** ($\text{SI} = 17.5$ versus 2.1, respectively), thanks to a better

antiplasmodial IC_{50} value ($0.2 \mu M$) [23]. Such interesting results noted with compound **J** prompted us to carry on the investigation of the antiplasmodial potential of the 3-substituted-2-trichloromethylquinoxaline scaffold, in continuation of the pharmacomodulation of the quinazoline moiety in region V (**Figure 5**).

Figure 5. Rational for the synthesis of 2-substituted-3-trichloromethylquinoxaline derivatives.

The first attempted synthetic route involved the reaction of nucleophiles **a-d** with 2-chloro-3-trichloromethylquinoxaline **K** [44] under S_NAr conditions, as previously reported in quinazoline series. Unfortunately, all tested conditions failed to afford the expected derivatives **32a-d**, where no conversion or degradation of starting material could occur.

To solve this problem, we next investigated another pathway by reacting 2-chloro-3-methylquinoxaline **L** [45] with the nucleophiles **a-d** via a S_NAr reaction, before operating the *gem*-trichloromethylation reaction (**Scheme 9**). The best reaction conditions found were to carry out the S_NAr reaction in DMF, under moderate heating, in the presence of Cs_2CO_3 leading to **31a**, **31b**, **31c** and **31d** in, respectively, 19%, 5%, 85% and 83% respectively. Another strategy to improve these low reaction yields was successful only with 3-trifluoromethylaniline **b**, by reacting **L** in aniline **b** (without any solvent) under microwave heating in a sealed vial (69%). Then, from intermediates **31a-d**, a final chlorination step was needed to transform the methyl group into a $-CCl_3$ one. This step was achieved using the classical route elaborated by our team using a $PCl_5/POCl_3$ mixture, under microwave heating. Unfortunately, only the ether and thioether derivatives **32c** and **32d** were obtained in good yields (70-72%) (**Scheme 9**). It seemed that the presence of the amine function of **31a** and **31b** inhibited the chlorination reaction. The results of the biological assays of this series are presented in **Table 3**.

Scheme 9. Synthesis of 2-substituted-3-trichloromethylquinoxaline series.

Reagents and conditions: (i) 2,4-dichloroaniline, 4-chlorophenol or 4-chlorothiophenol 1 equiv, Cs_2CO_3 1 equiv, anh. DMF, 70 °C, 12 h, sealed vial, N_2 ; (ii) 3-trifluoromethylaniline 7 equiv, 140 °C, MW, 45 min, sealed vial; (iii) PCl_5 6 equiv, POCl_3 as solvent, 100 °C, MW, 20 min, 800 W.

2.8. Modulation of region V: biological evaluation.

Thus, the ring variation of region V afforded two series of compounds: a quinoline and a quinoxaline series.

Among the four tested compounds in quinoline series, it appeared that all exerted an antiparasmodial activity globally similar to the ones of **hits A-D**. However, the cytotoxicity toward the HepG2 human cell line was increased for all of the quinoline series, in comparison to the Hits **A-D** and particularly for aniline **b** (6.5 μM vs 150 μM for **hit B**). It must be pointed out that for quinoline **3d**, a lack of solubility in the biological media was observed, as previously reported for **hit D**. Thus, the substitution of the quinazoline ring by a quinoline ring was globally detrimental.

Contrary to the quinoline series, the quinoxaline series appeared very promising. Thus, derivative **32c** showed both a better antiparasmodial activity than its analog **hit C** (0.4 μM vs 1.1 μM) and a great improvement in the cytotoxic profile, its SI reaching 100, compared to 45 for **hit C**. Similarly, the same profile was observed for compound **32d** with a 2-fold activity increase compared to **hit D** (0.45 μM vs 0.9 μM) and a SI of 77 without any solubility impairment, as observed for **hit D**.

To conclude, we noted that for the region V, the suppression of nitrogen *N*-3 of the quinazoline led to generally more cytotoxic compounds while the replacement of the quinazoline moiety by a quinoxaline one was a significant improvement of the biological profile, compound **32c** reaching a SI of 100.

Table 3. *In vitro* antiplasmodial and cytotoxicity evaluation of compounds **30a-d**, and **32c-d**: modulations at region V.

Molecule	Structure	HepG2 CC ₅₀ (μM)	K1 <i>P.falciparum</i> IC ₅₀ (μM)	Selectivity Index ^c
30a		11.7	1.3	9.0
30b		6.5	1.3	5.0
30c		31	1.5	20.7
30d		>15.6 ^d	1.1	>14.2
32c		40.2	0.4	100.5
32d		38.6	0.5	77.2
Hit A		16	0.4	40
Hit B		150	1.8	83
Hit C		50	1.1	45
Hit D		>25 ^d	0.9	>28
Doxorubicine^a		0.2	-	-
Atovaquone^b		> 15.6 ^d	0.001	15600
Chloroquine^b		30	0.6	50
Doxycycline^b		20	6.0	3.3

^aDoxorubicine was used as a cytotoxic reference-drug; ^bAtovaquone, Chloroquine and doxycycline were used as antimalarial reference-drugs; ^cSelectivity indices were calculated according to the formula : SI = HepG2 CC₅₀ / K1 IC₅₀. ^dHighest concentration tested due to a lack of solubility.

Finally, when comparing the biological activities of the synthetic intermediates prepared for studying region V, it was interesting to note that the intermediate **E** was not active ($IC_{50} = 54.5 \mu M$) while intermediates **H** and **K** displayed a significant IC_{50} (0.8 and 0.4 μM respectively). Noticeably, the intermediate **K** possessed a very good SI (265) thanks to a moderate value of its CC_{50} . Moreover, the presence of a substituent at position 3 appeared to be mandatory for displaying a good antiplasmodial profile by comparison of compound **I** which displayed a SI of only 2 (**Figure 6**). It is to note that both derivatives **32c** ($IC_{50} = 0.15 \mu g/mL$) and **K** ($IC_{50} = 0.11 \mu g/mL$) meet the hit to lead *in vitro* TDR criteria [46] for “selectively active antiplasmodial agents”: $IC_{50} < 0.2 \mu g/mL$, $SI > 100$.

Figure 6. Biological assessment of synthetic intermediates.

3. Conclusion

From previously identified antiplasmodial hits **A-D**, in 2-trichloromethyl-4-substituted-quinazoline series, new derivatives were synthesized in order to study the SARs in 5 precisely defined regions (I-V). The modulation of region I showed that the $-CCl_3$ group was mandatory for providing antiplasmodial activity in quinazoline series. In the same series, the study of regions II and III demonstrated that the optimal substitution of the aniline moiety by halogens atoms was 2,4-dichloro and that the *N*-methylation was detrimental toward activity. Then, by modulating regions IV and V, it respectively appeared that the replacement of the quinazoline ring by a quinoline ring led to activity levels similar to the ones of Hits **A-D** but also to increased cytotoxicities values, while the replacement of the quinazoline ring by a quinoxaline ring highlighted derivatives **32c-d**, displaying both a preserved activity and a reduced cytotoxicity, in comparison with hits **A-D**, as summarized in **Table 4**. These molecules now appear as new reference antiplasmodial hits. Finally, the great antiplasmodial potential of intermediate **K** was revealed. Both derivatives **32c** ($IC_{50} = 0.15 \mu g/mL$) and **K** ($IC_{50} = 0.11 \mu g/mL$) meet the hit to lead *in vitro* TDR criteria [46] for “selectively active antiplasmodial agents”: $IC_{50} < 0.2 \mu g/mL$, $SI > 100$, opening the way to the synthesis of novel lead-compounds.

Table 4. Summary of SAR data at regions IV and V.

R
		Previous hits A-D	21a-d	29a,b,d	30a-d	32c-d
IC ₅₀	0,4	0.6	0.5	1.3	-
	CC ₅₀	16	3.2	6.2	11.7	-
	SI	40	5.3	12.4	9.0	-
IC ₅₀	1,8	0.9	0.6	1.3	-
	CC ₅₀	150	0.7	4.0	6.5	-
	SI	83	0.8	6.7	5.0	-
IC ₅₀	1.1	0.6	-	1.5	0.4
	CC ₅₀	50	4.3	-	31	40.2
	SI	45	7.2	-	20.7	100.5
IC ₅₀	0.9	0.4	0.8	1.1	0.5
	CC ₅₀	>25	6.9	13.3	>15.6*	38.6
	SI	>28	17.2	16.6	>14.2	77.2

*Highest concentration tested due to a lack of solubility.

In bold, newly identified hit-molecules

4. Experimental

4.1. Chemistry

Commercial reagents were used as received without additional purification. Melting points were determined on a Kofler bench and are uncorrected. Elemental analysis and HRMS were carried out at the Spectropole, Faculté des Sciences et Techniques de Saint-Jérôme, Marseille, France. NMR spectra were recorded on a Bruker ARX 200 spectrometer or a Bruker AV 250 spectrometer at the Faculté de Pharmacie de Marseille or a BRUKER Avance III nanobay 400 at the the Spectropole, Faculté des Sciences et Techniques de Saint-Jérôme, Marseille (¹H-NMR: 200, 250 or 400 MHz, ¹³C-NMR: 50, 63 or 100 MHz). NMR references were the following: ¹H: CHCl₃ δ = 7.26, DMSO-*d*₆ δ = 2.50 and ¹³C: CHCl₃ δ = 76.9, DMSO-*d*₆ δ = 39.5. Solvents were dried by conventional methods. The following adsorbent was used for column chromatography: silica gel 60 (Merck, particle size 0.063–0.200 mm, 70–230 mesh ASTM). TLC was performed on 5 cm × 10 cm aluminium plates coated with silica gel 60F-

254 (Merck) in an appropriate eluent. Visualization was made with ultraviolet light (234 nm). HRMS spectra were recorded on QStar Elite (Applied Biosystems SCIEX) spectrometer. PEG was the matrix for HRMS. The experimental exact mass was given for the ion which has the maximum isotopic abundance. Purity of synthesized compounds was checked with LC-MS analyses which were realized at the Faculté de Pharmacie de Marseille with a Thermo Scientific Accela High Speed LC System[®] coupled with a single quadrupole mass spectrometer Thermo MSQ Plus[®]. The RP-HPLC column used is a Thermo Hypersil Gold[®] 50 × 2.1 mm (C18 bounded), with particles of 1.9 µm diameter. The volume of sample injected on the column was 1 µL. The chromatographic analysis, total duration of 8 min, is made with the gradient of following solvents: $t = 0$ min, water/methanol 50/50; $0 < t < 4$ min, linear increase in the proportion of water to a ratio water/methanol 95/5; $4 < t < 6$ min, water/methanol 95/5; $6 < t < 7$ min, linear decrease in the proportion of water to return to a ratio 50/50 water/methanol; $6 < t < 7$ min, water/methanol 50/50. The water used was buffered with 5 mM ammonium acetate. The retention times (t_R) of the molecules analyzed are indicated in min. The preparation of 4-chloro-2-trichloromethylquinazoline **E** [24], 2-methyl-*N*-(2,4-dichlorophenyl)quinazolin-4-amine **1** [10], *N*-(2,4-dichlorophenyl)quinazolin-4-amine **9** [27], 4-chloro-2-trichloromethylquinoline **H** [43], methyl 3-acetamidothiophene-2-carboxylate **15** [28], 4-chloro-2-chloromethylquinazoline **F** [24], 4-chloro-2-dichloromethylquinazoline **G** [26], 4-chloro-2-trifluoromethylquinazoline [47], 2-chloro-3-trichloromethylquinoxaline **K** [44] and 2-chloro-3-methylquinoxaline **L** [45] was achieved as described in the literature and characterization were consistent as reported in literature.

4.1.1. General procedure for the preparation of *N*-aryl-2-trichloromethylquinazolin-4-amine (2, 3, 7, 8 and 14)

To a mixture of appropriate 4-chloroquinazoline (1 equiv) in isopropanol (10 mL) and appropriate aniline (0.8 equiv) were added a few drops of concentrated HCl. The vial was then sealed and heated at 70 °C for 2 h. The mixture was allowed to cool to room temperature and was poured into 20 mL of iced water. The mixture was then extracted twice with CH₂Cl₂. The combined organic phases were washed twice with water, then dried (Na₂SO₄) and the solvent was evaporated under reduced pressure. The residue was then purified by silica gel column chromatography (Petroleum Ether/CH₂Cl₂ 1/1).

4.1.1.1. 2-Chloromethyl-*N*-(2,4-dichlorophenyl)quinazolin-4-amine (2)

Starting from 4-chloro-2-chloromethylquinazoline [25] (500 mg, 2.35 mmol) and 2,4-dichloroaniline (304 mg, 1.88 mmol).

Yield 71%. White powder. mp 171 °C. ^1H NMR (200 MHz, CDCl_3) δ = 8.97 (d, J = 8.9 Hz, 1H), 8.21 (bs, 1H), 7.99-7.82 (m, 3H), 7.68-7.60 (m, 1H), 7.47-7.35 (m, 2H), 4.75 (s, 2H). ^{13}C NMR (50 MHz, CDCl_3) δ = 160.9, 157.1, 149.9, 133.6, 128.9, 128.7, 128.0, 127.6, 127.5, 123.9, 122.9, 120.1, 113.9, 47.6. A quaternary C was not observed under these experimental conditions. Anal. Calcd for $\text{C}_{15}\text{H}_{10}\text{Cl}_3\text{N}_3$: C, 53.20; H, 2.98; N, 12.41. Found: C, 53.60; H, 3.12; N, 11.61. LC-MS (ESI, 35 eV): t_R = 4.21 min, m/z 338 $[\text{M}+\text{H}]^+$.

4.1.1.2. 2-Dichloromethyl-*N*-(2,4-dichlorophenyl)quinazolin-4-amine (3)

Starting from 4-chloro-2-dichloromethylquinazoline **G** [26] (500 mg, 2.02 mmol) and 2,4-dichloroaniline (259 mg, 1.62 mmol).

Yield 42%. White powder. mp 203 °C. ^1H NMR (200 MHz, CDCl_3) δ = 9.11 (d, J = 8.9 Hz, 1H), 8.30 (bs, 1H), 8.02-7.85 (m, 3H), 7.73-7.66 (m, 1H), 7.49-7.39 (m, 2H), 6.78 (s, 1H). ^{13}C NMR (50 MHz, CDCl_3) δ = 160.6, 157.6, 149.6, 133.9, 133.6, 129.5, 128.7, 128.3, 128.2, 123.7, 122.9, 120.1, 114.5, 71.8. A quaternary C was not observed under these experimental conditions. Anal. Calcd for $\text{C}_{15}\text{H}_9\text{Cl}_4\text{N}_3$: C, 48.29; H, 2.43; N, 11.26. Found: C, 48.41; H, 2.29; N, 11.03. LC-MS (ESI, 35 eV): t_R = 4.71 min, m/z 372 $[\text{M}+\text{H}]^+$.

4.1.1.3. 2-(chlorodifluoromethyl)-*N*-(2,4-dichlorophenyl)quinazolin-4-amine (7)

Starting from 4-chloro-2-chlorodifluoromethylquinazoline **6** (500 mg, 2.01 mmol) and 2,4-dichloroaniline (293 mg, 1.8 mmol).

Yield 30%. White powder. mp 195 °C. ^1H NMR (400 MHz, CDCl_3) δ = 8.93 (d, J = 9.0 Hz, 1H), 8.38 (bs, 1H), 8.11 (d, J = 8.3 Hz, 1H), 8.00-7.92 (m, 2H), 7.79-7.74 (m, 1H), 7.48 (d, J = 2.5 Hz, 1H), 7.42-7.35 (m, 1H). ^{13}C NMR (100 MHz, CDCl_3) δ = 157.3, 155.6 (t, J = 29 Hz), 149.4, 134.2, 133.2, 130.0, 129.2, 129.0, 128.8, 128.2, 123.9, 122.9 (t, J = 292 Hz), 122.8, 120.1, 114.9. HRMS (ESI): m/z calcd. for $\text{C}_{15}\text{H}_8\text{Cl}_3\text{F}_2\text{N}_3$ $[\text{M}+\text{H}]^+$: 374.9908. Found: 374.9909. LC-MS (ESI, 35 eV): t_R = 4.58 min, m/z 376 $[\text{M}+\text{H}]^+$.

4.1.1.4. *N*-(2,4-dichlorophenyl)-2-(trifluoromethyl)quinazolin-4-amine (8)

Starting from 4-chloro-2-trifluoromethylquinazoline [45] (500 mg, 2.15 mmol) and 2,4-dichloroaniline (279 mg, 1.72 mmol).

Yield 37%. White powder. mp 198 °C. ^1H NMR (200 MHz, CDCl_3) δ = 8.88 (d, J = 9.0 Hz, 1H), 8.29 (bs, 1H), 8.11-8.07 (m, 1H), 7.97-7.89 (m, 2H), 7.78-7.70 (m, 1H), 7.47-7.35 (m, 2H). ^{13}C NMR (50 MHz, CDCl_3) δ = 157.3, 152.1 (q, J = 36 Hz), 149.5, 134.2, 133.2, 130.1, 129.2, 129.1, 128.8, 128.2, 123.9, 122.8, 120.1, 119.8 (q, J = 275 Hz), 115.2. Anal. Calcd for $\text{C}_{15}\text{H}_8\text{Cl}_2\text{F}_3\text{N}_3$: C, 50.30; H, 2.25; N, 11.73. Found: C, 50.39; H, 2.21; N, 11.80.

4.1.1.5. *N*-(2,4-dichlorophenyl)-*N*-methyl-2-trichloromethylquinazolin-4-amine (14)

Starting from 4-chloro-2-(trichloromethyl)quinazoline **E** [24] (500 mg, 1.77 mmol) and *N*-methyl-2,4-dichloroaniline (188 μ L, 1.95 mmol).

Yield 15%. Colorless oil. ^1H NMR (200 MHz, CDCl_3) δ = 8.12-8.08 (m, 1H), 7.61-7.60 (m, 2H), 7.35-7.17 (m, 3H), 6.93-6.89 (m, 1H), 3.63 (s, 3H). ^{13}C NMR (50 MHz, CDCl_3) δ = 161.8, 160.1, 150.9, 143.1, 134.3, 133.1, 133.0, 131.3, 129.9, 129.5, 129.1, 127.1, 124.4, 114.9, 97.4, 41.3. HRMS (ESI): m/z calcd. for $\text{C}_{16}\text{H}_{10}\text{Cl}_5\text{N}_3$ $[\text{M}+\text{H}]^+$: 418.9317. Found: 418.9319. LC-MS (ESI, 35 eV): t_R = 5.18 min, m/z 419.7 $[\text{M}+\text{H}]^+$.

4.1.2. Synthesis of 4-chloro-2-chlorodifluoromethylquinazoline (6)

4.1.2.1. 2-Chloro-*N*-(2-cyanophenyl)-2,2-difluoroacetamide (4)

In a 250 mL round-bottom flask was successively introduced 2-aminobenzonitrile (2.0 g, 16.9 mmol), 4-(*N,N*-dimethylamino)pyridine (DMAP) (1.7 g, 13.3 mmol) and *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (EDCI.HCl) (3.4 g, 17.6 mmol) in CH_2Cl_2 (50 mL). Then, 2-chloro-2,2-difluoroacetic acid (1.4 mL, 16.9 mmol) was added into the mixture and it was stirred for 24 h at rt. The solvent was evaporated under reduced pressure and water (200 mL) was added to the residue. The mixture was gently heated at 40 $^\circ\text{C}$ until a homogeneous solution was obtained. After cooling down to rt, conc. HCl was added to adjust the pH = 1. The resulting precipitate was filtered, thoroughly washed with water and dried in an oven leading 2-chloro-*N*-(2-cyanophenyl)-2,2-difluoroacetamide.

Yield 15%. White powder. mp 84 $^\circ\text{C}$. ^1H NMR (400 MHz, CDCl_3) δ = 8.37 (d, J = 8.5 Hz, 2H), 7.71 (t, J = 7.6 Hz, 2H), 7.36 (t, J = 7.6 Hz, 1H). ^{13}C NMR (100 MHz, CDCl_3) δ = 157.1 (t, J = 30 Hz), 137.6, 134.5, 132.7, 126.3, 121.6, 121.5, (t, J = 300 Hz), 115.3, 103.7. Anal. Calcd for $\text{C}_9\text{H}_5\text{ClF}_2\text{N}_2\text{O}$: C, 46.88; H, 2.19; N, 12.15. Found: C, 46.98; H, 2.13; N, 12.35. LC-MS (ESI, 35 eV): t_R = 0.85 min, m/z 229 $[\text{M}-\text{H}]^-$.

4.1.2.2. 2-Chlorodifluoromethylquinazolin-4(3*H*)-one (5)

In a 100 mL round-bottom flask was successively introduced 2-chloro-*N*-(2-cyanophenyl)-2,2-difluoroacetamide **4** (500 mg, 2.17 mmol), 30% H_2O_2 (2 mL) and a mixture of EtOH (6 mL) and water (8 mL). Then, NaOH (140 mg, 3.47 mmol) was added at 0 $^\circ\text{C}$. The mixture was stirred at rt for 90 min and the volatiles were removed under vacuum. Water (100 mL) was added to the residue and the resulting mixture was acidified to pH = 4 with conc. HCl.

The precipitate obtained was filtered, thoroughly washed with water and dried in an oven gave the 2-chlorodifluoromethylquinazolin-4(3*H*)-one (**5**).

Yield 80%. Beige powder. mp 246 °C. ¹H NMR (400 MHz, CDCl₃) δ = 10.44 (bs, 1H), 8.37 (d, *J* = 8.5 Hz, 1H), 7.89 (d, *J* = 3.7 Hz, 2H), 7.67-7.61 (m, 1H). ¹³C NMR (100 MHz, CDCl₃) δ = 135.6, 129.2, 128.7, 126.9. Five quaternary C were not observed under these experimental conditions. LC-MS (ESI, 35 eV): *t*_R = 0.66 min, *m/z* 229 [M-H]⁻.

4.1.2.3. 4-Chloro-2-chlorodifluoromethylquinazoline (**6**)

In a 20 mL vial was successively introduced 2-chlorodifluoromethylquinazolin-4(3*H*)-one **5** (500 mg, 2.17 mmol) and POCl₃ (1 mL, 10.8 mmol). The mixture was stirred at 140 °C for 10 min under microwave irradiation (800 W). After cooling at rt, the crude was purified by silica gel column chromatography (CH₂Cl₂) to afford 4-chloro-2-chlorodifluoromethylquinazoline.

Yield 75%. Beige powder. mp 90 °C. ¹H NMR (400 MHz, CDCl₃) δ = 8.38 (d, *J* = 8.3 Hz, 1H), 8.23 (d, *J* = 8.3 Hz, 1H), 8.12-8.08 (dt, *J* = 8.2, 1.2 Hz, 1H), 7.89 (dt, *J* = 8.2, 1.2 Hz, 1H). ¹³C NMR (100 MHz, CDCl₃) δ = 164.2, 155.0 (t, *J* = 30 Hz), 150.4, 136.2, 130.9, 129.6, 126.1, 123.8, 122.0 (t, *J* = 292 Hz). Anal. Calcd for C₉H₄Cl₂F₂N₂: C, 43.40; H, 1.62; N, 11.25. Found: C, 43.29; H, 1.56; N, 11.42. LC-MS (ESI, 35 eV): *t*_R = 3.42 min, *m/z* 249 [M+H]⁺.

4.1.3. Representative procedure for the preparation of *N*-(2,4-dihalo-phenyl)-2-trichloromethylquinazolin-4-amine (**10-13**)

A 100 mL round-bottomed single-neck flask equipped with a condenser and a magnetic stir bar was charged with 4-chloro-2-trichloromethylquinazoline **E** [24] (500 mg, 1.77 mmol), 2 equiv. of the appropriated dihalogeno-aniline (3.54 mmol) and 30 mL of isopropanol. The mixture was refluxed for 48 h. After cooling, water was added (50 mL), and the resulting solution was extracted three times with CH₂Cl₂. Then, the combined organic layers were washed with water and dried (Na₂SO₄). Finally, the solvent was evaporated under reduced pressure. The residue was purified by trituration with CH₂Cl₂ and filtration yielding the *N*-(2,4-dihalo-phenyl)-2-trichloromethylquinazolin-4-amines (**10-13**).

4.1.3.1. *N*-(2,4-dibromophenyl)-2-trichloromethylquinazolin-4-amine (**10**)

Yield 55%. White powder. mp 243 °C. ¹H NMR (200 MHz, DMSO-*d*₆) δ = 10.32 (bs, 1H), 8.55 (d, *J* = 8.2 Hz, 1H), 8.02- 7.91 (m, 3H), 7.82-7.59 (m, 3H). ¹³C NMR (50 MHz, DMSO-*d*₆) δ = 160.0, 159.5, 148.9, 136.3, 134.7, 134.4, 131.1, 130.8, 128.5, 128.3, 123.1, 122.5, 119.5, 113.5, 97.7. Anal. Calcd for C₁₅H₈Br₂Cl₃N₃: C, 36.59; H, 1.62; N, 8.46. Found: C, 36.16; H, 1.53; N, 8.50. LC-MS (ESI, 35 eV): *t*_R = 5.41 min, *m/z* 494 [M+H]⁺.

4.1.3.2. *N*-(4-bromo-2-chlorophenyl)-2-trichloromethylquinazolin-4-amine (11)

Yield 63%. White powder. mp 222 °C. ¹H NMR (200 MHz, DMSO-*d*₆) δ = 10.38 (bs, 1H), 8.62 (d, *J* = 8.3 Hz, 1H), 8.07-7.97 (m, 2H), 7.93 (s, 1H), 7.87-7.80 (m, 1H), 7.75 - 7.65 (m, 2H). ¹³C NMR (50 MHz, DMSO-*d*₆) δ = 160.2, 159.7, 149.1, 135.0, 134.6, 132.1, 131.8, 130.8, 130.6, 128.7, 128.5, 123.4, 119.3, 113.7, 97.9. Anal. Calcd for C₁₅H₈BrCl₄N₃: C, 39.86; H, 1.78; N, 9.30. Found: C, 39.83; H, 1.70; N, 9.45. LC-MS (ESI, 35 eV): *t*_R = 5.41 min, *m/z* 450 [M+H]⁺.

4.1.3.3. *N*-(2-bromo-4-chlorophenyl)-2-trichloromethylquinazolin-4-amine (12)

Yield 70%. White powder. mp 223 °C. ¹H NMR (200 MHz, DMSO-*d*₆) δ = 9.09 (d, *J* = 9.0 Hz, 1H), 8.38 (s, 1H), 8.11 (d, *J* = 8.3 Hz, 1H), 8.00-7.89 (m, 2H), 7.77-7.70 (m, 1H), 7.63 (d, *J* = 2.3 Hz, 1H), 7.44 (dd, *J* = 9.0, 2.3 Hz, 1H). ¹³C NMR (50 MHz, DMSO-*d*₆) δ = 160.6, 157.2, 149.6, 134.5, 134.0, 131.7, 130.3, 129.2, 128.8, 128.7, 122.8, 120.1, 120.0, 114.1, 97.5. Anal. Calcd for C₁₅H₈BrCl₄N₃: C, 39.86; H, 1.78; N, 9.30. Found: C, 40.01; H, 1.69; N, 9.26. LC-MS (ESI, 35 eV): *t*_R = 4.82 min, *m/z* 450 [M+H]⁺.

4.1.3.4. *N*-(2,4-difluorophenyl)-2-trichloromethylquinazolin-4-amine (13)

Yield 70%. White powder. mp 165 °C. ¹H NMR (200 MHz, DMSO-*d*₆) δ = 9.00-8.87 (m, 1H), 8.12-8.08 (m, 1H), 7.95-7.66 (m, 4H), 7.07-6.92 (m, 2H). ¹³C NMR (50 MHz, DMSO-*d*₆) δ = 160.7, 157.4, 153.2 (d, *J* = 260 Hz), 152.9 (d, *J* = 260 Hz), 149.6, 133.9, 130.2, 128.5, 123.5 (d, *J* = 8.8 Hz), 123.1 (dd, *J* = 9.5, 4.1 Hz), 120.1, 113.8, 111.4 (dd, *J* = 21.6, 3.6 Hz), 104.3-103 (m, 1CH), 97.6. Anal. Calcd for C₁₅H₈Cl₃F₂N₃: C, 48.09; H, 2.15; N, 11.22. Found: C, 48.45; H, 1.99; N, 11.25. LC-MS (ESI, 35 eV): *t*_R = 4.58 min, *m/z* 374 [M+H]⁺.

4.1.4. 2-Methylthieno[3,2-*d*]pyrimidin-4(3*H*)-one (16)

A solution of methyl 3-acetamidothiophene-2-carboxylate **15** [29] (760 mg, 3.81 mmol) in NH₄OH 25% (9.5 mL) was heated, in a sealed vial, at 105 °C for 3 h. After cooling to room temperature, concentrated HCl was added until pH 8. The precipitate thus formed was filtered, washed with water and dried in a vacuum oven at 40 °C to afford 2-methylthieno[3,2-*d*]pyrimidin-4(3*H*)-one. No further purification necessary.

Yield 63%. White solid. mp 91 °C. ¹H NMR (200 MHz, DMSO-*d*₆) δ = 12.31 (bs, 1 H, NH), 8.12 (d, *J* = 5.2 Hz, 1 H), 7.29 (d, *J* = 5.2 Hz, 1 H), 2.36 (s, 3 H). ¹³C NMR (63 MHz, DMSO-*d*₆) δ = 181.0, 158.8, 158.0, 156.3, 133.9, 124.5, 21.1. LC-MS (ESI, 35eV): *t*_R = 0.64 min, *m/z* 167 [M+H]⁺.

4.1.5. 4-Chloro-2-methylthieno[3,2-*d*]pyrimidine (17) [30]

To a solution of compound **16** (200 mg, 1.20 mmol) in POCl₃ (5.0 mL) were added a few drops of pyridine. The mixture was then refluxed for 24 h. After cooling down to room temperature, the solution was poured on ice and a saturated Na₂CO₃ solution was added until the excess of POCl₃ was neutralized. The aqueous phase was then extracted with EtOAc, the combined organic phases were washed with brine, dried (Na₂SO₄) and the solvent was evaporated under reduced pressure. The residue was then purified by silica gel column chromatography (eluent: Petroleum Ether/EtOAc 8/2) to afford 4-chloro-2-methylthieno[3,2-*d*]pyrimidine.

Yield 97%. White solid. mp 84 °C. ¹H NMR (250 MHz, CDCl₃) δ = 7.96 (d, *J* = 5.5 Hz, 1H), 7.45 (d, *J* = 5.5 Hz, 1H), 2.78 (s, 3H). ¹³C NMR (63 MHz, CDCl₃) δ = 164.7, 162.6, 154.7, 136.7, 127.9, 124.8, 25.5. LC-MS (ESI, 35 eV): *t*_R = 1.74 min, *m/z* 185 [M+H]⁺.

4.1.6. Methyl 3-(2,2,2-trichloroacetamido)thiophene-2-carboxylate (18)

To a solution of methyl 3-aminothiophene-2-carboxylate (500 mg, 3.18 mmol) and triethylamine (440 μL, 3.18 mmol) in tetrahydrofuran (4.0 mL) at 5 °C was added trichloroacetyl chloride (350 μL, 3.18 mmol). The reaction was stirred at 5 °C for 30 min. Water (5.0 mL) was then added and the aqueous phase was extracted with EtOAc. The organic phase was then washed, dried (Na₂SO₄) and the solvent was evaporated to afford methyl 3-(2,2,2-trichloroacetamido)thiophene-2-carboxylate. No further purification necessary.

Yield 99%. White solid. mp 244 °C. ¹H NMR (200 MHz, DMSO-*d*₆) δ = 11.42 (bs, 1 H), 8.06 (d, *J* = 5.4 Hz, 1 H), 7.89 (d, *J* = 5.4 Hz, 1 H), 3.88 (s, 3 H). ¹³C NMR (63 MHz, DMSO-*d*₆) δ = 163.4, 158.1, 140.9, 133.5, 121.0, 113.6, 91.8, 52.2. LC-MS (ESI, 35 eV): *t*_R = 4.11 min, *m/z* 300 [M+H]⁺.

4.1.7. 2-(Trichloromethyl)thieno[3,2-*d*]pyrimidin-4(3*H*)-one (19)

HCl_(g) was bubbled into acetic acid (1.23 mL) for 30 minutes. Then, nitrogen was swept across the solution. This solution was added to a mixture of methyl 3-aminothiophene-2-carboxylate (3.0 g, 19.08 mmol) and trichloroacetonitrile (0.85 mL, 8.48 mmol) in acetic acid (1.86 mL). The reaction was then heated at 100 °C for 18 h. After cooling down to rt, the acetic acid was evaporated under reduced pressure and isopropanol (4.5 mL) was added. This solution was refluxed for 5 min, then refrigerated. The precipitate thus formed was filtered, and then purified by silica gel column chromatography (eluent: Petroleum Ether 100% + 10

mL of EtOAc every 100 mL of eluent) to afford 2-(trichloromethyl)thieno[3,2-*d*]pyrimidin-4(3*H*)-one.

Yield 63%. White solid. mp 237 °C. ¹H NMR (200 MHz, CDCl₃) δ = 10.50 (bs, 1 H), 7.93 (d, *J* = 5.3 Hz, 1 H), 7.51 (d, *J* = 5.3 Hz, 1 H). ¹³C NMR (50 MHz, CDCl₃) δ = 157.7, 155.7, 151.8, 135.7, 126.1, 123.7, 92.0. LC-MS (ESI, 35 eV): *t*_R = 0.78 min, *m/z* 267 [M-H]⁺.

4.1.8. 4-Chloro-2-(trichloromethyl)thieno[3,2-*d*]pyrimidine (20)

To a solution of 2-(trichloromethyl)thieno[3,2-*d*]pyrimidin-4(3*H*)-one **19** (200 mg, 0.74 mmol) in POCl₃ (3.0 mL) were added a few drops of pyridine. The reaction was refluxed for 1 h. After cooling to room temperature, the solution was poured on ice and the aqueous phase was extracted with EtOAc. The combined organic phases were washed with brine, dried (Na₂SO₄) and the solvent was evaporated under reduced pressure. The residue was then purified by silica gel column chromatography (eluent: Petroleum Ether/EtOAc 8/2) to yield 4-chloro-2-(trichloromethyl)thieno[3,2-*d*]pyrimidine.

Yield 99%. White solid. mp 99 °C. ¹H NMR (200 MHz, CDCl₃) δ = 8.21 (d, *J* = 5.5 Hz, 1 H), 7.74 (d, *J* = 5.5 Hz, 1 H). ¹³C NMR (50 MHz, CDCl₃) δ = 161.9, 161.7, 155.6, 139.0, 130.7, 125.6, 96.0. LC-MS (ESI, 35 eV): *t*_R = 4.05 min (no ionization).

4.1.9. *N*-(2,4-Dichlorophenyl)-2-(trichloromethyl)thieno[3,2-*d*]pyrimidin-4-amine (21a)

To a solution of 4-chloro-2-(trichloromethyl)thieno[3,2-*d*]pyrimidine **20** (190 mg, 0.66 mmol) and 2,4-dichloroaniline (106 mg, 0.66 mmol) in ethanol (3.0 mL) were added a few drops of concentrated HCl. The vial was sealed and the reaction was heated at 100 °C for 24 h. The solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (eluent: Petroleum Ether/CH₂Cl₂ 8/2 + 10 mL of CH₂Cl₂ every 100 mL of eluent) to afford *N*-(2,4-dichlorophenyl)-2-(trichloromethyl)thieno[3,2-*d*]pyrimidin-4-amine.

Yield 45%. White solid. mp 197 °C. ¹H NMR (200 MHz, CDCl₃) δ = 8.72 (d, *J* = 8.9 Hz, 1 H), 7.96 (d, *J* = 5.4 Hz, 1 H), 7.66 (d, *J* = 5.4 Hz, 1 H), 7.61 (s, 1 H), 7.48 (d, *J* = 2.3 Hz, 1 H), 7.37 (dd, *J* = 8.9, 2.3 Hz, 1 H). ¹³C NMR (200 MHz, CDCl₃) δ = 161.6, 161.0, 154.6, 134.1, 133.2, 130.0, 129.1, 128.3, 126.2, 125.1, 124.0, 116.2, 97.1. LC-MS (ESI, 35 eV): *t*_R = 4.90 min, *m/z* 410 [M-H]⁺. Anal. Calcd for C₁₃H₆Cl₅N₃S: C, 37.76; H, 1.46; N, 10.16. Found: C, 38.03; H, 1.26; N, 10.30.

4.1.10. 2-(Trichloromethyl)-*N*-[3-(trifluoromethyl)phenyl]thieno[3,2-*d*]pyrimidin-4-amine (21b)

To a solution of 4-chloro-2-(trichloromethyl)thieno[3,2-*d*]pyrimidine **20** (150 mg, 0.52 mmol) and 3-(trifluoromethyl)aniline (87 μ L, 0.52 mmol) in ethanol (2.5 mL) were added a few drops of concentrated HCl. The vial was sealed and the reaction was heated at 100°C for 24 h. The solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (eluent: Petroleum Ether/EtOAc 8/2) to yield 2-(trichloromethyl)-*N*-[3-(trifluoromethyl)phenyl]thieno[3,2-*d*]pyrimidin-4-amine.

Yield 55%. White solid. mp 141 °C. ^1H NMR (200MHz, CDCl_3) δ = 8.42 (s, 1 H), 7.88-7.83 (m, 2 H), 7.58-7.39 (m, 4 H). ^{13}C NMR (50 MHz, CDCl_3) δ = 161.9, 161.0, 155.0, 138.3, 133.9, 131.9, 131.3, 129.7, 125.7, 124.3, 121.3, 118.6, 115.7, 97.4. Anal. Calcd for $\text{C}_{14}\text{H}_7\text{Cl}_3\text{F}_3\text{N}_3\text{S}$: C, 40.75; H, 1.71; N, 10.18. Found: C, 40.99; H, 1.80; N, 10.39. LC-MS (ESI, 35 eV): t_R = 4.87 min, m/z 412 $[\text{M}+\text{H}]^+$

4.1.11. 4-(4-Chlorophenoxy)-2-(trichloromethyl)thieno[3,2-*d*]pyrimidine (21c)

To a solution of 4-chloro-2-(trichloromethyl)thieno[3,2-*d*]pyrimidine **20** (210 mg, 0.73 mmol) and 4-chlorophenol (281 mg, 2.19 mmol) in acetonitrile (5.0 mL) was added K_2CO_3 (302 mg, 2.19 mmol). The reaction was heated at 160 °C for 1h10 under microwave irradiation. The mixture was poured onto a NaOH 10% solution. The aqueous phase was extracted with EtOAc. The organic phases were combined, washed with brine, dried (Na_2SO_4) and the solvent was evaporated under reduced pressure. The residue was then purified by silica gel column chromatography (eluent: Petroleum Ether/ CH_2Cl_2 8/2 + 10 mL of CH_2Cl_2 every 100 mL of eluent) to yield 4-(4-chlorophenoxy)-2-(trichloromethyl)thieno[3,2-*d*]pyrimidine.

Yield 39%. White solid. mp 161 °C. ^1H NMR (200 MHz, CDCl_3) δ = 8.09 (d, J = 5.4 Hz, 1 H), 7.70 (d, J = 5.4 Hz, 1 H), 7.45-7.31 (m, 4 H). ^{13}C NMR (50 MHz, CDCl_3) δ = 163.5, 163.1, 161.5, 150.3, 136.9, 131.4, 129.9, 129.5, 125.2, 122.9, 117.7, 116.4, 96.4. Anal. Calcd for $\text{C}_{13}\text{H}_6\text{Cl}_4\text{N}_2\text{OS}$: C, 41.08; H, 1.59; N, 7.37. Found: C, 41.28; H, 1.72; N, 7.45. LC-MS (ESI, 35 eV): t_R = 5.08 min, m/z 379 $[\text{M}+\text{H}]^+$.

4.1.12. 4-[(4-Chlorophenyl)thio]-2-(trichloromethyl)thieno[3,2-*d*]pyrimidine (21d)

To a solution of 4-chlorothiophenol (100 mg, 0.69 mmol) in DMSO (0.5mL), NaH (33 mg, 1.39 mmol) was added portionwise under inert atmosphere. The mixture was stirred at room temperature for 20 mi. 4-Chloro-2-(trichloromethyl)thieno[3,2-*d*]pyrimidine **20** (200 mg, 0.69 mmol) was then added portionwise and the reaction was stirred for 24 h at rt. Water (1.0 mL) was added and the aqueous phase was then extracted with CH_2Cl_2 . The organic phase was washed with water, brine and then dried (Na_2SO_4). The solvent was evaporated and the residue was purified by silica gel column chromatography (eluent: Petroleum Ether 100% +

10 mL of CH₂Cl₂ every 100 mL of eluent) to afford 4-[(4-chlorophenyl)thio]-2-(trichloromethyl)thieno[3,2-*d*]pyrimidine.

Yield 7%. White solid. mp not determined. ¹H NMR (200 MHz, CDCl₃) δ = 8.02 (d, *J* = 5.5 Hz, 1 H), 7.67-6.64 (m, 3 H), 7.45 (d, *J* = 8.8 Hz, 2 H). ¹³C NMR (50 MHz, CDCl₃) δ = 164.5, 161.1, 159.9, 137.0, 136.6, 136.4, 129.5, 127.5, 125.0, 124.7, 96.8. Anal. Calcd for C₁₃H₆Cl₄N₂S₂: C, 39.41; H, 1.53; N, 7.07. Found: C, 39.75; H, 1.72; N, 7.20. LC-MS (ESI, 35 eV): *t*_R = 5.36 min, *m/z* 395 [M+H]⁺.

4.1.13. *N*-(3-Cyanothiophen-2-yl)acetamide (22)

To a solution of 2-aminothiophene-3-carbonitrile (1.020 g, 8.21 mmol) in acetic anhydride (3.0 mL) were added a few drops of acetic acid. The reaction was heated at 120 °C for 10 min under microwave irradiation. The brown solid thus obtained was filtered, washed with water and dried in a vacuum oven at 40 °C to afford *N*-(3-cyanothiophen-2-yl)acetamide. No further purification necessary.

Yield 90%. Brown solid. mp 210 °C (litt. 210-211 °C). ¹H NMR (200 MHz, DMSO-*d*₆) δ = 11.70 (bs, 1 H), 7.13-7.12 (m, 2 H), 2.20 (s, 3 H). ¹³C NMR (50 MHz, DMSO-*d*₆) δ = 168.4, 149.4, 124.7, 118.7, 114.9, 91.8, 22.5. LC-MS (ESI, 35 eV): *t*_R = 0.81 min, *m/z* 165 [M-H]⁻.

4.1.14. 2-Methylthieno[2,3-*d*]pyrimidin-4(3*H*)-one (23)

To a solution of *N*-(3-cyanothiophen-2-yl)acetamide **22** (1.232 g, 7.41 mmol) and H₂O₂ 30% (6.5 mL) in ethanol (22 mL) and water (4 mL), at 0 °C, was added NaOH in pellets (0.495 g, 12.4 mmol). The reaction was stirred at room temperature for 15 min and then heated at 55 °C until a homogeneous solution was obtained. After cooling to rt, the solvent was evaporated under reduced pressure. Water (10 mL) was added and the mixture was heated to 50 °C. After cooling to rt, a HCl 1N solution was added until pH 4. The precipitate was filtered, washed with water and dried in a vacuum oven at 40 °C to afford 2-methylthieno[2,3-*d*]pyrimidin-4(3*H*)-one. No further purification necessary.

Yield 72%. White solid. mp 210 °C. ¹H NMR (200 MHz, DMSO-*d*₆) δ = 7.90 (s, 1 H), 7.39 (d, *J* = 5.8 Hz, 1 H), 6.93 (d, *J* = 5.8 Hz, 1 H), 2.19 (s, 3 H). ¹³C NMR (50 MHz, DMSO-*d*₆) δ = 166.9, 166.8, 145.8, 123.0, 115.8, 114.8, 23.3. LC-MS (ESI, 35 eV): *t*_R = 0.65 min, *m/z* 167 [M+H]⁺.

4.1.15. 4-Chloro-2-methylthieno[2,3-*d*]pyrimidine (24)

To a solution of 2-methylthieno[2,3-*d*]pyrimidin-4(3*H*)-one **23** (300 mg, 1.80 mmol) in POCl₃ (5.0 mL) were added a few drops of pyridine. The mixture was refluxed for 24 h and, after

cooling to rt, was poured onto ice. A saturated Na₂CO₃ solution was added until complete neutralization of the excess of POCl₃. The aqueous phase was extracted with EtOAc. The combined organic phases were washed with brine, dried (Na₂SO₄) and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (eluent: Petroleum Ether/EtOAc 8/2) to yield 4-chloro-2-methylthieno[2,3-*d*]pyrimidine.

Yield 61%. Yellow solid mp 93 °C. ¹H NMR (200 MHz, CDCl₃) δ = 7.48 (d, *J* = 6.0 Hz, 1 H), 7.35 (d, *J* = 6.0 Hz, 1 H), 2.79 (s, 3 H). ¹³C NMR (50 MHz, CDCl₃) δ = 169.2, 163.0, 154.4, 126.8, 126.6, 119.5, 25.5. LC-MS (ESI, 35 eV): *t*_R = 1.84 min, *m/z* 185 [M+H]⁺.

4.1.16. 2,2,2-Trichloro-*N*-(3-cyanothiophen-2-yl)acetamide (26)

To a solution of 2-aminothiophene-3-carbonitrile (1.08 g, 8.0 mmol) and trichloroacetyl chloride (0.89 mL, 8.0 mmol) in dioxane (15 mL), triethylamine (1.67 mL, 12.0 mmol) was added dropwise. The reaction was stirred at room temperature for 36 h. The aqueous phase was extracted with diethyl ether and the combined organic phases were washed with brine and dried (Na₂SO₄). The solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (eluent: Petroleum Ether/EtOAc 9/1) to afford 2,2,2-trichloro-*N*-(3-cyanothiophen-2-yl)acetamide.

Yield 76%. White solid. mp 127 °C. ¹H NMR (200 MHz, CDCl₃) δ = 9.48 (s, 1 H), 7.08 (s, 2 H). ¹³C NMR (50 MHz, CDCl₃) δ = 158.7, 147.0, 124.8, 120.5, 113.4, 96.8, 90.8. LC-MS (ESI, 35 eV): *t*_R = 0.80 min, *m/z* 267 (M⁺). HRMS (ESI): *m/z* calcd. for C₇H₃Cl₃N₂OS [M+H]⁺: 268.9104. Found: 268.9110.

4.1.17. 2-(Trichloromethyl)thieno[2,3-*d*]pyrimidin-4(3*H*)-one (27)

Compound 2,2,2-trichloro-*N*-(3-cyanothiophen-2-yl)acetamide **26** (809 mg, 3.0 mmol) was added to a mixture of phosphoric acid and polyphosphoric acid (60 mL, 1:1). The solution was stirred at rt for 1 h and then heated at 70 °C for 3 h. After cooling to room temperature, the mixture was poured onto ice. The precipitate was filtered, washed with ice-cold water and dried in a vacuum oven at 40 °C to yield 2-(trichloromethyl)thieno[2,3-*d*]pyrimidin-4(3*H*)-one. No further purification necessary.

Yield 90%. White solid. mp 195 °C. ¹H NMR (200 MHz, DMSO-*d*₆) δ = 8.23 (s, 1 H), 7.56 (d, *J* = 5.8 Hz, 1 H), 7.25 (d, *J* = 5.8 Hz, 1 H). ¹³C NMR (50 MHz, DMSO-*d*₆) δ = 167.4, 158.8, 144.6, 124.1, 119.2, 118.5, 91.6. LC-MS (ESI, 35 eV): *t*_R = 0.69 min, *m/z* 267 [M-H]⁻.

4.1.18. *N*-(2,4-Dichlorophenyl)-2-(trichloromethyl)thieno[2,3-*d*]pyrimidin-4-amine (29a)

To a solution of 2,2,2-trichloro-*N*-(3-cyanothiophen-2-yl)acetamide **27** (200 mg, 0.74 mmol) and 2,4-dichloroaniline (118 mg, 0.74 mmol) in CH₂Cl₂ (0.3 mL), AlCl₃ (100 mg, 0.74 mmol) was added portionwise. The mixture was stirred at rt for 72 h. The solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (eluent: Petroleum Ether/EtOAc 7/3) to afford a white solid **28a** (LC-MS (ESI, 35 eV): *t*_R = 4.00 min, *m/z* 430 (M+H)⁺) which was dissolved in ethanol (2 mL) and the resulting solution was refluxed for 4 h. The solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (eluent: Petroleum Ether/CH₂Cl₂ 8/2 to 5/5) to afford *N*-(2,4-dichlorophenyl)-2-(trichloromethyl)thieno[2,3-*d*]pyrimidin-4-amine.

Yield 36%. White solid. mp 181 °C. ¹H NMR (200 MHz, CDCl₃) δ = 8.94 (d, *J* = 9.0 Hz, 1 H), 7.77 (bs, 1 H), 7.64 (d, *J* = 6.0 Hz, 1 H), 7.47 (d, *J* = 2.3 Hz, 1 H), 7.37 (dd, *J* = 5.4, 2.3 Hz, 2 H). ¹³C NMR (50 MHz, CDCl₃) δ = 167.3, 160.2, 153.8, 133.5, 128.9, 128.8, 128.2, 127.9, 123.4, 122.6, 116.8, 116.2, 96.3. Anal. Calcd for C₁₃H₆Cl₅N₃S: C, 37.76; H, 1.46; N, 10.16. Found: C, 37.95; H, 1.52; N, 10.39. LC-MS (ESI, 35 eV): *t*_R = 5.26 min, *m/z* 412 [M+H]⁺.

4.1.19. 2-(Trichloromethyl)-*N*-(3-(trifluoromethyl)phenyl)thieno[2,3-*d*]pyrimidin-4-amine (29b)

To a solution of 2,2,2-trichloro-*N*-(3-cyanothiophen-2-yl)acetamide **26** (320 mg, 1.18 mmol) and 3-(trifluoromethyl)aniline (200 μL, 1.18 mmol) in CH₂Cl₂ (0.3 mL), AlCl₃ (160 mg, 1.18 mmol) was added portionwise. The mixture was stirred at rt for 72 h. The solvent was evaporated under reduced pressure and the residue **28b** (LC-MS (ESI, 35 eV): *t*_R = 4.99 min, *m/z* 430 [M+H]⁺) was dissolved in ethanol (2.0 mL) and the solution was refluxed for 18 h. The solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (eluent: Petroleum Ether/CH₂Cl₂ 8/2 to 5/5) to afford 2-(Trichloromethyl)-*N*-(3-(trifluoromethyl)phenyl)thieno[2,3-*d*]pyrimidin-4-amine.

Yield 15%. Pale blue solid. mp 124 °C. ¹H NMR (200 MHz, CDCl₃) δ = 8.42 (s, 1 H), 7.92 (d, *J* = 8.1 Hz, 1 H), 7.60 (d, *J* = 5.9 Hz, 1 H), 7.52 (t, *J* = 7.9 Hz, 1 H), 7.48-7.40 (m, 2 H), 7.35 (d, *J* = 5.9 Hz, 1 H). ¹³C NMR (50 MHz, CDCl₃) δ = 167.4, 160.4, 154.3, 138.7, 132.0, 131.3, 129.7, 127.6, 123.4, 120.9, 117.7, 116.6, 116.4, 97.2. Anal. Calcd for C₁₄H₇Cl₃F₃N₃S: C, 40.75; H, 1.71; N, 10.18. Found: C, 41.03; H, 1.86; N, 10.24. LC-MS (ESI, 35 eV): *t*_R = 5.22 min, *m/z* 412 [M+H]⁺.

4.1.20. 4-((4-Chlorophenyl)thio)-2-(trichloromethyl)thieno[2,3-*d*]pyrimidine (29d)

HBr_(g) was bubbled into a solution of 2,2,2-trichloro-*N*-(3-cyanothiophen-2-yl)acetamide **26** (270 mg, 1.0 mmol) and 4-chlorothiophenol (145 mg, 1.0 mmol) in diethyl ether (2.5 mL), at -20 °C, for 30 min. The bubbling was stopped when the mixture became homogeneous (bright yellow solution). The reaction was then stirred at rt for 18 h. Water (2.0 mL) was then added and the aqueous phase was extracted with EtOAc. The organic phase was dried (Na₂SO₄) and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (eluent: Petroleum Ether/CH₂Cl₂ 8/2 + 5 mL of CH₂Cl₂ every 100 mL of eluent) to afford a white solid **28d** (86 mg) (LC-MS (ESI, 35 eV): *t*_R = 4.46 min, *m/z* 413 [M+H]⁺) which was dissolved in ethanol (1.0 mL) and the resulting solution was refluxed for 5 minutes. The solvent was evaporated under reduced pressure to yield 4-((4-chlorophenyl)thio)-2-(trichloromethyl)thieno[2,3-*d*]pyrimidine. No further purification necessary.

Yield 22%. White solid. mp 150 °C (dec.). ¹H NMR (200 MHz, CDCl₃) δ = 7.70 (d, *J* = 6.0 Hz, 1 H), 7.61 (d, *J* = 8.6 Hz, 2 H), 7.46-7.42 (m, 3 H). ¹³C NMR (50, MHz, CDCl₃) δ = 166.5, 164.6, 159.5, 136.8, 136.3, 129.8, 129.5, 126.7, 125.1, 118.8, 96.7. Anal. Calcd for C₁₃H₆Cl₄N₂S₂: C, 39.41; H, 1.53; N, 7.07. Found: C, 39.61; H, 1.61; N, 7.16. LC-MS (ESI, 35 eV): *t*_R = 5.45 min, *m/z* 395 [M+H]⁺.

4.1.21. *N*-(2,4-Dichlorophenyl)-2-trichloromethylquinoline-4-amine (30a)

To a solution of 4-chloro-2-(trichloromethyl)quinoline **H** [43] (281 mg, 1.0 mmol) and 2,4-dichloroaniline (160 mg, 1.0 mmol) in ethanol (5.0 mL) were added a few drops of concentrated HCl. The vial was then sealed and heated at 100 °C overnight. The mixture was allowed to cool to rt and the organic phase was then washed with a solution of NaOH 2N. The aqueous phase was extracted with CH₂Cl₂. The combined organic phases were dried (Na₂SO₄) and the solvent was evaporated under reduced pressure. The residue was then purified by silica gel column chromatography (gradient: Petroleum Ether 100% + 10 mL of EtOAc every 100 mL of eluent) to afford 158 mg of the expected *N*-(2,4-dichlorophenyl)-2-trichloromethylquinoline-4-amine.

Yield 39%. Orange powder. mp 121 °C. ¹H NMR (200 MHz, CDCl₃) δ = 8.22 (d, *J* = 8.3 Hz, 1 H), 8.02 (d, *J* = 8.3 Hz, 1 H), 7.84-7.76 (m, 1 H), 7.68-7.64 (m, 2 H), 7.54-7.51 (m, 1 H), 7.47 (s, 1 H), 7.35-7.29 (m, 1 H), 7.11 (s, 1 H). ¹³C NMR (50 MHz, CDCl₃) δ = 158.6, 147.5, 147.0, 135.4, 131.0, 130.3, 129.7, 128.3, 127.8, 126.9, 122.5, 119.8, 99.7, 98.1. Anal. Calcd for C₁₆H₉Cl₅N₂: C, 47.27; H, 2.23; N, 6.89. Found: C, 47.01; H, 2.28; N, 6.99. LC-MS (ESI, 35 eV): *t*_R = 5.10 min, *m/z* 403 [M-H]⁻.

4.1.22. 2-(Trichloromethyl)-N-[3-(trifluoromethyl)phenyl]quinoline-4-amine (30b)

To a solution of 4-chloro-2-(trichloromethyl)quinoline **H** [43] (281 mg, 1.0 mmol) and 3-(trifluoromethyl)aniline (170 μ L, 1.0 mmol) in ethanol (5.0 mL) were added a few drops of concentrated HCl. The vial was then sealed and heated at 100 °C overnight. The mixture was allowed to cool to rt and the organic phase was then washed with a solution of NaOH 2N. The aqueous phase was extracted with CH₂Cl₂. The combined organic phases were dried (Na₂SO₄) and the solvent was evaporated under reduced pressure. The residue was then purified by silica gel column chromatography (gradient: Petroleum Ether/CH₂Cl₂ 8/2 + 5 mL of CH₂Cl₂ every 100 mL of eluent) to afford 2-(trichloromethyl)-N-[3-(trifluoromethyl)phenyl]quinoline-4-amine.

Yield 73%. Yellow powder. mp 159 °C. ¹H NMR (200 MHz, CDCl₃) δ = 8.22 (d, *J* = 8.3 Hz, 1 H), 8.12 (d, *J* = 8.3 Hz, 1 H), 7.75 (t, *J* = 7.7 Hz, 1 H), 7.69 (s, 1 H), 7.57-7.42 (m, 6 H). ¹³C NMR (50 MHz, CDCl₃) δ = 157.8, 149.0, 146.4, 140.0, 132.8, 131.3, 130.6, 130.1, 127.6, 125.3, 121.7, 120.3, 119.3, 119.1, 119.0, 98.8, 97.5. Anal. Calcd for C₁₇H₁₀Cl₃F₃N₂: C, 50.34; H, 2.48; N, 6.91. Found: C, 50.73; H, 2.69; N, 6.66. LC-MS (ESI, 35 eV) *t*_R = 5.09 min, *m/z* 403 [M-H]⁻.

4.1.23. 4-(4-Chlorophenoxy)-2-(trichloromethyl)quinoline (30c)

To a solution of 4-chlorophenol (141 mg, 1.1 mmol) in DMSO (1.0 mL), NaH (48 mg, 2.0 mmol) was added portionwise under inert atmosphere. The mixture was stirred at rt for 20 min. 4-chloro-2-(trichloromethyl)quinoline **H** [43] (281 mg, 1.0 mmol) was then added portionwise and the reaction was stirred for 24 h at 50 °C. The aqueous phase was then extracted with CH₂Cl₂ and the organic phase was washed with water, dried (Na₂SO₄) and the solvent was evaporated. The residue was then purified by silica gel column chromatography (eluent Petroleum Ether/CH₂Cl₂ 6/4) to afford 4-(4-chlorophenoxy)-2-(trichloromethyl)quinoline.

Yield 72%. White solid. mp 166 °C. ¹H NMR (200 MHz, CDCl₃) δ = 8.34 (d, *J* = 8.3 Hz, 1 H), 8.20 (d, *J* = 8.6 Hz, 1 H), 7.89-7.81 (m, 1 H), 7.71-7.64 (m, 1 H), 7.48 (d, *J* = 8.8 Hz, 2 H), 7.22 (s, 1 H), 7.18 (d, *J* = 8.8 Hz, 2 H). ¹³C NMR (50 MHz, CDCl₃) δ = 162.6, 158.4, 152.6, 147.4, 131.3, 130.6, 130.0, 128.0, 122.2, 121.6, 121.0, 100.6, 97.7. Anal. Calcd for C₁₆H₉Cl₄NO: C, 51.51; H, 2.43; N, 3.75. Found: C, 51.69; H, 2.53; N, 3.81. LC-MS (ESI, 35 eV): *t*_R = 5.52 min, *m/z* 372 [M+H]⁺.

4.1.24. 4-((4-Chlorophenyl)thio)-2-(trichloromethyl)quinoline (30d)

To a solution of 4-chlorothiophenol (159 mg, 1.1 mmol) in DMSO (1.0 mL), NaH (48 mg, 2.0 mmol) was added portionwise under inert atmosphere. The mixture was stirred at rt for 20

min. 4-Chloro-2-(trichloromethyl)quinoline **H** [43] (281 mg, 1.0 mmol) was then added portionwise and the reaction was stirred for 24 h at rt. The aqueous phase was then extracted with CH₂Cl₂ and the organic phase was washed with water, dried (Na₂SO₄) and the solvent was evaporated. The residue was then purified by silica gel column chromatography (eluent: Petroleum Ether/CH₂Cl₂ 8/2) to afford 4-((4-chlorophenyl)thio)-2-(trichloromethyl)quinolone.

Yield 32%. White powder. mp 151 °C. ¹H NMR (200 MHz, CDCl₃) δ = 8.24-8.18 (m, 2 H), 7.87-7.78 (m, 1 H), 7.72-7.64 (m, 1 H), 7.57-7.46 (m, 5 H). ¹³C NMR (50 MHz, CDCl₃) δ = 156.8, 150.2, 145.4, 136.4, 136.1, 131.0, 130.8, 130.5, 129.3, 128.4, 127.7, 125.6, 123.2, 114.1, 97.7. Anal. Calcd for C₁₆H₉Cl₄NS: C, 49.39; H, 2.33; N, 3.60. Found: C, 49.49; H, 2.23; N, 3.53. LC-MS (ESI, 35 eV): *t*_R = 5.76 min, *m/z* 388 [M+H]⁺.

4.1.25. *N*-(2,4-Dichlorophenyl)-3-methylquinoxalin-2-amine (31a)

To a solution of 2-chloro-3-methylquinoxaline **L** [45] (1.0 g, 5.6 mmol) and 2,4-dichloroaniline (0.91 g, 5.6 mmol) in anhydrous DMF (10 mL), Cs₂CO₃ (1.82 g, 5.6 mmol) was added under inert atmosphere. The mixture was stirred at 70 °C for 24 h. After cooling, water then CH₂Cl₂ were added. The organic layer was then washed five times with water and dried with Na₂SO₄. After filtration and evaporation, the resulting solid was purified by silica gel column chromatography (eluent: Petroleum Ether/CH₂Cl₂ 7/3 then 1/1) to afford *N*-(2,4-dichlorophenyl)-3-methylquinoxalin-2-amine.

Yield 19%. White powder. mp 152 °C. ¹H NMR (250 MHz, CDCl₃) δ = 9.06 (d, *J* = 9.0 Hz, 1H), 7.90 (d, *J* = 8.2 Hz, 1H), 7.83 (d, *J* = 8.2 Hz, 1H), 7.67- 7.57 (m, 1H), 7.57- 7.30 (m, 4H), 2.78 (s, 3H). ¹³C NMR (63 MHz, CDCl₃) δ = 147.5, 145.2, 140.4, 138.0, 135.3, 129.6, 128.8, 128.4, 128.0, 127.6, 127.0, 126.4, 123.2, 121.4, 21.1. LC-MS (ESI, 35 eV): *t*_R = 5.49 min, *m/z* 304 [M+H]⁺.

4.1.26. 3-Methyl-*N*-(3-(trifluoromethyl)phenyl)quinoxalin-2-amine (31b)

A solution of 2-chloro-3-methylquinoxaline **L** [45] (500 mg, 2.8 mmol) and 3-trifluoromethylaniline (2.5 mL, 20 mmol) in anhydrous DMF (10 mL) was heated at 140 °C in a sealed vial under microwave irradiation. After completion of the reaction (45 min), CH₂Cl₂ was added, and the organic phase was washed successively with 1M HCl and brine. The organic layer was then dried with Na₂SO₄, filtered and evaporated. The resulting solid was purified by silica gel column chromatography (eluent: CH₂Cl₂) to afford 3-Methyl-*N*-(3-(trifluoromethyl)phenyl)quinoxalin-2-amine.

Yield 69%. Amber powder. mp 98 °C. ¹H NMR (250 MHz, CDCl₃) δ = 8.24 (s, 1H), 7.93 (d, *J* = 8.1 Hz, 1H), 7.87 (dd, *J* = 8.1, 1.1 Hz, 1H), 7.78 (dd, *J* = 8.2, 1.0 Hz, 1H), 7.64 –

7.53 (m, 1H), 7.52-7.39 (m, 2H), 7.37- 7.27 (m, 1H), 6.77 (bs, 1H, NH), 2.65 (s, 3H). ^{13}C NMR (63 MHz, CDCl_3) δ = 147.7, 144.6, 140.3 (d, J = 3.2 Hz), 137.9, 131.5 (q, J = 33.1 Hz), 129.4 (d, J = 6.0 Hz), 128.2, 126.9, 126.1, 124.3 (q, J = 272.9 Hz), 122.8, 119.6 (d, J = 3.8 Hz), 116.6 (d, J = 3.4 Hz), 21.03. LC-MS (ESI, 35 eV): t_{R} = 4.75 min, m/z 304 $[\text{M}+\text{H}]^+$.

4.1.27. 2-(4-Chlorophenoxy)-3-methylquinoxaline (31c)

To a solution of 2-chloro-3-methylquinoxaline **L** [45] (650 mg, 3.6 mmol) and 4-chlorophenol (0.37 mL, 3.6 mmol) in anhydrous DMF (15 mL), Cs_2CO_3 (1.19 g, 3.6 mmol) was added under inert atmosphere. The mixture was stirred at 70 °C overnight. After completion of the reaction, water was added, leading to a precipitate which was separated by filtration. The resulting precipitate was then thoroughly washed with water. The precipitate was dissolved in CH_2Cl_2 and dried with Na_2SO_4 . After filtration and evaporation, the resulting solid was purified by silica gel column chromatography (eluent: Petroleum Ether/ CH_2Cl_2 1/1) to afford 2-(4-chlorophenoxy)-3-methylquinoxaline.

Yield 85%. Off-white powder. mp 108 °C. ^1H NMR (250 MHz, CDCl_3) δ = 7.69 (dd, J = 6.1, 3.7 Hz, 1H), 7.57 (dd, J = 6.3, 3.5 Hz, 2H), 7.46-7.38 (m, 2H), 7.28 – 7.19 (m, 2H), 2.81 (s, 3H). ^{13}C NMR (63 MHz, CDCl_3) δ = 155.8, 151.5, 147.9, 139.6, 139.4, 130.6, 129.7, 129.4, 128.1, 127.6, 127.4, 123.3, 20.7. LC-MS (ESI, 35 eV): t_{R} = 4.35 min, m/z 271 $[\text{M}+\text{H}]^+$.

4.1.28. 2-((4-Chlorophenyl)thio)-3-methylquinoxaline (31d)

To a solution of 2-chloro-3-methylquinoxaline **L** [45] (500 mg, 2.8 mmol) and 4-chlorothiophenol (405 mg, 2.8 mmol) in anhydrous DMF (10 mL), Cs_2CO_3 (912 mg, 2.8 mmol) was added under inert atmosphere. The mixture was stirred at 70 °C overnight. After completion of the reaction, water was added, leading to a precipitate which was separated by filtration. The resulting precipitate was then thoroughly washed with water. The precipitate was dissolved in CH_2Cl_2 and dried with Na_2SO_4 . After filtration and evaporation, the resulting solid was purified by silica gel column chromatography (eluent: Petroleum Ether/ CH_2Cl_2 1/1) to afford 2-((4-chlorophenyl)thio)-3-methylquinoxaline.

Yield 83%. Beige powder. mp 118 °C. ^1H NMR (250 MHz, CDCl_3) δ = 7.96-7.92 (m, 1H), 7.72-7.68 (m, 1H), 7.60-7.54 (m, 4H), 7.44 (d, J = 6.6 Hz, 2H), 2.77 (s, 3H). ^{13}C NMR (101 MHz, CDCl_3) δ = 155.3, 151.4, 141.4, 140.0, 136.7, 135.6, 129.5, 129.2, 128.6, 128.3, 128.1, 127.2, 22.4. LC-MS (ESI, 35 eV): t_{R} = 5.51 min, m/z 287 $[\text{M}+\text{H}]^+$.

4.1.29. 2-(4-Chlorophenoxy)-3-(trichloromethyl)quinoxaline (32c)

To a solution of 2-(4-chlorophenoxy)-3-methylquinoxaline **31c** (624 mg, 2.8 mmol) and PCl_5 (2.88 g, 16.8 mmol), POCl_3 was added to make a slurry (ca 5 mL). The mixture was then

heated in a multimode microwave oven at 100 °C, 800 W for 20 min. After completion of the reaction, the mixture was poured into ice and was then neutralized with Na₂CO₃. The resulting solution was extracted with CH₂Cl₂ and dried with Na₂SO₄. After filtration and evaporation, the resulting solid was purified by silica gel column chromatography (eluent: Petroleum Ether/CH₂Cl₂ 7/3) to afford 2-(4-chlorophenoxy)-3-(trichloromethyl)quinoxaline.

Yield 72%. White powder. mp 108 °C. ¹H NMR (250 MHz, CDCl₃) δ = 8.11 (d, *J* = 8.0 Hz, 1H), 7.72-7.63 (m, 3H), 7.38 (dd, *J* = 6.6, 3.5 Hz, 2 H), 7.22 (dd, *J* = 6.6, 3.5 Hz, 2 H). ¹³C NMR (101 MHz, CDCl₃) δ = 152.5, 151.2, 142.4, 141.2, 137.0, 132.3, 131.1, 129.9, 129.8, 128.8, 127.1, 123.2, 94.9. LC-MS (ESI, 35 eV): *t*_R = 5.63 min, (no ionization). HRMS (ESI): *m/z* calcd. for C₁₅H₈Cl₄N₂O [M+H]⁺: 372.9464. Found: 372.9461. Anal. Calcd for C₁₅H₈Cl₄N₂O: C, 48.16; H, 2.16; N, 7.49. Found: C, 48.37; H, 2.32; N, 7.42.

4.1.30. 2-((4-Chlorophenyl)thio)-3-(trichloromethyl)quinoxaline (32d)

To a solution of 2-((4-chlorophenyl)thio)-3-methylquinoxaline **31d** (623 mg, 2.2 mmol) and PCl₅ (2.7 g, 13.0 mmol), POCl₃ was added to make a slurry (ca 6 mL). The mixture was then heated in a multimode microwave oven at 100 °C, 800 W for 20 min. After completion of the reaction, the mixture was poured into ice and was then neutralized with Na₂CO₃. The resulting solution was extracted with CH₂Cl₂ and dried with Na₂SO₄. After filtration and evaporation, the resulting solid was purified by silica gel column chromatography (eluent: Petroleum Ether/CH₂Cl₂ 9/1) to afford 2-((4-chlorophenyl)thio)-3-(trichloromethyl)quinoxaline.

Yield 72%. Yellow powder. mp 92 °C. ¹H NMR (250 MHz, CDCl₃) δ = 8.15- 8.04 (m, 1H), 7.76-7.64 (m, 3H), 7.56 (d, *J* = 8.6 Hz, 2H), 7.43 (d, *J* = 8.6 Hz, 2H). ¹³C NMR (63 MHz, CDCl₃) δ = 152.9, 147.7, 142.0, 137.2, 137.0, 136.0, 132.1, 129.9, 129.8, 129.5, 128.7, 127.8, 96.7. LC-MS (ESI, 35 eV): *t*_R = 6.59 min, *m/z* 389 [M+H]⁺. Anal. Calcd for C₁₅H₈Cl₄N₂S: C, 46.18; H, 2.07; N, 7.18. Found: C, 46.33; H, 2.26; N, 6.97.

4.2. Biology

4.2.1. *In vitro* Antiplasmodial evaluation

In this study, a K1 culture-adapted *P. falciparum* strain resistant to chloroquine, pyrimethamine and proguanil was used in an *in vitro* culture. Maintenance in continuous culture was done as described previously by Trager and Jensen [48]. Cultures were maintained in fresh A+ human erythrocytes at 2.5% hematocrit in complete medium (RPMI 1640 with 25 mM HEPES, 25 mM NaHCO₃, 10% of A+ human serum) at 37 °C under reduced O₂ atmosphere (gas mixture 10% O₂, 6% CO₂, and 84% N₂). Parasitaemia was maintained daily between 1% and 6%. The *P. falciparum* drug susceptibility test was carried

out by comparing quantities of DNA in treated and control cultures of parasite in human erythrocytes according to a SYBR Green I fluorescence-based method [49] using a 96-well fluorescence plate reader. Compounds, previously dissolved in DMSO (final concentration less than 0.5% v/v) were incubated in a total assay volume of 200 μ L of synchronized culture suspension (2% hematocrit and 0.4% parasitaemia) for 72 h in a humidified atmosphere (84% N₂, 10% O₂ and 6% CO₂) at 37 °C, in 96-well flat bottom plates. Duplicate assays were performed for each sample. After incubation, 125 μ L supernatant was discarded and cells were washed twice with 125 μ L 1X PBS. 15 μ L re-suspended cells were transferred to 96-well flat bottom nonsterile black plates (Greiner Bio-one) already containing 15 μ L of the SYBR Green I lysis buffer (2X SYBR Green I, 20 mM Tris base pH 7.5, 20 mM EDTA, 0.008% w/v saponin, 0.08% w/v Triton X-100). Negative control, treated by solvents (DMSO or H₂O) and positive controls (atovaquone, chloroquine and doxycycline) were added to each set of experiments. Plates were incubated for 15 min at 37 °C and then read on a TECAN Infinite F-200 spectrophotometer with excitation and emission wavelengths at 485 and 535 nm, respectively. The concentrations of compounds required to induce a 50% decrease of parasite growth (IC₅₀ K1) were calculated from three independent experiments.

4.2.2. *In vitro* Cytotoxicity evaluation

HepG2 cell line was maintained at 37 °C, 6% CO₂ with 90% humidity in RPMI supplemented with 10% foetal bovine serum, 1% L-glutamine (200 mM) and penicillin (100 U/mL) / streptomycin (100 μ g/mL) (complete RPMI medium). The evaluation of the tested molecules cytotoxicity on the HepG2 (hepatocarcinoma cell line purchased from ATCC, ref HB-8065) cell line was performed according to the method of Mosmann [50] with slight modifications. Briefly, 5.10³ cells in 100 μ L of complete medium were inoculated into each well of 96-well plates and incubated at 37 °C in a humidified 6% CO₂. After 24 h incubation, 100 μ L of medium with various product concentrations dissolved in DMSO (final concentration less than 0.5% v/v) were added and the plates were incubated for 72 h at 37 °C. Triplicate assays were performed for each sample. Each plate-well was then microscope-examined for detecting possible precipitate formation before the medium was aspirated from the wells. 100 μ L of MTT (3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl-2H-tetrazolium bromide) solution (0.5 mg/mL in medium without FCS) were then added to each well. Cells were incubated for 2 h at 37 °C. After this time, the MTT solution was removed and DMSO (100 μ L) was added to dissolve the resulting blue formazan crystals. Plates were shaken vigorously (700 rpm) for 10 min. The absorbance was measured at 570 nm with 630 nm as reference wavelength using a

BIO-TEK ELx808 Absorbance Microplate Reader. DMSO was used as blank and doxorubicin (purchased from Sigma Aldrich) as positive control. Cell viability was calculated as percentage of control (cells incubated without compound). The 50% cytotoxic concentration (CC₅₀) was determined from the dose–response curve by using the TableCurve software 2D v.5.0. CC₅₀ values represent the mean value calculated from three independent experiments.

Acknowledgement

This work was supported by Aix-Marseille Université and the CNRS. The authors thank Dr Vincent Remusat for the NMR spectra.

References

- [1] WHO World Malaria Report 2015, WHO. <http://www.who.int/malaria/publications/world-malaria-report-2015/report/en/> (accessed February 15, 2016).
- [2] A.P. Phyto, S. Nkhoma, K. Stepniewska, E.A. Ashley, S. Nair, R. McGready, M.C. Ler, S. Al-Saai, A.M. Dondorp, K.M. Lwin, P. Singhasivanon, N.P. Day, N.J. White, T.J. Anderson, F. Nosten, Emergence of artemisinin-resistant malaria on the western border of Thailand: a longitudinal study, *Lancet*. 379 (2012) 1960–1966.
- [3] M.D. Spring, J.T. Lin, J.E. Manning, P. Vanachayangkul, S. Somethy, R. Bun, Y. Se, S. Chann, M. Ittiverakul, P. Sia-ngam, W. Kuntawunginn, M. Arsanok, N. Buathong, S. Chaorattanakawee, P. Gosi, W. Ta-aksorn, N. Chanarat, S. Sundrakes, N. Kong, T.K. Heng, S. Nou, P. Teja-isavadharm, S. Pichyangkul, S.T. Phann, S. Balasubramanian, J.J. Juliano, S.R. Meshnick, C.M. Chour, S. Prom, C.A. Lanteri, C. Lon, D.L. Saunders, Dihydroartemisinin-piperaquine failure associated with a triple mutant including kelch13 C580Y in Cambodia: an observational cohort study, *Lancet Infect. Dis.* 15 (2015) 683–691.
- [4] L. Paloque, A.P. Ramadani, O. Mercereau-Puijalon, J.-M. Augereau, F. Benoit-Vical, *Plasmodium falciparum*: multifaceted resistance to artemisinins, *Malar. J.* 15 (2016).
- [5] B. St Laurent, B. Miller, T.A. Burton, C. Amaratunga, S. Men, S. Sovannaroeth, M.P. Fay, O. Miotto, R.W. Gwadz, J.M. Anderson, R.M. Fairhurst, Artemisinin-resistant

- Plasmodium falciparum* clinical isolates can infect diverse mosquito vectors of Southeast Asia and Africa, *Nat. Commun.* 6 (2015) 8614.
- [6] C. Mendoza-Martínez, J. Correa-Basurto, R. Nieto-Meneses, A. Márquez-Navarro, R. Aguilar-Suárez, M.D. Montero-Cortes, B. Noguera-Torres, E. Suárez-Contreras, N. Galindo-Sevilla, Á. Rojas-Rojas, A. Rodríguez-Lezama, F. Hernández-Luis, Design, synthesis and biological evaluation of quinazoline derivatives as anti-trypanosomatid and anti-plasmodial agents, *Eur. J. Med. Chem.* 96 (2015) 296–307.
- [7] N.A. Malmquist, T.A. Moss, S. Mecheri, A. Scherf, M.J. Fuchter, Small-molecule histone methyltransferase inhibitors display rapid antimalarial activity against all blood stage forms in *Plasmodium falciparum*, *Proc. Natl. Acad. Sci. U. S. A.* 109 (2012) 16708–16713.
- [8] S. Sundriyal, N.A. Malmquist, J. Caron, S. Blundell, F. Liu, X. Chen, N. Srimongkolpithak, J. Jin, S.A. Charman, A. Scherf, M.J. Fuchter, Development of diaminoquinazoline histone lysine methyltransferase inhibitors as potent blood-stage antimalarial compounds, *ChemMedChem*. 9 (2014) 2360–2373.
- [9] A. Mishra, K. Srivastava, R. Tripathi, S.K. Puri, S. Batra, Search for new pharmacophores for antimalarial activity. Part III: synthesis and bioevaluation of new 6-thioureido-4-anilinoquinazolines, *Eur. J. Med. Chem.* 44 (2009) 4404–4412.
- [10] L.A. Dunn, A.G. Burgess, K.G. Krauer, L. Eckmann, P. Vanelle, M.D. Crozet, F.D. Gillin, P. Upcroft, J.A. Upcroft, A new-generation 5-nitroimidazole can induce highly metronidazole-resistant *Giardia lamblia* in vitro, *Int. J. Antimicrob. Agents*. 36 (2010) 37–42.
- [11] P. Verhaeghe, N. Azas, S. Hutter, C. Castera-Ducros, M. Laget, A. Dumètre, M. Gasquet, J.-P. Reboul, S. Rault, P. Rathelot, P. Vanelle, Synthesis and in vitro antiplasmodial evaluation of 4-anilino-2-trichloromethylquinazolines, *Bioorg. Med. Chem.* 17 (2009) 4313–4322.
- [12] Y. Kabri, N. Azas, A. Dumètre, S. Hutter, M. Laget, P. Verhaeghe, A. Gellis, P. Vanelle, Original quinazoline derivatives displaying antiplasmodial properties, *Eur. J. Med. Chem.* 45 (2010) 616–622.
- [13] P. Verhaeghe, N. Azas, M. Gasquet, S. Hutter, C. Ducros, M. Laget, S. Rault, P. Rathelot, P. Vanelle, Synthesis and antiplasmodial activity of new 4-aryl-2-trichloromethylquinazolines, *Bioorg. Med. Chem. Lett.* 18 (2008) 396–401.
- [14] C. Castera-Ducros, N. Azas, P. Verhaeghe, S. Hutter, P. Garrigue, A. Dumètre, L. MBatchi, M. Laget, V. Remusat, F. Sifredi, S. Rault, P. Rathelot, P. Vanelle, Targeting

- the human malaria parasite *Plasmodium falciparum*: *in vitro* identification of a new antiplasmodial hit in 4-phenoxy-2-trichloromethylquinazoline series, *Eur. J. Med. Chem.* 46 (2011) 4184–4191.
- [15] P. Verhaeghe, A. Dumètre, C. Castera-Ducros, S. Hutter, M. Laget, C. Fersing, M. Prieri, J. Yzombard, F. Sifredi, S. Rault, P. Rathelot, P. Vanelle, N. Azas, 4-Thiophenoxy-2-trichloromethylquinazolines display *in vitro* selective antiplasmodial activity against the human malaria parasite *Plasmodium falciparum*, *Bioorg. Med. Chem. Lett.* 21 (2011) 6003–6006.
- [16] N. Primas, P. Verhaeghe, A. Cohen, C. Kieffer, A. Dumètre, S. Hutter, S. Rault, P. Rathelot, N. Azas, P. Vanelle, A New Synthetic Route to Original Sulfonamide Derivatives in 2-Trichloromethylquinazoline Series: A Structure-Activity Relationship Study of Antiplasmodial Activity, *Molecules*. 17 (2012) 8105–8117.
- [17] C. Kieffer, P. Verhaeghe, N. Primas, C. Castera-Ducros, A. Gellis, R. Rosas, S. Rault, P. Rathelot, P. Vanelle, Sonogashira cross-coupling reaction in 4-chloro-2-trichloromethylquinazoline series is possible despite a side dimerization reaction, *Tetrahedron*. 69 (2013) 2987–2995.
- [18] A. Gellis, C. Kieffer, N. Primas, G. Lanzada, M. Giorgi, P. Verhaeghe, P. Vanelle, A new DMAP-catalyzed and microwave-assisted approach for introducing heteroaryl amino substituents at position 4 of the quinazoline ring, *Tetrahedron* 70 (2014) 8257–8266.
- [19] A. Gellis, N. Primas, S. Hutter, G. Lanzada, V. Remusat, P. Verhaeghe, P. Vanelle, N. Azas, Looking for new antiplasmodial quinazolines: DMAP-catalyzed synthesis of 4-benzyloxy- and 4-aryloxy-2-trichloromethylquinazolines and their *in vitro* evaluation toward *Plasmodium falciparum*, *Eur. J. Med. Chem.* 119 (2016) 34–44.
- [20] H. Kikuchi, K. Yamamoto, S. Horoiwa, S. Hirai, R. Kasahara, N. Hariguchi, M. Matsumoto, Y. Oshima, Exploration of a New Type of Antimalarial Compounds Based on Febrifugine, *J. Med. Chem.* 49 (2006) 4698–4706.
- [21] A. Cohen, P. Suzanne, J.-C. Lancelot, P. Verhaeghe, A. Lesnard, L. Basmacıyan, S. Hutter, M. Laget, A. Dumètre, L. Paloque, E. Deharo, M.D. Crozet, P. Rathelot, P. Dallemagne, A. Lorthiois, C.H. Sibley, P. Vanelle, A. Valentin, D. Mazier, S. Rault, N. Azas, Discovery of new thienopyrimidinone derivatives displaying antimalarial properties toward both erythrocytic and hepatic stages of *Plasmodium*, *Eur. J. Med. Chem.* 95 (2015) 16–28.

- [22] D. González Cabrera, C. Le Manach, F. Douelle, Y. Younis, T.-S. Feng, T. Paquet, A.T. Nchinda, L.J. Street, D. Taylor, C. de Kock, L. Wiesner, S. Duffy, K.L. White, K.M. Zabiulla, Y. Sambandan, S. Bashyam, D. Waterson, M.J. Witty, S.A. Charman, V.M. Avery, S. Wittlin, K. Chibale, 2,4-Diaminothienopyrimidines as Orally Active Antimalarial Agents, *J. Med. Chem.* 57 (2014) 1014–1022.
- [23] N. Primas, P. Suzanne, P. Verhaeghe, S. Hutter, C. Kieffer, M. Laget, A. Cohen, J. Broggi, J.-C. Lancelot, A. Lesnard, P. Dallemagne, P. Rathelot, S. Rault, P. Vanelle, N. Azas, Synthesis and in vitro evaluation of 4-trichloromethylpyrrolo[1,2-*a*]quinoxalines as new antiplasmodial agents, *Eur. J. Med. Chem.* 83 (2014) 26–35.
- [24] P. Verhaeghe, P. Rathelot, A. Gellis, S. Rault, P. Vanelle, Highly efficient microwave assisted α -trichlorination reaction of α -methylated nitrogen containing heterocycles, *Tetrahedron*. 62 (2006) 8173–8176.
- [25] C.J. Shishoo, M.B. Devani, V.S. Bhadti, S. Ananthan, G.V. Ullas, Reaction of nitriles under acidic conditions: a novel, direct formation of condensed 4-chloropyrimidines, *Tetrahedron Lett.* 24 (1983) 4611–4612.
- [26] C.J. Shishoo, M.B. Devani, V.S. Bhadti, K.S. Jain, S. Ananthan, Reaction of nitriles under acidic conditions. Part VI. Synthesis of condensed 4-chloro- and 4-aminopyrimidines from ortho-aminonitriles, *J. Heterocycl. Chem.* 27 (1990) 119–126.
- [27] Z. Wang, C. Wang, Y. Sun, N. Zhang, Z. Liu, J. Liu, A novel strategy to the synthesis of 4-anilinoquinazoline derivatives, *Tetrahedron*. 70 (2014) 906–913.
- [28] V. Bavetsias, E.A. Henderson, E. McDonald, Cyclopenta[*g*]quinazolinone-based inhibitors of thymidylate synthase targeting α -folate receptor overexpressing tumours: synthetic approaches to 4-{*N*-[(6*RS*)-2-hydroxymethyl-4-oxo-3,4,7,8-tetrahydro-6*H*-cyclopenta[*g*]quinazolin-6-yl]-*N*-(prop-2-ynyl)amino}benzoic acid, *Tetrahedron*. 63 (2007) 1537–1543.
- [29] L. Englert, K. Silber, H. Steuber, S. Brass, B. Over, H.-D. Gerber, A. Heine, W.E. Diederich, G. Klebe, Fragment-Based Lead Discovery: Screening and Optimizing Fragments for Thermolysin Inhibition, *ChemMedChem*. 5 (2010) 930–940.
- [30] R.J. Gillespie, D.R. Adams, D. Bebbington, K. Benwell, I.A. Cliffe, C.E. Dawson, C.T. Dourish, A. Fletcher, S. Gaur, P.R. Giles, A.M. Jordan, A.R. Knight, L.J.S. Knutsen, A. Lawrence, J. Lerpiniere, A. Misra, R.H.P. Porter, R.M. Pratt, R. Shepherd, R. Upton, S.E. Ward, S.M. Weiss, D.S. Williamson, Antagonists of the human adenosine A2A receptor. Part 1: Discovery and synthesis of thieno[3,2-*d*]pyrimidine-4-methanone derivatives, *Bioorg. Med. Chem. Lett.* 18 (2008) 2916–2919.

- [31] G.D. Madding, M.D. Thompson, Regioselective syntheses of 2-amino-4,5-dialkylthiophene-3-carboxylates and their conversion to 3,4-dihydro-4-oxothieno[2,3-*d*]pyrimidine-2-carboxylates, *J. Heterocycl. Chem.* 24 (1987) 581–587.
- [32] W. Ried, R. Giesse, New 4-Hydroxythienopyrimidines, *Angew. Chem. Int. Ed. Engl.* 7 (1968) 136–136.
- [33] G. Hallas, A.D. Towns, Dyes derived from aminothiophenes. Part 4: Synthesis of some nitro-substituted thiophene-based azo disperse dyes, *Dyes Pigments.* 33 (1997) 319–336.
- [34] P. Herdewijn, J. De, L.-J. Gao, M.-Y. Jang, B. Vanderhoydonck, M.J.A. Waer, Y. Lin, J.F. Herman, T.A.M. Louat, Novel Bicyclic Heterocycles, WO2010103130, 2010. (accessed February 3, 2016).
- [35] M. Robba, J. Lecomte, M. Cugnon de Sevrécourt, Thienopyrimidines. VI. Study of Halothieno[2,3-*d*]pyrimidines, *Bull. Soc. Chim. Fr. Partie I.* (1975) 592–597.
- [36] M. Bobošíková, W. Clegg, S.J. Coles, M. Dandárová, M.B. Hursthouse, T. Kiss, A. Krutošíková, T. Liptaj, N. Prónayová, C.A. Ramsden, The oxidative rearrangement of furan-2-carboximidamides: preparation and properties of 2-acylamino-furans, *J. Chem. Soc. [Perkin 1]*. (2001) 680–689.
- [37] N. Senhorães, A. Dias, L. Conde, M. Proença, One-Pot Regioselective Synthesis of 2,6,9-Trisubstituted Adenines, *Synlett.* 2011 (2011) 181–186.
- [38] R. Baati, V. Gouverneur, C. Mioskowski, An Improved Method for the Preparation of Amidines via Thiophenylimidic Esters, *Synthesis.* 1999 (1999) 927–929.
- [39] S.V. Pan'kov, N.A. Belyakova, V.V. Vishnyakov, P.P. Purygin, Reactions of 1-cyanoimidazoles with phenols, *Russ. J. Org. Chem.* 37 (2001) 426–429.
- [40] K.W. Breukink, L.H. Krol, P.E. Verkade, B.M. Wepster, Preparation of quinazoline derivatives through ring-closure of aromatic *o*-cyano(acylamino) compounds in alkaline alcoholic or phenolic medium. I. 4-RO-substituted quinazolines, *Recl. Trav. Chim. Pays-Bas Belg.* 76 (1957) 401–14.
- [41] W.M. Koppes, H.G. Adolph, Pressure-induced cyclotrimerization of electron-deficient nitriles. Catalysis by acidic alcohols and phenols, *J. Org. Chem.* 46 (1981) 406–12.
- [42] C. Alonso-Alija, M. Michels, H. Schirok, K.-H. Schlemmer, J. Bell, M.F. Fitzgerald, S. Dodd, A. Gill, Preparation of amino(monocyclic aroyl)pyridinones that inhibit p38 map kinase for use as antiinflammatory agents., WO 2003076405, 2003.
- [43] T. Kato, N. Katagiri, A. Wagai, Trichloromethylquinolines: Synthesis and Reaction with Trimethyl Phosphite, *Chem Pharm Bull.* 29 (1981) 1069–1075.

- [44] S. Goswami, A.C. Maity, H.-K. Fun, Facile Synthesis of 6-Trichloromethylpterin and 2-Chloro-3-trichloromethylquinoxaline along with a Library of Trichloromethyl Heterocycles Using *N*-Chlorosuccinimide and Triphenyl Phosphine, *Chem. Lett.* 36 (2007) 552–553.
- [45] S. Wagle, A.V. Adhikari, N.S. Kumari, Synthesis of some new 4-styryltetrazolo[1,5-*a*]quinoxaline and 1-substituted-4-styryl[1,2,4]triazolo[4,3-*a*]quinoxaline derivatives as potent anticonvulsants, *Eur. J. Med. Chem.* 44 (2009) 1135–1143.
- [46] S. Nwaka, B. Ramirez, R. Brun, L. Maes, F. Douglas, R. Ridley, Advancing Drug Innovation for Neglected Diseases—Criteria for Lead Progression, *PLoS Negl Trop Dis.* 3 (2009) e440.
- [47] T. Storz, R. Heid, J. Zeldis, S.M. Hoagland, V. Rapisardi, S. Hollywood, G. Morton, Convenient and Practical One-Pot Synthesis of 4-Chloropyrimidines via a Novel Chloroimidate Annulation, *Org. Process Res. Dev.* 15 (2011) 918–924.
- [48] W. Trager, J.B. Jensen, Human malaria parasites in continuous culture, *Science.* 193 (1976) 673–675.
- [49] W.A. Guiguemde, A.A. Shelat, D. Bouck, S. Duffy, G.J. Crowther, P.H. Davis, D.C. Smithson, M. Connelly, J. Clark, F. Zhu, M.B. Jimenez-Diaz, M.S. Martinez, E.B. Wilson, A.K. Tripathi, J. Gut, E.R. Sharlow, I. Bathurst, M.F. El, J.W. Fowble, I. Forquer, P.L. McGinley, S. Castro, I. Angulo-Barturen, S. Ferrer, P.J. Rosenthal, J.L. Derisi, D.J. Sullivan, J.S. Lazo, D.S. Roos, M.K. Riscoe, M.A. Phillips, P.K. Rathod, W.C. Van Voorhis, V.M. Avery, R.K. Guy, Chemical genetics of *Plasmodium falciparum*, *Nature.* 465 (2010) 311–315.
- [50] T. Mosmann, Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays, *J. Immunol. Methods.* 65 (1983) 55–63.

Highlights:

► Antiplasmodial pharmacomodulation of CCl₃-substituted-nitrogen containing heterocycles was made. ► Thienopyrimidine derivatives appeared more cytotoxic. ► Original 3-substituted-2-trichloromethylquinoxaline analogs were prepared. ► Two quinoxaline derivatives displayed *in vitro* IC₅₀ values of 0.4 and 0.5 μ M on the K1 multi-resistant *P. falciparum* strain. ► Cytotoxicity was assessed on the human HepG2 cell line showing low cytotoxicity (CC₅₀ ~ 40 μ M) and improved selectivity indices (77-100).