The Laschamp geomagnetic dipole low expressed as a cosmogenic ^{10}Be atmospheric overproduction at 41 ka

L L Ménabréaz , Nicolas Thouveny, D L Boulès, P Deschamps, B Hamelin,
F Demory

► To cite this version:

L L Ménabréaz , Nicolas Thouveny, D L Boulès, P Deschamps, B Hamelin, et al.. The Laschamp geomagnetic dipole low expressed as a cosmogenic ^{10}Be atmospheric overproduction at 41 ka. *Earth and Planetary Science Letters*, 2011, 312 (3-4), pp.305-317. 10.1016/j.epsl.2011.10.037 . hal-01420387

HAL Id: hal-01420387

<https://hal.science/hal-01420387>

Submitted on 6 Feb 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The Laschamp geomagnetic dipole low expressed as a cosmogenic ^{10}Be atmospheric overproduction at ~41 ka.

L. Ménabréaz, N. Thouveny, D.L. Bourlès, P. Deschamps, B. Hamelin and F. Demory.

Aix-Marseille Univ., CEREGE, BP80, 13545 Aix-en-Provence Cedex 04, France.

CNRS, CEREGE, BP80, 13545 Aix-en-Provence Cedex 04, France.

IRD, CEREGE, BP80, 13545 Aix-en-Provence Cedex 04, France.

Abstract

Authigenic $^{10}\text{Be}/^9\text{Be}$ ratio measurements were performed at high resolution along a Portuguese Margin deep-sea core (37°48 N; 10°09 W) spanning the 20-50 ka time interval, in order to reconstruct variations in atmospheric cosmogenic ^{10}Be production rates and derive the related geomagnetic dipole moment modulation. A complementary approach consisting in $^{230}\text{Th}_{\text{xs}}$ -normalized ^{10}Be deposition rate determination on selected samples confirmed the reliability of the authigenic $^{10}\text{Be}/^9\text{Be}$ record. This study constitutes the first successful comparison of the two widely-used normalization techniques of ^{10}Be concentrations. For both methods, the presented results herein evidence a factor of ~2 cosmogenic nuclide overproduction linked to a minimum dipole moment associated with the Laschamp Excursion. The latter is stratigraphically constrained beneath Heinrich Event 4. Its age is estimated at ~41 ka on the basis of direct correlation between the series of rapid paleoclimatic events recorded in the Portuguese Margin sediments and in the Greenland ice sheet, and is confirmed by calibrated radiocarbon dating carried out on the same sediments.

The remarkable agreement between the authigenic $^{10}\text{Be}/^9\text{Be}$ and the Greenland Ice cores ^{10}Be deposition rate records attest to their global significance. This new authigenic $^{10}\text{Be}/^9\text{Be}$ record has been combined with that previously obtained at the same site to produce a stacked record that is calibrated using absolute values of Virtual Dipole Moment determined on lava flows. This provides a reconstruction of dipole geomagnetic moment variations over the 20-50 ka interval, independent from paleomagnetically-constrained methods, that documents the Laschamp dipole low but fails to express any dipole low related to the Mono Lake Excursion.

This high resolution record responds to the necessity of supplementing the knowledge of the atmospheric $\Delta^{14}\text{C}$ variations in the 30-45 ka interval during which the ^{14}C calibration curve suffers from a lack of accurate data, and during which a discrepancy of about 5500 yr

between the ^{14}C and U-Th ages is due to the Laschamp geomagnetic dipole low. Such new high resolution datasets from records obtained from different latitudes will be required to make significant advances in understanding the causes of atmospheric $\Delta^{14}\text{C}$ variations.

Key words: Cosmogenic nuclides; atmospheric ^{10}Be production rate; geomagnetic dipole moment; Laschamp Excursion; marine sediments.

1. Introduction

The cosmogenic nuclide Beryllium-10 (^{10}Be , half-life: 1.387 ± 0.012 Ma (Chmeleff et al., 2010; Korschinek et al., 2010)) is produced in the Earth's upper atmosphere by nuclear interactions between energetic primary and secondary cosmic ray particles with target elements O and N. As shown in the early work of Elsasser et al. (1956), and later refined by Lal (1988), ^{10}Be atmospheric production is directly modulated by the geomagnetic field strength over millennial time scales, following a negative power law. This relationship between the dipole moment and the atmospheric production rate of cosmogenic nuclides was initially inferred from archeomagnetic absolute paleointensities and neutron monitor data comparison (Elsasser et al., 1956). More recently, a physical model of cosmic ray particle interactions with atmospheric targets yielded compatible simulations (Masarik and Beer, 1999; Wagner et al., 2000). Regarding the classically-used paleomagnetic reconstructions, ^{10}Be -derived paleointensity records can therefore constitute an alternative global and independent reading of the dipole moment variations, and more particularly those that accompany geomagnetic excursions and polarity events. During the last few years, efforts have been made to extract a geomagnetic signal from both single and stacked ^{10}Be records in natural archives such as ice sheets (e.g. Muscheler et al., 2005) and marine sediments (e.g. Frank et al., 1997; Carcaillet et al., 2004a; Christl et al., 2007). For these latter archives, two correction techniques were used to account for oceanic transport: the authigenic (adsorbed from the water column) $^{10}\text{Be}/^9\text{Be}$ ratio (Bourlès et al., 1989; Henken-Mellies et al., 1990; Robinson et al., 1995; Carcaillet et al., 2003; Carcaillet et al., 2004a; Carcaillet et al., 2004b; Leduc et al., 2006) and $^{230}\text{Th}_{\text{xs}}$ -normalisation (e.g. Frank et al., 1997; Christl et al., 2003; Christl et al., 2007; Christl et al., 2010). The sole attempt to compare both correction techniques was presented by Knudsen et al. (2008), but large uncertainties and environmental

complications prevented them from isolating the ^{10}Be production component from the $^{10}\text{Be}/^{230}\text{Th}_{\text{xs}}$ signal.

The last 50 ka is a key period for the study and calibration of geomagnetic excursions and the associated cosmogenic responses, because this period contains the best-documented and best-dated geomagnetic instability, the Laschamp Excursion (LE). This excursion was first identified in the Laschamp and Olby lava flows of the Chaîne des Puys, France (Bonhommet and Babkine, 1967), and was later described in lava flows of Iceland (e.g. Kristjansson and Gudmundsson, 1980; Levi et al., 1990) and New Zealand (Cassata et al., 2008) notably. Other records of the Laschamp Excursion and/or associated dipole low were obtained from lacustrine and marine sediments (e.g. Thouveny and Creer, 1992; Vlag et al., 1996; Lund et al., 2005; Channell, 2006). The most recent chronological constraints are provided by Ar-Ar, K-Ar, and ^{230}Th -U dating of lava flows from France (Laschamp, Olby and Louchadière), and from New Zealand (McLennans Hill) (Guillou et al., 2004; Plenier et al., 2007; Singer et al., 2009). The statistical analyses of these age populations have led to the following average ages: 40.4 ± 2.0 ka (Guillou et al., 2004) 40.7 ± 1.0 ka (Singer et al., 2009), and 37.0 ± 0.7 ka (Plenier et al., 2007), the third thus introducing a significantly younger age estimation.

Another younger, but more controversial, excursion has also been documented: the Mono Lake (ML) Excursion that was first detected in the Wilson Creek formation, California, by Denham and Cox (1971) as they were seeking for a record of the Laschamp Excursion. It was later described in other northwestern American lacustrine (e.g. Negrini et al., 1984; Liddicoat, 1992; Coe and Liddicoat, 1994) as well as in marine sediments (e.g. Nowaczyk and Knies, 2000; Channell, 2006). The radiometric age of the ML Excursion has been revised from ~ 24 ka (Denham and Cox, 1971) to ~ 32 ka (Negrini et al., 2000; Benson et al., 2003). Surprisingly, other ^{39}Ar - ^{40}Ar dating of ash layers surrounding the excursion at Wilson Creek (Kent et al., 2002) has yielded ages between 38 and 41 ka, which suggests that the excursion recorded at Mono Lake could be the Laschamp Excursion. The correlation of the Mono Lake RPI stack (Zimmerman et al., 2006) to the GLOPIS record of Laj et al. (2004) supported the assignment of this excursion to the Laschamp. It remains true that most RPI records exhibit two successive RPI lows at 32-34 ka and at 41 ka (e.g. Channell et al., 2006; Lund et al., 2005), which is also supported by recent reports of excursions directions and low absolute paleointensities in lava flows at ages near 32 ka, i.e. distinct from the Laschamp age (Cassata et al., 2008; Singer et al., 2009; Kissel et al., 2011).

The dipole field reduction linked to the Laschamp Excursion has been recorded as enhancements of ^{10}Be deposition rate in ice cores (e.g. Yiou et al., 1997; Muscheler et al., 2004; Raisbeck et al., 2007) and marine sediments (e.g. Frank et al., 1997; Christl et al., 2003; Carcaillet et al., 2004b; Leduc et al., 2006). Regarding the ML Excursion, a Chlorine-36 atmospheric production enhancement detected at around 32 ka in the GRIP ice core (Wagner et al., 2000) constitutes the only documented evidence for a significant increase of the cosmogenic production. Atmospheric cosmogenic nuclide production variations induced by these geomagnetic instabilities need to be well constrained, notably because they impact ^{14}C age calibration methods.

Here we present a highly resolved reconstruction of the ^{10}Be atmospheric production variations over the 20-50 ka interval, obtained from a well-dated 10 m sediment sequence deposited off the coast of Portugal. Authigenic $^{10}\text{Be}/^9\text{Be}$ ratios were measured, and compared to $^{230}\text{Th}_{\text{xs}}$ -derived ^{10}Be fluxes which were investigated on the same samples. The ^{10}Be production record was then transformed into a ^{10}Be -derived geomagnetic dipole moment record.

2. Material and methods

2.1 Coring site and sediment description

Climate and oceanographic settings as well as sedimentation processes within the area have been described notably in Baas et al. (1997) and Pailler and Bard (2002). The sediment deposition regime is related to glacial-interglacial sea-level variations and to the Tagus River inputs. The contribution of large and variable continental material gave rise to a sediment carbonate content of ~10% during glacials and ~50% during interglacials (Thomson et al., 1999; Hall and McCave, 2000), and with less than 1% organic matter (Pailler and Bard, 2002). Hemipelagic sediments along the Portuguese Margin also contain scarce and thin Ice Rafted Debris (IRD) material diluted in the clay matrix (feldspars, quartz, magnetite and hematite-coated grains). These layers detected by their IRD abundance and Magnetic Susceptibility (MS) signatures have been shown to be contemporaneous with Heinrich Events (Bard et al., 2000; Thouveny et al., 2000; Moreno et al., 2002).

Using different methodologies, several studies have demonstrated that the upper part of the giant cores collected with the Calypso corer of the R/V *Marion Dufresne* are affected

by significant oversampling (e.g. Thouveny et al., 2000; Skinner and McCave, 2003 and Széréméta et al., 2004). This artefact has led to overestimations of apparent sedimentation rates (e.g. Thompson et al., 1999). Despite their perfect stratigraphic preservation, the upper 12 m of core MD95-2042 are affected by this artefact, which hampered the evaluation of accurate sedimentation rates and the interpretation of paleomagnetic and cosmogenic nuclides results.

During the MD-140 PRIVILEGE campaign of the R/V *Marion Dufresne* (2004) on the Portuguese Margin, a giant box corer CASQ (0.25m²x12m) allowed to collect a high resolution, undisturbed sediment sequence (Core MD04-2811; Lat.: 37°48 N; Long.: 10°09 W; 3162m water-depth) at the site of the previously studied SU81-18 and MD95-2042 cores (Fig. 1; see references in caption). These cores were extensively studied for radiocarbon calibration (section 3.2) and therefore benefits from an excellent chronological control. Core MD04-2811 was thus selected in order to document a lack of the MD95-2042 record in the critical time interval (20-35 ka) and to assess accurate sediment fluxes using the ²³⁰Th_{xs} normalization.

MD04-2811 sediments are composed of oxidized carbonate-rich silty clays in the upper Holocene section (1.5m), and homogeneous hemipelagic silty clays deposited during the last glacial period.

2.2 Paleomagnetic record and sampling strategy

Paleomagnetic investigations were performed on 1.5 m long U-channels collected along the MD04-2811 core. Low field Magnetic Susceptibility (MS) was measured using a Bartington MS2C probe. As described in former studies (e.g. Thouveny et al., 2000), the major structures of the MS profile consist in a succession of peaks corresponding to the ferrimagnetic responses of IRD layers deposited during Heinrich Events 1 to 5 superimposed on the paramagnetic contribution of the clayey fraction and a diamagnetic contribution of carbonates (Fig. 2).

Natural Remanent Magnetization (NRM) and artificially induced magnetizations were measured using a 2G cryogenic magnetometer 760-SRM model and were demagnetized by Alternating Fields (AF). The Relative Paleointensity (RPI) curve was established by normalizing the NRM intensity to the Anhyseretic Remanent Magnetization (ARM) intensity. The NRM/ARM ratio obtained for the 30mT AF demagnetization step (NRM_{30mT}

/ARM_{30mT}) represents the best estimate of the RPI. Three main RPI lows are recorded in the 435-520 cm, 575-630 cm and 740-810 cm intervals, the latter displaying the lowermost values (Fig. 2). This structure is classically attributed to the dipole low related to the Laschamp Excursion (e.g. Laj et al., 2004; Thouveny et al., 2004). The mean inclination of the Characteristic Remanent Magnetizations equals $\sim 50^\circ$, i.e. it is weaker than the inclination (56.8°) of the field created by a geomagnetic axial dipole field (GAD) at the site latitude. Unlike other records of the area (Thouveny et al., 2004), there is no anomalous direction related to the RPI low.

Based on the interpretation of the RPI profile, 52 sediment samples were taken at a 30 cm resolution, increased to 10 cm within paleointensity lows intervals, for ^{10}Be and ^9Be measurements.

2.3 Analytical techniques

Leaching procedure: dissolution of the authigenic phase.

In this study, authigenic $^{10}\text{Be}/^9\text{Be}$ ratios, interpreted as a proxy of ^{10}Be fluxes, are presented. Following the procedure thoroughly described in Bourlès et al. (1989), the authigenic phase is extracted by leaching $\sim 1\text{g}$ of dried and homogenized sediment using a 0.04M hydroxylamine in 25% acetic acid solution. A 2ml aliquot of the resulting leaching solution is separated for natural ^9Be measurements. The remaining solution, spiked with 300 μl of a 10^{-3} g/g ^9Be -carrier, is used for chemical extraction and Accelerator Mass Spectrometry (AMS) measurements of the resulting $^{10}\text{Be}/^9\text{Be}$ ratio. This procedure permits to overcome changes in extraction efficiency.

Total sample digestion and bulk sediment ^{10}Be content.

$^{230}\text{Th}_{\text{xs}}$ -normalized ^{10}Be flux quantification requires a total dissolution of sediment samples, assuming that lattice-bound ^{10}Be (from in-situ production) concentrations are negligible with respect to authigenic content. Sample dissolution was achieved at 70°C in a mixture of HF, HNO_3 and HCl (Bourlès et al., 1989), and spiked with 300 μl of a 10^{-3} g/g ^9Be -carrier to perform chemical extraction and AMS measurements.

Beryllium isotopes chemical extraction and measurements.

Once the sample is in solution, whatever the method used, Beryllium is then extracted in the form of Be acetylacetonate using organic solvent. After decomposition of these

complexes, Be oxy-hydroxides are precipitated at pH 8 and oxidized to perform AMS measurements at ASTER, the French AMS national facility (installed at the CEREGE) which operates at 5MV. ^{10}Be concentrations are deduced from spiked $^{10}\text{Be}/^9\text{Be}$ measured ratios calibrated against the NIST 4325 Standard Reference Material with an assigned $^{10}\text{Be}/^9\text{Be}$ value of 2.79×10^{-11} (Nishiizumi et al., 2007), and are decay-corrected using the ^{10}Be half-life of 1.387 ± 0.012 Ma (Chmeleff et al., 2010; Korschinek et al., 2010):

$$[^{10}\text{Be}]_{\text{g/g}}^{\text{sample}} = \left(\frac{^{10}\text{Be}}{^9\text{Be}} \right)_M \cdot \frac{m_{\text{spike}} \cdot [^9\text{Be}]_{\text{spike}}}{m_{\text{sample}}} \cdot e^{\lambda_{10} \cdot t}$$

where $(^{10}\text{Be}/^9\text{Be})_M$ is the measured Be ratio, λ_{10} is the decay constant of ^{10}Be and m_{sample} is the weight of the leached sediment powder sample.

Measured ratios as well as the resulting ^{10}Be concentrations are listed in Table 1. Chemistry blanks cluster around 9×10^{-15} , up to 1000 times lower than the $^{10}\text{Be}/^9\text{Be}$ ratio of the samples.

The external reproducibility obtained on the NIST SRM 4325 standard (estimated from 1σ standard deviation of long-term repeated measurements and integrating all effects contributing to the variability of ASTER) is limited to 0.6% (Arnold et al., 2010). Uncertainties in the measured $^{10}\text{Be}/^9\text{Be}$ ratios and ^{10}Be concentrations were calculated from counting statistics and instrumental error propagation, according to the standard equation (e.g. Taylor, 1997).

Natural ^9Be concentrations were measured on the assigned 2ml aliquot using a graphite-furnace Atomic Absorption Spectrophotometer with a Zeeman effect background correction (Thermo Scientific ICE 3400) and the standard additions method (Table 1). Uncertainties in ^9Be concentrations depend on: (1) the reproducibility of the measured absorbances at each added ^9Be concentrations (standard deviation less than 1%), and (2) the least-square fitting between measured absorbances and added ^9Be concentrations ($R^2 > 0.999$). Uncertainties (1σ) in the ^9Be concentrations are generally significantly lower than 10% (average value: 4.3%).

The final $^{10}\text{Be}/^9\text{Be}$ uncertainty results from standard propagation of the above-cited ^{10}Be and ^9Be concentrations uncertainties.

Replicate authigenic ^{10}Be and ^9Be analyses were performed in order to check the reproducibility of the Be isotopes chemical extraction (Table 1).

Uranium and Thorium isotopic analyses: chemical procedure and TIMS measurements.

U-series measurements were performed by Thermo-ionisation Mass Spectrometry (TIMS) at the CEREGE) by using a ^{236}U - ^{233}U - ^{229}Th mixed spike calibrated according to the procedure described in Deschamps et al. (2003).

The chemical procedure used to separate the uranium and thorium fractions for determination of $^{230}\text{Th}_{\text{xs}}$ is similar to that described in Deschamps et al. (2004). Before total digestion, samples were spiked. Total digestion was achieved by sequential treatments with aqua regia and concentrated nitric and hydrofluoric acids. Following co-precipitation on iron oxyhydroxides, U and Th fractions were chemically separated and purified using standard ion exchange resins (AG-1X8 and U-Teva) and separation protocols modified from Deschamps et al. (2004). Typical chemical blanks attained over the course of this study were about 0.25 ppb for uranium and 0.28 ppb for thorium.

U and Th analyses were performed using a VG-Sector 54-30 mass spectrometer equipped with a 30-cm electrostatic analyzer and an ion-counting Daly detector. The instrumental abundance sensitivity is greater than 0.15 ppm at 1 amu (proportion of the ^{238}U ion beam measured at mass 237). The detailed procedure is described in Bard et al. (1990) with further modifications described in Deschamps et al. (2011). U and Th fraction was loaded on the side filament of a triple zone-refined Re filament assembly. Reported errors (2σ) for the $^{234}\text{U}/^{238}\text{U}$ and $^{230}\text{Th}/^{232}\text{Th}$ ratios are about 0.14 ‰ and 0.9 ‰, respectively. The internal analytical reproducibility achieved in the course of this study on replicate measurements of the NBS-960 international standard yielded a mean $\delta^{234}\text{U}$ value of $36.5 \pm 0.8\text{‰}$ (2σ , $n=23$) which is in excellent agreement with values reported in the literature (Deschamps et al., 2003; Andersen et al., 2004).

To calculate excess ^{230}Th activity ($^{230}\text{Th}_{\text{xs}}$), measured ^{230}Th concentration is corrected for detrital and authigenic ^{230}Th components. The fraction of ^{230}Th supported by the decay of detrital ^{238}U is estimated from the ^{232}Th sample concentration and an assumed average lithogenic $^{238}\text{U}/^{232}\text{Th}$ activity ratio of $R = 0.6 \pm 0.1$ for the Atlantic Basin (Henderson and Anderson, 2003). Ingrowth of ^{230}Th from decay of authigenic uranium is estimated by

246 assuming that authigenic uranium has an initial $^{234}\text{U}/^{238}\text{U}$ activity ratio equivalent to that of
 247 sea-water (i.e. 1.147, Andersen et al., 2010). $^{230}\text{Th}_{\text{xs}}$ activities were calculated as:

$$\begin{aligned} {}^{230}\text{Th}_{\text{xs}} = & {}^{230}\text{Th}_{\text{Total}} \\ & + R_{\text{Crust}} \times {}^{232}\text{Th}_{\text{Total}} \left(\left[\left(\frac{{}^{234}\text{U}}{{}^{238}\text{U}} \right)_0 - 1 \right] \times \frac{\lambda_{230}}{\lambda_{230} - \lambda_{234}} \left[e^{-\lambda_{234} \times t} - e^{-\lambda_{230} \times t} \right] - e^{-\lambda_{230} \times t} \right) \\ & - {}^{238}\text{U}_{\text{Total}} \left(1 - e^{-\lambda_{230} \times t} + \left[\left(\frac{{}^{234}\text{U}}{{}^{238}\text{U}} \right)_0 - 1 \right] \times \frac{\lambda_{230}}{\lambda_{230} - \lambda_{234}} \left[e^{-\lambda_{234} \times t} - e^{-\lambda_{230} \times t} \right] \right) \end{aligned}$$

252 where $^{230}\text{Th}_{\text{Total}}$, $^{232}\text{Th}_{\text{Total}}$ and $^{238}\text{U}_{\text{Total}}$ are the measured ^{230}Th , ^{232}Th and ^{238}U activities,
 253 respectively, and $^{238}\text{U}_{\text{Total}}$ is the sedimentary ^{238}U activity. $(^{234}\text{U}/^{238}\text{U})_0$ stands for the initial
 254 (decay-corrected) authigenic activity ratio and the decay constants of radioisotopes i are
 255 expressed by λ_i .

256 The preserved, decay-corrected, $^{230}\text{Th}_{\text{xs}}$ -normalized ^{10}Be deposition rates are then
 257 calculated following equations (11) and (12) as given in François et al. (2004), using a water
 258 depth of 3162 m.

259 Analytical errors and uncertainty related to assumption concerning the lithogenic
 260 $^{238}\text{U}/^{232}\text{Th}$ activity ratio, R, were properly propagated through $^{230}\text{Th}_{\text{xs}}$ and flux calculations.
 261 Note that owing to small analytical errors achieved by the TIMS method, most of the final
 262 errors on $^{230}\text{Th}_{\text{xs}}$ and flux calculations are related to the arbitrary fixed uncertainty on R.

263

264 **2.4 Using authigenic $^{10}\text{Be}/^{\rho}\text{Be}$ and $^{230}\text{Th}_{\text{xs}}$ -normalized ^{10}Be fluxes as tracers of geomagnetic** 265 **moment variations**

266 The particle-reactive ^{10}Be is mainly produced in the atmosphere through nuclear
 267 reactions (spallation reactions) on oxygen (O) and nitrogen (N), and is transferred to the
 268 Earth's surface in soluble form by precipitation (Raisbeck et al., 1981) within ~3 yrs (Baroni
 269 et al., 2011). Atmospheric ^{10}Be is ultimately removed from water via settling particles and is
 270 either deposited in marine and lacustrine sediments, or is efficiently retained by continental
 271 sediment components, and finally enters the oceans through various paths and deposits in
 272 deep sea sediments. Due to its short atmospheric residence time and chemical reactivity, ^{10}Be
 273 records of production enhancements should be synchronous in the geophysical reservoirs. In
 274 marine sediments, the ^{10}Be concentration results from a complex interplay between several

processes: e.g. cosmogenic production, redistribution by atmospheric, riverine and oceanic transport, adsorption and deposition processes. Considering the adsorption processes, for example, the absolute ^{10}Be concentration is controlled by the scavenging efficiency and the specific surface of the settling sedimentary particles and thus depends on environmental conditions affecting their chemical and grain size composition. Consequently, ^{10}Be concentrations are meaningless and in order to account for these dependencies, a correction procedure is required. Soluble forms of ^{10}Be and ^9Be have different sources: while ^{10}Be is cosmogenic, the stable isotope ^9Be originates from partial dissolution of detrital, aeolian and riverine inputs (Brown et al., 1992). Once homogenized in seawater, both isotopes are scavenged with the same efficiency. Authigenic $^{10}\text{Be}/^9\text{Be}$ ratio of marine surface sediments reflects the Be isotope composition of the overlying deep waters, and its spatial variability is mainly controlled by the proximity to continental ^9Be inputs (Bourlès et al., 1989). Although the authigenic ^9Be normalization method has provided promising results in specific environments (Bourlès et al., 1989; Carcaillet et al., 2003; Carcaillet et al., 2004a; Carcaillet et al., 2004b; Lebatard et al., 2010), quantitative reconstructions of the ^{10}Be fluxes to the sediment requires taking into account syndepositional lateral transport of adsorbed Be. Because residence time of Be in the water column is about 500-1000 yrs, boundary scavenging and deepwater circulation must also contribute to the removal of dissolved Be and thus influence the $^{10}\text{Be}/^9\text{Be}$ ratio. Moreover, lateral transport of adsorbed Be can be significant in sediments heavily affected by focusing, such as drift deposits. However, at ocean margins where particle flux is high, the short Be residence time allows to record the ^{10}Be atmospheric flux variations with almost no signal attenuation.

Assuming a scavenged flux equivalent to its known production rate in the overlying water column (Bacon, 1984), ^{230}Th can be used to quantify rates of particle rain to the seafloor and to correct for syndepositional sediment redistribution by bottom currents. The principles and limitations of this method are fully reviewed in Henderson and Anderson (2003) and François et al. (2004). Like authigenic $^{10}\text{Be}/^9\text{Be}$, the $^{10}\text{Be}/^{230}\text{Th}_{\text{xs}}$ ratio may also be potentially influenced by advection of dissolved Be. Moreover, biases resulting from nuclide characteristics (different chemical affinities) are introduced: one illustration of this problem is reported by Chase et al. (2002), who showed a composition-dependant differential scavenging of Be and Th, suggesting that opal/carbonate ratio of settling particles may partly explain the sedimentary $^{10}\text{Be}/^{230}\text{Th}_{\text{xs}}$ variability.

Here, authigenic $^{10}\text{Be}/^9\text{Be}$ ratios were measured for 51 sampled layers, and an additional $^{230}\text{Th}_{\text{xs}}$ -normalization was performed for 9 of them, which were selected using the Be isotopes ratio profile. This allows direct comparison and cross-calibration on the same homogenized material.

3. Results

3.1 The authigenic $^{10}\text{Be}/^9\text{Be}$ ratios and RPI on the same sequence

Authigenic ^9Be concentrations vary from 4.35×10^{-7} g/g to 7.22×10^{-7} g/g (Table 1), refining and confirming the previous range obtained on adjacent core MD95 2042 by Carcaillet et al. (2004). Decay-corrected authigenic ^{10}Be concentrations vary around a mean value of 7.32×10^{-15} g/g with a prominent increase to 13.6×10^{-15} g/g at 712 cm. The range of variation (5.33 to 13.6×10^{-15} g/g) and the average value are identical to those of the adjacent core MD95 2042 (Carcaillet et al., 2004b). The resulting authigenic $^{10}\text{Be}/^9\text{Be}$ evolution is presented in Figure 2, and compared to the RPI and MS profiles. The main feature of the $^{10}\text{Be}/^9\text{Be}$ record is a unique significant enhancement at 712 cm, with ^{10}Be concentrations leading to a near doubling of the long-term average $^{10}\text{Be}/^9\text{Be}$ value of 1.44×10^{-8} to 2.69×10^{-8} .

Along this sequence, both records of the ^{10}Be atmospheric production and its geomagnetic modulation display inverse long-term trends. While the RPI record exhibits three minima, only one significant enhancement appears in the $^{10}\text{Be}/^9\text{Be}$ record, with values above the “mean+1 σ ” occurring between 662 and 722 cm depth. This interval overlaps the depth interval recording the lowest RPI values located between ~690 and ~820 cm. The depth shift thus evidenced between these two expressions of the dipole low is further discussed in section 4.1

3.2 Chronostratigraphy

The first step in establishing the chronostratigraphy of core MD04-2811 is to construct a correlation matrix based on the MS records (Fig. 3) with the dated neighboring core MD95 2042 (Lat.: 37°45'N; Long.: 10°10'W; 3146 m water depth) in order to translate the available chronological data onto the MD04-2811 depth scale (Fig. 4).

Two major studies have established radiocarbon dates series on MD95-2042 (Bard et al., 2004; Shackleton et al., 2004, Electronic Appendix, Table A.1) that were compared with ages obtained from correlations with annually laminated Greenland ice cores (GRIP and GISP2). Among the paleoclimatic proxies measured along the MD95-2042 sequence, sea-

surface temperature proxies (alkenones and planktonic $\delta^{18}\text{O}$) show marked variations which are in robust mutual stratigraphic agreement, and which correlate to Dansgaard-Oeschger cycles and Heinrich Events. For this study, the age control on the core MD95-2042 was updated by tuning the alkenone-derived sea-surface temperature record with the NGRIP oxygen isotopes record using a minimum number of tie points ($R = 0.876$, Fig. 3 and 4).

Uncertainties affecting the age model are derived from the two-sigma GICC05 time-scale uncertainties associated with multi parameter counting of annual layers (Andersen et al., 2006; Rasmussen et al., 2006): the glacial part of this time scale has an estimated associated error of 2% back to 40 ka and of 5–10% back to 57 ka. Although multiple correlation procedures - limited by precision and resolution of the individual time series - and the accuracy of the cross-correlation procedure represent a source of uncertainty which is difficult to quantify using simple statistical parameters, the computed age model is validated by the fact that the ages assigned to the Heinrich Events (Fig. 4) are indistinguishable from the calibrated ^{14}C ages obtained on North Atlantic basin sediment sequences (Thouveny et al., 2000). Moreover, the ^{14}C ages obtained on the studied sequence calibrated using the INTCAL09 dataset (Reimer et al., 2009) relying mainly on $^{234}\text{U}/^{230}\text{Th}$ absolute dating of Hulu Cave stalagmites (China, Wang et al., 2001) over the 35-45 ka period (Electronic Appendix, Table A.1) are fully compatible with this ice core-derived chronology.

3.3 ^{10}Be deposition rates (excess ^{230}Th -normalized ^{10}Be fluxes)

Bulk ^{10}Be concentrations obtained after total sample dissolution are comprised between 1.1×10^{-14} g/g and 1.8×10^{-14} g/g (see Table 2). Total mass flux ranges between 4.79 and 6.09 g/cm²/ka, and is fully compatible with results obtained further northward on the Portuguese Margin by Thomson et al. (1999), who determine a total sediment accumulation flux of ~ 5 g/cm²/ka during glacial times over the last 140 ka. Preserved vertical ^{10}Be deposition rates vary from 3.18×10^9 to 6.59×10^9 atoms/cm²/ka (Fig. 5a). The ^{10}Be deposition rate downcore evolution is strictly similar to that of the authigenic $^{10}\text{Be}/^9\text{Be}$ ratios measured on the same samples: the characteristic increase with a maximum at 712 cm depth is indeed reflected in the ^{10}Be deposition rates record. The comparison between authigenic $^{10}\text{Be}/^9\text{Be}$ ratios and $^{230}\text{Th}_{\text{xs}}$ -normalized ^{10}Be fluxes is illustrated in Figure 5b. The linear relationship between these two variables is highly significant ($R = 0.912$), albeit mainly carried by the maximum variability recorded during the main enhancement.

4. Discussion

4.1 Identification of the Laschamp dipole low and chronological implications

During low dipole field intervals, the magnetic torques are insufficient to align magnetic grains and so ensure acquisition of a well-defined magnetization. Nevertheless, after recovery of the field strength, re-alignments of magnetic grains and acquisition of viscous magnetization (VRM) result in a biased record of directions and intensities (e.g. Coe and Liddicoat, 1994). In core MD04-2811, as in other neighboring cores (see Figures 4 and 9 in Thouveny et al., 2004), the structure of paleomagnetic records (directions and RPI) does not allow an accurate and precise stratigraphic positioning of the dipole field minimum. Comparison of the positions of the respective boundaries of the main RPI low (at ~700 and ~820 cm) with the main $^{10}\text{Be}/^9\text{Be}$ enhancement (at ~660 cm and ~725 cm) calls attention to a ~50 cm lag between the two signals (Fig. 2). This lag integrates a maximum 10 cm shift due to the residence time with respect to scavenging of Be, given a ~1 cm/50 yrs average sedimentation rate. The remaining 40 cm lag is greater than the 15-30 cm lags revealed from neighboring cores by Carcaillet et al. (2004); however given the sources of distortions cited above, it is not possible to assign this lag to the magnetization lock-in depth only.

In core MD95-2042 core, the declination anomaly and the RPI low related to the Laschamp excursion are recorded 50 cm beneath the MS peak associated with the Heinrich Event 4 (Thouveny et al., 2004). In North Atlantic Ocean cores (Kissel et al., 1999), excursions peaks related to the Laschamp are recorded at the same stratigraphic position as rock-magnetic signatures of the Dansgaard-Oeschger (D-O) Interstadial 10, i.e. prior to H4. Nevertheless, the best stratigraphic indicator of the occurrence of the dipole low related to the Laschamp is the cosmogenic nuclide signature which is not affected by delays and distortions imparted by post-depositional processes.

Along core MD04-2811 the highest $^{10}\text{Be}/^9\text{Be}$ ratio peak - recorded ~40 cm beneath the MS peak corresponding to H4 - can thus undoubtedly be interpreted as the ^{10}Be atmospheric overproduction due to the geomagnetic dipole reduction associated with the Laschamp Excursion (LE). This 40 cm depth interval suggests that the LE occurred ~2 ka before Heinrich Event 4, which is dated at ~39 ka in North Atlantic sedimentary sequences (e.g. Thouveny et al., 2000). Indeed, the chronological reconstruction positions the $^{10}\text{Be}/^9\text{Be}$ Laschamp signature at 41.2 ± 1.6 ka (minimal uncertainty derived from GICC05 chronology). This age is coherent with the most recent independent radiometric age determinations

obtained on volcanic material (40.4 ± 2.0 ka, Guillou et al., 2004; 40.7 ± 0.9 ka, Singer et al., 2009). It also coincides tightly with the age of the maximum ^{10}Be flux in Summit ice cores (Muscheler et al., 2005; Svensson et al., 2008). This suggests that the ^{10}Be residence time in the water column in such settings is short enough to preserve the simultaneity of the record in both geological reservoirs, considering age uncertainties and resolution.

The ^{10}Be production and recording processes in sedimentary sequences that are totally independent from: 1) the local geomagnetic vector, and 2) magnetization processes in weak and highly variable local fields, enable to evaluate the duration of the dipole field anomaly. Both the high resolution and the accurate dating of this record highlight the persistence of a weak dipole field defined by authigenic $^{10}\text{Be}/^9\text{Be}$ values above “mean+1 σ ” during ~ 2.5 ka. The minimum field intensity phase defined by authigenic $^{10}\text{Be}/^9\text{Be}$ values above “mean+2 σ ” lasted about 1 ka. This duration estimate is similar to that provided by the ice core record of Muscheler et al. (2005).

4.2 Significance of the $^{10}\text{Be}/^9\text{Be}$ record

$^{10}\text{Be}/^9\text{Be}$ ratios and $^{230}\text{Th}_{\text{xs}}$ -normalized ^{10}Be fluxes: a cross evaluation

In this study, ^{10}Be deposition rates reinforce the authigenic $^{10}\text{Be}/^9\text{Be}$ signal validity. Comparison between the amplitude of $^{10}\text{Be}/^{230}\text{Th}_{\text{xs}}$ and authigenic $^{10}\text{Be}/^9\text{Be}$ variations shows that both records agree within uncertainties (Fig. 6a) and document an increase by a factor of ~ 2 coincidental with the Laschamp geomagnetic dipole low. This is the first time that such a match has been reported, demonstrating that both normalization methods provide equivalent results in this case, and that normalizing authigenic ^{10}Be concentrations by authigenic ^9Be concentrations allows a reliable correction for total particle flux variations. Moreover, quantification of the vertical ^{10}Be fluxes allows to underline the effects of boundary scavenging: the ^{10}Be deposition rates determined in this study (mean value: $4.09 \pm 0.31 \times 10^9$ at./cm²/ka) are (1) both slightly higher than an estimate of the present global production rate of $1.21 \pm 0.26 \times 10^9$ at./cm²/ka (Monaghan et al., 1986) and higher than Greenland ^{10}Be fluxes (0.25 to 0.6×10^9 at./cm²/ka (Muscheler et al., 2004)); and (2) within the range reported for Atlantic drift deposits of the last 75 ka (~ 1 to 7×10^9 at./cm²/ka, Christl et al., 2007; Knudsen et al., 2008; Christl et al., 2010). Although the resolution of the $^{10}\text{Be}/^{230}\text{Th}_{\text{xs}}$ record must be improved, the observed close agreement suggests that the effects of rapid millennial-scale changes in deep water circulation (advection of dissolved Be and Th) could be either similar in both approaches, or insignificant.

Comparison with ^{10}Be production rate ice core records and with relative geomagnetic field intensity reconstructed from paleomagnetic records

Over the 20-50 ka interval, there is a remarkable match between the MD04-2811 authigenic $^{10}\text{Be}/^9\text{Be}$ record and the ^{10}Be flux variations recorded along the GRIP and GISP2 ice cores (Fig. 6b; Finkel and Nishiizumi, 1997; Yiou et al., 1997; Muscheler et al., 2005). It must be emphasized that even small variations of the two records match which strongly supports the high quality of the authigenic $^{10}\text{Be}/^9\text{Be}$ record presented. Observed for such different geophysical reservoirs as marine sediments and polar ice, this agreement demonstrates that this ratio most likely mainly reflects the changes in global atmospheric ^{10}Be production rate. This confirms that the normalization of authigenic ^{10}Be concentrations to authigenic ^9Be concentrations minimizes the impact of secondary mechanisms, among which ocean system effects (see section 2.4). In addition, the fact that these variations are recorded with the same amplitude in both archives, indicates that the ^{10}Be residence time in the atmosphere-ocean reservoirs should be short enough to minimize the attenuation of the production signal. The Laschamp signature corresponds to the near-doubling of the global atmospheric ^{10}Be production expected from the physically constrained algorithm (Elsasser et al., 1956; Lal, 1988) which expresses the control of global atmospheric ^{10}Be production rate exerted by the magnetospheric shielding as modulated by geomagnetic moment variations,. This also suggests that a significant component of the Greenland ^{10}Be flux signal results from an atmospheric transport from lower latitudes. Another important result is that these marine and ice core records of the atmospheric ^{10}Be production rates both fail to display any other significant ^{10}Be deposition enhancement. Wagner et al. (2000a) identified in the GRIP ice core an increased ^{36}Cl flux between D-O events 6 and 7, which they attributed to the RPI minimum observed at ~34 ka in the NAPIS-75 stack (based on the GISP-2 chronology; Laj et al., 2000), and referred to as the Mono Lake Excursion. After smoothing of the dataset with a $1/2000 \text{ yrs}^{-1}$ low-pass filter, the ^{36}Cl peak amplitude appears equivalent to that of the Laschamp. Surprisingly, in Wagner et al. (2000b), smoothing of the dataset with a $1/3000 \text{ yrs}^{-1}$ low-pass filter reduces the ^{36}Cl Mono Lake peak previously noted in Wagner et al. (2000a) to the level of an insignificant variation. Together with this observation, the ^{10}Be fluxes in Summit ice cores and the presented high-resolution $^{10}\text{Be}/^9\text{Be}$ record question the validity of a ^{36}Cl signature linked to the Mono Lake dipole field low.

Lal's algorithm (1988) has been used to convert the authigenic $^{10}\text{Be}/^9\text{Be}$ record into a relative Virtual Dipole Moment (VDM) record that is compared to the GLOPIS-75 stacked

paleomagnetic record (Fig. 6c). This reconstruction and the GLOPIS-75 curve agree relatively well over the 30-48 ka time interval, although the cosmogenic record points out possible higher intensities for the 20-30 ka period and beyond 48 ka than indicated in the paleomagnetic records of the GLOPIS-75 stack.

4.3 Reconstruction of the geomagnetic dipole moment variations from the $^{10}\text{Be}/^9\text{Be}$ ratios

A stack of the Portuguese Margin $^{10}\text{Be}/^9\text{Be}$ records

In order to achieve a statistically significant record of the authigenic $^{10}\text{Be}/^9\text{Be}$ ratio, the MD04-2811 record was combined with the record previously obtained by Carcaillet et al. (2004) on core MD95-2042, recovered at the same site. These data were relocated on the depth scale of core MD04-2811 using the MS correlation matrix (reciprocal to the one used to transfer chronological data (Fig. 7a)). Be ratios measured on both cores vary in a similar range and present the same structure characterized both by a sharp maximum linked to the LE dipole low and by the absence of any other significant peak. Differences between MD95-2042 and MD04-2811 data sets result from local sedimentary processes, differential post-depositional processes (i.e. compaction), sampling and analytical uncertainties. After stacking the two $^{10}\text{Be}/^9\text{Be}$ ratios records, a smoothed composite record was produced by computing weighted moving averages over a 1000 years sliding window offset by 500 years (except for the 20-25 ka interval, for which MD04-2811 data are directly reported). Associated uncertainties combine both the sample analytical uncertainty and standard deviation around each weighted average, and are calculated using the unbiased weighted estimator of the sample variance (see Electronic Appendix Table A.2). Fig. 7b shows this composite record and the associated standard error (1σ). The 1000 yrs-smoothed authigenic $^{10}\text{Be}/^9\text{Be}$ composite record displays a ~ 1.45 time increase during the LE time interval compared to the long-term average (1.32×10^{-8}). Although the studied time interval covers the age range of the Mono Lake Excursion, the slight increase of the $^{10}\text{Be}/^9\text{Be}$ composite curve at ~ 33 -34 ka is not significant regarding the uncertainties associated with the adjacent data points. This reveals the absence of any ^{10}Be overproduction linked to a dipole reduction associated with the Mono lake Excursion. This observation -if confirmed at other sites- would have strong implications on the interpretation of the magnitude and duration of the “geomagnetic dipole low” associated with the Mono lake Excursion.

The previously described ^{10}Be -stack was converted into Virtual Dipole Moment (VDM) variations using absolute paleointensities reconstructed from lava flows. In order to reproduce the full range of the ^{10}Be production variation, the $^{10}\text{Be}/^9\text{Be}$ ratios measured along core MD04-2811 were associated with VDM values selected from the GEOMAGIA-50 (Korhonen et al., 2008) absolute paleointensity database (see Electronic Appendix Fig. A.3 and Table A.3). ^{10}Be production rates are inversely proportional to VDM values, thus (1) the $^{10}\text{Be}/^9\text{Be}$ maximum value was assigned to the minimum VDM value linked to the LE, (2) the intermediate $^{10}\text{Be}/^9\text{Be}$ ratios were assigned to the intermediate VDM values, and (3) the minimum $^{10}\text{Be}/^9\text{Be}$ values were assigned to the maximum VDM values.

The relationship thus obtained between authigenic $^{10}\text{Be}/^9\text{Be}$ ratios and VDMs is, at best, fitted by a polynomial of order 2 (Fig. A.3). When applied to the $^{10}\text{Be}/^9\text{Be}$ stack this polynomial produces a “ ^{10}Be -derived VDM” record for the 20-50 ka interval, which can be compared to GLOPIS-75 and SINT-800 paleointensity stacks (Fig. 8). This ^{10}Be based VDM record provides a reconstruction in better agreement with GLOPIS-75 than does Lal’s algorithm conversion (Fig. 6c). It documents a stronger geomagnetic moment before (maximum $\sim 10.5 \times 10^{22} \text{ A.m}^2$) than after (maximum $\sim 7.0 \times 10^{22} \text{ A.m}^2$) the LE, and allows to estimate a decreasing rate of $\sim 1 \times 10^{22} \text{ A.m}^2/\text{ka}$ for the dipole collapse which initiates the Laschamp (minimum $\sim 2.6 \times 10^{22} \text{ A.m}^2$ reached at $\sim 41 \text{ ka}$). Although this record exhibits the same main variations as GLOPIS-75, it fails to reproduce a sharp VDM reduction at the age of the Mono Lake Excursion: at 34-32 ka relatively low VDM values ($\sim 4.7 \times 10^{22} \text{ A.m}^2$) are in opposition with the sharp increase recorded at 32-33 ka in the GLOPIS-75.

5. Conclusion

Authigenic $^{10}\text{Be}/^9\text{Be}$ ratios measured along a rapidly accumulating sedimentary sequence from the North-East Atlantic (core MD04-2811) provide a new record of dipole geomagnetic moment variations over the 20-50 ka time interval, independent from paleomagnetic methodological constraints.

This first successful geomagnetic reconstruction using both authigenic $^{10}\text{Be}/^9\text{Be}$ and $^{10}\text{Be}/^{230}\text{Th}_{\text{xs}}$ normalization techniques of ^{10}Be concentrations in marine sediments evidence an almost doubling of the atmospheric ^{10}Be production which documents the occurrence of the

Laschamp geomagnetic dipole low prior to the Heinrich Event 4, at an age of ~ 41 ka. These ratios are fully compatible with the Summit ice core ^{10}Be record, which further supports their significance in terms of ^{10}Be atmospheric production. Identical ^{10}Be flux responses recorded in a polar ice core and in a mid-latitude marine core strongly suggest that the cosmogenic nuclide overproduction signal in Greenland ice results from the global geomagnetic moment modulation.

The authigenic $^{10}\text{Be}/^9\text{Be}$ ratio records of neighboring cores MD04-2811 and MD95-2042 cores were compiled and averaged over a 1 ka sliding window. The obtained stack was calibrated using absolute virtual dipole moment (VDM) values, determined from lava flow paleointensities, to provide a ^{10}Be derived VDM record. This allows to characterize the Laschamp dipole low by a decrease from $\sim 10.5 \times 10^{22} \text{ A.m}^2$ at ~49 ka to $\sim 2.6 \times 10^{22} \text{ A.m}^2$ at ~41 ka. The latter estimate is concordant with radiometric ages obtained on excursions lava flows. This record fails to document a significant VDM reduction at the age of the Mono Lake Excursion (32-34 ka).

By providing a new high-resolution ^{10}Be dataset at mid-latitude, the results presented here are of great interest for better constraining the influence of geomagnetic field intensity variations on cosmogenic nuclide production rates. Among these cosmogenic nuclides, atmospheric ^{14}C is of particular importance considering that the ^{14}C calibration curve lacks accurate data during the time period covered by this study. Such new high-resolution datasets from different latitudes are required to make significant advances in quantifying all processes involved in atmospheric $\Delta^{14}\text{C}$ variations. Similarly, reconstructing past VDM variations is of fundamental importance in quantifying in-situ cosmogenic nuclide production rate variations and thus in accurately measuring Earth's surface processes. Further investigations, notably at low-latitude sites where the shielding effect from the geomagnetic dipole field is maximized, are currently underway.

Acknowledgements:

This study was funded by the MAG-ORB project of the ANR (Agence Nationale de la Recherche) program BLANC #ANR-09-BLAN-0053-01. The ASTER French AMS national facility (CEREGE, Aix en Provence) is supported by the INSU/CNRS, the French Ministry of Research and Higher Education, IRD and CEA. We gratefully thank Maurice Arnold, Georges Aumaître and Karim Keddadouche for their invaluable help and support during the AMS measurements, and Wulfran Barthelemy for his assistance during the TIMS measurements. We are also grateful to Eva Moreno, for helping in U-channel sampling and in preliminary paleomagnetic measurements, as well as R. Muscheler who kindly provided the ^{10}Be ice cores data. We acknowledge the scientific and technical crew of R.V. *Marion Dufresne* cruise 140 who collected core MD04-2811. We thank Professor Martin Frank and an anonymous reviewer who helped to improve this manuscript.

References:

Andersen, K.K., Svensson, A., Rasmussen, S.O., Steffensen, J.P., Johnsen, S.J., Bigler, M., Röthlisberger, R., Ruth, U., Siggaard-Andersen, M.-L., Dahl-Jensen, D., Vinther, B.M., Clausen, H.B., 2006. The Greenland Ice Core Chronology 2005, 15–42 ka. Part 1: constructing the time scale. *Quat. Sci. Rev.* 25, 246-257.

Andersen, M.B., Stirling, C.H., Potter, E.-K., Halliday, A.N., 2004. Toward epsilon levels of measurement precision on $^{234}\text{U}/^{238}\text{U}$ by using MC-ICPMS. *Int. J. of Mass Spectrom.* 237, 107-118.

Andersen, M.B., Stirling, C.H., Zimmermann, B., Halliday, A.N., 2010. Precise determination of the open ocean $^{234}\text{U}/^{238}\text{U}$ composition. *Geochem., Geophys., Geosyst.* 11, Q12003, doi: 10.1029/2010GC003318.

587 Baas, J.H., Mienert, J., Abrantes, F., Prins, M.A., 1997. Late Quaternary sedimentation on the
588 Portuguese continental margin: climate-related processes and products. *Palaeogeogr.*,
589 *Palaeoclimatol.*, *Palaeoecol.* 130, 1-23.

590

591 Bacon, M.P., 1984. Glacial to interglacial changes in carbonate and clay sedimentation in the
592 Atlantic Ocean estimated from ^{230}Th measurements. *Isot. Geosci.* 2, 97-111.

593

594 Bard, E., Arnold, M., Maurice, P., Duprat, J., Moyes, J., Duplessy, J.-C., 1987. Retreat
595 velocity of the North Atlantic polar front during the last deglaciation determined by ^{14}C
596 accelerator mass spectrometry. *Nature* 328, 791-794.

597

598 Bard, E., Hamelin, B., Fairbanks, R.G., Zindler, A., Mathieu, G., Arnold, M., 1990. U/Th and
599 ^{14}C ages of corals from Barbados and their use for calibrating the ^{14}C time scale beyond 9000
600 years BP. *Nucl. Instr. Meth. in Phys. Res. B*-52, 461-468.

601

602 Bard, E., Rostek, F., Turon, J.-L., and Gendreau, S., 2000. Hydrological Impact of Heinrich
603 Events in the Subtropical Northeast Atlantic. *Science* 289, 1321-1324.

604

605 Bard, E., Rostek, F., Ménot-Combes, G., 2004. Radiocarbon calibration beyond 20,000 ^{14}C yr
606 B.P. by means of planktonic foraminifera of the Iberian Margin. *Quat. Res.* 61, 204-214.

607

608 Baroni, M., Bard, E., Petit, J.R., Magnan, O., Bourlès, D.L., 2011. Volcanic and solar activity,
609 and atmospheric circulation influences on cosmogenic ^{10}Be fallout at Vostok and Concordia
610 (Antarctica) over the last 60 years. *Geochim. Cosmochim. Acta*, ISSN 0016-7037, doi:
611 10.1016/j.gca.2011.09.002.

612

613 Benson, L., Liddicoat, J., Smoot, J., Sarna-Wojcicki, A., Negrini, R., Lund, S., 2003. Age of
 614 the Mono Lake excursion and associated tephra. *Quat. Sci. Rev.* 22, 135-140.
 615
 616 Bonhommet, N., Babkine, J., 1967. Sur la présence d'aimantations inverses dans la Chaîne
 617 des Puys — Massif Central, France. *C.R.A.S. Paris* 264, 92.
 618
 619 Bourlès, D., Raisbeck, G.M., Yiou, F., 1989. ^{10}Be and ^9Be in marine sediments and their
 620 potential for dating. *Geochim. Cosmochim. Acta* 53, 443-452.
 621
 622 Brown, E.T., Measures, C.I., Edmond, J.M., Bourlès, D.L., Raisbeck, G.M., Yiou, F., 1992.
 623 Continental inputs of beryllium to the oceans. *Earth Planet. Sci. Lett.* 114, 101-111.
 624
 625 Carcaillet, J., Thouveny, N., Bourlès, D.L., 2003. Geomagnetic moment instability between
 626 0.6 and 1.3 Ma from cosmonuclide evidence. *Geophys. Res. Lett.* 30, 1792.
 627
 628 Carcaillet, J., Bourlès, D.L., N.Thouveny, 2004a. Geomagnetic dipole moment and ^{10}Be
 629 production rate intercalibration from authigenic $^{10}\text{Be}/^9\text{Be}$ for the last 1.3 Ma. *Geochem.*
 630 *Geophys. Geosyst.* 5, 397-412, Q05006, doi: 10.1029/2003GC000641.
 631
 632 Carcaillet, J., Bourlès, D.L., Thouveny, N., Arnold, M., 2004b. A high resolution authigenic
 633 $^{10}\text{Be}/^9\text{Be}$ record of geomagnetic moment variations over the last 300 ka from sedimentary
 634 cores of the Portuguese margin. *Earth Planet. Sci. Lett.* 219, 397-412.
 635
 636 Cassata, W.S., Singer, B.S., Cassidy, J., 2008. Laschamp and Mono Lake geomagnetic
 637 excursions recorded in New Zealand. *Earth Planet. Sci. Lett.* 268, 76-88.
 638

639 Channell, J., 2006. Late Brunhes polarity excursions (Mono Lake, Laschamp, Iceland Basin
640 and Pringle Falls) recorded at ODP Site 919 (Irminger Basin). *Earth Planet. Sci. Lett.* 244,
641 378-393.

642

643 Chase, Z., Anderson, R.F., Fleisher, M.Q., Kubik, P.W., 2002. The influence of particle
644 composition and particle flux on scavenging of Th, Pa and Be in the ocean. *Earth Planet. Sci.*
645 *Lett.* 204, 215-229.

646

647 Chmeleff, J., von Blanckenburg, F., Kossert, K., Jakob, D., 2010. Determination of the ^{10}Be
648 half-life by multicollector ICP-MS and liquid scintillation counting. *Nuclear Instruments and*
649 *Methods in Physics Research Section B: Beam Interactions with Materials and Atoms* 268,
650 192-199.

651

652 Christl, M., Strobl, C., Mangini, A., 2003. Beryllium-10 in deep-sea sediments: a tracer for
653 the Earth's magnetic field intensity during the last 200,000 years. *Quat. Sci. Rev.* 22, 725-739.

654

655 Christl, M., Mangini, A., Kubik, P.W., 2007. Highly resolved Beryllium-10 record from ODP
656 Site 1089--A global signal? *Earth Planet. Sci. Lett.* 257, 245-258.

657

658 Christl, M., Lippold, J., Steinhilber, F., Bernsdorff, F., Mangini, A., 2010. Reconstruction of
659 global ^{10}Be production over the past 250 ka from highly accumulating Atlantic drift
660 sediments. *Quat. Sci. Rev.* 29, 2663-2672.

661

662 Coe, R.S., Liddicoat, J.C., 1994. Overprinting of natural magnetic remanence in lake
663 sediments by subsequent high intensity field. *Nature* 367, 57-59.

664

665 Denham, C.R., Cox, A., 1971. Evidence that the Laschamp polarity event did not occur
666 13300–30400 years ago. *Earth Planet. Sci. Lett.* 13, 181-190.

667

668 Deschamps, P., Doucelance, R., Ghaleb, B., Michelot, J.L., 2003. Further investigations on
669 optimized tail correction and high-precision measurement of Uranium isotopic ratios using
670 Multi-Collector ICP-MS. *Chem. Geol.* 201, 141-160.

671

672 Deschamps, P., Hillaire-Marcel, C., Michelot, J.L, Doucelance, R., Ghaleb, B., Buschaert, S.,
673 2004. $^{234}\text{U}/^{238}\text{U}$ Disequilibrium along stylolitic discontinuities in deep Mesozoic limestone
674 formations of the Eastern Paris basin: evidence for discrete uranium mobility over the last 1-2
675 million years. *Hydrol. Earth Syst. Sci.* 8, 35-46.

676

677 Deschamps, P., Durand, N., Bard, E., Hamelin, B., Camoin, G., Thomas, A.L., Henderson,
678 G.M., Okuno, J., Yokoyama, Y., Submitted. Dramatic ice sheet collapse 14,600 years ago at
679 onset of Bølling warming. *Nature*.

680

681 Elsasser, W., Ney, E.P., Winckler, J.R., 1956. Cosmic-ray intensity and geomagnetism.
682 *Nature* 178, 1226-1227.

683

684 Finkel, R.C., Nishiizumi, K., 1997. Beryllium 10 concentrations in the Greenland Ice Sheet
685 Project 2 ice core from 3–40 ka. *J. Geophys. Res.* 102 26699-26706.

686

687 François, R., Frank, M., Rutgers van der Loeff, M.M., Bacon, M.P., 2004. ^{230}Th
688 normalization: An essential tool for interpreting sedimentary fluxes during the late
689 Quaternary. *Paleoceanography* 19, 10-18.

690

691 Frank, M., Schwarz, B., Baumann, S., Kubik, P.W., Suter, M., Mangini, A., 1997. A 200 kyr
692 record of cosmogenic radionuclide production rate and geomagnetic field intensity from ^{10}Be
693 in globally stacked deep-sea sediments. *Earth Planet. Sci. Lett.* 149, 121-129.

694

695 Frank, M., 2000. Comparison of cosmogenic radionuclide production and geomagnetic field
696 intensity over the last 200,000 years. *Phil. Trans. R. Soc. Lond. A* 358, 1089-1107.

697

698 Guillou, H., Singer, B.S., Laj, C., Kissel, C., Scaillet, S., Jicha, B.R., 2004. On the age of the
699 Laschamp geomagnetic excursion. *Earth Planet. Sci. Lett.* 227, 331-343.

700 Guyodo, Y., Valet, J.P., 1999. Global changes in intensity of the Earth's magnetic field during
701 the past 800 kyr. *Nature* 399, 249-252.

702

703 Hall, I.R., McCave, I.N., 2000. Palaeocurrent reconstruction, sediment and thorium focussing
704 on the Iberian margin over the last 140 ka. *Earth Planet. Sci. Lett.* 178, 151-164.

705

706 Henderson, G.M., Anderson, R.F., 2003. The U-series toolbox for paleoceanography,. *Rev.*
707 *Mineral. Geochem.* 52, 493-531.

708

709 Henken-Mellies, W.U., Beer, J., Heller, F., Hsü, K.J., Shen, C., Bonani, G., Hofmann, H.J.,
710 Suter, M., Wölfli, W., 1990. ^{10}Be and ^9Be in South Atlantic DSDP Site 519: Relation to
711 geomagnetic reversals and to sediment composition. *Earth Planet. Sci. Lett.* 98, 267-276.

712

713 Kent, D.V., Hemming, S.R., Turrin, B.D., 2002. Laschamp Excursion at Mono Lake? *Earth*
714 *Planet. Sci. Lett.* 197, 151-164.

715

716 Knudsen, M.F., Henderson, G.M., Frank, M., Mac Niocaill, C., Kubik, P.W., 2008. In-phase
 717 anomalies in Beryllium-10 production and palaeomagnetic field behaviour during the Iceland
 718 Basin geomagnetic excursion. *Earth Planet. Sci. Lett.* 265, 588-599.
 719
 720 Kissel, C., Laj, C., Labeyrie, L., Dokken, T., Voelker, A., Blamart, D., 1999. Rapid climatic
 721 variations during marine isotopic stage 3: magnetic analysis of sediments from Nordic Seas
 722 and North Atlantic. *Earth Planet. Sci. Lett.* 171, 489-502.
 723
 724 Korhonen, K., Donadini, F., Riisager, P., Pesonen, L., 2008. GEOMAGIA50: an
 725 archeointensity database with PHP and MySQL. *Geochem. Geophys. Geosyst.* 9 (4), Q04029,
 726 doi: 10.1029/2007GC001893
 727
 728 Korschinek, G., Bergmaier, A., Faestermann, T., Gerstmann, U.C., Knie, K., Rugel, G.,
 729 Wallner, A., Dillmann, I., Dollinger, G., von Gostomski, C.L., Kossert, K., Maiti, M.,
 730 Poutivtsev, M., Remmert, A., 2010. A new value for the half-life of ^{10}Be by Heavy-Ion
 731 Elastic Recoil Detection and liquid scintillation counting. *Nucl. Instr. Meth. in Phys. Res.*
 732 *Sect. B: Beam Interact. Mater. At.* 268, 187-191.
 733
 734 Kristjansson, L., Gudmundsson, A.A., 1980. Geomagnetic excursions in late-glacial basalt
 735 outcrop in south-western Iceland. *Geophys. Res. Lett.* 7, 337–340.
 736
 737 Laj, C., Kissel, C., Beer, J., 2004. High resolution global paleointensity stack since 75 kyr
 738 (GLOPIS-75) calibrated to absolute values. In: Channell, J.E.T., Kent, D.V., Lowrie, W.,
 739 Meert, J.G. (Eds.), *Timescales of the Paleomagnetic Field.* . *Geophys. Monogr.* 145.
 740
 741 Lal, D., 1988. Theoretically expected variations in the terrestrial cosmic-ray production rates
 742 of isotopes. *Soc. Italia. di Fisica-Bologna-Italy XCV corso*, 216-233.

743

744 Lebatard, A.-E., Bourlès, D.L., Braucher, R., Arnold, M., Düringer, P., Jolivet, M., Moussa,
745 A., Deschamps, P., Roquin, C., Carcaillet, J., Schuster, M., Lihoreau, F., Likius, A., Mackaye,
746 H.T., Vignaud, P., Brunet, M., 2010. Application of the authigenic $^{10}\text{Be}/^9\text{Be}$ dating method to
747 continental sediments: Reconstruction of the Mio-Pleistocene sedimentary sequence in the
748 early hominid fossiliferous areas of the northern Chad Basin. *Earth Planet. Sci. Lett.* 297, 57-
749 70.

750

751 Leduc, G., Thouveny, N., Bourlès, D.L., Blanchet, C.L., Carcaillet, J.T., 2006. Authigenic
752 $^{10}\text{Be}/^9\text{Be}$ signature of the Laschamp excursion: A tool for global synchronisation of
753 paleoclimatic archives. *Earth Planet. Sci. Lett.* 245, 19-28.

754

755 Levi, S., Gudmunsson, H., Duncan, R.A., Kristjansson, L., Gillot, P.V., Jacobsson, S.P.,
756 1990. Late Pleistocene geomagnetic excursion in Icelandic lavas: Confirmation of the
757 Laschamp excursion. *Earth Planet. Sci. Lett.* 96, 443–457.

758

759 Liddicoat, J.C., 1992. Mono Lake Excursion in Mono Basin, California, and at Carson Sink
760 and Pyramid Lake, Nevada. *Geophys. J. Intern.* 108, 442-452.

761

762 Lund, S.P., Schwartz, M., Keigwin, L., Johnson, T., 2005. Deep-sea sediment records of the
763 Laschamp geomagnetic field excursion (<41,000 calendar years before present. *J. Geophys.*
764 *Res.* 110, Q12006.

765

766 Masarik, J., Beer, J., 1999. Simulation of particle fluxes and cosmogenic nuclide production
767 in the Earth's atmosphere. *J. Geophys. Res.* 104, 12099-12111.

768

769 Monaghan, M.C., Krishnaswami, S., Turekian, K.K., 1986. The global-average production
770 rate of ^{10}Be . *Earth Planet. Sci. Lett.* 76, 279-287.

771

772 Moreno, E., Thouveny, N., Delanghe, D., McCave, I.N., Shackleton, N.J., 2002. Climatic and
773 oceanographic changes in the Northeast Atlantic reflected by magnetic properties of
774 sediments deposited on the Portuguese Margin during the last 340 ka. *Earth Planet. Sci. Lett.*
775 202, 465-480.

776

777 Muscheler, R., Beer, J., Wagner, G., Laj, C., Kissel, C., Raisbeck, G.M., Yiou, F., Kubik,
778 P.W., 2004. Changes in the carbon cycle during the last deglaciation as indicated by the
779 comparison of ^{10}Be and ^{14}C records. *Earth Planet. Sci. Lett.* 219, 325-340.

780

781 Muscheler, R., Beer, J., Kubik, P.W., Synal, H.A., 2005. Geomagnetic field intensity during
782 the last 60,000 years based on ^{10}Be and ^{36}Cl from the Summit ice cores and ^{14}C . *Quat. Sci.*
783 *Rev.* 24, 1849-1860.

784

785 Negrini, R.M., Davis, J.O., Verosub, K.L., 1984. Mono Lake geomagnetic excursion found at
786 Summer Lake, Oregon. *Geology* 12, 643-646.

787

788 Negrini, R., Erbes, K.F., Herrera, A.M., Roberts, A.P., Cohen, A., Palacios-Fest, M., Wigand,
789 P.E., Foit, F., 2000. A paleoclimate record for the past 250,000 years from Summer Lake,
790 Oregon, USA: I. Age control and magnetic lake level proxies. *J. Paleolimn.* 24, 125-149.

791

792 Nishiizumi, K., Imamura, M., Caffee, M.W., Southon, J.R., Finkel, R.C., McAninch, J., 2007.
793 Absolute calibration of ^{10}Be AMS standards. *Nucl. Instr. Meth. in Phys. Res. Sect. B: Beam*
794 *Interact. Mater. At.* 258, 403-413.

795

796 NGRIP-dating-group, 2006. Greenland Ice Core Chronology 2005 (GICC05). IGBP
797 PAGES/World Data Center for Paleoclimatology Data Contribution Series # 2006-118.
798 NOAA/NCDC Paleoclimatology Program, Boulder CO, USA.
799
800 NGRIP-Members, 2004. High-resolution record of Northern Hemisphere climate extending
801 into the last interglacial period. *Nature* 431, 147–151.
802
803 Nowaczyk, N.R., Knies, J., 2000. Magnetostratigraphic results from the eastern Arctic ocean:
804 AMS ^{14}C ages and relative paleointensity data of the mono lake and Laschamp geomagnetic
805 reversal excursions. *Geophys. J. Intern.* 140, 185-197.
806
807 Paillard, D., Labeyrie, L., Yiou, P., 1996. Macintosh program performs time-series analysis.
808 *Eos Trans. AGU* 77, 379.
809
810 Pailler, D., Bard, E., 2002. High frequency palaeoceanographic changes during the past 140
811 000 yr recorded by the organic matter in sediments of the Iberian Margin. *Palaeogeogr.*
812 *Palaeoclimatol. Palaeoecol.* 181, 431-452.
813
814 Plenier, G., Valet, J.-P., Guérin, G., Lefèvre, J.-C., LeGoff, M., Carter-Stiglitz, B., 2007.
815 Origin and age of the directions recorded during the Laschamp event in the Chaîne des Puys
816 (France). *Earth Planet. Sci. Lett.* 259, 414-431.
817
818 Raisbeck, G.M., Yiou, F., Fruneau, M., Loiseaux, J.M., Lieuvain, M., Ravel, J.C., 1981.
819 Cosmogenic Be-10/Be-7 as a probe of atmospheric transport processes. *Geophys. Res. Lett.*
820 8, 1015-1018.
821

Raisbeck, G.M., Yiou, F., Jouzel, J., Stocker, T.F., 2007. Direct north-south synchronization of abrupt climate change record in ice cores using Beryllium 10. *Climate of the Past* 3, 541-547.

Rasmussen, S.O., Seierstad, I.K., Andersen, K.K., Bigler, M., D. Dahl-Jensen, S.J. Johnsen, 2006. NGRIP, GRIP, and GISP2 Synchronization Match Points. IGBP PAGES/World Data Center for Paleoclimatology. Data Contribution Series # 2006-120. NOAA/NCDC Paleoclimatology Program, Boulder CO, USA.

Reimer, P.J., Baillie, M.G.L., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Bronk Ramsey, C., Buck, C.E., Burr, G.S., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Hajdas, I., Heaton, T.J., Hogg, A.G., Hughen, K.A., Kaiser, K.F., Kromer, B., McCormac, G., Manning, S., Reimer, R.W., Richards, D.A., Southon, J.R., Talamo, S., Turney, C.S.M., van der Plicht, J., Weyhenmeyer, C.E., 2009. IntCal09 and Marine09 radiocarbon age calibration curves, 0-50000 Years cal BP. *Radiocarbon* 51, 1111-1150.

Robinson, C., Raisbeck, G.M., Yiou, F., Lehman, B., Laj, C., 1995. The relationship between ^{10}Be and geomagnetic field strength records in central North Atlantic sediments during the last 80 ka. *Earth Planet. Sci. Lett.* 136, 551-557.

Shackleton, N.J., Fairbanks, R.G., Chiu, T.-C., Parrenin, F., 2004. Absolute calibration of the Greenland time scale: implications for Antarctic time scales and for $\delta^{14}\text{C}$. *Quat. Science Rev.* 23, 1513-1522.

Singer, B.S., Guillou, H., Jicha, B.R., Laj, C., Kissel, C., Beard, B.L., Johnson, C.M., 2009. $^{40}\text{Ar}/^{39}\text{Ar}$, K-Ar and ^{230}Th - ^{238}U dating of the Laschamp Excursion: A radioisotopic tie-point for ice core and climate chronologies. *Earth Planet. Sci. Lett.* 286, 80-88.

Skinner, L.C., and McCave, I.N., 2003. Analysis and modelling of gravity- and piston coring based on soil mechanics. *Mar. Geol.* 199, 181-204.

Svensson, A., Andersen, K.K., Bigler, M., Clausen, H.B., Dahl-Jensen, D., Davies, S.M., Johnsen, S.J., Muscheler, R., Parrenin, F., Rasmussen, S.O., Röthlisberger, R., Seierstad, I.,

Steffensen, J.P., Vinther, B.M., 2008. A 60 000 year Greenland stratigraphic ice core chronology. *Climate of the Past* 4, 47-57.

Széréméta, N., Bassinot, F., Balut, Y., Labeyrie, L., Pagel, M., 2004. Oversampling of sedimentary series collected by giant piston corer: Evidence and corrections based on 3.5-kHz chirp profiles. *Paleoceanography* 19, PA1005, doi: 10.1029/2002PA000795.

Taylor, J.R., 1997. *An Introduction to Error Analysis. The study of Uncertainties in Physical Measurements*, Second ed., University Science Books, Sausalito, USA.

Thomson, J., Nixon, S., Summerhayes, C.P., Schönfeld, J., Zahn, R., Grootes, P., 1999. Implications for sedimentation changes on the Iberian margin over the last two glacial/interglacial transitions from ($^{230}\text{Th}_{\text{excess}}$)₀ systematics. *Earth Planet. Sci. Lett.* 165, 255-270.

Thouveny, N., Creer, K.M., 1992. On the brevity of the Laschamp Excursion. *Bull. Soc. Géol. Fr.* 6, 771–780.

Thouveny, N., Moreno, E., Delanghe, D., Candon, L., Lancelot, Y., Shackleton, N.J., 2000. Rock magnetic detection of distal ice-rafted debries: clue for the identification of Heinrich layers on the Portuguese margin. *Earth Planet. Sci. Lett.* 180, 61-75.

Thouveny, N., Carcaillet, J., Moreno, E., Leduc, G., Nérini, D., 2004. Geomagnetic moment variation and paleomagnetic excursions since 400 kyr BP: a stacked record from sedimentary sequences of the Portuguese margin. *Earth Planet. Sci. Lett.* 219, 377-396.

Thouveny, N., Bourlès, D.L., Saracco, G., Carcaillet, J.T., Bassinot, F., 2008. Paleoclimatic context of geomagnetic dipole lows and excursions in the Brunhes, clue for an orbital influence on the geodynamo? *Earth Planet. Sci. Lett.* 275, 269-284.

Vlag, P., Thouveny, N., Williamson, D., Rochette, P., Ben-Atig, F., 1996. Evidence for a geomagnetic excursion recorded in the sediments of Lac St. Front, France: a link with the Laschamp Excursion? *J. Geophys. Res.* 101, 28211–28230.

Wagner, G., Beer, J., Laj, C., Kissel, C., Masarik, J., Muscheler, R., Synal, H.A., 2000a. Chlorine-36 evidence for the Mono Lake event in the Summit GRIP ice core. *Earth Planet. Sci. Lett.* 181, 1-6.

Wagner, G., Masarik, J., Beer, J., Baumgartner, S., Imboden, D., Kubik, P.W., Synal, H.A., Suter, M., 2000b. Reconstruction of the geomagnetic field between 20 and 60 kyr BP from cosmogenic radionuclides in the GRIP ice core. *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms* 172, 597-604.

Wang, Y.J., Cheng, H., Edwards, R.L., An, Z.S., Wu, J.Y., Shen, C.-C., and Dorale, J.A., 2001. A High-Resolution Absolute-Dated Late Pleistocene Monsoon Record from Hulu Cave, China *Science* 294, 2345-2348.

Yiou, F., Raisbeck, G.M., Baumgartner, S., Beer, J., Hammer, C., Johnsen, S., Jouzel, J., Kubik, P.W., Lestringuez, J., Stiévenard, M., Sutter, M., Yiou, P., 1997. Beryllium 10 in the Greenland Ice Core Project ice core at Summit, Greenland. *J. Geophys. Res.* 102, 26783-26794.

Zimmerman, S.H., Hemming, S.R., Kent, D.V., Searle, S.Y., 2006. Revised chronology for late Pleistocene Mono Lake sediments based on paleointensity correlation to the global reference curve. *Earth Planet. Sci. Lett.* 252, 94-106.

917

918

Figure Captions :

919

“The Laschamp geomagnetic dipole low expressed as a cosmogenic ^{10}Be atmospheric

920

overproduction at ~41 ka.”

921

By L. Ménabréaz *, N. Thouveny, D.L. Bourlès, P. Deschamps, B. Hamelin and F. Demory.

922

* corresponding author: menabreaz@cerege.fr

923

Revised version submitted to Earth and Planetary Science Letters

924

Fig. 1. Location map of the studied MD04-2811 core. Other cores cited in the text

925

(MD95-2042 and SU81-18 were recovered at the same site.

926

Fig. 2. Magnetic properties and cosmogenic nuclide ^{10}Be record as a function of depth in core

927

MD04-2811. (a) Magnetic susceptibility; susceptibility peaks identify distal ferrimagnetic

928

IRD layers deposited during Heinrich Events 1 to 5 (labeled H1 to H5). (b) Relative

929

paleointensity (NRM/ARM ratios at 30mT alternating field step). (c) Authigenic $^{10}\text{Be}/^9\text{Be}$

930

ratios (10^{-8}) with 1σ uncertainty. Solid and dotted lines correspond to the mean value over the

931

20-50 ka time period and the associated standard deviation (1σ), respectively (see Table 1).

932

Fig. 3. (a-b) Paleoclimatic variations over Greenland and the Portuguese margin: (a)

933

variations of the air paleotemperature proxy over Greenland: $\delta^{18}\text{O}$ of the NGRIP ice core

934

(NGRIP-Members, 2004 and references therein) plotted on the GICC05 time-scale (NGRIP

935

dating group, 2006 and references therein); (b) variations of the Portuguese margin sea-

936

surface paleotemperature proxy: UK'37 alkenone index in core MD95-2042 (raw data of

937

Pailler and Bard, 2002) on a time scale tuned to NGRIP $\delta^{18}\text{O}$ using the Analyserie software

938

(Paillard et al., 1996). Correlation tie-points are marked by vertical dashes. Triangles indicate

939

the position of the 23 samples used for AMS dating: raw data (Electronic Appendix, Table

940

A.1), are from Bard et al., 2004 and Shackleton et al., 2004); (c-d) Magnetic Susceptibility

941

(MS) records of Portuguese margin cores as a function of depth: (c) core MD95-2042

942

(specific MS in $10^{-9} \text{ m}^3/\text{kg}$) (data from Thouveny et al., 2000 transferred on the NGRIP

943

chronology; (d) core MD04-2811 (volume susceptibility 10^{-5} SI). Susceptibility peaks (S0 to

944

S5) mark IRD layers deposited during the Younger Dryas (YD) and Heinrich events (H1 to

945

H5) events.

Fig. 4. Age-depth relationship in core MD04-2811. Open circles stand for the age model derived from the GICC05 chronology: the MD95-2042 age control obtained tuning UK'37 alkenone index and NGRIP $\delta^{18}\text{O}$ (see Fig.3) was applied to core MD04-2811 by correlation of MS profiles using the Analyse series software (Paillard et al., 1996). Ages of the MD04-2811 susceptibility peaks (grey bands, ages reported on the left) are undistinguishable from ages of Heinrich layers computed by Thouveny et al. (2000): H2: 24778 ± 297 yr BP; H3: 31163 ± 509 yr BP; H4: 39379 ± 948 yr BP.

Fig. 5. (a) Downcore variation of the $^{230}\text{Th}_{\text{xs}}$ -normalized ^{10}Be fluxes and (b) relation with authigenic $^{10}\text{Be}/^9\text{Be}$ ratios measured on the same homogenized sediment samples.

Fig. 6. (a and b) Relative temporal variations of ^{10}Be atmospheric production rate proxies over the 20-50 ka interval. All records are normalized to their respective mean values over the studied time interval. (a) Black dots represent the authigenic $^{10}\text{Be}/^9\text{Be}$ record and the associated 1σ uncertainty. Grey dots represent the $^{230}\text{Th}_{\text{xs}}$ -normalized ^{10}Be fluxes. (b) Black dots represent the authigenic $^{10}\text{Be}/^9\text{Be}$ record and the associated 1σ uncertainty, and grey line represents the Greenland ^{10}Be flux in Summit ice cores on GICC05 time scale (Finkel et al., 1997; Yiou et al., 1997; composite GRIP-GISP2: Muscheler et al., 2005). (c) Relative temporal variations of geomagnetic field intensity over the 20-50 ka interval. Solid lines represent the 1σ uncertainty envelop of the GLOPIS-75 stack: raw data from Laj et al., (2004) are normalized to $8 \times 10^{22} \text{ A.m}^2$ (most recent GLOPIS VADM value) and plotted on the GICC05 time scale using the NGRIP, GRIP and GISP2 synchronization match points dataset (Rasmussen et al., 2006). Dots are the authigenic $^{10}\text{Be}/^9\text{Be}$ -derived VDM record (M/M_0) calculated using the Lal's algorithm (1988): $P/P_0 = (M/M_0)^{-0.5}$, where P/P_0 is the authigenic $^{10}\text{Be}/^9\text{Be}$ ratio divided by the surface sample value of $(1.36 \pm 0.35) \times 10^{-8}$, undistinguishable within uncertainties from the long-term average $(1.44 \pm 0.043) \times 10^{-8}$.

Fig. 7. (a) Authigenic $^{10}\text{Be}/^9\text{Be}$ ratios measured along MD04-2811 (solid circles) and MD95-2042 cores (open circle, Carcaillet et al., 2004) on the MD04-2811 depth scale. (b) Calculated authigenic $^{10}\text{Be}/^9\text{Be}$ composite record over the 20-50 ka interval. This record is low-pass filtered ($1/1000 \text{ yr}^{-1}$). Associated uncertainties are 1σ .

Fig. 8. Geomagnetic dipole moment variation during the 20-50 ka interval. The black curve is the Virtual Axial Dipole Moment (VADM) variations reconstructed from SINT800 (Guyodo and Valet, 1999). The grey zone is the 1σ envelop of the GLOPIS-75 based VADM

977 reconstruction (Laj et al., 2004). Bold black curve is the authigenic $^{10}\text{Be}/^9\text{Be}$ -based VDM
978 (Virtual Dipole Moment) record and 1σ associated uncertainties.

979

980

981 **Highlights to the manuscript:** “The Laschamp geomagnetic dipole low expressed as a
982 cosmogenic ^{10}Be atmospheric overproduction at ~ 41 ka.”

983

984 by L. Menabreaz *, N. Thouveny, D.L. Bourles, P. Deschamps, B. Hamelin and F. Demory.

985

Earth and Planetary Science Letters

986

987 ☐ Cosmogenic ^{10}Be concentration was measured along a marine sediment core (20 -50ka)

988

☐ Authigenic ^9Be and excess ^{230}Th were used to normalize ^{10}Be concentrations

989

☐ A doubling of the cosmogenic production is revealed at ~ 41 ka (NGRIP age)

990

☐ Geomagnetic field intensity variations are reconstructed from ^{10}Be production rates

991

☐ The dipole low of Laschamp excursion is confirmed at 41 ka.

992