The Generalized Linear Sampling Method for limited aperture measurements

Lorenzo Audibert* and Houssem Haddar †

Abstract. We extend the so-called Generalized Linear Sampling Method (GLSM) to the case of limited aperture data at a fixed frequency. In this case the factorization of the sampling operator does not obey the symmetry required in the justification of the GLSM introduced in Audibert-Haddar [Inverse Problems, 2014]. We propose a new formulation by adding an extra penalty term that asymptotically correct the non symmetry of the GLSM original penalty term. The analysis of the new formulation is first presented in an abstract framework. We then show how to apply our setting to the scalar problem with far field measurements or near field measurements on a limited aperture. We finally validate the method through some numerical tests in two dimensions and for far field measurements.

Key words. Inverse scattering problems, Linear Sampling Method, Generalized Linear Sampling Method, Factorization Method, Qualitative methods

AMS subject classifications. 35R60, 35R30, 65M32

1. Introduction. This work is concerned with the design of so-called sampling methods [7, 6, 8, 13, 4] for inverse scattering problems where one would like to determine the shape of extended targets from fixed frequency multi-static data. More precisely we extend and analyze the recently introduced Generalized Linear Sampling Method [3] (GLSM) to limited aperture data. The GLSM framework developed in [3] provides an exact characterization of the target shape in terms of the so-called far field operator (at fixed frequency and for full aperture). This characterization is based on two factorizations of the far field operator. The first one is used to justify the Linear Sampling Method (LSM) and the second one is at the heart of the Factorization Method (FM). Considering general limited aperture data break the symmetry of the second factorization and prevent the application of the results of [3] or [13] on the FM. The characterization of the GLSM is based on constructing nearby solution to the far field equation as minimizing sequences of a special cost functional. In this cost functional the symmetric factorization is important to ensure that the regularization term has suitable properties. In this article we propose a modification of the regularization term and analyze this modification in order to prove exact characterization even for non symmetric factorization.

The main idea behind our method is that without symmetric factorization it is not possible to control directly the norm of the Herglotz wave that approximately solves the far field equation. However we have access to a term that is close to this quantity and we can bound the error we made, therefore controlling the norm of the associated Herglotz wave. Due to this splitting the control is coarser and therefore it reflects the fact that this situation is less favorable for imaging. The fact that the regularization involves compact operators or the case of noisy operators are covered using the idea already proposed in [3]. However the interesting property of strong convergence of the minimizing sequence of the cost functional demonstrated

*Department STEP, EDF R&D, Chatou, France and CMAP, INRIA, Ecole Polytechnique, CNRS and Université Paris Saclay, Palaiseau, France. (lorenzo.audibert@edf.fr).

†CMAP, INRIA, Ecole Polytechnique, CNRS and Université Paris Saclay, Palaiseau, France. (houssem.haddar@inria.fr).

in [2] could not be simply extended. The second main contribution of this article is to add a regularization term to lower the hypothesis of [2] on the regularization term. This new results extend the validity of the results of [2] and enable an extension to non symmetric factorization. In order to introduce those ideas we choose to present the case of scalar inverse scattering from inhomogeneous inclusions for limited aperture far field measurements. We also indicate how the method can be easily extended to near field data.

On the numerical side we introduce a second order method to minimize the cost functional, this method prove to be more efficient than the one use in [3]. The superiority of our indicator function is demonstrated for symmetric factorization. The theory does not say how to choose the regularization parameter for symmetric factorization, the method does not seem to be very sensitive and an heuristic choice give good result. For non symmetric factorization this choice is by far more important and we propose three heuristics to set this parameter.

The article is organized as follows. In Section 2 a model problem is introduced to motivate the GLSM for non symmetric factorization. Theoretical extension for the symmetric factorization is given in section 3.1 and the case of non-symmetric factorization is treated in section 3.2. Section 4 provides an example of application by completely treating the model problem introduced in section 2. Section 5 show how nearfield data easily fit into the theory developed in Section 3. The last section (Section 6) is devoted to numerical algorithms issued from section 4 along with validating numerical results and discussion on the difference between symmetric and non-symmetric cases.

2. A model problem for limited aperture data. We choose to present our method for the simple model of inverse time harmonic scattering problem from inhomogeneous targets. For a wave number $k > 0$, the total field solve the following scalar wave equation:

$$\Delta u + k^2 n u = 0 \text{ in } \mathbb{R}^d$$

with $d = 2$ or 3 and with $n \in L^\infty(\mathbb{R}^d)$ denoting the refractive index such that the support of $n - 1$ is included inside $\bar{D}$ with D a bounded domain with Lipschitz boundary and connected complement and such that $\Im(n) \geq 0$.

We consider the cases where the total field is generated by incident plane waves, $u^i(\theta, x) := e^{ikx \cdot \theta}$ with $x \in \mathbb{R}^d$ and $\theta \in \Gamma_s$ ($\Gamma_s \subset \mathbb{S}^{d-1}$ the unit sphere) and we denote by u^s the scattered field defined by

$$u^s(\theta, \cdot) = u - u^i(\theta, \cdot) \quad \text{in } \mathbb{R}^d,$$

which is assumed to be satisfying the Sommerfeld radiation condition,

$$\lim_{r \rightarrow \infty} \int_{|x|=r} \left| \frac{\partial u^s}{\partial r} - iku^s \right|^2 ds = 0.$$

The data for the inverse problem is formed by noisy measurements of the so called far field pattern $u^\infty(\theta, \hat{x})$ defined by

$$u^s(\theta, x) = \frac{e^{ik|x|}}{|x|^{(d-1)/2}} (u^\infty(\theta, \hat{x}) + O(1/|x|))$$

65 as $|x| \rightarrow \infty$ for all $(\theta, \hat{x}) \in \Gamma_s \times \Gamma_m$, where Γ_m is a subset of $\mathbb{S}^{d-1}$ possibly different from Γ_s .
 66 The goal is to be able to reconstruct D from these measurements (without knowing n). We
 67 introduce the far field operator $F : L^2(\Gamma_s) \rightarrow L^2(\Gamma_m)$, defined by

$$Fg(\hat{x}) := \int_{\Gamma_s} u^\infty(\theta, \hat{x})g(\theta)ds(\theta), \quad \hat{x} \in \Gamma_m.$$

68 Let us define, for $\psi \in L^2(D)$, the unique function $w \in H_{\text{loc}}^1(\mathbb{R}^d)$ satisfying

$$(1) \quad \begin{cases} \Delta w + nk^2 w = -k^2(n-1)\psi \text{ in } \mathbb{R}^d, \\ \lim_{r \rightarrow \infty} \int_{|x|=r} \left| \frac{\partial w}{\partial r} - ikw \right|^2 ds = 0. \end{cases}$$

By linearity of the forward scattering problem, Fg is nothing but the far field pattern of w solution of (1) with $\psi = v_g$ in D , where

$$v_g(x) := \int_{\Gamma_s} e^{ikx \cdot \theta} g(\theta)ds(\theta), \quad g \in L^2(\Gamma_s), \quad x \in \mathbb{R}^d.$$

70 Now consider the (compact) operator $H_s : L^2(\Gamma_s) \rightarrow L^2(D)$ defined by

$$(2) \quad H_s g := v_g|_D,$$

72 and the (compact) operator $G_m : \overline{\mathcal{R}(H_s)} \subset L^2(D) \rightarrow L^2(\Gamma_m)$ defined by

$$(3) \quad G_m \psi := w^\infty|_{\Gamma_m}$$

where w^∞ is the far field of $w \in H_{\text{loc}}^1(\mathbb{R}^d)$ solution of (1) and where $\overline{\mathcal{R}(H_s)}$ denotes the closure of the range of H_s in $L^2(D)$. Then clearly

$$F = G_m H_s$$

74 One can still decompose F to get the second factorisation of the far field operator. More
 75 precisely, for the case under consideration, since the far field pattern of w has the following
 76 expression ([4])

$$77 \quad w^\infty(\hat{x}) = - \int_D e^{-iky \cdot \hat{x}} (1-n)k^2(\psi(y) + w(y))dy,$$

one simply has $G_m = H_m^* T \psi$, where $H_m^* : L^2(D) \rightarrow L^2(\Gamma_m)$ is the adjoint of the operator H_m (defined similarly to H_s but with Γ_s replaced by Γ_m) and whose expression is given by

$$H_m^* \varphi(\hat{x}) := \int_D e^{-iky \cdot \hat{x}} \varphi(y)dy, \quad \varphi \in L^2(D), \quad \hat{x} \in \Gamma_m,$$

78 and where the operator $T : L^2(D) \rightarrow L^2(D)$ is defined by

$$79 \quad (4) \quad T\psi := -k^2(1-n)(\psi + w),$$

80 with $w \in H_{\text{loc}}^1(\mathbb{R}^d)$ being the solution of (1). Finally we end up with

81 (5)
$$F = H_m^* T H_s.$$

82 This factorization is called "non symmetric" in the cases $H_m \neq H_s$ which correspond to
 83 $\Gamma_s \neq \Gamma_m$. The GLSM as formulated in [3] applies to the "symmetric" cases, i.e. $\Gamma_s = \Gamma_m$.
 84 Physically the latter correspond with sources and receivers on symmetric opposite sides of the
 85 target (as shown in Figure 1).

Figure 1. Sources-receivers configurations that correspond with symmetric factorizations of the far field operator.

86 Our focus in the following is to extend the GLSM to non symmetric factorizations of the
 87 measurement operator.

3. Theoretical foundation of the GLSM for limited aperture. In this section we shall give the theoretical foundation of the extension of the GLSM to non symmetric factorizations. We will adopt an abstract framework that can be applied to other settings than the one presented in the previous section (See for instance Section 5 where the case of near field data is considered). As pointed above, the "symmetry" in the factorization of the far field operator is of primary importance in the GLSM framework of [3] where the following cost functional (for noise-free data) was introduced:

$$J_\alpha(\phi; g) := \alpha |\langle Bg, g \rangle| + \alpha^{1-\eta} |\langle Fg - \phi, g \rangle| + \|Fg - \phi\|^2$$

88 with B being an operator constructed from F and that has a "symmetric" factorization. The
 89 latter seems to be hard to ensure in general when F itself has not a "symmetric" factorization.
 90 In some special cases this can be done as for heterogeneous backgrounds [10] or special settings
 91 of the near field data [5]. However, in the case of limited aperture presented above with
 92 $\Gamma_m \neq \Gamma_s$, this type of construction seems to be impossible to achieve. This is why we shall
 93 consider in the following only the case $B = F$.

94 As has been pointed out in [2], for the case $B = F$, one cannot guarantee in general the
 95 strong convergence of Herglotz waves associated with the minimizing sequences of $J_\alpha(\phi; g)$
 96 (when the sampling point is inside D). Since this convergence is an important property for
 97 some imaging algorithms (as in [2] for the case of differential measurements), we shall first
 98 modify the setting of GLSM so that one obtain this convergence result even in the case $B = F$.
 99 The idea is to add an extra (carefully chosen) penalty term that is inspired from difficulties
 100 encountered in establishing the over mentioned convergence result in the classical setting of
 101 GLSM.

3.1. A new formulation of the GLSM for symmetric factorizations.

3.1.1. Analysis of the noise free case. We denote by X and Y two (complex) reflexive Banach spaces with duals X^* and Y^* respectively and shall denote by $\langle \cdot, \cdot \rangle$ a duality product that refers to $\langle X^*, X \rangle$ or $\langle Y^*, Y \rangle$ duality. We consider the linear operator $F : X \rightarrow X^*$. Moreover we shall assume that the following factorization holds

$$(6) \quad F = H^*TH$$

where the operators $H : X \rightarrow Y$ and $T : Y \rightarrow Y^*$ are bounded. We denote by $G : \overline{\mathcal{R}(H)} \subset Y \rightarrow X^*$ the linear operator H^*T restricted to $\overline{\mathcal{R}(H)}$.

Let $\alpha > 0$ be a given parameter and $\phi \in X^*$. The new GLSM (for noise free measurements) is based on considering minimizing sequences of the functional $J_\alpha(\phi; \cdot) : X \rightarrow \mathbb{R}$

$$(7) \quad J_\alpha(\phi; g) := \alpha |\langle Fg, g \rangle| + \alpha^{1-\eta} |\langle Fg - \phi, g \rangle| + \|Fg - \phi\|^2 \quad \forall g \in X,$$

where $\eta \in]0, 1]$ is a fixed parameter. Following [3], we first observe that

$$(8) \quad j_\alpha(\phi) := \inf_{g \in X} J_\alpha(\phi; g) \rightarrow 0 \text{ as } \alpha \rightarrow 0.$$

for all $\phi \in X^*$ if one assumes that F has dense range. Indeed in this case, for a given $\varepsilon > 0$ there exists g_ε such that $\|Fg_\varepsilon - \phi\| < \frac{\varepsilon}{2}$. Then one can choose $\alpha_0(\varepsilon)$ such for all $\alpha \leq \alpha_0(\varepsilon)$, $\alpha |\langle Fg_\varepsilon, g_\varepsilon \rangle| + \alpha^{1-\eta} |\langle Fg_\varepsilon - \phi, g_\varepsilon \rangle| < \frac{\varepsilon}{2}$ so that $j_\alpha(\phi) < \varepsilon$, which proves (8). One then can prove the following characterization of the range of G in terms of F :

Theorem 1. *We assume that H is compact, G is injective and F is injective with dense range. We also assume that T satisfies the coercivity property*

$$(9) \quad |\langle Th, h \rangle| > \mu \|h\|^2 \quad \forall h \in \mathcal{R}(H),$$

where $\mu > 0$ is a constant independent of h . Consider for $\alpha > 0$ and $\phi \in X^*$, $g_\alpha \in X$ such that

$$(10) \quad J_\alpha(\phi; g_\alpha) \leq j_\alpha(\phi) + p(\alpha)$$

where $\frac{p(\alpha)}{\alpha}$ is bounded with respect to α . Then

$$\phi \in \mathcal{R}(G) \quad \text{iff} \quad \lim_{\alpha \rightarrow 0} |\langle Fg_\alpha, g_\alpha \rangle| < \infty.$$

In the case $\phi = G\varphi$, the sequence Hg_α converges strongly to φ in Y as α goes to zero.

Proof. Assume that $\phi \in \mathcal{R}(G)$ and let $\varphi \in \overline{\mathcal{R}(H)}$ such that $G\varphi = \phi$. For $\alpha > 0$ one can choose $g_0 \in X$ such that $\|Hg_0 - \varphi\|^2 < \alpha^2$. Then by continuity of G , $\|Fg_0 - \phi\|^2 < \|G\|^2 \alpha^2$. On the other hand the continuity of T implies

$$|\langle Fg_0, g_0 \rangle| = |\langle THg_0, Hg_0 \rangle| \leq \|T\| \|Hg_0\|^2 < 2 \|T\| (\alpha^2 + \|\varphi\|^2)$$

and

$$|\langle Fg_0 - \phi, g_0 \rangle| = |\langle T(Hg_0 - \varphi), Hg_0 \rangle| \leq \|T\| \|Hg_0 - \varphi\| \|Hg_0\| < 2 \|T\| \alpha (\alpha + \|\varphi\|).$$

From the definitions of $j_\alpha(\phi)$ and g_α we have

$$\alpha|\langle Fg_0, g_0 \rangle| + \alpha^{1-\eta}|\langle Fg_0 - \phi, g_0 \rangle| + \|Fg_0 - \phi\|^2 > j_\alpha(\phi) > J_\alpha(\phi, g_\alpha) - p(\alpha).$$

126 We then deduce from the definition of J_α , the fact that $\eta \in]0, 1]$ and previous inequalities

$$127 \quad (11) \quad \alpha|\langle Fg_\alpha, g_\alpha \rangle| \leq J_\alpha(\phi, g_\alpha) \leq p(\alpha) + 2\alpha\|T\|(\alpha^2 + \|\varphi\|^2) + \alpha^2\|G\|^2 + 2\|T\|\alpha^{2-\eta}(\alpha + \|\varphi\|).$$

Therefore $\limsup_{\alpha \rightarrow 0} |\langle Fg_\alpha, g_\alpha \rangle| < \infty$. The coercivity property of T implies that $\|Hg_\alpha\|^2$ is bounded. From (8) and (10) and the injectivity of G we infer that the only possible weak limit of (any subsequence of) Hg_α is φ . Thus the whole sequence Hg_α weakly converges to φ in Y . On the other hand we have that:

$$\begin{aligned} \|Hg_\alpha - \varphi\|^2 &\leq |\langle T(Hg_\alpha - \varphi), Hg_\alpha - \varphi \rangle| \\ &\leq |\langle T(Hg_\alpha - \varphi), Hg_\alpha \rangle| + |\langle T(Hg_\alpha - \varphi), \varphi \rangle| \\ &\leq |\langle Fg_\alpha - \phi, g_\alpha \rangle| + |\langle T(Hg_\alpha - \varphi), \varphi \rangle| \end{aligned}$$

The last term goes to zero due to the weak convergence of Hg_α . The first term goes to zero since the second inequality in (11) implies in particular that $|\langle Fg_\alpha - \phi, g_\alpha \rangle| \leq \alpha^\eta$. Therefore we conclude that Hg_α strongly converges to φ and consequently

$$\lim_{\alpha \rightarrow 0} |\langle Fg_\alpha, g_\alpha \rangle| = |\langle T\varphi, \varphi \rangle|.$$

128 We now consider the case $\phi \notin \mathcal{R}(G)$. Assume that $\liminf_{\alpha \rightarrow 0} |\langle Fg_\alpha, g_\alpha \rangle| < \infty$. Then, (for some
129 extracted subsequence g_α) $|\langle Fg_\alpha, g_\alpha \rangle| < A$ for some constant A independent of $\alpha \rightarrow 0$. The
130 coercivity of T implies that $\|Hg_\alpha\|$ is also bounded and therefore one can assume that, up to
131 an extracted subsequence, Hg_α weakly converges to some $\varphi \in \overline{\mathcal{R}(H)}$. Since G is compact,
132 we obtain that GHg_α strongly converges to $G\varphi$ as $\alpha \rightarrow 0$. On the other hand, (8) and the
133 definition of $J_\alpha(\phi, g_\alpha)$ imply that $\|Fg_\alpha - \phi\| \leq J_\alpha(\phi, g_\alpha) \leq j_\alpha(\phi) + C\alpha \rightarrow 0$ as $\alpha \rightarrow 0$. Since
134 $Fg_\alpha = GHg_\alpha$ we obtain that $G\varphi = \phi$ which is a contradiction. We then conclude that if
135 $\phi \notin \mathcal{R}(G)$ then $\lim_{\alpha \rightarrow 0} |\langle Fg_\alpha, g_\alpha \rangle| = \infty$. ■

136 **Remark 1.** The extension proposed in Theorem 1 requires indeed less assumptions to ensure
137 strong convergence than the one proposed in [2] for the case of symmetric factorizations.
138 However the result from [2] is still interesting for practical applications (when applicable)
139 since it uses a convex cost functional which is easier to minimize numerically.

3.1.2. Analysis for the case of noisy measurements. Let $F^\delta : X \rightarrow X^*$ be the operator associated with noisy far field measurements such that

$$\|F^\delta - F\| \leq \delta$$

140 for some $\delta > 0$. We assume that the operators F^δ and F are compact. Again let $\eta \in]0, 1]$ be
141 a fixed parameter. We define for $\alpha > 0$ and $\phi \in X^*$ the regularized functional

$$142 \quad (12) \quad J_\alpha^\delta(\phi; g) := \alpha|\langle F^\delta g, g \rangle| + \alpha^{1-\eta}|\langle F^\delta g - \phi, g \rangle| + \alpha^{1-\eta}\delta\|g\|^2 + \|F^\delta g - \phi\|^2$$

for $g \in X$. This functional has a minimizer

$$(13) \quad g_\alpha^\delta := \arg \min_{g \in X} J_\alpha^\delta(\phi; g)$$

and we also have, assuming that F has a dense range,

$$(14) \quad \lim_{\alpha \rightarrow 0} \limsup_{\delta \rightarrow 0} J_\alpha^\delta(\phi; g_\alpha^\delta) = 0.$$

The latter can be proved exactly the same way as in [3, Lemma 4] or Lemma 5 below and is based on the estimate

$$J_\alpha^\delta(\phi; g) \leq J_\alpha(\phi; g) + (\alpha\delta + \alpha^{1-\eta}\delta + \alpha^{1-\eta}\delta + \delta^2) \|g\|^2$$

and (8). We now state and prove the main result of this section.

Theorem 2. Assume that the hypothesis of Theorem 1 hold true. Let g_α^δ be the minimizer of $J_\alpha^\delta(\phi; \cdot)$ for $\alpha > 0$, $\delta > 0$ and $\phi \in X^*$. Then

$$\bullet \phi \in \mathcal{R}(G) \text{ implies } \lim_{\alpha \rightarrow 0} \limsup_{\delta \rightarrow 0} \left(|\langle F^\delta g_\alpha^\delta, g_\alpha^\delta \rangle| + \delta \alpha^{-\eta} \|g_\alpha^\delta\|^2 \right) < \infty.$$

$$\bullet \phi \notin \mathcal{R}(G) \text{ implies } \lim_{\alpha \rightarrow 0} \liminf_{\delta \rightarrow 0} \left(|\langle F^\delta g_\alpha^\delta, g_\alpha^\delta \rangle| + \delta \alpha^{-\eta} \|g_\alpha^\delta\|^2 \right) = \infty.$$

Moreover, when $\phi \in \mathcal{R}(G)$ we also have

$$\lim_{\alpha \rightarrow 0} \limsup_{\delta \rightarrow 0} \delta \|g_\alpha^\delta\|^2 = 0.$$

If $G\varphi = \phi$, then there exists $\delta_0(\alpha)$ such that for all $\delta(\alpha) \leq \delta_0(\alpha)$, $Hg_\alpha^{\delta(\alpha)}$ converges strongly to φ as α goes to zero.

Proof. The proof follows the lines of the proof of Theorem 1. Assume that $\phi = G(\varphi)$ for some $\varphi \in \overline{\mathcal{R}(H)}$. We consider g_0 (that depends on α but is independent from δ) such that $\|Hg_0 - \varphi\|^2 < \alpha^2$. Choosing δ sufficiently small such that

$$(\alpha\delta + \alpha^{1-\eta}\delta + \alpha^{1-\eta}\delta + \delta^2) \|g_0\|^2 \leq \alpha$$

we get

$$(15) \quad J_\alpha^\delta(\phi; g_\alpha^\delta) \leq J_\alpha^\delta(\phi; g_0) \leq J_\alpha(\phi; g_0) + \alpha.$$

Consequently, following the same arguments as for the second inequality in (11), we arrive at

$$\alpha \left(|\langle F^\delta g_\alpha^\delta, g_\alpha^\delta \rangle| + \alpha^{-\eta} \delta \|g_\alpha^\delta\|^2 \right) \leq J_\alpha^\delta(\phi; g_\alpha^\delta) \leq C\alpha,$$

for sufficiently small α with C a constant independent from α . This proves $\limsup_{\alpha \rightarrow 0} \limsup_{\delta \rightarrow 0} \left(|\langle F^\delta g_\alpha^\delta, g_\alpha^\delta \rangle| + \alpha^{-\eta} \delta \|F^\delta g_\alpha^\delta\|^2 \right) < \infty$. We also have, as a consequence of the inequalities above, that

$$\delta \|g_\alpha^\delta\|^2 \leq C\alpha^\eta,$$

which proves $\limsup_{\alpha \rightarrow 0} \lim_{\delta \rightarrow 0} \delta \|g_\alpha^\delta\|^2 = 0$. We also have

$$|\langle F^\delta g_\alpha^\delta - \phi, g_\alpha^\delta \rangle| \leq C\alpha^\eta$$

which proves, with the estimate on $\delta \|g_\alpha^\delta\|^2$ given above, that

$$\lim_{\alpha \rightarrow 0} \limsup_{\delta \rightarrow 0} |\langle F g_\alpha^\delta - \phi, g_\alpha^\delta \rangle| = 0.$$

We then get that $\lim_{\alpha \rightarrow 0} \limsup_{\delta \rightarrow 0} |\langle F^\delta g_\alpha^\delta, g_\alpha^\delta \rangle| < \infty$ and can conclude as in the proof of Theorem 1 that $Hg_\alpha^{\delta(\alpha)}$ converges strongly to φ as α goes to zero for $\delta(\alpha)$ sufficiently small. This also proves that $\lim_{\alpha \rightarrow 0} \limsup_{\delta \rightarrow 0} (|\langle F^\delta g_\alpha^\delta, g_\alpha^\delta \rangle| + \delta\alpha^{-\eta} \|g_\alpha^\delta\|^2) < \infty$.

Now assume that $\phi \notin \mathcal{R}(G)$ and $\liminf_{\alpha \rightarrow 0} \liminf_{\delta \rightarrow 0} (|\langle F^\delta g_\alpha^\delta, g_\alpha^\delta \rangle| + \alpha^{-\eta} \delta \|g_\alpha^\delta\|^2)$ is finite. The coercivity of T and $\alpha < 1$ implies that

$$\mu \|Hg_{\alpha(\delta)}^\delta\|^2 \leq |\langle Fg_\alpha^\delta, g_\alpha^\delta \rangle| \leq |\langle F^\delta g_\alpha^\delta, g_\alpha^\delta \rangle| + \alpha^{-\eta} \delta \|g_\alpha^\delta\|^2.$$

Therefore $\liminf_{\alpha \rightarrow 0} \liminf_{\delta \rightarrow 0} \|Hg_\alpha^\delta\|^2$ is also finite. This means the existence of a subsequence $(\alpha', \delta(\alpha'))$ such that $\alpha' \rightarrow 0$ and $\delta(\alpha') \rightarrow 0$ as $\alpha' \rightarrow 0$ and $\|Hg_{\alpha'}^{\delta(\alpha')}\|^2$ is bounded independently from α' . One can also choose $\delta(\alpha')$ such that $\delta(\alpha') \leq \alpha'^{1-\eta}$. On the other hand Equation (14) indicates that one can choose this subsequence such that $J_{\alpha'}^{\delta(\alpha')}(g_{\alpha'}^{\delta(\alpha')}) \rightarrow 0$ as $\alpha' \rightarrow 0$ and therefore $\|F^\delta g_{\alpha'}^{\delta(\alpha')} - \phi\| \rightarrow 0$ as $\alpha' \rightarrow 0$ and $\alpha'^{1-\eta} \delta(\alpha') \|g_{\alpha'}^{\delta(\alpha')}\|^2 \rightarrow 0$ as $\alpha' \rightarrow 0$. By a triangular inequality and $\delta(\alpha') \leq \alpha'^{1-\eta}$ we then deduce that $\|Fg_{\alpha'}^{\delta(\alpha')} - \phi\| \rightarrow 0$ as $\alpha' \rightarrow 0$. The compactness of G implies that a subsequence of $GHg_{\alpha'}^{\delta(\alpha')}$ converges for some $G\varphi$ in X^* . The uniqueness of the limit implies that $G\varphi = \phi$, which is a contradiction.

3.2. The GLSM for non symmetric factorizations. In this section we shall extend GLSM formalism presented in the previous section to the case of non symmetric factorisations. The general framework is given by the following assumptions. We shall denote by X_1 , X_2 and Y three (complex) reflexive Banach spaces with duals X_1^* , X_2^* and Y^* respectively and shall denote by $\langle, \rangle$ a duality product that refers to $\langle X_1^*, X_1 \rangle$, $\langle X_2^*, X_2 \rangle$ or $\langle Y^*, Y \rangle$ duality. We also set $X := X_1 \times X_2$.

We consider a linear operator $F : X_2 \rightarrow X_1^*$ that is assumed to be bounded and has the following factorization

$$(16) \quad F = U^*TV$$

where the operators $V : X_2 \rightarrow Y$, $T : Y \rightarrow Y^*$ and $U : X_1 \rightarrow Y$ are bounded. We set $G : \overline{\mathcal{R}_Y(V)} \subset Y \rightarrow X_1^*$ the restriction of U^*T to $\overline{\mathcal{R}_Y(V)}$ where $\overline{\mathcal{R}_Y(V)}$ is the closure of the

range of V in Y . We shall assume in addition the existence of a space $\hat{Y}$ such that U and V can be extended to bounded operators $V : X_2 \rightarrow \hat{Y}$ and $U : X_1 \rightarrow \hat{Y}$ such that

$$(17) \quad \|Vg_2 + Ug_1\|_Y \leq \|Vg_2 + Ug_1\|_{\hat{Y}}, \quad \forall (g_1, g_2) \in X.$$

We finally assume that

$$(18) \quad \overline{\mathcal{R}_Y(V)} = \overline{\mathcal{R}_Y(U)} \quad \text{and} \quad \overline{\mathcal{R}_{\hat{Y}}(V)} = \overline{\mathcal{R}_{\hat{Y}}(U)}.$$

A typical example is the case of limited aperture presented above with $X_2 = L^2(\Gamma_s)$, $X_1 = L^2(\Gamma_m)$, $Y = L^2(D)$ and $\hat{Y} = L^2(\Sigma)$ with Σ being any domain such that $D \subset \Sigma$. The domain Σ is assumed to be known a priori (which can coincide with the whole probed domain) and therefore the operators $V : X_2 \rightarrow \hat{Y}$ and $U : X_1 \rightarrow \hat{Y}$ are also known a priori. In the case of limited aperture presented above these operators are defined by

$$Vg(x) = \int_{\Gamma_s} e^{ikx \cdot \theta} g(\theta) ds(\theta) \quad \text{and} \quad Ug(x) = \int_{\Gamma_m} e^{ikx \cdot \theta} g(\theta) ds(\theta), \quad x \in \Sigma.$$

3.2.1. Analysis of the noise free case. Let $\alpha > 0$ be a given parameter and $\phi \in X_1^*$. We redefine the functional J_α as $J_\alpha(\phi; \cdot) : X = X_1 \times X_2 \rightarrow \mathbb{R}$

$$(19) \quad J_\alpha(\phi; g) := \alpha |\langle Fg_2, g_1 \rangle| + \alpha^{1-\eta} \|Vg_2 - Ug_1\|_Y^2 + \alpha^{1-\eta} |\langle Fg_2 - \phi, g_1 \rangle| + \|Fg_2 - \phi\|^2$$

for all $g = (g_1, g_2) \in X$ where $\eta \in]0, 1[$ is again a fixed parameter. We also set

$$(20) \quad j_\alpha(\phi) := \inf_{g \in X} J_\alpha(\phi; g).$$

Indeed the role of the extra term $\|Vg_2 - Ug_1\|$ is to formally ensure $Vg_2 \simeq Ug_1$ which cannot be done exactly since the ranges of the operators V and U are different in general. We then observe that the penalty term is of the form

$$|\langle Fg_2, g_1 \rangle| = \langle TVg_2, Ug_1 \rangle \simeq \langle TVg_2, Vg_2 \rangle$$

and therefore formally behaves as in the case of symmetric factorizations. The goal of the following analysis is to show that this is indeed asymptotically the case as $\alpha \rightarrow 0$. We first prove that with the additional penalty term, the inf still goes to 0 as $\alpha \rightarrow 0$ which guarantee that we can construct nearby solutions of the $Fg \simeq \phi$.

Lemma 3. Assume that F has dense range. Then for all $\phi \in X_1^*$, $j_\alpha(\phi) \rightarrow 0$ as $\alpha \rightarrow 0$.

Proof. Since F has dense range, for a given $\varepsilon > 0$ there exists g_2^ε such that

$$(21) \quad \|Fg_2^\varepsilon - \phi\| \leq \varepsilon/3.$$

Using (18) and (17) we can choose g_1^ε such that:

$$(22) \quad \|Vg_2^\varepsilon - Ug_1^\varepsilon\|_Y^2 < \|Vg_2^\varepsilon - Ug_1^\varepsilon\|_{\hat{Y}}^2 < \varepsilon/3$$

One then can choose α small enough such that

$$(23) \quad \alpha |\langle Fg_2^\varepsilon, g_1^\varepsilon \rangle| + \alpha^{1-\eta} |\langle Fg_2^\varepsilon - \phi, g_1^\varepsilon \rangle| \leq \varepsilon/3.$$

Together with inequalities (21) and (22) the latter inequality implies

$$j_\alpha(\phi) \leq J_\alpha(\phi; g^\varepsilon) \leq \varepsilon$$

for sufficiently small α where $g^\varepsilon := (g_1^\varepsilon, g_2^\varepsilon)$. ■

We now can state and prove the main theorem of this section that provides a characterization of the range of G in terms of F and U and V as operators with values in $\hat{Y}$.

Theorem 4. *We assume that $G : \overline{\mathcal{R}_Y(V)} \subset Y \rightarrow X_1^*$ is injective and that F has dense range. We also assume that T satisfies the coercivity property*

$$(24) \quad |\langle T\varphi, \varphi \rangle| > \mu \|\varphi\|^2 \quad \forall \varphi \in \overline{\mathcal{R}(U)} = \overline{\mathcal{R}(V)},$$

where $\mu > 0$ is a constant independent of φ . Let $p(\alpha)$ be a given function such that $\frac{p(\alpha)}{\alpha} = O(1)$ and consider for $\alpha > 0$ and $\phi \in X_1^*$, $g^\alpha = (g_1^\alpha, g_2^\alpha) \in X$ such that

$$(25) \quad J_\alpha(\phi; g^\alpha) \leq j_\alpha(\phi) + p(\alpha).$$

Then we have the following:

• $\phi \in \mathcal{R}(G)$ implies $\limsup_{\alpha \rightarrow 0} \left(|\langle Fg_2^\alpha, g_1^\alpha \rangle| + \alpha^{-\eta} \|Vg_2^\alpha - Ug_1^\alpha\|_{\hat{Y}}^2 \right) < \infty$.

• $\phi \notin \mathcal{R}(G)$ implies $\lim_{\alpha \rightarrow 0} \left(|\langle Fg_2^\alpha, g_1^\alpha \rangle| + \alpha^{-\eta} \|Vg_2^\alpha - Ug_1^\alpha\|_{\hat{Y}}^2 \right) = \infty$

In the case $\phi = G\varphi$, the two sequences Vg_2^α and Ug_1^α strongly converge to φ in Y .

Proof. The proof follows roughly the same steps and ideas as the proof for the case of symmetric factorizations. We start with the case $\phi \in \mathcal{R}(G)$. We consider $\varphi \in \overline{\mathcal{R}_Y(V)}$ such that $G\varphi = \phi$ and $h_2^\alpha \in X_2$ such that $\|Vh_2^\alpha - \varphi\|_Y^2 \leq \alpha^2$. According to (18) and (17), there exists $h_1^\alpha \in X_1$ such that:

$$(26) \quad \|Vh_2^\alpha - Uh_1^\alpha\|_Y^2 < \|Vh_2^\alpha - Uh_1^\alpha\|_{\hat{Y}}^2 < \alpha^\eta.$$

We also have

$$(27) \quad \begin{aligned} |\langle Fh_2^\alpha, h_1^\alpha \rangle| &= |\langle TVh_2^\alpha, Uh_1^\alpha \rangle| \\ &\leq |\langle TVh_2^\alpha, Vh_2^\alpha \rangle| + |\langle TVh_2^\alpha, Uh_1^\alpha - Vh_2^\alpha \rangle| \\ &\leq \|T\| \|Vh_2^\alpha\|_Y^2 + \|T\| \|Vh_2^\alpha\|_Y \sqrt{\alpha^\eta} \end{aligned}$$

and

$$\begin{aligned} |\langle Fh_2^\alpha - \phi, h_1^\alpha \rangle| &= |\langle T(Vh_2^\alpha - \varphi), Uh_1^\alpha \rangle| \leq \|T\| \|Vh_2^\alpha - \varphi\| \|Uh_1^\alpha\| \\ &< 2 \|T\| \alpha(\alpha + \|\varphi\| + \sqrt{\alpha^\eta}). \end{aligned}$$

The two previous inequalities and the definitions g^α and $j_\alpha(\phi)$ lead to

$$\alpha(|\langle Fg_2^\alpha, g_1^\alpha \rangle| + \alpha^{-\eta} \|Vg_2^\alpha - Ug_1^\alpha\|_{\hat{Y}}^2 + \alpha^{-\eta} |\langle Fh_2^\alpha - \phi, h_1^\alpha \rangle|) \leq j_\alpha(\phi) + p(\alpha) \leq C\alpha,$$

where C is bounded independently of α . This implies in particular that $(|\langle Fg_2^\alpha, g_1^\alpha \rangle| + \alpha^{-\eta} \|Vg_2^\alpha - Ug_1^\alpha\|_Y^2)$ remains bounded as $\alpha \rightarrow 0$. We also get

$$(28) \quad \|Vg_2^\alpha - Ug_1^\alpha\|_Y^2 \leq C\alpha^\eta$$

and

$$(29) \quad |\langle Fg_2^\alpha - \phi, g_1^\alpha \rangle| \leq C\alpha^\eta. \quad \blacksquare$$

We shall prove now the convergence of Vg_2^α strongly converges to φ in Y where $G\varphi = \phi$. The coercivity of T implies

$$\mu \|Vg_2^\alpha\|_Y^2 \leq |\langle TVg_2^\alpha, Vg_2^\alpha \rangle| \leq |\langle TVg_2^\alpha, Vg_2^\alpha \rangle| + |\langle TVg_2^\alpha, Ug_1^\alpha - Vg_2^\alpha \rangle| + |\langle TVg_2^\alpha, Ug_1^\alpha - Vg_2^\alpha \rangle|$$

On the one hand

$$|\langle TVg_2^\alpha, Vg_2^\alpha \rangle| + |\langle TVg_2^\alpha, Ug_1^\alpha - Vg_2^\alpha \rangle| = |\langle Fg_2^\alpha, g_1^\alpha \rangle| \leq C$$

and on the other hand

$$|\langle TVg_2^\alpha, Ug_1^\alpha - Vg_2^\alpha \rangle| \leq \|T\| \|Vg_2^\alpha\|_Y \|Vg_2^\alpha - Ug_1^\alpha\|_Y \leq \|T\| C\alpha^\eta \|Vg_2^\alpha\|_Y$$

These inequalities show that $\|Vg_2^\alpha\|_Y$ is bounded. Second, from Lemma 3 and (25) and the injectivity of G we infer that the only possible weak limit of (any subsequence of) Vg_2^α in Y is φ . Thus the whole sequence Vg_2^α weakly converges to φ in Y . Following the idea of proof of Theorem 1 we use the formula:

$$|\langle T(Vg_2^\alpha - \varphi), Vg_2^\alpha - \varphi \rangle| \leq |\langle T(Vg_2^\alpha - \varphi), \varphi \rangle| + \underbrace{|\langle T(Vg_2^\alpha - \varphi), Vg_2^\alpha - Ug_1^\alpha \rangle|}_{\leq \|T\|(\|Vg_2^\alpha\| + \|\varphi\|)\|Vg_2^\alpha - Ug_1^\alpha\|_Y} + |\langle Fg_2^\alpha - \phi, g_1^\alpha \rangle|$$

The first term on the right hand side goes to zero thanks to the weak convergence, the second term goes to zero thanks to (28) and the third term goes to zero thanks to (29). The coercivity property of T implies that Vg_2^α converges strongly to φ in Y . The strong convergence of Ug_1^α to φ in Y is a direct consequence of (28).

We now consider the case $\phi \notin \mathcal{R}(G)$ and assume that $\liminf_{\alpha \rightarrow 0} (|\langle Fg_2^\alpha, g_1^\alpha \rangle| + \alpha^{-\eta} \|Vg_2^\alpha - Ug_1^\alpha\|_Y^2) < \infty$. Then, (for some extracted subsequence g^α) $|\langle Fg_2^\alpha, g_1^\alpha \rangle| + \alpha^{-\eta} \|Vg_2^\alpha - Ug_1^\alpha\|_Y^2 \leq A$ for some A independent of α as α goes to 0. Using the same reasoning as in the first part of the theorem this implies that $\|Vg_2^\alpha\|_Y$ is bounded. We then obtain a contradiction exactly in the same way as in the proof of the second part of Theorem 1.

3.2.2. Analysis of the case of perturbed operators. We now consider the case of noisy data and/or non exact models. The noise in the data is modelled with an operator F^δ such that

$$\|F^\delta - F\| \leq \delta$$

for some $\delta > 0$. We can also assume error in the "model" by considering perturbed operators U^δ, V^δ

$$\|U^\delta - U\| \leq \delta \quad \text{and} \quad \|V^\delta - V\| \leq \delta.$$

236 The noisy operators F^δ , U^δ and V^δ are assumed to be compact. We introduce the counterpart
 237 of (19) in the noisy case (for a constant $\eta \in]0, 1[$) as

$$238 \quad (30) \quad J_\alpha^\delta(\phi; g) := \alpha \left(|\langle F^\delta g_2, g_1 \rangle| + \delta \alpha^{1-\eta} (\|g_1\|_{X_1}^2 + \|g_2\|_{X_2}^2) + \alpha^{1-\eta} |\langle F^\delta g_2 - \phi, g_1 \rangle| \right. \\ \left. + \alpha^{1-\eta} \|V^\delta g_2 - U^\delta g_1\|_{\tilde{Y}}^2 + \|F^\delta g_2 - \phi\|_{X_1^*}^2 \right)$$

for $g = (g_1, g_2) \in X$. We can also treat the case of noisy incorrect knowledge of ϕ by assume that one would consider $\phi^\delta \in X$ such that

$$\|\phi^\delta - \phi\| \leq \delta.$$

239 The analysis of the noisy case will then be mainly based on the following simple estimate

$$240 \quad (31) \quad J_\alpha^\delta(\phi^\delta; g) \leq J_\alpha(\phi; g) + n(\delta, \alpha, g),$$

241 where

$$242 \quad (32) \quad n(\delta, \alpha, g) := \delta(\alpha + \alpha^{1-\eta})(\|g_1\|_{X_1}^2 + \|g_2\|_{X_2}^2) \\ + \delta^2 \left(\|g_2\|_{X_2}^2 + \alpha^{1-\eta}(\|g_1\|_{X_1}^2 + \|g_2\|_{X_2}^2) + 1 \right).$$

Lemma 5. *For for all $\alpha, \delta > 0$ the functional $J_\alpha^\delta(\phi^\delta; \cdot)$ has a minimizer $g^{\alpha, \delta}$. Assume in addition that F has dense range. Then we have*

$$\lim_{\alpha \rightarrow 0} \lim_{\delta \rightarrow 0} J_\alpha^\delta(\phi^\delta; g^{\alpha, \delta}) = 0.$$

Proof. The existence of a minimizer is clear: for a fixed $\alpha > 0$, $\delta > 0$ and ϕ^δ , any minimizing sequence g^n of $J_\alpha^\delta(\phi^\delta; \cdot)$ is bounded and therefore there exists a weakly convergent subsequence to some $g^{\alpha, \delta}$. The lower semi-continuity of the norm with respect to the weak convergence and the compactness property of the operators then imply:

$$J_\alpha^\delta(\phi^\delta; g^{\alpha, \delta}) \leq \liminf_{n \rightarrow +\infty} J_\alpha^\delta(\phi^\delta; g^n) \leq \inf_g J_\alpha^\delta(\phi^\delta; g)$$

which proves that $g^{\alpha, \delta}$ is a minimizer of $J_\alpha^\delta(\phi; \cdot)$. Let $\epsilon > 0$ be given. We consider g^ϵ as introduced in the proof of Lemma 3 and choose δ sufficiently small ($\delta \leq \delta_0(\alpha, \epsilon)$) such that

$$n(\delta, \alpha, g^\epsilon) \leq \epsilon.$$

We then deduce from (31) and the definition of $g^{\alpha, \delta}$ that

$$J_\alpha^\delta(\phi^\delta; g^{\alpha, \delta}) \leq J_\alpha(\phi; g^\epsilon) + \epsilon$$

and conclude as in Lemma 3 that

$$J_\alpha^\delta(\phi; g^{\alpha, \delta}) \leq 2\epsilon$$

243 for sufficiently small α , which proves the second claim of the lemma. ■

We now can prove the following asymptotic characterization of the range of G (as $\delta 0$). In order to shorten the notaion we define

$$R(g, \alpha, \delta) := |\langle F^\delta g_2, g_1 \rangle| + \delta \alpha^{-\eta} (\|g_1\|_{X_1}^2 + \|g_2\|_{X_2}^2) + \alpha^{-\eta} \|V^\delta g_2 - U^\delta g_1\|_{\hat{Y}}^2.$$

244

245 **Theorem 6.** Assume that the hypothesis of Theorem 4 hold true. Let $g^{\alpha, \delta} = (g_1^{\alpha, \delta}, g_2^{\alpha, \delta})$ be
 246 the minimizer of $J_\alpha^\delta(\phi^\delta; \cdot)$. Then:

247 • $\phi \in \mathcal{R}(G)$ implies $\limsup_{\alpha \rightarrow 0} \limsup_{\delta \rightarrow 0} R(g^{\alpha, \delta}, \alpha, \delta) < \infty$.

248 • $\phi \notin \mathcal{R}(G)$ implies $\liminf_{\alpha \rightarrow 0} \liminf_{\delta \rightarrow 0} R(g^{\alpha, \delta}, \alpha, \delta) = \infty$.

249 Moreover, if $G\varphi = \phi$, then we also have

$$250 \quad (33) \quad \limsup_{\alpha \rightarrow 0} \limsup_{\delta \rightarrow 0} \delta \left(\|g_1^{\alpha, \delta}\|_{X_1}^2 + \|g_2^{\alpha, \delta}\|_{X_2}^2 \right) = 0$$

251 and there exists $\delta_0(\alpha)$ such that for all $\delta(\alpha) \leq \delta_0(\alpha)$, $Vg_2^{\alpha, \delta(\alpha)}$ and $Ug_1^{\alpha, \delta(\alpha)}$ converge strongly
 252 to φ in Y as α goes to zero.

Proof. Consider first the case $\phi \in \mathcal{R}(G)$. We shall make use of same function $h^\alpha = (h_1^\alpha, h_2^\alpha)$ as in the first part of the proof of Theorem 4 (that only depends on α). If we choose $\delta(\alpha)$ such that :

$$n(\delta(\alpha), \alpha, h^\alpha) \leq \alpha$$

(where n is defined in (32)) then we get (as in first part of the proof of Theorem 4)

$$J_\alpha^\delta(\phi; g^{\alpha, \delta}) \leq C\alpha + \alpha.$$

253 Consequently

$$254 \quad (34) \quad R(g^{\alpha, \delta}, \alpha, \delta) \leq C$$

which proves the first assertion of the theorem. We also get, as a consequence of the inequalities above, that

$$\delta \left(\|g_1^{\alpha, \delta}\|_{X_1}^2 + \|g_2^{\alpha, \delta}\|_{X_2}^2 \right) \leq C\alpha^\eta$$

255 which proves

$$256 \quad (35) \quad \lim_{\alpha \rightarrow 0} \limsup_{\delta \rightarrow 0} \delta \left(\|g_1^{\alpha, \delta}\|_{X_1}^2 + \|g_2^{\alpha, \delta}\|_{X_2}^2 \right) = 0.$$

257 For the same reasons, since $\eta > 0$ we have

$$258 \quad (36) \quad \lim_{\alpha \rightarrow 0} \limsup_{\delta \rightarrow 0} \|V^\delta g_2^{\alpha, \delta} - U^\delta g_1^{\alpha, \delta}\|_{\hat{Y}}^2 = 0.$$

Now choose $\delta_0(\alpha)$ small enough such that, $\limsup_{\alpha \rightarrow 0} n(\delta_0(\alpha), \alpha, h^\alpha) = 0$, consider $\delta(\alpha) \leq \delta_0(\alpha)$ and denote by $\tilde{g}^\alpha := g^{\alpha, \delta(\alpha)}$. Then, from (34) and (36) we clearly obtain that the quantity

$\langle TV^{\delta(\alpha)}\tilde{g}_2^\alpha, V^{\delta(\alpha)}\tilde{g}_2^\alpha \rangle$ is bounded. To conclude as in the proof of Theorem 4 that $V\tilde{g}_2^\alpha$ and $U\tilde{g}_1^\alpha$ converge strongly to φ in Y as α goes to zero, one just need to remark that

$$\|V\tilde{g}_2^\alpha - U\tilde{g}_1^\alpha\|_Y^2 \leq \|V^{\delta(\alpha)}\tilde{g}_2^\alpha - U^{\delta(\alpha)}\tilde{g}_1^\alpha\|_Y^2 + \delta(\alpha)^2 \|\tilde{g}_1^\alpha\|_{X_1}^2 + \delta(\alpha)^2 \|\tilde{g}_2^\alpha\|_{X_2}^2 \rightarrow 0$$

as $\alpha \rightarrow 0$ and

$$|\langle F\tilde{g}_2^\alpha - \phi, \tilde{g}_1^\alpha \rangle| \leq |\langle F^{\delta(\alpha)}\tilde{g}_2^\alpha - \phi, \tilde{g}_1^\alpha \rangle| + \delta(\|\tilde{g}_1^\alpha\|_{X_1}^2 + \|\tilde{g}_2^\alpha\|_{X_2}^2) \rightarrow 0.$$

as $\alpha \rightarrow 0$.

Consider now the case $\phi \notin \mathcal{R}(G)$ and assume that $\liminf_{\alpha \rightarrow 0} \liminf_{\delta \rightarrow 0} R(g^{\alpha, \delta}, \alpha, \delta)$ is finite.

Then, from

$$(37) \quad \left| \langle Fg_2^{\alpha, \delta}, g_1^{\alpha, \delta} \rangle \right| \leq \left| \langle F^{\delta}g_2^{\alpha, \delta}, g_1^{\alpha, \delta} \rangle \right| + \frac{\delta}{2} \|g_1^{\alpha, \delta}\|_{X_1}^2 + \frac{\delta}{2} \|g_2^{\alpha, \delta}\|_{X_2}^2$$

we deduce that $\left| \langle Fg_2^{\alpha, \delta}, g_1^{\alpha, \delta} \rangle \right|$ is bounded for some subsequence $\delta(\alpha)$. One also get that $\|Vg_2^{\alpha, \delta} - Ug_1^{\alpha, \delta}\|_Y^2$ is bounded for the same sequence $\delta(\alpha)$ meaning that, similarly to the second part of the proof of Theorem 4, the sequence $\|Vg_2^{\alpha, \delta(\alpha)}\|_Y$ is bounded as $\alpha \rightarrow 0$. We then can obtain a contradiction exactly the same way as in the proof of Theorem 4. ■

4. Application to inverse scattering. The purpose of this section is to apply the result of section 3 to limited aperture data (described in section 2). This will be possible if $D \subset \Sigma$ where Σ is some bounded known domain. We then define $\hat{Y}$ from section 3 to be $L^2(\Sigma)$ and set $V = H_s$ and $U = H_m$.

The basis of the GLSM is the characterization of the obstacle D in term of the range of G_m . This characterization is based on the solvability of the following interior transmission problem for $u, v \in L^2(D)$ such that $u - v \in H^2(D)$,

$$(38) \quad \begin{cases} \Delta u + k^2 n u = 0 & \text{in } D, \\ \Delta v + k^2 v = 0 & \text{in } D, \\ (u - v) = f & \text{on } \partial D, \\ \frac{\partial}{\partial \nu}(u - v) = g & \text{on } \partial D, \end{cases}$$

for a given $f \in H^{\frac{3}{2}}(\partial D)$ and $g \in H^{\frac{1}{2}}(\partial D)$. We should make the following assumption

Hypothesis 1. We assume that $k^2 \in \mathbb{R}_+$ is such that for all $f \in H^{\frac{3}{2}}(\partial D)$ and $g \in H^{\frac{1}{2}}(\partial D)$ problem (38) has a unique solution in $(u, v) \in L^2(D) \times L^2(D)$ and $u - v \in H^2(D)$.

We recall that it is known [15] that if $n - 1$ positive definite or negative definite in a neighborhood of ∂D , Hypothesis 1 is verified for all $k \in \mathbb{R}$ except a countable set without finite accumulation point.

Defining

$$\phi_z(\hat{x}) := e^{-ik\hat{x} \cdot z} \text{ for } \hat{x} \in \Gamma_m$$

we have:

Theorem 7. Under Hypothesis 1, $\phi_z \in \mathcal{R}(G_m)$ (for G_m defined in (3)) if and only if $z \in D$.

Lemma 8. $\overline{\mathcal{R}(H_s)} = \{v \in L^2(D) \text{ s.t. } \Delta v + k^2 v = 0 \text{ in } D\} = \overline{\mathcal{R}(H_m)}$

The proof of this theorem is rather straightforward using the result of Lemma 8 (see [13]) and the fact that ϕ_z is the far field of $\Phi(\cdot, z)$, the fundamental solution of the Helmholtz equation satisfying the Sommerfeld radiation condition.

The central additional theorem needed in order to apply the theory developed in Section 3 is the following coercivity property of the operator T . This theorem holds true under the following assumptions:

Hypothesis 2. We assume that $n \in L^\infty(\mathbb{R}^d)$ with $\Im(n) \geq 0$ and there exist constants $n_0, \alpha > 0$ such that $1 - \Re(n) + \alpha \Im(n) \geq 0$ for a.e. x in a neighborhood of ∂D or $\Re(n) - 1 + \alpha \Im(n) \geq 0$ for a.e. x in a neighborhood of ∂D .

The following lemma has been proved in [1].

Lemma 9. Assume that Hypothesis 2 holds and that k^2 is not a transmission eigenvalue. Then the operator T satisfies the coercivity property (9).

Let $C > 0$ be a given constant (independent of α) and consider $\alpha > 0$ and $z \in \mathbb{R}^d$, $g^{z,\alpha} = (g_1^{z,\alpha}, g_2^{z,\alpha}) \in L^2(\Gamma_m) \times L^2(\Gamma_s)$ such that :

$$\begin{aligned} J_\alpha(\phi_z, g^{z,\alpha}) &= \alpha |\langle F g_2^{z,\alpha}, g_1^{z,\alpha} \rangle| + \alpha^{1-\eta} \|H_s g_2^{z,\alpha} - H_m g_1^{z,\alpha}\|_{L^2(\Sigma)}^2 \\ &\quad + \alpha^{1-\eta} |\langle F g_2^{z,\alpha} - \phi_z, g_1^{z,\alpha} \rangle| + \|F g_2^{z,\alpha} - \phi_z\|^2 \\ &\leq j_\alpha(\phi_z) + C\alpha, \end{aligned}$$

where $\eta \in]0, 1[$ and

$$j_\alpha(\phi_z) = \inf_{g \in L^2(\Gamma_m) \times L^2(\Gamma_s)} J_\alpha(\phi_z, g).$$

Combining the results of Theorems 4 and 7 we obtain the following theorem:

Theorem 10. Assume that Hypotheses 1 and 2 hold. Then $z \in D$ if and only if $\limsup_{\alpha \rightarrow 0} |\langle F g_2^{z,\alpha}, g_1^{z,\alpha} \rangle| + \alpha^{-\eta} \|H_s g_2^{z,\alpha} - H_m g_1^{z,\alpha}\|_{L^2(\Sigma)}^2 < \infty$. Moreover, if $z \in D$ then the sequence of Herglotz wave functions associated to $g^{z,\alpha}$ converges strongly to the solution v of (38) with $(f, g) = (\Phi_z, \frac{\partial \Phi_z}{\partial \nu})$ as α goes to zero.

For applications, it is important to rather use the criterion provided in Theorem 6. Consider $F^\delta : L^2(\Gamma_s) \rightarrow L^2(\Gamma_m)$ a compact operator such that:

$$\|F^\delta - F\| \leq \delta.$$

Then consider for $\alpha > 0$ and $\phi \in L^2(\Gamma_m)$ the functional $J_\alpha^\delta(\phi, \cdot) : L^2(\Gamma_s) \times L^2(\Gamma_m) \rightarrow \mathbb{R}$,

$$\begin{aligned} J_\alpha^\delta(\phi_z, g) &= \alpha |\langle F^\delta g_2, g_1 \rangle| + \alpha^{1-\eta} \|H_s g_2 - H_m g_1\|_{L^2(\Sigma)}^2 + \alpha^{1-\eta} \delta \|g\|^2 \\ &\quad + \alpha^{1-\eta} |\langle F^\delta g_2 - \phi_z, g_1 \rangle| + \|F^\delta g_2 - \phi_z\|^2 \end{aligned}$$

where $\eta \in]0, 1[$. Then as a direct consequences of Theorem 6 we obtain the following characterization of D ,

Theorem 11. Assume that the hypothesis of Theorem 10 hold true. For $z \in \mathbb{R}^d$ denote by $g^{z,\alpha,\delta}$ the minimizer of $J_\alpha^\delta(\phi, \cdot)$ over $L^2(\Gamma_s) \times L^2(\Gamma_m)$. Then $z \in D$ if and only if

$$\limsup_{\alpha \rightarrow 0} \limsup_{\delta \rightarrow 0} |\langle F^\delta g_2^{z,\alpha,\delta}, g_1^{z,\alpha,\delta} \rangle| + \alpha^{-\eta} \|H_s g_2^{z,\alpha,\delta} - H_m g_1^{z,\alpha,\delta}\|_{L^2(\Sigma)}^2 + \alpha^{-\eta} \delta \|g^{z,\alpha,\delta}\|^2 < \infty.$$

If $z \in D$, there exists $\delta_0(\alpha)$ such that for all $\delta(\alpha) \leq \delta_0(\alpha)$, $Hg^{z,\alpha,\delta(\alpha)}$ converges strongly to the solution v of (38) with $(f, g) = (\Phi_z, \frac{\partial \Phi_z}{\partial \nu})$ as α goes to zero.

5. Extension to near field data. We concentrated in the previous sections on incident plane waves and far field measurement and raise the problem of "non symmetric factorization" in the case of limited apertures. We here show how the theory of Section 3 can be applied to other configurations of non symmetric factorization. This is the case for instance of near field data that we shall present in this section.

The total field is generated by point sources and the scattered field is recorded on a surface of $\mathbb{R}^d$ (usually where the point source lies). If we denote by $\partial\Omega$ the surface where the sources lie, we consider an incident field $u^i(y, x) := \Phi(y, x)$ with $x \in \mathbb{R}^d$ and $y \in \partial\Omega$. We introduce $N : H^{-\frac{1}{2}}(\partial\Omega) \rightarrow H^{\frac{1}{2}}(\partial\Omega)$ defined by

$$(39) \quad Ng := \int_{\partial\Omega} u^s(y, x) g(y) ds(y), \quad g \in H^{-\frac{1}{2}}(\partial\Omega), \quad x \in H^{\frac{1}{2}}(\partial\Omega),$$

where $u^s(y, \cdot) = w$ solution of (1) with an incident field $\psi = u^i(y, \cdot)$. We introduce the compact operator $S : H^{-\frac{1}{2}}(\partial\Omega) \rightarrow L^2(D)$ (which plays the role of H_s) defined by

$$(40) \quad Sg := \int_{\partial\Omega} \Phi(y, x) g(y) ds(y), \quad g \in H^{-\frac{1}{2}}(\partial\Omega), \quad x \in D$$

and the (compact) operator $G : \overline{\mathcal{R}(S)} \subset L^2(D) \rightarrow H^{\frac{1}{2}}(\partial\Omega)$ defined by

$$G\psi := w|_{\partial\Omega},$$

where $\overline{\mathcal{R}(S)}$ denotes the closure of the range of S in $L^2(D)$ and w is defined as in (1). Then clearly

$$(41) \quad N = GS.$$

In the case under consideration, since the scattered field has the following expression :

$$w(x) = - \int_D \Phi(y, x) (1 - n) k^2 (\psi(y) + w(y)) dy,$$

one simply has $G = \bar{S}^* T \psi$ where $\bar{S}^* : L^2(D) \rightarrow L^2(\partial\Omega)$ is the conjugate of the adjoint of S given by:

$$\bar{S}^* \varphi(x) = \int_D \Phi(y, x) \varphi(y) dy, \quad x \in \Gamma,$$

and T is defined by (4). Finally we get

$$(42) \quad N = \bar{S}^* T S.$$

As for the limited aperture case this factorization is "non symmetric".

5.1. Point sources and point measurements on the same surface. The case where the point sources and the measurements are on the same surface can be solved without relying on the theory developed in Section 3.2. At the cost of computing an operator C (introduced in the following) such that :

$$B = CF = H^*TH,$$

one can rely on the theory of Section 3.1 or on the theory proposed in [3]. In [13] an inf-criterion is proposed to tackle the case of near field full aperture, through the use of the corresponding far field operator. We refer to [12] for similar ideas using near field measurement. We propose to adapt this idea to the setting of the GLSM and to revisit its analysis to avoid the use of the corresponding far field operator. To do so we need to introduce the following operator, which is closely connected to S and a technical lemma.

$$(43) \quad S_{\partial\Omega} : H^{-\frac{1}{2}}(\partial\Omega) \rightarrow H^{\frac{1}{2}}(\partial\Omega), S_{\partial\Omega}(f)(x) = \int_{\partial\Omega} \Phi(x, y) f(y) ds(y), \quad x \in \partial\Omega$$

Lemma 12. *If k^2 is not a Dirichlet eigenvalue of the Laplace operator in Ω , we have that:*

$$S_{\partial\Omega}^* S_{\partial\Omega}^{-1} \bar{S}^* = S^*.$$

Proof. If k^2 is not a Dirichlet eigenvalue of the Laplace operator in Ω , for $\varphi \in H^{-\frac{1}{2}}(\partial\Omega)$ we have that $\bar{S} S_{\partial\Omega}^{-1,*} S_{\partial\Omega} \varphi$ and $S\varphi$ solves the Helmholtz equation in Ω . Straightforward calculations provide that $\bar{S}_{\partial\Omega} S_{\partial\Omega}^{-1,*} S_{\partial\Omega}^* \varphi = \bar{S}_{\partial\Omega} \varphi$ therefore the two solutions share the same boundary values on $\partial\Omega$. By taking the adjoint we conclude the proof. ■

Using (42) we arrived at

$$B = S_{\partial\Omega}^* S_{\partial\Omega}^{-1} N = S^* T S$$

From this factorization one can either use the framework developed in [3] or the factorization method developed in [13]. One can also apply the results from Section 3.1 by substituting $\langle Fg - \phi, g \rangle$ with $\langle C(Fg - \phi), g \rangle$.

5.2. Point sources and measurements lying on different surfaces. One can consider a limited aperture nearfield measurement by considering that the point sources are located on $\Gamma_s \subset \partial\Omega$ and the measurements are done on $\Gamma_m \subset \partial\Omega$ and assume that Γ_s and Γ_m are analytic surfaces. In this case similarly to the far field case we obtain a factorization :

$$(44) \quad N = \bar{S}_m^* T S_s,$$

where S_m and S_s are defined similarly to $S_{\partial\Omega}$ with $\partial\Omega$ replaced by Γ_m and Γ_s respectively. Similarly to Section 2, we define the compact operator G from $\overline{\mathcal{R}(S_s)}$ to $H^{\frac{1}{2}}(\Gamma_m)$ by $G := \bar{S}_m^* T$. As for the far field case we have the following result which is proven in [13],

Lemma 13. *If Hypothesis 1 is verified, $\Phi_z \in \mathcal{R}(G)$ if and only if $z \in D$.*

Lemma 14. *If k is not a dirichlet eigenvalue of Ω we have that S_s and S_m are dense in $\{v \in L^2(D) \text{ s.t. } \Delta v + v = 0 \text{ in } D\}$*

As already pointed out the operator T is not changed by the type of incident wave and measurement therefore it keeps the coercivity property given in section 4.

The two previous lemmas, the coercivity of T and (44) are all the required ingredients to apply the framework of section 3.2 with $V = S_s$, $U = \bar{S}_m$ and $F = N$. We therefore obtain the following corollary for the GLSM with nearfield measurements.

Corollary 15. *Assume that Hypotheses 1 and 2 hold and that $D \subset \Sigma$. Then $z \in D$ if and only if*

$|\langle N g_2^{z,\alpha}, g_1^{z,\alpha} \rangle| + \alpha^{-\eta} \|S_s g_2^{z,\alpha} - \bar{S}_m g_1^{z,\alpha}\|_{L^2(\Sigma)}^2$ remains bounded for $g_1^{z,\alpha}$ and $g_2^{z,\alpha}$ defined as in Section 3 with $\phi = \phi_z$

$R(g^{z,\alpha,\delta}, \alpha, \delta)$ (defined in Theorem 6) remains bounded for $g^{z,\alpha,\delta}$ defined as in Section 3 with $\phi = \phi_z$, $V = S_s$ and $U = \bar{S}_m$.

Moreover we have that one can extract a subsequence from the sequence of herglotz wave functions associated to $g^{z,\alpha}$ (resp. $g^{z,\alpha,\delta}$) which will converge strongly to the solution v of (38) with $(f, g) = (\Phi_z, \frac{\partial \Phi_z}{\partial \nu})$ as α goes to zero (resp. as α and δ go to zero for $\delta \leq \delta_0$).

6. Numerical Algorithm and results. In order to fix the ideas, we shall restrict ourselves in a two dimensional setting with far field measurement. We identify $\mathbb{S}^1$ with the interval $[0, 2\pi[$. In order to collect the data of the inverse problem we solve numerically (1) for N incident fields using the surface integral equation forward solver available in [11]. The discrete version of F is then the matrix F_N . We add some noise to the data to build a noisy far field matrix F_N^δ where $(F_N^\delta)_{j,k} = (F_N)_{j,k}(1 + \sigma N_{ij})$ for $\sigma > 0$ and N_{ij} an uniform complex random variable in $[-1, 1]^2$. We denote $\Phi_{z,N} \in \mathbb{C}^N$, the vector defined by $\Phi_{z,N}(j) = \phi_z(\frac{2\pi j}{N})$ for $0 \leq j \leq N-1$. In all our experiments we take $\eta = 0$ as we do not find a significant influence for this parameter.

6.1. Symmetric case. First we will look at the result given when $\Gamma_m = \Gamma_s$. This setting could be seen as a reference image as it does not introduce any new regularization term based on a priori knowledge on D (the choice of Σ). Moreover it can be formulated [3] as a convex functional if one introduces $F_\#^\delta = |\Re(F^\delta)| + |\Im(F^\delta)|$, we introduce:

$$g_\#^{z,\alpha,\delta} = \arg \min_{g \in \mathbb{C}^N} \alpha \left\| (F_\#^\delta)^{\frac{1}{2}} g \right\|^2 + \alpha^{1-\eta} \delta \|g\|^2 + \|F^\delta g - \phi_z\|^2$$

This minimization is solved using the normal equation:

$$g_\#^{z,\alpha,\delta} = (\alpha F_\# + \alpha^{1-\eta} \delta Id + F^{\delta,*} F^\delta)^{-1} F^{\delta,*} \phi_z$$

And finally we use the following indicator function to retrieve the D

$$\mathcal{I}_\#(z) = \frac{1}{\left\| (F_\#^\delta)^{\frac{1}{2}} g_\#^{z,\alpha,\delta} \right\|^2 + \alpha^{-\eta} \delta \|g_\#^{z,\alpha,\delta}\|^2}$$

To compare with setting where $\Gamma_m \neq \Gamma_s$ we also introduced :

$$g^{z,\alpha,\delta} = \arg \min_{g \in \mathbb{C}^N} \alpha |\langle F^\delta g, g \rangle| + \alpha^{1-\eta} \delta \|g\|^2 + \alpha^{1-\eta} |\langle F^\delta g - \phi_z, g \rangle| + \|F^\delta g - \phi_z\|^2$$

and consider the following indicator function:

$$\mathcal{I}(z) = \frac{1}{|\langle F^\delta g^{z,\alpha,\delta}, g^{z,\alpha,\delta} \rangle| + \alpha^{-\eta} \delta \|g^{z,\alpha,\delta}\|^2}.$$

Computing $g^{z,\alpha,\delta}$ is much more challenging as the functional is non convex nor differentiable in general. In [3], a first order gradient method is used. We here improve the efficiency of this scheme by using a second order method. We give the formula of the gradient and the hessian explicitly in the more general case where $\Gamma_m \neq \Gamma_s$. The iteration are initialized by using the original LSM [4] with Tikhonov regularization :

$$g_0^{z,\beta,\delta} = \arg \min_{g \in \mathbb{C}^N} \beta \|g\|^2 + \|F^\delta g - \phi_z\|^2$$

where we choose β such that $\delta \|g_0^{z,\beta,\delta}\| = \|F_0^{z,\beta,\delta} - \phi_z\|$. From this choice of β we set $\alpha = \frac{\beta}{\|F_\# \|}$ or $\frac{\beta}{\|F\|}$.

We consider two examples one with two ellipses and one with a kite shape obstacle both penetrable obstacle with index of refraction of 0.2. The axis are labelled as multiple of the wavelength $\lambda = 2\pi/k$. We consider three apertures $[\pi/2, 3\pi/2[$, $[3\pi/4, 5\pi/4[$ and $[7\pi/8, 9\pi/8[$ with a noise $\delta = 1\%$. In figure 2, we show the results of $\mathcal{I}_\#$ and $\mathcal{I}$.

6.2. NonSymetric case. We consider the case where $\Gamma_m \neq \Gamma_s$. In this case we have to define $g^{z,\alpha,\delta}$ as the minimizer of a (non convex nor differentiable) cost functional,

$$\begin{aligned} g^{z,\alpha,\delta} = \arg \min_{g \in \mathbb{C}^N \times \mathbb{C}^N} & \alpha |\langle F^\delta g_2, g_1 \rangle| + \alpha^{1-\eta} \delta \|g\|^2 + \alpha^{1-\eta} |\langle F^\delta g_2 - \phi_z, g_1 \rangle| \\ & + \alpha^{1-\eta} \|H_s g_2 - H_m g_1\|^2 + \|F^\delta g_2 - \phi_z\|^2, \end{aligned}$$

and we introduced the indicator function:

$$\mathcal{I}(z) = \frac{1}{|\langle F^\delta g_2^{z,\alpha,\delta}, g_1^{z,\alpha,\delta} \rangle| + \alpha^{-\eta} \delta \|g^{z,\alpha,\delta}\|^2 + \alpha^{-\eta} \|H_s g_2^{z,\alpha,\delta} - H_m g_1^{z,\alpha,\delta}\|^2}.$$

To minimize the cost functional we will rely on a second order descent method. We will choose the starting point of the descent, g_0 , as

$$g_{0,2}^{z,\beta_2,\delta} = \arg \min_{g \in \mathbb{C}^N} \beta_2 \|g\|^2 + \|F^\delta g - \phi_z\|^2$$

$$(45) \quad g_{0,1}^{z,\beta_1,\delta} = \arg \min_{g \in \mathbb{C}^N} \beta_1 \|g\|^2 + \|H_m g - H_s g_{0,2}^{z,\beta_2,\delta}\|^2$$

where we choose β_2 such that $\delta \|g_{0,2}^{z,\beta_2,\delta}\| = \|F_{0,2}^{z,\beta_2,\delta} - \phi_z\|$ and β_1 such that $\|g_{0,1}^{z,\beta_1,\delta}\| = \|g_{0,2}^{z,\beta_2,\delta}\|$. This second choice is purely arbitrary, our purpose in setting β_1 is to avoid, $g_{0,1}^{z,\beta_1,\delta}$ to have large norm which would dominate numerically all other quantities.

The minimization of J_α^δ causes numerical problem. Indeed first numerically H_m is a compact operator and even if it is not important for the theory we are implicitly inverting it by minimizing J_α^δ therefore we have to be careful on the balance between the terms $\|g_1\|^2$ and $\|H_m g_1 - H_s g_2\|$. This is even more important as $H_s g_2$ is not in the range of H_m . Since the theory does not give a strategy to set α , we proposed and tested three strategies that give similar results. Those strategies are based on the idea (we also use to pick an initial guess g_0) that $\|g_1^{z, \beta_1, \delta}\|$ and $\|g_2^{z, \beta_1, \delta}\|$ should have the same order of magnitude.

First one should remark that we have used the same parameter, η in front of all the terms but it could have been chosen with a different value for each term (as long as it stays between 0 and 1 for the theory). Using different α instead of η to keep simple notation we introduce:

$$J_\alpha^\delta(g_1, g_2) = \alpha |\langle F^\delta g_2, g_1 \rangle| + \alpha_1 \delta \|g_1\|^2 + \alpha_2 \delta \|g_2\|^2 + \alpha^{1-\eta} |\langle F^\delta g_2 - \phi_z, g_1 \rangle| + \alpha_3 \|H_s g_2 - H_m g_1\|^2 + \|F^\delta g_2 - \phi_z\|^2$$

We have actually increase the number of parameter in order to get some freedom to balance the term involving g_1 and $H_m g_1$. To set α we use again our heuristic: $\alpha = \frac{\beta_2}{\|F\|}$. We propose to choose $\alpha_1 = \alpha_2 = \alpha$ and $\alpha_3 = \alpha \delta / \beta_1$ and therefore keep the regularizing power used to find the initial guess. The parameters set, we used a newton method to minimize J_α^δ .

A second solution we have experienced is to alternatively minimize J_α^δ as a function of g_2 with $\alpha_3 = \alpha_1 = 0$ and to minimize the same Tikhonov functional (45) we used to find the initial guess $g_{0,1}$. This will impose $\|g_1^{z, \beta_1, \delta}\| = \|g_2^{z, \beta_2, \delta}\|$ and limit the number of parameters to set, however it is not a scheme that is cover by the theory.

A third solution closely related to our heuristic for symmetric factorization, we have set α_3 to 1 and $\alpha_1 = \alpha_2 = \alpha$ where α is chosen to be equal to $\max(\beta_1, \beta_2) / \|F^\delta\|$.

All those three methods give similar result. In the following we will show only the results of the first method. In order to perform the Newton method we need to compute the gradient and the Hessian which we explicit in the following for the original cost functional, both gradient and Hessian can be easily derive from those formulas. If $\cdot$ is the dot product without conjugate, t the transposition and by * the classical transpose-conjugate, we can rewrite $J_\alpha^\delta(\phi, \cdot)$:

$$\alpha |g^* \cdot (F^\delta g)| + \alpha^{1-\eta} |g^* \cdot (F^\delta g - \phi)| + \delta \alpha^{1-\eta} \|F^\delta\| |g^* \cdot g| + \alpha^{1-\eta} (Hg)^* \cdot (Hg) + (F^\delta g - \phi)^* \cdot (F^\delta g - \phi)$$

where we use the matrix:

$$F^\delta = \begin{bmatrix} 0 & F_N^\delta \\ 0 & 0 \end{bmatrix} \text{ and } H = \begin{bmatrix} H_m & -H_s \end{bmatrix} \text{ and } g = \begin{bmatrix} g_2 \\ g_1 \end{bmatrix} \text{ and } \phi = \begin{bmatrix} \Phi_z \\ 0 \end{bmatrix}$$

Using this notation we can compute following the framework of [14] the gradient

$$\delta \alpha^{1-\eta} \|F^\delta\| |g + F^{\delta*} (F^\delta g - \phi) + \alpha^{1-\eta} H^* H g| + \alpha \frac{\overline{g^* \cdot (F^\delta g)} F^\delta g + (g^* \cdot (F^\delta g)) F^{\delta*} g}{|g^* \cdot (F^\delta g)|} + \alpha^{1-\eta} \frac{(g^* \cdot (F^\delta g - \phi)) (F^\delta g - \phi) + (g^* \cdot (F^\delta g - \phi)) F^{\delta*} g}{|g^* \cdot (F^\delta g - \phi)|}$$

and the Hessian,

$$\begin{aligned} & \delta\alpha^{1-\eta} \left\| F^\delta \right\| \left\| Id + F^{\delta*} F^\delta + \alpha^{1-\eta} H^* H + \alpha \frac{\overline{g^* \cdot (F^\delta g)} F^\delta + (g^* \cdot (F^\delta g)) F^{\delta*} + F^\delta g g^* F^{\delta*} + F^{\delta*} g g^* F^\delta}{|g^* \cdot (F^\delta g)|} \right. \\ & - \alpha \frac{\overline{(g^* \cdot (F^\delta g)) F^\delta g + (g^* \cdot (F^\delta g)) F^{\delta*} g} (g^* \cdot (F^\delta g)) F^\delta g + (g^* \cdot (F^\delta g)) F^{\delta*} g)^*}{2|g^* \cdot (F^\delta g)|^{\frac{3}{2}}} \\ & + \alpha^{1-\eta} \left(\frac{\overline{(g^* \cdot (F^\delta g - \phi)) F^\delta} + (g^* \cdot (F^\delta g - \phi)) F^{\delta*} + (F^\delta g - \phi)(g^* F^{\delta*} - \phi^*) + F^{\delta*} g g^* F^\delta}{|g^* \cdot (F^\delta g - \phi)|} \right. \\ & \left. - \frac{((g^* \cdot (F^\delta g - \phi))(F^\delta g - \phi) + (g^* \cdot (F^\delta g - \phi)) F^{\delta*} g)((g^* \cdot (F^\delta g - \phi))(F^\delta g - \phi) + (g^* \cdot (F^\delta g - \phi)) F^{\delta*} g)^*}{2|g^* \cdot (F^\delta g - \phi)|^{\frac{3}{2}}} \right). \end{aligned}$$

We apply those techniques to the case of back scattering data which is when $\Gamma_m = -\Gamma_s$, for apertures $\Gamma_s = [\pi/2, 3\pi/2[$, $[3\pi/4, 5\pi/4[$ and $[7\pi/8, 9\pi/8[$. The result are shown in figure 3 for the kite example for a domain Σ which occupies the whole image and the smallest rectangle that contains D . We also consider the case of Γ_s being either $[\pi/2, 3\pi/2[$, $[3\pi/4, 5\pi/4[$ and $[7\pi/8, 9\pi/8[$ and Γ_m being either $[0, \pi[$, $[\pi/4, 3\pi/4[$ and $[3\pi/8, 5\pi/8[$. The results are shown in figure 4, again for a kite example and an original setting of sources and measurements. On those simulation the size of Σ has no clear impact therefore we will only show simulation for the large grid.

Figures 5 and 6 consider backscattering data from aperture of the same size as previously, but rotated around the obstacle. We see the strong dependency with the mean direction of the aperture. The fact that the results are coherent with the aperture we consider lets us think that non symmetric aperture is intrinsically worst than symmetric one. Connected to that subject in [9] they study invisibility for a finite number of incident direction and demonstrate that imposing invisibility in symmetric direction is equivalent to impose invisibility in all direction. Meaning that there is more information inside symmetric-factorization like far field operator than any other setting of sources and measurements.

REFERENCES

- [1] Lorenzo Audibert. Sampling method for sign changing contrast. 2016. Preprint.
- [2] Lorenzo Audibert, Alexandre Girard, and Houssem Haddar. Identifying defects in an unknown background using differential measurements. *Inverse Problems and Imaging*, 9(3):625–643, 2015.
- [3] Lorenzo Audibert and Houssem Haddar. A generalized formulation of the linear sampling method with exact characterization of targets in terms of farfield measurements. *Inverse Problems*, 30(3):035011, 2014.
- [4] Fioralba Cakoni and David Colton. *Qualitative methods in inverse scattering theory*. Interaction of Mechanics and Mathematics. Springer-Verlag, Berlin, 2006. An introduction.
- [5] Fioralba Cakoni and Isaac Harris. The factorization method for a defective region in an anisotropic material. *Inverse Problems*, 31(2):025002, 22, 2015.
- [6] David Colton, Houssem Haddar, and Michele Piana. The linear sampling method in inverse electromagnetic scattering theory. *Inverse Problems*, 19(6):S105–S137, 2003. Special section on imaging.
- [7] David Colton and Andreas Kirsch. A simple method for solving inverse scattering problems in the resonance region. *Inverse Problems*, 12(4):383–393, 1996.
- [8] David Colton and Rainer Kress. *Inverse acoustic and electromagnetic scattering theory*, volume 93 of *Applied Mathematical Sciences*. Springer, New York, third edition, 2013.
- [9] Anne-Sophie Bonnet-Ben Dhia, Lucas Chesnel, and Sergei A Nazarov. Non-scattering wavenumbers and far field invisibility for a finite set of incident/scattering directions. *Inverse Problems*, 31(4):045006, 2015.
- [10] Y. Grisel, V. Mouysset, P.-A. Mazet, and J.-P. Raymond. Determining the shape of defects in non-

- absorbing inhomogeneous media from far-field measurements. *Inverse Problems*, 28(5):055003, 19, 2012.
- [11] H. Haddar. Sampling 2d, Mars 2013. <http://sourceforge.net/projects/samplings-2d/>.
- [12] Guanghui Hu, Jiaqing Yang, Bo Zhang, and Haiwen Zhang. Near-field imaging of scattering obstacles with the factorization method. *Inverse Problems*, 30(9):095005, 25, 2014.
- [13] Andreas Kirsch and Natalia Grinberg. *The factorization method for inverse problems*, volume 36 of *Oxford Lecture Series in Mathematics and its Applications*. Oxford University Press, Oxford, 2008.
- [14] L. Sorber, M. Barel, and L. Lathauwer. Unconstrained optimization of real functions in complex variables. *SIAM J. Optim.*, 22(3):879–898, 2012.
- [15] John Sylvester. Discreteness of transmission eigenvalues via upper triangular compact operators. *SIAM J. Math. Anal.*, 44(1):341–354, 2012.

Figure 2. On the left $\mathcal{I}_\#$ and on the right $\mathcal{I}$. From up to down the aperture is : $[\pi/2, 3\pi/2[$, $[3\pi/4, 5\pi/4[$ and $[7\pi/8, 9\pi/8[$ (as depicted on the right column).

Figure 3. $\mathcal{I}$ computed on the left with a large Σ and with on the right with a small one. From up to down the apertures are : $\Gamma_s = [\pi/2, 3\pi/2[$, $[3\pi/4, 5\pi/4[$ and $[7\pi/8, 9\pi/8[$ (as depicted in the right column).

Figure 4. $\mathcal{I}$ computed on the left with a large Σ and with on the right with a small one. From up to down the apertures are $\Gamma_s = [\pi/2, 3\pi/2[$, $[3\pi/4, 5\pi/4[$ and $[7\pi/8, 9\pi/8[$ and $\Gamma_m = [0, \pi[$, $[\pi/4, 3\pi/4[$ and $[3\pi/8, 5\pi/8[$ (as depicted in the right column).

Figure 5. $\mathcal{I}$ computed with Σ equals the full grid. From left to right and up to down the aperture are : $\Gamma_s[3\pi/4, 7\pi/4[$, $[\pi, 2\pi[$ and $[-\pi/2, \pi/2[$ and $\Gamma_m = \Gamma_s + \pi$ (the sensor setting are depicted following the same order in the last image).

Figure 6. $\mathcal{I}$ computed with Σ equals the full grid. From left to right and up to down the aperture are : $\Gamma_s = [3\pi/4, 7\pi/4[$, $[\pi, 2\pi[$ and $[-\pi/2, \pi/2[$ and $\Gamma_m = \Gamma_s + \pi$ (the sensor setting are depicted following the same order in the last image).