



Projected pH reductions by 2100 might put deep North Atlantic biodiversity at risk

Marion Gehlen, Roland S  f  rian, D. O. B. Jones, Tilla Roy, R. Roth, J. Barry, Laurent Bopp, Scott C. Doney, John P. Dunne, Christoph Heinze, et al.

► To cite this version:

Marion Gehlen, Roland S  f  rian, D. O. B. Jones, Tilla Roy, R. Roth, et al.. Projected pH reductions by 2100 might put deep North Atlantic biodiversity at risk. *Biogeosciences*, 2014, 11 (23), pp.6955 - 6967. 10.5194/bg-11-6955-2014 . hal-01436121

HAL Id: hal-01436121

<https://hal.science/hal-01436121>

Submitted on 27 Oct 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin  e au d  p  t et    la diffusion de documents scientifiques de niveau recherche, publi  s ou non,   manant des   tablissements d'enseignement et de recherche fran  ais ou   trangers, des laboratoires publics ou priv  s.



Projected pH reductions by 2100 might put deep North Atlantic biodiversity at risk

M. Gehlen¹, R. Séférian², D. O. B. Jones³, T. Roy⁴, R. Roth⁵, J. Barry⁶, L. Bopp¹, S. C. Doney⁷, J. P. Dunne⁸, C. Heinze^{9,11,12}, F. Joos⁵, J. C. Orr¹, L. Resplandy¹, J. Segschneider¹⁰, and J. Tjiputra^{11,12}

¹LSCE/IPSL, Laboratoire des Sciences du Climat et de l'Environnement, Orme des Merisiers, CEA/Saclay 91198 Gif-sur-Yvette Cedex, France

²CNRM-GAME, Centre National de Recherche Météorologique-Groupe d'Etude de l'Atmosphère Météorologique, Météo-France/CNRS, 42 Avenue Gaspard Coriolis, 31100 Toulouse, France

³National Oceanography Centre, University of Southampton Waterfront Campus, European Way, Southampton, SO14 3ZH, UK

⁴LOCEAN/IPSL, 4, place Jussieu 75252 PARIS Cedex 05, France

⁵Climate and Environmental Physics, Physics Institute and Oeschger Centre for Climate Change Research, University of Bern, 3012 Bern, Switzerland

⁶Monterey Bay Aquarium Research Institute, Moss Landing, CA 95039, USA

⁷Marine Chemistry and Geochemistry, Woods Hole Oceanographic Institution, Woods Hole, MA 02543, USA

⁸National Oceanic and Atmospheric Administration/Geophysical Fluid Dynamics Laboratory, Princeton, NJ 08540, USA

⁹Geophysical Institute, University of Bergen and Bjerknes Centre for Climate Research, 5007 Bergen, Norway

¹⁰Max Planck Institute for Meteorology, Bundesstr. 53, 20146 Hamburg, Germany

¹¹Uni Research Climate, Bergen, Norway

¹²Bjerknes Centre for Climate Research, Bergen, Norway

Correspondence to: M. Gehlen (marion.gehlen@lsce.ipsl.fr)

Received: 6 May 2014 – Published in Biogeosciences Discuss.: 11 June 2014

Revised: 28 October 2014 – Accepted: 3 November 2014 – Published: 11 December 2014

Abstract. This study aims to evaluate the potential for impacts of ocean acidification on North Atlantic deep-sea ecosystems in response to IPCC AR5 Representative Concentration Pathways (RCPs). Deep-sea biota is likely highly vulnerable to changes in seawater chemistry and sensitive to moderate excursions in pH. Here we show, from seven fully coupled Earth system models, that for three out of four RCPs over 17 % of the seafloor area below 500 m depth in the North Atlantic sector will experience pH reductions exceeding -0.2 units by 2100. Increased stratification in response to climate change partially alleviates the impact of ocean acidification on deep benthic environments. We report on major pH reductions over the deep North Atlantic seafloor (depth > 500 m) and at important deep-sea features, such as seamounts and canyons. By 2100, and under the high CO_2 scenario RCP8.5, pH reductions exceeding -0.2 (-0.3) units

are projected in close to 23 % (~ 15 %) of North Atlantic deep-sea canyons and ~ 8 % (3 %) of seamounts – including seamounts proposed as sites of marine protected areas. The spatial pattern of impacts reflects the depth of the pH perturbation and does not scale linearly with atmospheric CO_2 concentration. Impacts may cause negative changes of the same magnitude or exceeding the current target of 10 % of preservation of marine biomes set by the convention on biological diversity, implying that ocean acidification may offset benefits from conservation/management strategies relying on the regulation of resource exploitation.

1 Introduction

Global ocean anthropogenic carbon inventories suggest that the ocean took up a cumulative $\sim 155 \pm 31$ Pg C (10^{15} g of carbon) in 2010 (Khaliwala et al., 2013). This uptake of CO₂ is causing profound changes in seawater chemistry resulting from increased hydrogen ion concentration (decrease in pH, $\text{pH} = -\log_{10}[\text{H}^+]$), referred to as ocean acidification (IPCC, 2011). Experimental and modelling studies provide compelling evidence that ocean acidification will put marine ecosystems at risk (e.g. Orr et al., 2005; Kroeker et al., 2013). However, with the exception of assessments focusing on cold-water coral systems (Barry et al., 2005, 2013; Fleeger et al., 2006; Guinotte et al., 2006; Tittensor et al., 2010), quantifications of biological consequences of ocean acidification mostly targeted surface ocean or coastal environments (Kroeker et al., 2010). The aim of this study is to extend our understanding of broad scale impacts of ocean acidification from the existing shallow water studies to focus specifically on deep-sea ecosystems. The deep sea is under increasing anthropogenic pressure as technological advances allow for exploitation of formerly inaccessible regions (Clauss and Hoog, 2002). While waste disposal, fishing and, in the future, mineral extraction are well recognized as dominant human pressures (Ramirez-Llodra et al., 2011), expert assessments urge consideration of climate change and ocean acidification impacts in future ecosystem conservation/management strategies (Taranto et al., 2012; Billé et al., 2013).

While previous studies quantified changes in carbonate mineral saturation state as a measure for potential detrimental impacts on deep calcifying communities (Guinotte et al., 2005, 2006; Turley et al., 2007; Fautin et al., 2009), this model-based assessment uses pH. The tight control of pH at the cellular scale is an important prerequisite of proper cell functioning, and mechanisms of pH control are ubiquitous across many taxa (Seibel and Walsh, 2003, and references therein). Deep-sea organisms might be particularly vulnerable to changes in seawater chemistry, at least in part owing to limitations on rate processes, caused by low temperature (Childress, 1995; Seibel and Walsh, 2001) and possibly food availability (Ramirez-Llodra, 2002), as well as the environmental stability of their habitat in the past (Barry et al., 2011; Seibel and Walsh, 2003). A recent review by Somero (2012) highlights the link between environmental stability and the capacity to acclimate to future changes in environmental variables such as pH. According to this study, environmental stability might impair the potential for acclimation. This stands in sharp contrast to shallow water or intertidal organisms, which are adapted to a dynamic environment with large changes in temperature and seawater chemistry (Hofmann et al., 2011; Duarte et al., 2013).

A model sensitivity study by Gehlen et al. (2008) suggested the potential for large pH reductions (up to -0.6 pH units) in the deep North Atlantic. Regions of large pH reductions coincided with areas of deep-water formation. Deep-

water formation drives the rapid propagation of surface-derived changes in carbonate chemistry to depth as underlined by high vertically integrated water column inventories of anthropogenic carbon (Sabine et al., 2004), as well as tritium and chlorofluorocarbon distributions (Doney and Jenkins, 1994). Gehlen et al. (2008) used output from a single model and for a scenario following an atmospheric CO₂ increase of 1 % per year over 140 years starting from an atmospheric CO₂ level of 286 ppm. This rate of increase is about twice as large as the rate typical for a high-end IPCC concentration pathway. The study did not include circulation changes in response to climate change.

Here we extend the study by Gehlen et al. (2008) by analysing pH projections from seven Earth system models that contributed to the Coupled Model Intercomparison Project Phase 5 (CMIP5) and for four different Representative Concentration Pathways (RCPs; Van Vuuren et al., 2011) ranging from a strong emission mitigation scenario (RCP2.6) to the high-CO₂ scenario RCP8.5. We assess the magnitude of deep-water pH reductions in the North Atlantic (35–75° N, 0–90° W) over this century in response to atmospheric CO₂ increase and climate change. The North Atlantic is a well-ventilated region of the world ocean and, despite a projected increase in stratification, will remain well oxygenated in the future (Bopp et al., 2013). The study complements assessments by Bopp et al. (2013) and Mora et al. (2013), which evaluated large-scale average pH reductions in response to the same RCP pathways, but without a detailed discussion of spatial patterns and their link to circulation. We define a critical threshold for pH reductions based on evidence from palaeo-oceanographic studies, contemporary observations and model results. Future multi-model projections of pH changes over the seafloor are analysed with reference to this threshold and without discrimination of particular habitats first. Next, model results are put into the perspective of ecosystem conservation by evaluating changes in pH against the distribution of seamounts and deep-sea canyons. These features are known as sites of high-biodiversity deep-sea ecosystems, such as cold-water corals and sponge communities (ICES, 2007; Clark et al., 2010; De Leo et al., 2010) and are selected as representative examples of deep-sea environments.

2 Material and methods

2.1 Earth system models

Our study draws on results from two types of Earth system models: (1) the Bern3-D-LPJ carbon-cycle/climate model (Steinacher et al., 2013; Roth and Joos, 2013) and (2) seven fully coupled three-dimensional atmosphere ocean climate models that participated in the Coupled Model Intercomparison Project Phase 5 (CMIP5; Taylor et al., 2011) and contributed to the Fifth Assessment Report of the Inter-

governmental Panel on Climate Change (IPCC AR5). The Bern3-D-LPJ is a model of intermediate complexity featuring a 3-D geostrophic-balance ocean and 2-D atmospheric energy and moisture-balance model. The cycle of carbon and related tracers is represented including prognostic formulations for marine production, a seafloor sediment and a dynamic global vegetation model. This model is relatively cost-efficient compared to CMIP5 models. It is used to evaluate the order of magnitude of pH reductions associated with past abrupt climate change by analysing results from freshwater hosing experiments (Bryan, 1986; Marchal et al., 1999; Matsumoto and Yokoyama, 2013).

Concerning the subset of CMIP5 models, we selected models for which 3-D pH fields were available and that had been part of a published multi-model evaluation (Bopp et al., 2013). We analyse output for four future atmospheric CO₂ concentration scenarios (Representative Concentration Pathways, RCPs), along with the corresponding pre-industrial control simulations, piControl. The nomenclature follows CMIP5 recommendations. Historical simulations cover the period between 1870 and 2005 and are followed by climate change scenarios according to RCP8.5, RCP6.0, RCP4.5 and RCP2.6 from 2006 to 2100 (Van Vuuren et al., 2011; Moss et al., 2010). RCP identifiers refer to the additional radiative forcing in 2100 relative to pre-industrial (or before 2100 in case of the peak-and-decline scenario RCP2.6). These additional radiative forcings correspond to atmospheric CO₂ levels in 2100 of 421, 538, 670 and 936 ppm for RCP2.6, RCP4.5, RCP6.0 and RCP8.5. Individual RCPs differ with respect to the temporal evolution of atmospheric CO₂ and range from a stringent emission mitigation RCP2.6 to the high-CO₂ scenario RCP8.5. The complete set of RCPs was not available for all models. Please refer to Table S1 (Supplement) for model name, scenario and references.

2.2 Deep-sea ecosystems

This study uses data sets of seamounts (Yesson et al., 2011) and canyons (Harris and Whiteway, 2011). For seamounts, these data include location, height and surface assuming a conical shape. For canyons, the data consist of a high-resolution vector database of canyon centre lines that was converted into a raster data set of canyon presence (using ArcGIS v10) for analysis. Data were projected on a 1° × 1° regular grid.

2.3 Post-treatment of model output and data

2.3.1 Post-treatment of CMIP5 model output

Model output is interpolated on a regular grid of 1° × 1° resolution. Anomalies are computed as the difference between the decade 2090–2099 and the long-term mean of the pre-industrial state. As the focus of this study is on impacts on benthic communities, we quantify pH changes in the deepest

model box over a topography range from 500 to > 4500 m water depth.

2.3.2 Computation of the area of seamounts for impact assessment

The area of North Atlantic seafloor impacted by ocean acidification is estimated on the basis of individual grid cells for which the reduction in pH exceeded ≥ 0.2 or 0.3 units. The impacted area follows as the integral of the area of these 1° × 1° grid cells. The area of seamounts with a pH reduction ≥ 0.2 or 0.3 units is computed based on distribution and height assuming a conical shape (Danovaro et al., 2008; Yesson et al., 2011). The database provides height above seafloor and base area. The area of the seamount (A) is given by

$$A = \pi r \sqrt{r^2 + (h + h')^2}, \quad (1)$$

where r is the base radius of the seamount and $h + h'$ is the height. The height impacted by a pH reduction exceeding the threshold (h') is diagnosed from the depth of the pH anomaly corresponding to the threshold. The radius of the seamount at the depth of the anomaly (r') is obtained from the Thales theorem:

$$\frac{r'}{r} = \frac{h'}{h}, \quad (2)$$

$$\text{as } r' = \frac{h'}{h} r. \quad (3)$$

The final expression of r' is the positive analytical solution of the fourth-order polynomial

$$\frac{A^2}{\pi^2} = \frac{h^2}{h'^2} r'^2 \left(\frac{h^2}{h'^2} r'^2 + (h + h')^2 \right), \quad (4)$$

$$\text{as } r' = \pm \frac{h'}{h} \left[\frac{1}{2} \left(-1 \pm \sqrt{\frac{4A^2}{\pi^2 h^2}} \right) \right]^{\frac{1}{2}}. \quad (5)$$

The impacted area of the seamount (A^*) follows from the depth of pH anomaly as a function of seamount height:

$$A^* = \pi \frac{h'}{h} \left[\frac{1}{2} \left(-1 + \sqrt{\frac{4A^2}{\pi^2 h^2}} \right) \right]^{\frac{1}{2}} \left(h'^2 + \frac{h'^2}{h^2} \left[\frac{1}{2} \left(-1 + \sqrt{\frac{4A^2}{\pi^2 h^2}} \right) \right] \right), \quad (6)$$

where A is the total surface area of the seamount.




Figure 1. North Atlantic freshwater hosing experiment. **(a)** Time series of strength of Atlantic Meridional Overturning Circulation (Sv); freshwater release occurred over 300 years (grey bar). **(b)** Times series of pH change relative to pre-industrial averaged over the deep (below 2000 m) North Atlantic (45–65° N). **(c)** Spatial distribution of the pH reduction averaged over experiment years 400–450 (grey bar in panel **b**) in terms of pH anomalies relative to pre-industrial at the seafloor and **(d)** in a section through the Atlantic at 38.5° W.

3 Results and discussion

3.1 Environmental stability and critical threshold for pH reduction

Considering that environmental stability might impair the potential for acclimation, we assessed pH changes over glacial–interglacial timescales and for past events of rapid climate changes recognized for having driven major reorganizations in North Atlantic circulation and carbonate chemistry.

The pH is defined as the negative logarithm of the hydrogen ion concentration ($[H^+]$). From the basic properties of logarithms it follows that the difference in pH equals the logarithm of the ratio of hydrogen ion concentrations. For a given pH change, the change in $[H^+]$, $\Delta[H^+]$, is a linear function of the initial hydrogen ion concentration ($[H^+]_i$) as $\Delta[H^+] = [H^+]_i \cdot ((1/10^{\Delta pH}) - 1)$. Hence, the larger the initial $[H^+]$, the larger the perturbation (Supplement Fig. S1). Contrasting shallow and deep environments highlights that absolute changes in $[H^+]$ are amplified at depth for any threshold, that is for environments of low natural variability.

3.1.1 Glacial–interglacial timescales

The palaeo-record permits evaluation of environmental variability of the deep ocean over the past million years. Avail-

able evidence indicates a low variability over this time interval (Elderfield et al., 2012; Yu et al., 2010, 2013). Changes in carbonate chemistry were small in the deep ocean compared to surface layers (Yu et al., 2010). Recent studies re-evaluated deep-water pH changes between glacial and present (Sanyal et al., 1995), arguing that carbonate compensation kept deep-water pH close to constant (Hönisch et al., 2008). We use data available in Yu et al. (2010) (and associated supplement) and follow their reasoning to infer dissolved inorganic carbon (DIC) changes from $[CO_3^{2-}]$ and hence alkalinity in order to compute associated changes in pH for sediment core BOFS 8K (52.5° N, 22.1° W, 4045 m). This pH change is computed using CO₂sys (<http://cdiac.ornl.gov/oceans/co2rprt.html>) with alkalinity and DIC as input variables, along with temperature, depth, phosphate and silicate (Yu et al., 2010). We estimate a pH reduction of ~0.1 pH units for North Atlantic deep water over the early deglacial (17 500 to 14 500 years before present).

3.1.2 Rapid events associated with freshwater release: Heinrich and Dansgaard Oeschger events

Model experiments yield maximum pH reductions in North Atlantic deep water below 0.15 pH units in response to a shutdown of the North Atlantic Meridional Overturning Circulation (AMOC, Fig. 1). To realize an abrupt shutdown of

the AMOC, different durations of freshwater perturbations in the North Atlantic on top of a pre-industrial steady state have been tested releasing a total of $3 \times 10^{15} \text{ m}^3$ freshwater ($\sim 9 \text{ m}$ sea level equivalent). In terms of pH changes in the North Atlantic region, the experiment with a 300-year freshwater forcing of 0.33 Sv results in the strongest response (Fig. 1a). In these experiments, the cause of the pH decrease is not high atmospheric CO_2 (CO_2 only increases a few ppm during the freshwater experiment), but is mainly a result of the decrease in deep-ocean ventilation. This leads to the additional accumulation of DIC by the respiration of organic matter. Although alkalinity is also increased in the deep by the dissolution of carbonate particles settling through the water column, it does not compensate for the increase in DIC leading to more acidic waters in the deep. The most extreme negative excursion of the pH averaged over the deep (below 2000 m) North Atlantic ($45\text{--}65^\circ \text{ N}$) occurs ~ 150 years after the end of the freshwater forcing with a decrease of $\sim 0.13 \text{ pH}$ units relative to the unperturbed pre-industrial state (Fig. 1b). The pH decrease does not exceed -0.18 pH units in any of the individual grid boxes. In Fig. 1c and d the spatial distribution of the pH-reduction averaged over years 400–450 (i.e. during the maximum of the pH decrease) is shown in terms of pH anomalies at the seafloor and in a section through the Atlantic at 38.5° W .

3.1.3 Critical threshold for pH reductions

For the purpose of evaluating the potential for negative impacts on deep-sea benthic environments, a critical threshold for pH reduction needs to be identified. Reductions of pH exceeding the envelope set by past and present natural variability are considered as critical. Palaeo-evidence suggests that the deep-sea fauna has evolved under conditions of environmental variability confined to a narrow range over the past million years (Yu et al., 2010; Elderfield et al., 2012). Many past episodes of climate change occurred over significantly longer timescales than the current anthropogenic perturbation of the climate system, allowing carbonate compensation to keep deep-water pH close to constant (Hönisch et al., 2008). This is corroborated by computing pH reduction over glacial–interglacial cycles for a North Atlantic site. Decadal-to-centennial changes are addressed by freshwater hosing model experiments to simulate effects of circulation changes associated with rapid Heinrich and Dansgaard Oeschger events. In both cases, pH reductions are below 0.15 pH units. Similarly, small amplitude natural temporal pH variability at depth emerges from a multi-annual time series station (González-Dávila et al., 2010) and the analysis of the long pre-industrial simulation “piControl” (Fig. S2). In summary, natural pH variations on multi-annual, decadal-to-century, and longer timescales were likely smaller than 0.2 pH units on the regional-to-basin scale in the deep Atlantic and at least for the past million years.

This suggests that pH variations of up to 0.1 to 0.2 pH units do not present a risk for marine life.

This leads us to define two thresholds for pH reduction between pre-industrial and the end of the 21st century: -0.2 and -0.3 pH units. Both stand for pH reductions exceeding palaeo-record-based estimates of changes in North Atlantic deep-water chemistry over the past $10\,000$ years, as well as being much larger than the amplitude of natural temporal variability of pH in the deep North Atlantic (González-Dávila et al., 2010). The first threshold (-0.2) is in line with recommendations by environmental agencies (Schubert et al., 2006) following the precautionary principle, and is reported to increase mortality of deep-sea benthic organisms during in situ exposure experiments (Barry et al., 2005). The second threshold (-0.3) allows for bracketing of a range of changes spanning from an $\sim 58\%$ increase in hydrogen ion concentration up to $\sim 100\%$.

3.2 Projections of pH reductions over the 21st century

Time series of atmospheric CO_2 (ppm) for three out of four IPCC RCP scenarios between 2006 and 2100 show an increase in CO_2 over this century; only RCP2.6 does not show a general increase with time (Fig. 2a). The corresponding simulated pH reductions for surface and deep North Atlantic waters are presented in Fig. 2b and c. Projected pH changes are indicated as the multi-model mean along with the between-model spread. Monitoring at time series stations reveals that the observed surface ocean pH decrease tracks increasing atmospheric $p\text{CO}_2$ (Orr, 2011). This trend is confirmed by the decline in simulated surface ocean pH (Fig. 2b) with a small between-model spread. In the surface ocean the extent of ocean acidification is set by the atmospheric CO_2 trajectory, along with physical climate change, namely warming and associated changes in ocean circulation and CO_2 thermodynamic properties. Surface waters, with high levels of dissolved anthropogenic CO_2 and characterized by low pH values, are entrained to the interior ocean during seasonal mixed layer deepening and deep convection episodes. As a result, deep pH changes (Fig. 2c) reflect atmospheric CO_2 to a lesser extent. Because the deep-water formation differs between models, the inter-model spread is significantly larger in deep waters than for the surface ocean.

The spatial pattern of pH reductions is exemplified for RCP4.5 and RCP8.5 in Fig. 3 (Fig. S3 for RCP2.6 and RCP6.0). Under RCP4.5 (Fig. 3a) and RCP8.5 (Fig. 3b), pH reductions crossing the -0.2 threshold are projected for continental slopes and a latitudinal band extending from 55 to 65° N . Since the pH perturbation originates at the sea surface, the continental slope and topographic heights (e.g. mid-Atlantic ridge) experience the largest pH reductions. Increasing impact on the sea floor between RCP4.5 and RCP8.5 for a threshold of -0.2 reflects the depth exposure to the pH perturbation of continental slopes and the mid-Atlantic ridge. In summary, the spatial pattern is set by a combination of




Figure 2. Time series of (a) atmospheric CO₂ (ppm) for RCP2.6, RCP4.5, RCP6.0 and RCP8.5 scenarios between 2006 and 2100 and corresponding simulated average North Atlantic pH changes relative to the pre-industrial mean for (b) surface waters and (c) deep waters. Hatching indicates the 2.5–97.5 % confidence interval of multi-model averages.

topography and North Atlantic circulation pathways. It reflects the transfer of the surface born anomaly of pH to the ocean interior during deep-water formation and downstream transport away from convection sites by the deep western boundary current.

By the end of the 21st century, projected pH reductions (Table 1) cross the -0.2 threshold for all scenarios except RCP2.6. For RCP2.6, deep-water pH reductions remain be-

low thresholds with likely limited impact on benthic environments. Under moderate RCP4.5, a decrease in pH beyond -0.2 units is projected for large areas of the North Atlantic, with about 16.7 ± 4.2 % of the sea floor area below 500 m being impacted. This estimate increases to 21.0 ± 4.4 % of the North Atlantic sea floor area under the most severe scenario (RCP8.5) and is still 14.0 ± 3.3 % of the sea floor for a threshold of -0.3 . The area impacted does not scale linearly

Table 1. Fraction of North Atlantic seafloor (35–75° N, 0–90° W) below 500 m experiencing a reduction in pH ≥ 0.2 , as well as ≥ 0.3 , at the end of the 21st century. Fractions for multi-model mean and standard deviation are given in percentage of impacted surface area relative to the total surface seafloor area of the North Atlantic sector. n is the number of simulations available at time of analysis for each RCP.

	n	pH reduction ≥ 0.2		pH reduction ≥ 0.3	
		mean (%)	SD (%)	mean (%)	SD (%)
RCP2.6	6	1.2	1.1	0.0	0.1
RCP4.5	7	16.7	4.2	0.6	0.5
RCP4.5/fixclim	2	18.1	n.a	0.8	n.a
RCP6.0	4	19.9	5.0	4.4	1.5
RCP8.5	7	21.0	4.4	14.0	3.3

with atmospheric CO₂ (Table 1) but levels off at higher RCPs for the -0.2 threshold. The -0.3 pH unit threshold (a 100 % increase in $[H^+]$) is not reached for RCP4.5, and only modest impacts are projected for RCP6.0 (Table 1). We expect, however, an increase in impacted area for all scenarios and pH thresholds beyond 2100 in response to legacy effects of CO₂ emissions and ongoing downward propagation of the pH perturbation (Frölicher and Joos, 2010).

3.2.1 Opposing effects of climate change and ocean acidification

The progression from RCP2.6 to RCP8.5 corresponds to a series of increasing geochemical (atmospheric pCO_2) and physical (climate change, defined here as changes in ocean dynamics in response to atmospheric warming) forcing with opposing effects on deep-ocean acidification.

In order to distinguish between the physical and geochemical drivers of North Atlantic deep-water acidification, we assessed two contrasting simulations available for two Earth system models (GFDL-ESM2M and IPSL-CM5A-LR) for RCP4.5. The first simulation (Fig. 4a) includes climate change effects on ocean circulation and geochemical effects on the seawater CO₂ system in response to atmospheric pCO_2 increase (RCP4.5). In the second experiment (Fig. 4b), the circulation and ocean physics are kept at pre-industrial conditions, but atmospheric CO₂ levels following RCP4.5 are used to force ocean acidification (RCP4.5/fixclim). The difference in pH between RCP4.5 and RCP4.5/fixclim (Fig. 4c) allows, at first order and within the limits of non-linearities (Schwinger et al., 2014), for isolation of the effect of climate change on pH changes. The negative differences in pH on panel c indicate stronger acidification in RCP4.5/fixclim, and suggest a slight alleviation of ocean acidification at depth and over the timescale of this study by climate change. In the experiment where ocean circulation was held at pre-industrial conditions (RCP4.5/fixclim), there was a small increase in the area impacted by pH re-

ductions for all thresholds (Table 1). Largest differences in projected pH values between RCP4.5 and RCP4.5/fixclim co-occur with large negative anomalies in winter mixed layer depth maxima in the Labrador Sea and negative pH anomalies downstream of convection sites following the deep western boundary current (Doney and Jenkins, 1994). This is in line with the projected enhancement of stratification across the North Atlantic in response to increasing temperatures and freshening. It will result in changes in winter mixed layer depth and deep convection and a decrease in the Atlantic Meridional Overturning Circulation (Meehl et al., 2007; Cheng et al., 2013). While increasing atmospheric CO₂ reduces pH, increasing climate change reduces surface-to-deep water exchange. In addition, topography modulates the extent of deep-water acidification. The combination of climate change, the non-linearities of the carbonate system, and topography explains the levelling-off of impacts in Table 1 for pH reductions exceeding -0.2 .

3.2.2 Projected impacts on ecosystems

In order to evaluate the risk for specific benthic ecosystems to be affected by pH reductions, we co-located seamounts (Fig. 3, black dots) and deep-sea canyons (Fig. 3, red dots) – both of which are key habitats of high biodiversity – and pH changes for RCP4.5 and RCP8.5 separately computed from the multi-model mean (see Supplement for RCP2.6 and RCP6.0). To further the evaluation of potential impacts of pH reductions beyond pH thresholds, we computed the area of seamounts for which a corresponding decrease is projected. A significant proportion of these habitats will be impacted by pH reductions exceeding -0.2 units by the end of the 21st century under moderate to high emission scenarios (Fig. 5). The geographic pattern results in close to 22.5 ± 5.3 % (14.7 ± 4.1 %) of North Atlantic deep-sea canyons and 7.7 ± 3.6 % (2.7 ± 0.9 %) of seamount ecosystems being exposed to pH reductions exceeding -0.2 (-0.3) units under RCP8.5. Under the moderate scenario, RCP4.5, model projections indicate that 19.0 ± 5.7 % of deep-sea canyons and 3.5 ± 1.6 % of seamounts still will experience pH reductions exceeding the -0.2 threshold. The close to constant impact reflects the use of a diagnostic that is based on counts of features being impacted, in addition to the depth distribution and propagation of the pH anomaly.

Seamounts and deep-sea canyons are known as hotspots of biodiversity and harbour a variety of distinct communities including reef-building cold-water corals, soft coral gardens and deep-sea sponge aggregations (Buhl-Mortensen et al., 2010, 2012; Clark et al., 2010; De Leo et al., 2010; ICES, 2007). Recent assessments reveal a high level of anthropogenic pressures on these ecosystems (Clark et al., 2010; Ramirez-Llodra et al., 2011). While fishing and resource extraction are recognized as the dominant human pressures at present and in the near future, expert assessments highlight the need for an appropriate quantification




Figure 3. Projected changes in pH between pre-industrial and the experiments forced by IPCC RCP scenarios by 2100. The panels show ensemble-mean differences in pH between the pre-industrial and the 2090–2100 average for (a) RCP4.5 and (b) RCP8.5. Locations of deep-sea canyons and seamounts are indicated by red and black symbols, respectively. The -0.2 pH contour line is plotted to delineate areas experiencing pH reductions beyond this threshold.

of the impacts of climate change and ocean acidification (Taranto et al., 2012). Present international conservation targets aim at preserving 10 % of marine biomes by 2020 (Convention of Biodiversity, 2011). Although not directly comparable to the outcome of model projections, it is nevertheless of interest to confront this preservation target with model results suggesting that ~ 8 % of North Atlantic seamounts and 23 % of canyons will experience a decrease in pH ex-

ceeding 0.2 pH units by the year 2100 for the most severe scenario. Seamounts identified as marine protection areas in the OSPAR region and excluding active venting sites (e.g. Josephine Seamount, $36^{\circ}40.02'N$, $14^{\circ}15.00'W$; Sedlo seamount, $40^{\circ}12.8'N$, $26^{\circ}15.8'W$) fall within the area for which these pH reductions are projected.

Our knowledge of the ecology of deep benthic communities is still limited, and impacts of pH changes on these




Figure 4. Projected changes in deep-ocean pH between pre-industrial and experiments forced with RCP scenarios by 2100: **(a)** RCP4.5, **(b)** RCP4.5/fixclim, and **(c)** difference in pH between **(a)** and **(b)** together with changes in maximum winter mixed layer depth (contour lines). The change in pH is computed as the difference in mean pH between the pre-industrial and the 2090–2100 average.

communities are difficult to evaluate owing to the lack of experimental and observational data. Rapid changes in pH will likely lead to disruption of extracellular acid–base balance, impedance of calcification and other physiological effects in

deep-water organisms, and whatever acclimation is required may have increased energetic costs (Widdicombe and Spicer, 2008) – e.g. for metabolism/maintenance, growth, reproduction – and could extend to increases in mortality of both




Figure 5. Projected impacts on seamounts (circles) and canyons (diamonds) as a function of atmospheric CO₂ levels by year 2100 for pH reductions exceeding (a) -0.2 and (b) -0.3 . Impact is computed as the fraction of the surface area affected by a reduction exceeding the threshold for seamounts, as well as the number of canyons surrounded by waters for which the reduction in pH exceeding the threshold is projected. Model pH is the decadal mean (2090–2100). Note that the seamount and canyon multi-model averages for the RCP2.6 scenario overlie each other. Light coloured circles: values obtained for each Earth system model; dark coloured circles: multi-model average for each scenario. Vertical and horizontal bars: 2.5–97.5 % confidence interval of multi-model averages.

adults and juveniles. Changes at the individual and population level will inevitably lead to more widespread ecosystem and community level changes and potential shifts in biodiversity (Hendriks et al., 2010) and ecosystem functioning (Danovaro et al., 2008). Biodiversity reductions could arise from a loss of species, or even functional or taxonomic groups, sensitive to pH change. The ecological implications of pH change could be more severe if keystone or habitat-forming species are impacted (Widdicombe and Spicer, 2008), which seems likely (Guinotte et al., 2006). These effects may be likely exacerbated in the presence of other stressors (Walther et al., 2009), such as global warming and projected reductions in deep-sea food supply (Bopp et al., 2013), as well as elevated resource exploitation and pollution. In particular, reductions in food supply to deep benthic communities are projected to result in a decrease in biomass and a shift towards smaller sized organisms (Jones et al., 2014). These changes will modify energy transfer rates through benthic food webs and may leave communities more

susceptible to pH reductions. We propose these and future model projections be taken into account when defining long-term preservation and management approaches to deep-sea ecosystems.

4 Conclusions

This study assesses the potential for detrimental pH reduction to occur across the deep North Atlantic by the end of the 21st century. It evaluates results from seven fully coupled Earth system models and for four Representative Concentration Pathways ranging from RCP2.6 to RCP8.5. In three out of the four scenarios, the multi-model analysis suggests that by 2100 over 17 % of the seafloor area below 500 m depth in the North Atlantic sector will experience pH reductions exceeding -0.2 units. Enhanced stratification in response to warming and freshening of surface waters slightly counteracts deep-water acidification. pH reductions co-occur

with sites of high deep-sea biodiversity such as seamounts and canyons. Model projections indicate that by the end of this century and for the high CO₂ scenario RCP8.5, close to 23 % (~ 15 %) of North Atlantic deep-sea canyons and ~ 8 % (3 %) of seamounts will experience pH reductions exceeding −0.2 (−0.3) units. Seamounts proposed as sites of marine protected areas are potentially threatened by these pH reductions. The spatial pattern of impacts reflects the depth of the pH perturbation and does not scale linearly with atmospheric CO₂ concentration. Impacts may cause negative changes of the same magnitude or exceeding the current biodiversity target of 10 % of preservation of marine biomes, implying that ocean acidification may offset benefits from conservation/management strategies relying on the regulation of resource exploitation.

The Supplement related to this article is available online at doi:10.5194/bg-11-6955-2014-supplement.

Acknowledgements. This work was supported through EU FP7 projects EPOCA (grant no. 211384) and CARBOCHANGE (grant no. 264879). D. O. B. Jones was funded by the UK Natural Environment Research Council as part of the Marine Environmental Mapping Programme (MAREMAP). S. C. Doney acknowledges support from the National Science Foundation (AGS-1048827). F. Joos acknowledges support from the Swiss National Science Foundation. This is a contribution from the BIOFEEDBACK project of the Centre for Climate Dynamics at Bjerknes Centre. C. Heinze and J. Tjiputra are grateful for support through NOTUR projects NN2980K and NN2345K as well as NorStore projects NS2980K and NS2345K for HPC CPU time and data storage. To analyse the CMIP5 data, this study benefited from the IPSL Prodiguer-Ciclad facility, which is supported by CNRS and UPMC, as well as Labex L-IPSL, which is funded by the ANR (grant no. ANR-10-LABX-0018) and by the European FP7 IS-ENES2 project (grant no. 312979).

Edited by: G. Herndl

References

- Barry, J. P., Buck, K. R., Lovera, C., Kuhn, L., and Whaling, P. J.: Utility of deep sea CO₂ release experiments in understanding the biology of a high-CO₂ ocean: Effects of hypercapnia on deep sea meiofauna, *J. Geophys. Res.*, 110, C09S12, doi:10.1029/2004JC002629, 2005.
- Barry, J. P., Widdicombe, S., and Hall-Spencer, J. M.: Effects of ocean acidification on marine biodiversity and ecosystem function, in: *Ocean Acidification*, edited by: Gattuso, J.-P. and Hansson, L., Oxford University Press, 192–209, 2011.
- Barry, J. P., Buck, K. R., Lovera, C., Brewer, P. G., Seibel, B. A., Drazen, J. C., Tamburri, M. N., Whaling, P. J., Kuhn, L., and
- Pane, E. F.: The response of abyssal organisms to low pH conditions during a series of CO₂-release experiments simulating deep-sea carbon sequestration, *Deep-Sea Res. Pt. II*, 92, 249–260, 2013.
- Billé, R., Kelly, R., Biastoch, A., Harrould-Kolieb, E., Herr, D., Joos, F., Kroeker, K., Laffoley, D., Oschlies, A., and Gattuso, J.-P.: Taking action against ocean acidification: A review of management and policy options, *Environ. Manag.*, 52, 761–779, 2013.
- Bopp, L., Resplandy, L., Orr, J. C., Doney, S. C., Dunne, J. P., Gehlen, M., Halloran, P., Heinze, C., Ilyina, T., Séférian, R., Tjiputra, J., and Vichi, M.: Multiple stressors of ocean ecosystems in the 21st century: projections with CMIP5 models, *Biogeosciences*, 10, 6225–6245, doi:10.5194/bg-10-6225-2013, 2013.
- Bryan, F.: High-latitude salinity effects and interhemispheric thermohaline circulations, *Nature*, 323, 301–304, 1986.
- Buhl-Mortensen, L., Vanreusel, A., Gooday, A. J., Levin, L. A., Priede, I. G., Buhl-Mortensen, P., Gheerardyn, H., King, N. J., and Raes, M.: Biological structures as a source of habitat heterogeneity and biodiversity on the deep ocean margins, *Mar. Ecol.*, 31, 21–50, 2010.
- Buhl-Mortensen, L., Buhl-Mortensen, P., Dolan, M. F. J., Dannheim, J., Bellec, V., and Holte, B.: Habitat complexity and bottom fauna composition at different scales on the continental shelf and slope of northern Norway, *Hydrobiologia*, 685, 191–219, 2012.
- Cheng, W., Chiang, J., and Zhang, D.: Atlantic Meridional Overturning Circulation (AMOC) in CMIP5 models: RCP and Historical Simulations, *J. Climate*, 26, 7187–7197, 2013.
- Childress, J. J.: Are there physiological and biochemical adaptations of metabolism in deep-sea animals, *Trends Ecol. Evol.*, 10, 1–36, 1995.
- Clark, M. R., Rowden, A. A., Schlacher, T., Williams, A., Consalvey, M., Stocks, K. I., Rogers, A. D., O'Hara, T. D., White, M., Shank, T. M., and Hall-Spencer, J. M.: The ecology of seamounts: structure, function, and human impacts, *Annu. Rev. Mar. Sci.*, 2, 253–278, 2010.
- Clauss, G. and Hoog, S.: *Deep Sea Challenges of Marine Technology and Oceanographic Engineering*, Developments in Marine Technology 12, Science-Technology Synergy for Research in the Marine Environment: Challenges for the XXI Century, Elsevier, Oxford, England, 2002.
- Convention on Biodiversity (CBD): target 11, document COP/10/INF/12/Rev.1, <http://www.cbd.int/sp/targets/rationale/target-11/>, 2011.
- Danovaro, R., Gambi, C., Dell'Anno, A., Corinaldesi, C., Fraschetti, S., Vanreusel, A., Vincx, M., and Gooday, A. J.: Exponential decline of deep-sea ecosystem functioning linked to benthic biodiversity loss, *Curr. Biol.*, 18, 1–8, 2008.
- De Leo, F. C., Smith, C. R., Rowden, A. A., Bowden, D. A., and Clark, M. R.: Submarine canyons: hotspots of benthic biomass and productivity in the deep sea, *Proc. R. Soc. B.*, 277, 2783–2792, 2010.
- Doney, S. C. and Jenkins, W. J.: Ventilation of the deep western boundary current and the abyssal western North Atlantic: estimates from tritium and ³He distributions, *J. Phys. Oceanogr.*, 24, 638–659, 1994.

- Duarte, C. M., Hendriks, I. E., Moore, T. S., Olsen, Y. S., Steckbauer, A., Ramajo, L., Carstensen, J., Trotter, J. A., and McCulloch, M.: Is Ocean Acidification an Open-Ocean Syndrome? Understanding Anthropogenic Impacts on Seawater pH, *Estuar. Coast.*, 36, 221–236, 2013.
- Elderfield, H., Ferretti, P., Greaves, M., Crowhurst, S., McCave, I. N., Hodell, D., and Piotrowski, A. M.: Evolution of Ocean Temperature and Ice Volume Through the Mid-Pleistocene Climate Transition, *Science*, 337, 704–709, 2012.
- Fautin, D. G., Guinotte, J. M., and Orr, J. C.: Comparative depth distribution of corallimorpharians and scleractinians (Cnidaria: Anthozoa), *Mar. Ecol. Prog. Ser.*, 397, 63–70, 2009.
- Fleeger, J. W., Carman, K. R., Weisenhorn, P. B., Sofranko, H., Marshall, T., Thistle, D., and Barry, J. P.: Simulated sequestration of anthropogenic carbon dioxide at a deep-sea site: Effects on nematode abundance and biovolume, *Deep-Sea Res. Pt. I*, 53, 1135–1147, 2006.
- Frölicher, T. L. and Joos, F.: Reversible and irreversible impacts of greenhouse gas emissions in multi-century projections with the NCAR global coupled carbon cycle-climate model, *Clim. Dyn.*, 35, 1439–1459, 2010.
- Gehlen, M., Bopp, L. and Aumont, O.: The short-term dissolution response of pelagic carbonate sediments to the invasion of anthropogenic CO₂: A model study, *Geochem. Geophys. Geosyst.*, 9, Q02012, doi:10.1029/2007GC001756, 2008.
- González-Dávila, M., Santana-Casiano, J. M., Rueda, M. J., and Llinás, O.: The water column distribution of carbonate system variables at the ESTOC site from 1995 to 2004, *Biogeosciences*, 7, 3067–3081, doi:10.5194/bg-7-3067-2010, 2010.
- Guinotte, J. M., Orr, J., Cairns, S., Freiwald, A., Morgan, L., and George, R.: Will human-induced changes in seawater chemistry alter the distribution of deep-sea scleractinian corals? *Front. Ecol. Environ.*, 4, 141–146, 2006.
- Harris, P. T. and Whiteway, T.: Global distribution of large submarine canyons: Geomorphic differences between active and passive continental margins, *Mar. Geol.*, 285, 69–86, 2011.
- Hendriks, I. E., Duarte, C. M., and Álvarez, M.: Vulnerability of marine biodiversity to ocean acidification: A meta-analysis, *Est. Coast. Shelf Sci.*, 86, 157–164, 2010.
- Hofmann, G. E., Smith, J. E., Johnson, K. S., Send, U., Levin, L. A., Micheli, F., Paytan, A., Price, N. N., Peterson, B., Takeshita, Y., Matson, P. G., Crook, E. D., Kroeker, K. J., Gambi, M. C., Rivest, E. B., Frieder, C. A., Yu, P. C., and Martz, T. R.: High-Frequency Dynamics of Ocean pH: A Multi-Ecosystem Comparison, *PLoS ONE*, 6, e28983, doi:10.1371/journal.pone.0028983, 2011.
- Hönisch, B., Ridgwell, A., Schmidt, D. N., Thomas, E., Gibbs, S. J., Sluijs, A., Zeebe, R., Kump, L., Martindale, R. C., Greene, S. E., Kiessling, W., Ries, J., Zachos, J. C., Royer, D. L., Barker, S., Marchitto, T. M., Moyer, R., Pelejero, C., Ziveri, P., Foster, G. L., and Williams, B.: The Geological Record of Ocean Acidification, *Science*, 335, 1058–1063, 2008.
- ICES, Report of the Working Group on Deep-water Ecology 26–28, ICES CM 2007/ACE:01 Ref. LRC, International Council for the Exploration of the Sea, Copenhagen, Denmark, 61 pp., 2007.
- IPCC: Workshop Report of the Intergovernmental Panel on Climate Change Workshop on Impacts of Ocean Acidification on Marine Biology and Ecosystems, eds Field CB et al. (IPCC Working Group II Technical Support Unit, Carnegie Institution, Stanford, California, United States of America), 164 pp., 2011.
- Jones, D. O. B., Yool, A., Wei, C.-L., Henson, S. A., Ruhl, H. A., Watson, R. A., and Gehlen, M.: Global reductions in seafloor biomass in response to climate change, *Glob. Change Biol.*, 20, 1864–1873, doi:10.1111/gcb.12480, 2014.
- Khaliwala, S., Tanhua, T., Mikaloff-Fletcher, S., Gerber, M., Doney, S. C., Graven, H. D., Gruber, N., McKinley, G. A., Murata, A., Ríos, A. F., and Sabine, C. L.: Global ocean storage of anthropogenic carbon, *Biogeosciences*, 10, 2169–2191, doi:10.5194/bg-10-2169-2013, 2013.
- Kroeker, K. J., Kordas, R. L., Crim, R. N., and Singh, G. G.: Meta-analysis reveals negative yet variable effects of ocean acidification on marine organisms, *Ecol. Lett.*, 13, 1419–1434, 2010.
- Marchal, O., Stocker, T. F., Joos, F., Indermühle, A., Blunier, T., and Tschumi, J.: Modeling the concentration of atmospheric CO₂ during the Younger Dryas climate event, *Clim. Dyn.*, 15, 341–354, 1999.
- Matsumoto, K. and Yokoyama, Y.: Atmospheric $\Delta^{14}\text{C}$ reduction in simulations of Atlantic overturning circulation shutdown, *Global Biogeochem. Cy.*, 27, 296–304, 2013.
- Meehl, G. A., Stocker, T. F., Collins, W. D., Friedlingstein, P., Gaye, A. T., Gregory, J. M., Kitoh, A., Knutti, R., Murphy, J. M., Noda, A., Raper, S. C. B., Watterson, I. G., Weaver, A. J., and Zhao, Z.-C.: Global Climate Projections, in: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2007.
- Mora, C., Wei, C.-L., Rollo, A., Amaro, T., Baco, A. R., et al.: Biotic and Human Vulnerability to Projected Changes in Ocean Biogeochemistry over the 21st Century, *PLoS Biol.*, 11, e1001682, doi:10.1371/journal.pbio.1001682, 2013.
- Moss, R. H., Edmonds, J. A., Hibbard, K. A., Manning, M. R., Rose, S. K., van Vuuren, D. P., Carter, T. R., Emori, S., Kainuma, M., Kram, T., Meehl, G. A., Mitchell, J. F. B., Nakicenovic, N., Riahi, K., Smith, S. J., Stouffer, R. J., Thomson, A. M., Weyant, J. P., and Wilbanks, T. J.: The next generation of scenarios for climate change research and assessment, *Nature*, 463, 747–756, 2010.
- Orr, J. C.: Recent and future changes in ocean carbonate chemistry, in: *Ocean Acidification*, edited by: Gattuso, J.-P. and Hansson, L., Oxford University Press, Oxford, 41–66, 2011.
- Orr, J. C., Fabry, V. J., Aumont, O., Bopp, L., Doney, S. C., Feely, R. A., Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., Key, R. M., Lindsay, K., Maier-Reimer, E., Matear, R., Monfray, P., Mouchet, A., Najjar, R. G., Plattner, G.-K., Rodgers, K. B., Sabine, C. L., Sarmiento, J. L., Schlitzer, R., Slater, R. D., Totterdell, I. J., Weirig, M.-F., Yamanaka, Y., and Yool, A.: Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms, *Nature*, 437, 681–686, 2005.
- Ramirez-Llodra, E.: Fecundity and life-history strategies in marine invertebrates, *Advances in Marine Biology*, 43, 87–170, 2002.
- Ramirez-Llodra, E., Tyler, P. A., Baker, M. C., Bergstad, O. A., Clark, M. R., Escobar, E., Levin, L. A., Menot, L., Rowden, A. A., Smith, C. R., and Van Dover, C. Ø.: Man and the Last Great Wilderness: Human Impact on the Deep Sea, *PLoS ONE*, 6, e22588, doi:10.1371/journal.pone.0022588, 2011.
- Roth, R. and Joos, F.: A reconstruction of radiocarbon production and total solar irradiance from the Holocene ^{14}C and CO₂

- records: implications of data and model uncertainties, *Clim. Past*, 9, 1879–1909, doi:10.5194/cp-9-1879-2013, 2013.
- Sabine, C. L., Feely, R. A., Gruber, N., Key, R. M., Lee, K., Bullister, J. L., Wanninkhof, R., Wong, C. S., Wallace, D. W. R., Tilbrook, B., Millero, F. J., Peng, T.-H., Kozyr, A., Ono, T., and Rios, A. F.: The oceanic sink for anthropogenic CO₂, *Science*, 305, 367–371, 2004.
- Sanyal, A., Hemming, N. G., Hanson, G. N., and Broecker, W. S.: Evidence for a higher pH in the glacial ocean from boron isotopes in foraminifera, *Nature*, 373, 234–236, 1995.
- Schubert, R., Schellnhuber, H.-J., Buchmann, N., Epiney, A., Griesshammer, R., Kulessa, M., Messner, D., Rahmstorf, S., and Schmid, J.: The future oceans – Warming up, rising high, turning sour, Special Report by the German Advisory Council on Global Change (WBGU), 123 pp., 2006.
- Schwinger, J., Tjiputra, J. F., Heinze, C., Bopp, L., Christian, J. R., Gehlen, M., Ilyina, T., Jones, C. D., Salas-Mélia, D., Segsneider, J., Séférian, R., and Totterdell, I.: Non-linearity of ocean carbon cycle feedbacks in CMIP5 earth system models, *J. Climate*, 27, 3869–3888, 2014.
- Seibel, B. A. and Walsh, P. J.: Potential Impacts of CO₂ Injection on Deep-Sea Biota, *Science*, 294, 319–320, 2001.
- Seibel, B. A. and Walsh, P. J.: Biological impacts of deep-sea carbon dioxide injection inferred from indices of physiological performance, *J. Experim. Biol.*, 206, 641–650, 2003.
- Somero, G. N.: The Physiology of Global Change: Linking Patterns to Mechanisms, *Annu. Rev. Mar. Sci.*, 4, 39–61, 2012.
- Steinacher, M., Joos, F., and Stocker, T. F.: Allowable carbon emissions lowered by multiple climate targets, *Nature*, 499, 197–201, 2013.
- Taranto, G. H., Kvile, K. Ø., Pitcher, T. J., and Morato, T.: An Ecosystem Evaluation Framework for Global Seamount Conservation and Management, *PLoS ONE*, 7, e42950, doi:10.1371/journal.pone.0042950, 2012.
- Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An overview of CMIP5 and the experiment design, *Bull. Am. Meteor. Soc.*, 93, 485–498, doi:10.1175/BAMS-D-11-00094, 2011.
- Tittensor, D. P., Baco, A. R., Hall-Spencer, J. M., Orr, J. C., and Rogers, A. D.: Seamounts as refugia from ocean acidification for coldwater stony corals, *Mar. Ecol.*, 31, 212–225, 2010.
- Turley, C. M., Roberts, J. M., and Guinotte, J. M.: Corals in deep-water: Will the unseen hand of ocean acidification destroy cold-water ecosystems?, *Coral Reefs*, 26, 445–448, 2007.
- Van Vuuren, D. P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G. C., Kram, T., Krey, V., Lamarque, J.-F., Masui, T., Meinshausen, M., Nakicenovic, N., Smith, S. J., and Rose, S. K.: The representative concentration pathways: an overview, *Clim. Change*, 109, 5–31, 2011.
- Walther, K., Sartoris, F. J., Bock, C., and Pörtner, H. O.: Impact of anthropogenic ocean acidification on thermal tolerance of the spider crab *Hyas araneus*, *Biogeosciences*, 6, 2207–2215, doi:10.5194/bg-6-2207-2009, 2009.
- Widdicombe, S. and Spicer, J. I.: Predicting the impact of ocean acidification on benthic biodiversity: What can physiology tell us?, *J. Exp. Mar. Biol. Ecol.*, 366, 187–197, 2008.
- Yesson, C., Clark, M. R., Taylor, M. L., and Rogers, A. D.: The global distribution of seamounts based on 30 arc seconds bathymetry data, *Deep-Sea Res. I*, 58, 442–453, 2011.
- Yu, J., Broecker, W. S., Elderfield, H., Jin, Z., McManus, J., and Zhang, F.: Loss of Carbon from the Deep Sea Since the Last Glacial Maximum, *Science*, 330, 1084–1087, 2010.
- Yu, J., Anderson, R. F., Jin, Z., Rae, J. W. B., Opdyke, N., and Eggin, S. M.: Responses of the deep ocean carbonate system to carbon reorganization during the Last Glacial–interglacial cycle, *Quat. Sci. Rev.*, 76, 39–52, 2013.