Otolith shape analysis and mitochondrial DNA markers distinguish three sand smelt species in the *Atherina boyeri* species complex in western Mediterranean

A. S. Boudinar, L. Chaoui, J. P. Quignard, D. Aurelle, M. H. Kara

► To cite this version:

A. S. Boudinar, L. Chaoui, J. P. Quignard, D. Aurelle, M. H. Kara. Otolith shape analysis and mitochondrial DNA markers distinguish three sand smelt species in the *Atherina boyeri* species complex in western Mediterranean. *Estuarine, Coastal and Shelf Science*, 2016, 182 (A), pp.202–210. 10.1016/j.ecss.2016.09.019 . hal-01444074

HAL Id: hal-01444074

<https://hal.science/hal-01444074>

Submitted on 1 Mar 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Otolith shape analysis and mitochondrial DNA markers distinguish three sand smelt species in the *Atherina boyeri* species complex in western Mediterranean.

Boudinar A.S.^a, Chaoui L.^a, Quignard J.P.^b, Aurelle D.^c, Kara M.H.^{a*}

^aLaboratoire Bioressources marines, Université d'Annaba Badji Mokhtar, Annaba, Algérie

^bLaboratoire d'ichtyologie, Université Montpellier II, Montpellier, France.

^cAix-Marseille Université, CNRS, IMBE UMR 7263, 13007, Marseille, France.

*Corresponding author: M. H. Kara, Laboratoire Bioressources marines, Université d'Annaba Badji Mokhtar, Algérie. Tel: +213 77 0312458. Fax number: +213 38 868510. E-mail: kara_hichem@yahoo.com

BOUDINAR AHMED SOFIANE : boudinar-2009@live.fr

CHAOUI LAMYA : chaouilamy@hotmail.com

QUIGNARD JEAN-PIERRE : qjp2@wanadoo.fr

AURELLE DIDIER : didier.aurelle@univ-amu.fr

KARA MOHAMED HICHEM : kara_hichem@yahoo.com

Abstract

Atherina boyeri is a common euryhaline teleost fish in the Mediterranean and adjacent areas, which inhabits coastal and estuarine waters, including coastal lagoons and more rarely inland waters. Several recent studies have pointed the possible existence of three distinct groups or species, one lagoon/freshwater group and two 'punctuated and unpunctuated on the flanks' marine groups, within an *A. boyeri* species complex. This study is a combined approach using otolith shape and molecular markers to better define the structure of the species in the western Mediterranean.

Genetic differentiation and species delimitation among nine *Atherina boyeri* populations from several marine and lagoon/brackish habitat sites in Algeria, Tunisia and France were investigated using three mitochondrial (control region, *Cyt b* and 16S) and one nuclear markers (2nd intron of *S7*). For further phylogenetic and phylogeographic study, we added sequences from Genbank covering more areas (Ionian Sea, Adriatic Sea, Tyrrhenian Sea, Black Sea, Atlantic). Five groups were found. Two of them perfectly corresponded to two species already recognized *Atherina presbyter* and *Atherina hepsetus*, both living in marine waters; and three additional, including *Atherina boyeri* (brackish and freshwater environments) and two independent groups of marine punctuated and unpunctuated individuals. Those findings are corroborated by the study of the otolith contour shape of 362 individuals of seven populations from different habitats using Fourier analysis. Individuals could be discriminated into five groups based on the first two functions (Wilk's lambda = 0.07, $p < 0.001$). Samples from Ziam inlet, marine punctuated individuals and unpunctuated marine specimens from Annaba's Gulf formed three well separated groups. Specimens from Mellah and Manguio lagoons formed another group. The last one includes individuals from Bizerte and Thau lagoons. The divergences between them strongly support the potential species within the *A. boyeri* species complex.

Keywords: *Atherina boyeri* species complex; mt-DNA; Mediterranean Sea; Fourier analysis; genetic divergence

1. Introduction

The Mediterranean Sea had a complex history, marked by the Messinian salinity crisis and climatic fluctuations of the recent Quaternary, characterized with the succession of glacial and interglacial periods. These events induced habitat fragmentation and alteration of connections between populations, and resulted in the establishment of its current biodiversity and of different biogeographical zones (Krijgsman et al., 1999; Patarnello et al., 2007). The Mediterranean Sea is now characterized by a high biodiversity with a quarter of endemic species (Bianchi et al., 2012). Despite decades of dedicated studies, knowledges on biodiversity still evolving, even for well known groups such as fishes (Bianchi et al., 2012). The development of genetic approaches allows a better analysis of species limits, for example by identifying cryptic species (Borsa et al., 2004; Pante et al., 2015). The comparison of genetic and morphological data can bring useful information on the evolution of characters used for species delimitation, on morphological plasticity, and on species interactions (differentiation vs gene flow).

The *Atherina* Linnaeus, 1758 genus has been a subject of several major systematic revisions (Kiener and Spillmann, 1969), where two genera and 20 species were condensed into a single genus. In Europe the Atherinidae family is represented by a single genus and three species: *Atherina boyeri* Risso, 1810, *Atherina hepsetus* Linnaeus, 1758 and *Atherina presbyter* Cuvier, 1829. *Atherina hepsetus* is restricted to the Mediterranean, although it has also been reported in the eastern Atlantic coasts of Spain, Morocco, Madeira and Canary Islands (Quignard and Pras, 1986). The distribution of *A. presbyter* ranges from the British Isles and southern North Sea to the Canary Islands, Mauritania and Cape Verde (Quignard and Pras, 1986). It has also been reported from the Azores archipelago (Santos et al., 1997) and rarely from the western Mediterranean (Kiener and Spillmann, 1969; Quignard and Pras, 1986). *Atherina boyeri* has been described as an extremely euryhaline teleost fish which can take a wide range of salinities, from freshwater to hypersaline conditions (110 ‰ maximum recorded) (Gon and Ben-Tuvia, 1983). It mainly inhabits coastal waters of the East Atlantic Ocean as well as in the Mediterranean, Black, Caspian and Aral seas, including estuarine waters, coastal lagoons and more rarely inland waters (Gon and Ben-Tuvia, 1983; Henderson and Bamber, 1987). This small fish forms local semi-isolated populations each with a characteristic morphology and life history traits (Henderson and Bamber, 1987). This species could be a good model to study differentiation and speciation in the Mediterranean Sea with a range spanning important phylogeographical limits and with important morphological diversity according to geography and habitats. Kottelat (1997) pointed out the possible

1 existence of two different groups in *A. boyeri s.l.*, one belonging to the lagoon / freshwater
2 environment and the other to the marine one. This author limited the name of *A. boyeri* to the
3 lagoonal form and proposed *A. mochon* Cuvier, 1829 for the marine one.

4 Several investigations based on morphometry (Kartas and Trabelsi 1990; Trabelsi et al.,
5 2002a, 2004, Boudinar et al., 2015), and allozymic analyses (Focant et al., 1992, 1993, 1999;
6 Cammarata et al., 1996) supported the hypothesis that *A. boyeri* could be subdivided into
7 separate sub-populations or species. Data based on allozymic analysis alone (Mauro et al.,
8 2007), or combined with morphometry (Trabelsi et al., 2002a, 2002b, 2004) pointed out the
9 existence of two (Klossa-Kilia et al., 2002, 2007; Kraitsek et al., 2008) or even three
10 Mediterranean species (Trabelsi et al., 2002a, 2002b, 2004; Astolfi et al., 2005; Francisco et
11 al., 2008, 2011; Milana et al., 2008), including one lagoon form (*A. lagunae*) and one or two,
12 (punctuated on the flanks "*A. punctata*" and non-punctuated "*A. boyeri*") marine forms. Since
13 the genetic divergence values estimated between marine and lagoon *A. boyeri* populations
14 were found to be similar to those existing between well-defined species of the same genus, the
15 authors elevated them to the species rank. All of these studies pointed out the existence of an
16 *A. boyeri* species complex, but for now there is no formal nor extensive description of the
17 biology and distribution range of these putative species. The under-representation of southern
18 populations from the western Mediterranean might have hindered the detection of additional
19 genetic structure (e.g. Chaoui et al., 2009) or a good estimation of the potential species range,
20 as the only data on this species in the Algerian coast are the study of *A. boyeri* species
21 complex using otolith shape and somatic morphology at a local scale (Boudinar et al. 2015),
22 and the age, growth and reproduction of Mellah lagoon population (Boudinar et al., 2016).
23 Additionally the use of a single genetic marker, mostly mitochondrial, is not sufficient for
24 precise species delimitation nor the identification of putative hybridization events. Finally an
25 evaluation of the morphological differentiation of these taxa has not been performed in all
26 cases. Indeed an integrative study including large geographical scope, multilocus and non
27 molecular approaches appears as the best approach for species delimitation (Puillandre et al.,
28 2012; Pante et al., 2015).

29 The purpose of the present work was to use otolith shape analysis and mitochondrial (Control
30 Region: CR, Cytochrome b: Cyt *b*, ribosomal DNA 16S) and nuclear DNA (S7 intron 2)
31 markers to study the morphological and genetic diversity of *A. boyeri* species complex from
32 different habitat (lagoon, inlet, sea) in the western Mediterranean. This combination of

investigative tools was attempted for the first time in this species complex and allowed
precising the species repartition and species limits in western Mediterranean.

2. Materials and methods

2.1. Sampling sites

Nine populations of *Atherina boyeri* including freshwater/brackish, marine punctuated (Fig. 1a) and marine unpunctuated fishes (Fig. 1b) were sampled from eight locations in the northern and southern banks of the western Mediterranean Sea (Fig. 2 and Table 1) representing different geographical areas (Algerian, Tunisian and French coasts and lagoons) and different environments: inshore brackish lagoons and the nearshore marine environment (hereafter called lagoon and sea habitats respectively) and inlets. All specimens were morphologically identified, according to Marfin (1981) and stored in 96% ethanol.

2.2. Otoliths morphometry

Otolith shape analysis was performed on six of the nine populations, the same specimens being used for molecular analyses. Samples from Lion's Gulf and Mafragh inlet were discarded: as sampling size was too small for statistical analyses ($N = 7$) and also because the specimens were fingerlings, which did not permit comparisons with adult fish. In order to eliminate fish size effects on otolith shape, all individuals used in this analysis belonged to the same size interval (5 to 8.5 cm Total Length, TL). A total of 362 fish were used for otolith shape analysis (Table 1). The sagittal otoliths were taken as pairs, washed in clean water, air-dried and stored in plastic tubes. Both right and left otoliths were examined under a stereomicroscope fitted with a numerical camera (Lumenera's INFINITY lite) linked to a computer monitor. Otoliths were orientated in a consistent manner, with the sulcus side up. Numerical pictures were then acquired with TNPC software (Mahé et al., 2011). For each numerical image, the software Shape 1.2 (Iwata and Ukai, 2002) calculated the Fourier coefficients in order to make them invariants to the otolith size, orientation and position, regarding the beginning of the outline, which is arbitrarily defined (Crampton, 1995). Also, the Fourier Power (FP) spectrum was calculated to estimate the sufficient number of harmonics accounting for the best reconstruction of the otolith outline (Pothin et al., 2006). A subsample of 90 otoliths (10 samples $\times$ 7 populations) was randomly created for this purpose. As the first 13 harmonics reached 99.99% of the mean cumulated power, the Fourier analysis indicated that the otolith shape could be summarized by these thirteen harmonics. A Factorial Discriminant Analysis (FDA) was performed with Fourier descriptors to compare otolith

shape variations among the sampling areas using the SPSS software version 13 (SPSS, 2004). Using Ward's method, a hierarchical clustering was performed with the FactoMineR package with R[®] software (R Core Team 2014). The classification success of the discriminant analysis was checked using jackknifed cross-validation.

2.3. Molecular analyses

For molecular analysis, nine populations of *A. boyeri* species complex were collected from eight locations in the northern and southern banks of the western Mediterranean Sea representing variety in environment kinds of habitats (Fig. 1). Total DNA was extracted from caudal fin using 10% Chelex® with a 1 h proteinase K digestion followed by 15 min at 95°C. Mitochondrial fragments of the control region (CR), 16rRNA, cytochrome *b* (Cyt *b*) genes and nuclear fragment of the S7 intron 2 were amplified by polymerase chain reaction (PCR); the PCR conditions and primers sequences and corresponding references are given in Table 2. For each marker, other available sequences from *Atherina* were retrieved from GenBank. The PCR products were sent for Sanger sequencing at Eurofins MWG. The numbers of sequences used for each marker in each locality are given in Table 1.

2.4. Species delimitation and population genetics analyses

The sequences were aligned with ClustalW (Thompson et al., 1994). In addition to the alignments for each gene, a CR and Cyt *b* concatenated alignment was performed as these markers presented the highest number of common sequences. Sequences of two Atheriniformes were used as outgroups: *Leptatherina wallacei* Prince, Ivantsoff & Potter, 1982 belonging to the family Atherinidae (GenBank accession numbers AY326785.1 for CR, AY313118.1 for Cyt *b*) and *Iso natalensis* Regan, 1919 belonging to the family Isonidae (GenBank accession numbers HM855075.1 for 16S). We also added sequences of the only two other congeneric species in the area, *A. presbyter* (GenBank accession numbers JF309618.1) and *A. hepsetus* (GenBank accession numbers JF309620.1), in order to obtain the closest outgroups of the *A. boyeri* species complex. For further phylogenetic and phylogeographic analyses we added sequences from Genbank covering more localities (Table 3, Fig. 3, Table S1-S3; supplementary material). We used Jmodeltest 3.7 (Guindon and Gascuel, 2003; Posada, 2008) with the Akaike Information Criterion to identify the best-fit model of evolution for each locus to be used in Maximum-Likelihood (ML) analysis: TVM+G for CR (gamma distribution shape parameter $G = 0.584$), TVM+I+G for Cyt *b* ($I = 0.252$, $G = 0.649$), TPM3uf+G for 16S ($G = 0.171$), and TIM3+G for S7 ($G = 1.221$). ML

analyses were performed with PhyML 3.0 (ref) and we tested the robustness of the inferred trees with 1000 bootstrap replicates (Felsenstein, 1985). Bayesian analyses (BA) were implemented in MrBayes 3.2 (Huelsenbeck and Ronquist, 2001; Ronquist and Huelsenbeck, 2003). The best-fit model of evolution for each locus was set using MrModeltest 2.3 (Nylander, 2004): HKY+G for CR (gamma distribution shape parameter $G = 0.522$), GTR+I+G for Cyt *b* ($I = 0.332$, $G = 0.786$), SYM+G for 16S ($G = 0.171$), and SYM+G for S7 ($G = 1.016$). We used one independent run of two million generations of Markov chain Monte Carlo (MCMC) simulations for each marker to approximate posterior probabilities of phylogenetic trees. Topologies were sampled every 1000 generations, and a majority-rule consensus tree was estimated after discarding the first 2500 sampled generations.

The obtained sequences were used for a phylogeographic analysis. For each marker data set, we analyzed the total amount of genetic variance into three variance components: among groups, among populations within groups and within populations, using an analysis of molecular variance (AMOVA) in Arlequin v3.5 (Excoffier and Lisher, 2010). Samples were grouped by geographic proximity, habitat type or a combination of both, resulting in five different groupings (Table 4). According to the analyzed marker, some test combination may be absent due to the lack of sequence in those localities. The analysis was based on Φ_{st} (Excoffier et al., 1992), and its significance was tested through 1000 permutations. The significance of tests involving multiple comparisons was adjusted with a Bonferroni correction. Average number of nucleotide substitutions per site, Dxy (Nei, 1987) were used to estimate differentiation between populations using DNAsp v5 (Librado and Rozas, 2009).

3. Results

3.1. *Otoliths morphometry*

The first two discriminant functions of the FDA performed with Fourier descriptors accounted for 89.40% of the variance. Individuals could be discriminated into five groups based on the first two functions (Wilk's lambda = 0.07, $p < 0.001$). Samples from Ziama inlet, marine punctuated individuals and unpunctuated marine specimens from Annaba's Gulf formed well separated groups. Specimens from Mellah and Manguio lagoons formed another group. The last one includes individuals from Bizerte and Thau lagoons (Fig. 5). Overall area-classification success was 84.30%. The best reclassification rate is 84.03% for Annaba's Gulf punctuated marine specimens, then specimens from Ziama Inlet 75%, Bizerte Lagoon 71.92%, Manguio Lagoon 66.07%, Annaba's Gulf unpunctuated marine specimens 64.75%, Mellah Lagoon specimens 55%, and the Thau Lagoon individuals 53.48% (Table 4). As for

the molecular analysis, some of the Thau Lagoon specimens are scattered between lagoon and marine environment.

3.2. Species delimitation

In total, 373 bp (379 sequences) were sequenced from the CR, 464 bp (81 sequences) from the 16S, 315 bp (217 sequences) from the Cyt *b* and 187 from the S7 (149 sequences) (including indels), yielding 170, 35, 72 and 27 haplotypes in *A. boyeri* species complex, respectively. The number of sequences used for the analysis includes the Genbank sequences. BA and ML species delimitation trees gave similar results in all cases.

For the Cyt *b*, CR, 16S, and the concatenated fragments CR and Cyt *b*, converged on a similar tree and showed the same topology. Seven clades were observed, *A. presbyter* and *A. hepsetus*, appeared as a sister group of the *A. boyeri* species complex. The five clades of the *A. boyeri* species complex: marine punctuated form, marine unpunctuated form, and two or three clades of the lagoonal form, are well supported by high bootstrap values.

The different clades of the lagoonal form correspond to different geographical areas: for the Cyt *b* the first one grouped specimens from Greece (Nea Peramos-Kavala), the second grouped lagoon and insular Tunisian specimens, and the third clade grouped other of lagoonal populations. Four specimens from Thau lagoon, were grouped with lagoonal samples while the other specimens were grouped with the marine unpunctuated individuals (Figure S1; supplementary material).

For the 16S, three clades inside the lagoonal form, the first group exclusively Tunisian specimens from Bizerte lagoon, the second specimens from Bulgaria and Turkey, and the third one grouped other of lagoonal populations (Figure S1; supplementary material).

For the CR, three clades were observed, the first one grouped specimens from Tunisia, and Italy (Marsala and one specimen from Cagliari), the second grouped specimens from Turkey (Danube, Lake Sapanca) and one specimen from Croatia (Spalato). The third clade grouped other lagoonal samples.

The tree for CR includes the most localities and sample size is shown in Fig. 5, along with bootstrap and posterior probabilities values.

For the concatenated fragments CR and Cyt *b*, the lagoonal form is separated into two clades, one with the tunisian specimens and the other with those sampled from Algerian and French lagoons (Figure S1; supplementary material).

The nuclear S7 marker did not lead to a well resolved phylogeny, but it allowed us to confirm the grouping of some specimens from Thau with the lagoonal samples (presented in a rectangle on the tree) (Figure S2; supplementary material).

3.3. Population genetics

AMOVA results (Table 5) confirmed the existence of a significant genetic structure based on the three mt-DNA markers and showed a highly significant global structure when comparing lagoonal, marine unpunctuated and marine punctuated samples (1), CR ($F_{SC} = 0.66$; $F_{ST} = 0.92$; $F_{CT} = 0.78$; $P < 0.001$); Cyt *b* ($F_{SC} = 0.77$; $F_{ST} = 0.94$; $F_{CT} = 0.76$; $P < 0.001$); 16S ($F_{SC} = 0.77$; $F_{ST} = 0.96$; $F_{CT} = 0.82$; $P < 0.001$).

The analysis based on Cyt *b* between marine punctuated samples (2) shows a significant divergence among populations within groups ($F_{SC} = 0.06$; $P < 0.001$). No differences are observed within populations ($F_{ST} = 0.61$; $P = 0.06$) or among groups ($F_{CT} = 0.58$; $P = 0.1$).

Comparison based on Cyt *b*, between marine unpunctuated samples (3) show a significant differences among populations within groups ($F_{SC} = 0.14$; $P < 0.001$) and within populations ($F_{ST} = 0.38$; $P = 0.06$), no significant differences are observed among groups ($F_{CT} = 0.28$; $P = 0.16$).

The comparison between North, South Ouest and South East Mediterranean lagoonal samples (4) also revealed significant differences, CR ($F_{SC} = 0.43$; $F_{ST} = 0.82$; $P < 0.001$; $F_{CT} = 0.68$; $P = 0.01$); Cyt *b* ($F_{SC} = 0.16$; $F_{ST} = 0.83$; $F_{CT} = 0.80$; $P < 0.001$); 16S ($F_{SC} = 0.17$; $P < 0.001$; $F_{ST} = 0.83$; $P = 0.02$), based on 16S no significant differences were observed among groups ($F_{CT} = 0.80$; $P = 0.38$).

Significant differences were also found while comparing lagoonal samples as observed in the tree topology, based on CR ($F_{SC} = 0.70$; $F_{ST} = 0.82$; $P < 0.001$; $F_{CT} = 0.41$; $P = 0.01$).

The average number of nucleotide substitutions per site between populations for the Cyt *b* was respectively $D_{xy} = 0.14$, $D_{xy} = 0.15$ when comparing lagoonal sequences with marine unpunctuated and punctuated ones, and $D_{xy} = 0.11$ when comparing marine unpunctuated and punctuated sequences. When we compare respectively *A. hepsetus* with lagoonal, marine unpunctuated and punctuated samples, the average number of nucleotide substitutions is $D_{xy} = 0.17$, $D_{xy} = 0.13$, $D_{xy} = 0.14$. Those values are lower when comparing *A. presbyter* with lagoonal ($D_{xy} = 0.14$), marine unpunctuated ($D_{xy} = 0.10$) and punctuated ($D_{xy} = 0.13$) samples. The average number of nucleotide substitutions is the lowest when comparing *A. hepsetus* and *A. presbyter* ($D_{xy} = 0.06$).

For the CR, when comparing lagoonal sequences with marine unpunctuated and punctuated ones Dxy is respectively equal to 0.15 and 0.18, and Dxy = 0.17 when comparing marine unpunctuated and punctuated sequences. When we compare respectively *A. hepsetus* with lagoonal, marine unpunctuated and punctuated samples, the average number of nucleotide substitutions is Dxy = 0.21, Dxy = 0.16, Dxy = 0.18. In the other hand when we compare *A. presbyter* with lagoonal, marine unpunctuated and punctuated samples the average number of nucleotide substitutions is respectively, Dxy = 0.20, Dxy = 0.19, Dxy = 0.16, its lowest value is set when comparing *A. hepsetus* and *A. presbyter* (Dxy = 0.11).

For the 16S, when comparing lagoonal sequences with marine unpunctuated and punctuated ones Dxy is respectively equal to 0.08 and 0.07, and Dxy = 0.07 when comparing marine unpunctuated and punctuated sequences. When we compare respectively *A. hepsetus* or *A. presbyter* with lagoonal, marine unpunctuated and punctuated samples, the average number of nucleotide substitutions is Dxy = 0.08, Dxy = 0.07, Dxy = 0.07, and Dxy = 0.02 when comparing *A. hepsetus* and *A. presbyter*.

4. Discussion

Otoliths are widely used in different studies, such as species identification (Aguirre and Lombarte, 1999; Parmentier et al., 2001) and may vary within species according to fish size, site, depth and/or other environmental factors (Begg et al., 2001; Stransky and MacLellan, 2005; Hüseyin, 2008). Several studies have shown that shape analysis allowed discrimination of local fish stocks according to ecological factors (Smith, 1992; Friedland and Reddin, 1994; Bolles and Begg, 2000; Cardinale et al., 2004). But this raises the question of the genetic versus environmental determinism (i.e. plasticity) of otolith shape.

In the present study, otolith shape analysis showed five groups: The first one included Mellah and Mauguio lagoons samples, when the second gathered individuals from Bizerte and Thau lagoons, the remaining samples (Ziama inlet, marine punctuated specimens and marine unpunctuated specimens from Annaba's gulf) constitute each a well separated group. Those results correspond to what have been found by Boudinar et al. (2015) in a local study on the same Algerian populations, where differences were observed between Mellah lagoon, Ziama inlet, marine punctuated specimens and marine unpunctuated specimens from Annaba's gulf, using otolith shape and somatic morphology. Three different mtDNA markers (CR, Cyt b, 16S) corroborated otolith shape findings. Both gave us information about how much the punctuated and unpunctuated marine forms are different. Besides, an intra-specific variation into the lagoonal form of *A. boyeri* species complex using both methods was apparent. Molecular

results for the three mtDNA markers using different tree construction methods are consistent and allow us to distinguish seven separated groups. Two of them perfectly corresponded to two species already recognized as valid (*A. hepsetus* ① and *A. presbyter* ②) and three additional corresponded to the marine punctuated *A. punctata* ③, marine unpunctuated *A. boyeri* ④ and lagoonal *A. lagunae* ⑤ traditionally included in the *A. boyeri* species complex (Trabelsi et al., 2002b). The main clade grouped lagoon samples (Bizerte, Mauguio, Mellah) and fresh water samples (Ziama and Mafragh inlets). The remaining two clades correspond to punctuated and unpunctuated marine groups, in which we found respectively marine punctuated samples from Annaba Gulf and marine unpunctuated samples from Annaba and Lion's Gulf. Those clades are supported with high bootstraps and posterior probabilities. The divergence between them was comparable to that separating *A. presbyter* from *A. hepsetus* thereby supporting the potential species status of the different clades inside the *A. boyeri* species complex. The observation of misplaced individuals in the species tree was interpreted as the possibility of rare migration events and supported the hypothesis of gene flow among populations. The inter-lagoon difference has already been reported by Trabelsi et al., (2002b) who found an important genetic divergence between Tunisian and French lagoon sand smelts which is supported by high bootstrap values, suggesting that there has been a long-term barrier to the gene flow between these two populations. This was also reported by Astolfi et al., (2005) in a larger scale considering populations from Atlantic, Mediterranean and Black Sea lagoon, where an important and significant genetic structure was observed, with some groupings displaying low bootstraps values. Milana et al., (2012) along the Italian and Albanian coasts of the Mediterranean Sea where a pronounced spatial genetic structure has been detected by microsatellites among lagoon populations of *A. boyeri* species complex with different statistical approaches used.

The distinctiveness of the three species of the *A. boyeri* species complex using both molecular and otoliths shape analysis is observed across the eastern and western basins of the Mediterranean Sea. These results support the findings based on morphometric data (Kartas and Trabelsi 1990; Trabelsi et al., 2002a, 2004), Cyt *b* gene (Trabelsi et al., 2002b), and mtDNA control region (Astolfi et al., 2005). Francisco et al., (2008) confirmed these results with further mtDNA fragments (control region and 12S), while Milana et al., (2008) in their analysis of portion of the mitochondrial cytochrome *b*, detected the presence of an insertion common to lagoon and punctuated forms of *A. boyeri* species complex. (88% sequence identity of the insertion was found for marine punctuated form. 93% sequence identity of the

1 insertion for the lagoon form. And 47% similarity when comparing the insertions for the two
2 forms). Using three mtDNA markers (CR, 12S, 16S) and two nuclear markers (Rhodopsin and
3 S7 intron 2) Francisco et al., (2011) sustained the previous the previous results.

4 These differences may have a genetic determinism linked to a local selective process
5 potentially and / or habitat choice or (Lemaire et al., 2000; Chaoui et al., 2012). This could
6 also correspond to morphological plasticity (e.g. Capoccioni et al., 2011) or to a combination
7 of environmental and genetic effects. Most likely, the reason for *A. boyeri* to appear as a
8 complex clade is related to its distinct life cycle which is, among the other species, the one
9 most associated with more fragmented environments (lagoons, estuaries and freshwater
10 bodies). The combination between this fragmentation in its habitats linked to the low dispersal
11 ability of the species is probably responsible for the highly phylogeographic pattern observed.

12 Our results reinforce the species status of the three Mediterranean forms traditionally included
13 in the *A. boyeri* species complex using both of otolith shape and mitochondrial DNA markers.
14 A formal description and range distribution definition should be considered as a priority for
15 this species, with further phylogeographic investigations including wider sampling and
16 additional molecular genetic markers.

17 18 **Acknowledgements**

19 This work is funded in part by PHC-Maghreb research program (Partenariat Hubert Curien N°
20 12MDU880M). The authors thank also the Algerian Ministry for Higher education and
21 scientific research (General directorate for scientific research and technology development,
22 GDRSDT) which financially supported this study, within the framework of the National
23 Funds of Research (NFR).

References

- Aguirre, H., Lombarte, A., 1999. Ecomorphological comparisons of sagittae in *Mullus barbatus* and *M. surmuletus*. *Journal of Fish Biology* 55, 105–114.
- Almada, F., Almada, V., Guillemaud, T., Wirtz, P., 2005. Phylogenetic relationships of the north-eastern Atlantic and Mediterranean blenniids. *Biological Journal of the Linnean Society* 86, 283–295.
- Astolfi, L., Dupanloup, I., Rossi, R., Bisol, P., Faure, E., Congiu, L., 2005. Mitochondrial variability of sand smelt *Atherina boyeri* populations from north Mediterranean coastal lagoons. *Marine Ecology Progress Series* 297, 233–243.
- Begg, G.A., Overholtz, W.J., Munroe, N.J., 2001. The use of internal otolith morphometrics for identification of haddock (*Melanogrammus aeglefinus*) stocks on Georges Bank. *Fishery Bulletin* 99, 1–14.
- Bianchi, C.N., Morri, C., Chiantore, M., Montefalcone, M., Parravicini, V., Rovere, A., 2012. Mediterranean Sea biodiversity between the legacy from the past and a future of change. *Life in the Mediterranean Sea: a look at habitat changes*. New York: Nova Science Publishers, 1–55.
- Bolles, K., Begg, G., 2000. Distinction between silver hake (*Merluccius bilinearis*) stocks in US waters of the northwest Atlantic based on whole otolith morphometrics. *Fishery Bulletin* 98, 451–462.
- Borsa, P., Collet, A., Durand, J.D., 2004. Nuclear-DNA markers confirm the presence of two anchovy species in the Mediterranean. *Comptes rendus biologies* 327, 1113–1123.
- Boudinar, A.S., Chaoui, L., Mahe, K., Cachera, M., Kara, M.H., 2015. Habitat discrimination of big scale sand smelt *Atherina boyeri* Risso, 1810 (Atheriniformes: Atherinidae) in eastern Algeria using somatic morphology and otolith shape. *Ital. J. Zool.* 82, 446–453.
- Boudinar, A.S., Chaoui, L., M., Kara, M.H., 2016. Age, growth and reproduction of the sand smelt *Atherina boyeri* Risso, 1810 in Mellah Lagoon (eastern Algeria). *J. Appl. Ichthyol.* 32, 302–309.
- Camarata, M., Mauro, A., Mazzola, A., Scilipoti, D., Arculeo, M., Parrinello, N., 1996. A biochemical genetic study of isoenzyme polymorphism within and between two populations of *Atherina boyeri* Risso. *Genetika* 32, 1220–1224.
- Cantatore, P., Roberti, M., Pesole, G., Ludovico, A., Milella, F., Gadaleta, M.N., Saccone, C., 1994. Evolutionary analysis of cytochrome b sequences in some Perciformes

- evidence for a slower rate of evolution than in mammals. *Journal of Molecular Evolution* 39, 589–597.
- Capoccioni, F., Costa, C., Aguzzi, J., Menesatti, P., Lombarte, A., Ciccotti, E., 2011. Ontogenetic and environmental effects on otolith shape variability in three Mediterranean European eel (*Anguilla anguilla*, L.) local stocks. *Journal of Experimental Marine Biology and Ecology* 397, 1–7.
- Cardinale, M., Doering-Arjes, P., Kastowsky, M., Mosegaard, H., 2004. Effects of sex, stock, and environment on the shape of known-age Atlantic cod (*Gadus morhua*) otoliths. *Canadian Journal of Fisheries and Aquatic Sciences* 61, 158–167.
- Chaoui, L., Quignard, J.P., Kara, M.H., 2001. Différenciation morphologique de deux populations marine et lagunaire de daurade *Sparus aurata* (Linné, 1785). Rapport de la Commission Internationale pour l'Exploration Scientifique de la Mer Méditerranée 36, 371.
- Chaoui, L., Kara, M. H., Quignard, J. P., Faure, E., Bonhomme, F., 2009. Forte différenciation génétique de la daurade *Sparus aurata* (L., 1758) entre les deux rives de la Méditerranée occidentale. *Comptes Rendus Biologies* 332, 329–335.
- Chaoui, L., Gagnaire, P.A., Guinand, B., Quignard, J.P., Tsigenopoulos, C., Kara, M.H., 2012. Microsatellite length variation in candidate genes correlates with habitat in the gilthead sea bream *Sparus aurata*. *Molecular ecology* 21, 5497–5511.
- Chow, S., Hazama, K., 1998. Universal PCR primers for S7 ribosomal protein gene introns in fish. *Molecular Ecology* 7, 1247–1263.
- Crampton, J.S., 1995. Elliptic Fourier shape analysis of fossil bivalves, practical considerations. *Lethaia* 28, 179–186.
- Excoffier, L., Lischer, H.E.L., 2010. Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Mol. Ecol. Resour.* 10, 564–567.
- Excoffier, L., Smouse, P.E., Quattro, J.M., 1992. Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data. *Genetics* 131, 479–491.
- Felsenstein, J., 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39, 783–791.
- Focant, B., Trabelsi, M., Kartas, F., Quignard, J.P., 1992. Caractérisation biochimique de trois populations d'*Atherina boyeri* des milieux lagunaires méditerranéens. Rapport de la

- Commission Internationale pour l'Exploration Scientifique de la Mer Méditerranée 33, 94.
- Focant, B., Trabelsi, M., Vandewalle, P., Kartas, F., Quignard, J.P., 1993. Apport de l'analyse biochimique à la caractérisation de populations d'*Atherina boyeri* des milieux lagunaires méditerranéens Actes du colloque Okeanos, «Le système littoral méditerranéen», Montpellier, 145–148.
- Focant, B., Rosocchi, E., Crivelli, A.J., 1999. Attempt at biochemical characterization of sand smelt *Atherina boyeri* Risso, 1810 (Pisces, Atherinidae) populations from the Camargue (Rhône delta, France). Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology 122, 261–267.
- Francisco, S.M., Congiu, L., Stefanni, S., Castilho, R., Brito, A., Ivanova, P.P., Levy, A., Cabral, H., Kiliyas, G., Doadrio, I., Almada, V.C., 2008. Phylogenetic relationships of the North-eastern Atlantic and Mediterranean forms of *Atherina* (Pisces, Atherinidae). Molecular Phylogenetics and Evolution 48, 782–788.
- Francisco, S.M., Congiu, L., von der Heyden, S., Almada, V.C., 2011. Multilocus phylogenetic analysis of the genus *Atherina* (Pisces: Atherinidae). Molecular Phylogenetics and Evolution 61, 71–78.
- Friedland, K., Reddin, D., 1994. Use of otolith morphology in stock discriminations of Atlantic salmon (*Salmo salar*). Canadian Journal of Fisheries and Aquatic Sciences 51, 91–98.
- Gon, O., Ben-Tuvia, A., 1983. The biology of Boyer's sand smelt, *Atherina boyeri* Risso in the Bardawil Lagoon on the Mediterranean coast of Sinai. Journal of Fish Biology 22, 537–547.
- Guindon, S., Gascuel, O.A., 2003. Simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. Systematic Biology 52, 696–704.
- Henderson, P., Bamber, R., 1987. On the reproductive biology of the sand smelt *Atherina boyeri* Risso (Pisces: Atherinidae) and its evolutionary potential. Biological Journal of the Linnean Society 32, 395–415.
- Huelsenbeck, J.P., Ronquist, F., 2001. MRBAYES: Bayesian inference of phylogeny. Bioinformatics 17, 754–755.
- Hüssy, K., 2008. Otolith shape in juvenile cod (*Gadus morhua*): Ontogenetic and environmental effects. Journal of Experimental Marine Biology and Ecology 364, 35–41.

- 1 Iwata, H., Ukai, Y., 2002. SHAPE: a computer program package for quantitative evaluation of
2 biological shapes based on elliptic Fourier descriptors. *Journal of Heredity* 93, 384–
3 385.
- 4 Kartas, F., Trabelsi, M., 1990. Sur le polymorphisme de l'athérine, *Atherina boyeri* Risso,
5 1810 (*Atherinidae*) des eaux littorales tunisiennes. *Cybium* 14, 295–311.
- 6 Kiener, A., Spillmann, C.J., 1969. Contribution à l'étude systématique et écologique des
7 athérines des côtes françaises. *Mémoires du Muséum national d'Histoire naturelle -*
8 *Série A : Zoologie* 60, 33–74.
- 9 Klossa-Kilia, E., Prassa, M., Papatziropoulos, V., Alahiotis, S., Kiliass, G., 2002.
10 Mitochondrial DNA diversity in *Atherina boyeri* populations as determined by RFLP
11 analysis of three mtDNA segments. *Heredity* 89, 363–370.
- 12 Klossa-Kilia, E., Papatziropoulos, V., Tryfonopoulos, G., Alahiotis, S., Kiliass, G., 2007.
13 Phylogenetic relationships of *Atherina hepsetus* and *Atherina boyeri* (Pisces:
14 Atherinidae) populations from Greece, based on mtDNA sequences. *Biological*
15 *Journal of the Linnean Society* 92, 151–161.
- 16 Kottelat, M., 1997. European freshwater fishes. *Biologia* 52, 1–271.
- 17 Kraitsek, S., Klossa-Kilia, E., Papatziropoulos, V., Alahiotis, S.N., Kiliass, G., 2008. Genetic
18 divergence among marine and lagoon *Atherina boyeri* populations in Greece using
19 mtDNA analysis. *Biochemical genetics* 46, 781–798.
- 20 Krijgsman, W., Hilgen, F.J., Raffi, I., Sierro, F.J., Wilson, D.S., 1999. Chronology, causes and
21 progression of the Messinian salinity crisis. *Nature* 400, 652–655.
- 22 Lemaire, C., Allegrucci, G., Naciri, M., Bahri-Sfar, L., Kara, M.H., Bonhomme, F., 2000. Do
23 discrepancies between microsatellite and allozyme variation reveal differential
24 selection between sea and lagoon in the sea bass (*Dicentrarchus labrax*)?. *Molecular*
25 *ecology* 9, 457–467.
- 26 Librado, P., Rozas J., 2009. DnaSP v5: a software for comprehensive analysis of DNA
27 polymorphism data. *Bioinformatics* 25, 1451–1452.
- 28 Mahé, K., Fave, S., Couteau, J., 2011. TNPC : User guide. Version 5. Boulogne-sur-mer:
29 Ifremer Institute. Computer Program.
- 30 Marfin, J. I., 1981. Biologie de l'Atherine: *Atherina boyeri* Risso, 1810 (Poissons
31 téléostéen) dans trois milieux saumâtres du Roussillon (Leucate, Canet,
32 Bourdigou). Ph.D. Thesis. University of Perpignan, Montpellier.

- 1 Mauro, A., Arculeo, M., Mazzola, A., Parrinello, N., 2007. Are there any distinct genetic sub-
2 populations of sand smelt, *Atherina boyeri* (Teleostei: Atherinidae) along Italian
3 coasts? Evidence from allozyme analysis. *Folia Zoologica* 56, 194–200.
- 4 Meyer, A., Kocher, T.D., Basasibawaki, P., Wilson, A.C., 1990. Monophyletic origin of Lake
5 Victoria's cichlid fishes suggested by mitochondrial DNA sequences. *Nature* 347,
6 550–553.
- 7 Milana, V., Sola, L., Congiu, L., Rossi, A., 2008. Mitochondrial DNA in *Atherina* (Teleostei,
8 Atheriniformes): differential distribution of an intergenic spacer in lagoon and marine
9 forms of *Atherina boyeri*. *Journal of Fish Biology* 73, 1216–1227.
- 10 Milana, V., Franchini, P., Sola, L., Angiulli, E., Rossi, A.R., 2012. Genetic structure in
11 lagoons: the effects of habitat discontinuity and low dispersal ability on populations of
12 *Atherina boyeri*. *Marine biology* 159, 399–411.
- 13 Nylander, J.A.A., 2004. MrModeltest v2. Program distributed by the author. Evolutionary
14 Biology Centre, Uppsala University.
- 15 Nei, M., 1987. *Molecular Evolutionary Genetics*. Columbia University Press, New York.
- 16 Ostellari, L., Barglloni, L., Penzo, E., Patarnello, P., Patarnello, T., 1996. Optimization of
17 single-strand conformation polymorphism and sequence analysis of the
18 mitochondrial control region in *Pagellus bogaraveo* (Sparidae, Teleostei):
19 rationalized tools in fish population biology. *Animal Genetics* 27, 423–427.
- 20 Pante, E., Puillandre, N., Viricel, A., Arnaud-Haond, S., Aurelle, D., Castelin, M.,
21 Chenuil, A., Destombe, C., Forcioli, D., Valero, M., Viard, F., Samadi, S., 2015.
22 Species are hypotheses: avoid connectivity assessments based on pillars of sand.
23 *Molecular ecology* 24, 525–544.
- 24 Parmentier, E., Vandewalle, P., Lagardère, F., 2001. Morpho-anatomy of the otic region in
25 carapid fishes: eco-morphological study of their otoliths. *Journal of Fish Biology* 58,
26 1046–1061.
- 27 Patarnello, T., Volckaert, F.A.M., Castilho, R., 2007. Pillars of Hercules: is the Atlantic-
28 Mediterranean transition a phylogeographical break? *Mol. Ecol.* 16, 4426–4444.
- 29 Posada, D., 2008. JModeltest: phylogenetic model averaging. *Molecular Biology and*
30 *Evolution* 25, 1253–1256.
- 31 Pothin, K., Gonzalez-Salas, C., Chabanet, P., Lecomte-Finiger, R., 2006. Distinction
32 between *Mulloidichthys flavolineatus* juveniles from Reunion Island and Mauritius
33 Island (south-west Indian Ocean) based on otolith morphometrics. *Journal of Fish*
34 *Biology* 69, 38–53.

- Puillandre, N., Modica, M.V., Zhang, Y., Sirovich, L., Boisselier, M.C., Cruaud, C., Samadi S., 2012. Large-scale species delimitation method for hyperdiverse groups. *Molecular ecology* 21, 2671–2691.
- Quignard, J.P., Pras, A., 1986. Atherinidae. In: Whitehead, P., Bauchot, M.L., Hureau, J.C., Nielsen, J., Tortonese, E. (Eds.), *Fishes of the North-eastern Atlantic and the Mediterranean*. UNESCO, Paris, pp. 1207–1210.
- R Core Team. 2014. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Available: <http://www.R-project.org/>. Accessed Apr 2014 22.
- Ronquist, F., Huelsenbeck, J.P., 2003. MRBAYES 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19, 1572–1574.
- Santos, R.S., Porteiro, F.M., Barreiros, J.P., 1997. Marine Wshes of the Azores: annotated checklist and bibliography. *Bull Univ Azores*, 1, 244
- Smith, M.K., 1992. Regional differences in otolith morphology of the deep slope red snapper *Etelis carbunculus*. *Canadian Journal of Fisheries and Aquatic Sciences* 49, 795–804.
- SPSS, 2004. SPSS for Windows, Version 13.0. Chicago, SPSS Inc. Computer Program.
- Stransky, C., MacLellan, S.E., 2005. Species separation and zoogeography of redfish and rockfish (genus *Sebastes*) by otolith shape analysis. *Canadian Journal of Fisheries and Aquatic Sciences* 62, 2265–2276.
- Thompson, J.D., Gibson, T.J., Plewniak, F., Jeanmougin, F., Higgins, D.G., 1997. The Clustal X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Research* 25, 4876–4882.
- Trabelsi, M., Faure, E., Quignard, J.P., Boussaid, M., Focant, B., Maamouri, F., 2002a. *Atherina punctata* and *Atherina laguna* (Pisces, Atherinidae) new species found in the Mediterranean sea, Article 1: morphological investigations of three Atherinid species. *Comptes Rendus de l'Académie des Sciences Paris Life Science* 325, 967–975.
- Trabelsi, M., Gilles, A., Fleury, C., Quignard, J.P., Maamouri, F., Faure, E., 2002b. *Atherina punctata* and *Atherina lagunae* (Pisces, Atherinidae), new species found in the Mediterranean Sea. Article 2: molecular investigations of three Atherinid species. *Comptes Rendus de l'Académie des Sciences Paris Life Science* 325, 1119–1128.
- Trabelsi, M., Maamouri, F., Quignard, J.P., Boussaïd, M., Faure, E., 2004. Biometric and molecular investigations evidenced allopatric speciation in western Mediterranean lagoons within the *Atherina lagunae* species (*Teleostei*, *Atherinidae*). *Estuarine, Coastal and Shelf Science* 61, 713–723.

Fig. 1. Marine punctuated (A) and marine unpunctuated (B) forms belonging to *Atherina boyeri* species complex.

1

2

3 Fig. 2. Map of the study area. The eight sampling locations are indicated by their codes as
 4 defined in Table 2. Circles indicate inshore lagoonal habitat, triangles marine habitat and
 5 square inlet/brakish habitat. The upper map shows the north-western Mediterranean sampling
 6 area, and the lower map shows the south-western Mediterranean sampling area.

7

Fig. 3. Map of sampling locations for the extended analysis. The location numbers details are presented in table 3.

Fig. 4. Hierarchical clustering using Fourier's descriptors on otolith shape of *A. boyeri* species complex. Sampling locations are indicated by their codes as defined in Table 2.

Fig. 5. Phylogenetic relationships within *Atherina* for the Control region (CR). Each main node is represented by circled capital letters. ① : *A. hepsetus*, ② : *A. presbyter*, ③ : *A. punctata*, ④ : *A. boyeri* and ⑤ : *A. lagunae*. Percentages of bootstrap for maximum likelihood are in bold followed by the posterior probability for Bayesian analysis.

Table 1. Description of *A. boyeri* species complex samples including sampling locations, samples codes, number of used sequences for each marker and number of otoliths for the shape analysis in the western Mediterranean Sea region.

Sampling locations	Country	Sample code	CR	Cyt <i>b</i>	16S	S7	Otoliths
Annaba Gulf	Algeria	MP	16	12	7	7	61
Annaba Gulf		MNP	18	16	4	8	60
Mellah lagoon		ML	15	10	4	9	50
Mafragh Inlet		MI	22	24	9	10	
Ziama Inlet		Z	24	28	6	8	56
Bizerte lagoon	Tunisia	BZ	17	8	4	7	57
Thau lagoon	France	TH	15	7	6	11	22
Mauguio lagoon		MG	21	20	8	9	56
Lion's Gulf		M	7	5	7	6	

1 Table 2. Primers and polymerase chain reaction conditions for each fragment.

Fragment	Primer name	Sequence	Sequence Source	PCR conditions
Control region	L-pro1	5'-ACTCTCACCCCTAGCTCCCAAAG-3'	(Ostellari et al., 1996)	94 °C 2' 35 x (94 °C 30", 55 °C 30", 72 °C 1') 72 °C 5'
	H-DL1	5'-CCTGAAGTAGGAACCAGATGCCAG-3'		
Cytochrome b	New-For	5'-AGCCTACGAAAACCCACCC-3'	(Meyer et al., 1990)	92 °C 2" 30 x (92 °C 15', 52 °C 45', 72 °C 1.5") 72 °C 8"
	34-Rev	5'-AAACTGCAGCCCCTCAGAATGATATTTGTCCTCA-3'	(Cantatore et al., 1994)	
16S	16SFor	5'-AAGCCTCGCCTGTTTACCAA-3'	(Almada et al., 2005)	94 °C 3' 35 x (94 °C 45", 50 °C 1', 72 °C 1') 72 °C 10'
	16SRev	5'-CTGAACTCAGATCACGTAGG-3'		
S7 intron 2	S7RPEX2F	5'-AGCGCCAAAATAGTGAAGCC-3'	(Chow and Hazama, 1998)	94 °C 2' 40 x (95 °C 45", 52 °C 1', 72 °C 45') 72 °C 10'
	S7RPEX3R	5'-GCCTTCAGGTCAGAGTTCAT-3'		

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

1 Table 3. Sampling location and number of *Atherina boyeri* species complex individuals used
 2 in the extended phylogenetic analyses.

N° in Fig.	Collection site		N	lagoonal fish			unpunctuated fish			punctuated fish		
				Cyt <i>b</i>	16S	CR	Cyt <i>b</i>	16S	CR	Cyt <i>b</i>	16S	CR
1	Portugal	Mondego	28			28						
2		Tagus	21		3	21						
3		Boquilobo	1			1						
4		Tago	2			2						
5		Sado	4			4						
6		Lagoa Albufeira	22			22						
7		Ilha de Faro	6			6						
8		Castro Marim	1			1						
9	Spain	Zonar	2			2						
10		Ebro	5			5						
		Barcelona	4		4							
11	France	Ayrolle	6	6								
12		Agde	3							3	3	2
13		Thau	20				20	3	17			
14		Mauguio	23	18	8	23						
15		Or	9			9						
16		Marseille	7						7			
17		Scandola	7				7					
18		Scandola	2							2	2	2
19		Biguglia	9	9		2						
20		Lavezzi	3							3	2	2
21		Corsica										2
22	Algeria	Ziama Inlet	27	25	5	27						
23		Annaba Gulf	17							10	4	17
24		Annaba Gulf	13				2	1	13			
25		Mafragh Inlet	23	22	9	23						
26		Mellah lagoon	16	6	5	16						
27	Tunisia	Bizerte lagoon	12	5	3	12						
28		Cap Zebib	4							4		
29		Cap Zebib	8				8	3	2			
30		Tunis	3			3						
31		Hergla	10				10					
32		Hergla	3							3		2
33		Monastir	10				10					
34		Karkennah Sidi Fraj	9	9								
35		Karkennah El Attaya	7	7								
36		Karkennah Sidi Youssef	6	6								
37		Djerba	8	8								
38	Italy	Cagliari	4			4						
39		Muravera	1	1		1						
40		Fucecchio	7			7						
41		Marsala	4			4						
42		Faro	8			8						
43		Scardovari	7			7						
44		Chioggia	21		4	21						
45		Varano	1	1		1						
46		Acquatina	6			6						
47	Croatia	Spalato	10		4	10						
48	Greece	Nea Peramos-Kavala	2						1			1
49	Romania	Danube	9			9						
50	Bulgaria	Bulgaria	5		5							
51	Turkey	Turkey	3		3							
		Lake Sapanca	5			5						
52	Israel	Israel	3					2	3			
53	Morocco	Loukkos	4		4							
54		Sebou	16			16						

1 Table 4. Classification success rate using based on otolith shape of *A. boyeri* species complex
2 using jackknifed cross-validation. Sampling locations are indicated by their codes as defined
3 in Table 2.

Sites	ML	MNP	MP	BZ	Z	MG	TH
ML	55	8	1	11	6	14	5
MNP	5	79	9	0	11	10	8
MP	1	8	100	3	3	1	3
BZ	8	1	6	82	4	5	8
Z	5	5	2	1	84	12	3
MG	10	7	3	8	5	74	5
TH	3	3	2	7	0	5	23

1 Table 5. AMOVA results for five different groupings, as a function of geographic proximity,
 2 habitat type or a combination of both.

	CR				Cyt <i>b</i>				16S			
	ddl	Variation	F-statistics	P-value	ddl	Variation	F-statistics	P-value	ddl	Variation	F-statistics	P-value
(1) MP/MNP/Lagoon												
Among populations within groups	2	78.67	FSC=0.66	0	2	76.67	FSC=0.77	0	2	82.81	FSC=0.77	0
Within populations	38	14.16	FST=0.92	0	24	18.1	FST=0.94	0	16	13.31	FST=0.96	0
Among groups	333	7.18	FCT=0.78	0	181	5.23	FCT=0.76	0	54	3.88	FCT=0.82	0
(2) Within MP												
Among populations within groups					2	58.94	FSC=0.06	0				
Within populations					3	2.52	FST=0.61	0.0645				
Among groups					19	38.54	FCT=0.58	0.109				
(3) Within MNP												
Among populations within groups					2	28.77	FSC=0.14	0				
Within populations					3	10.07	FST=0.38	0.0009				
Among groups					51	61.16	FCT=0.28	0.1632				
(4) Lagoons: Med North/Med South Ouest/Med South East												
Among populations within groups	2	68.17	FSC=0.43	0	2	80.02	FSC=0.16	0	2	80.12	FSC=0.17	0
Within populations	7	13.95	FST=0.82	0	8	3.25	FST=0.83	0	2	3.49	FST=0.83	0.028
Among groups	114	17.88	FCT=0.68	0.014	110	16.73	FCT=0.80	0	24	16.39	FCT=0.80	0.385
(5) Lagoon populations as in tree topology												
Among populations within groups	2	41.63	FSC=0.70	0								
Within populations	27	41.16	FST=0.82	0								
Among groups	275	17.21	FCT=0.41	0.017								

3

4