Habitat availability and geographic isolation as potential drivers of population structure in an oceanic dolphin in the Southwest Indian Ocean

Amélia Viricel, Benoit Simon-Bouhet, Laura Ceyrac, Violaine Dulau-Drouot, Per Berggren, Omar A. Amir, Narriman S. Jiddawi, Philippe Mongin, Jeremy J. Kiszka

► To cite this version:

Amélia Viricel, Benoit Simon-Bouhet, Laura Ceyrac, Violaine Dulau-Drouot, Per Berggren, et al.. Habitat availability and geographic isolation as potential drivers of population structure in an oceanic dolphin in the Southwest Indian Ocean. *Marine Biology*, 2016, 163 (219), 10.1007/s00227-016-2999-3 . hal-01444114

HAL Id: hal-01444114

<https://hal.science/hal-01444114>

Submitted on 23 Jan 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Habitat availability and geographic isolation as potential drivers of population structure in an oceanic dolphin in the Southwest Indian Ocean

Amélia Viricel¹, Benoit Simon-Bouhet ^{1,7}, Laura Ceyrac², Violaine Dulau-Drouot³, Per Berggren⁴, Omar A. Amir⁵, Narriman S. Jiddawi⁶, Philippe Mongin⁸, Jeremy J. Kiszka^{1,2,7,9*}

¹ Littoral ENvironnement et Sociétés (LIENSs), UMR 7266 CNRS-Université de la Rochelle. Institut du Littoral et de l'Environnement, 2 rue Olympe de Gouges, F-17000 La Rochelle, France.

² Institut de Recherche pour le Développement (IRD), UMR 212 EME (Exploited Marine Ecosystems). Centre de Recherche Halieutique Méditerranéenne et Tropicale, BP 171, F-34203, Sète cedex, France.

³ Globice Réunion, 30 chemin Parc Cabris, Grand Bois, F-97410 Saint Pierre, La Réunion, France.

⁴ School of Marine Science and Technology, Newcastle University, NE1 7RU, UK

⁵Ministry of Livestock and Fisheries, P. O. Box 295, Nyangumi House, Maruhubi Street, Zanzibar, Tanzania.

⁶ Institute of Marine Sciences, P.O. Box 668, Zanzibar, Tanzania.

⁷ Centre d'Etudes Biologiques de Chizé (CEBC), UMR 7372 CNRS-Université de la Rochelle. Institut du Littoral et de l'Environnement, 2 rue Olympe de Gouges, F-17000 La Rochelle, France

⁸ Brigade Nature Ocean Indien, Parc de la Providence, 97400 Saint-Denis, La Réunion, France

⁹ Current affiliation: Department of Biological Sciences, Florida International University, 3000
NE 151st Street, North Miami, FL33181, USA

*corresponding author: Jeremy Kiszka, jeremy.kiszka@gmail.com, Tel: +1 305 919 4104

ABSTRACT

Delphinid populations show highly variable patterns of genetic diversity and population structure. Previous studies indicate that habitat discontinuities and geographic isolation are major drivers of population division in cetaceans. Spinner dolphins (*Stenella longirostris*) are distributed in all tropical oceans, but they are particularly common around islands and atolls. This species occurs in shallow waters at daytime to rest and socialise, and feeds on offshore mesopelagic prey overnight. Here we investigated the genetic population structure of spinner dolphins in the Southwest (SW) Indian Ocean along a west-east geographical gradient, from eastern Africa to the Mascarene archipelago. We combined analyses of 12 microsatellite loci, mtDNA control region sequences, and sighting data to assess genetic differentiation and characterise habitat preferences of these populations. Significant genetic structure among the three sampled sites (Zanzibar, Mayotte and La Réunion) was observed using both types of molecular markers. Overall, our results indicate that geographic isolation, and potentially other factors such as shallow water habitats to rest and socialise, may be important drivers of the genetic population structure of insular spinner dolphins in this region.

Keywords: spinner dolphin, *Stenella longirostris*, genetic structure, microsatellites, mtDNA, East Africa.

INTRODUCTION

Understanding factors influencing population connectivity has been a central and long-standing research avenue in marine ecology (e.g. Cowen et al. 2000; 2007; Selkoe et al 2008). Past studies have shown that the genetic structure of marine populations is driven by a number of processes, including water currents, sea floor topography, water temperature and life history (e.g., Fullard et al. 2000; Fontaine et al. 2007; Pelc et al. 2009; Ciannelli et al. 2010; Mendez et al. 2011; 2013). In cetaceans, factors that can lead to discontinuous relationships between genetic and geographic distance include habitat characteristics, intra-species niche partitioning (e.g. foraging specialization) and kinship, combined with demographic processes (e.g., Hoelzel 2009; Möller et al. 2007, 2011, Louis et al. 2014a,b; Viricel and Rosel 2014).

Beside its fundamental importance in marine ecology, understanding the spatial structure and genetic connectivity of marine populations is also critical for conservation and management purposes. Indeed, while no marine ecosystem is completely unaffected by human activities, threats to populations of marine organisms vary geographically (Halpern et al. 2008). In the Southwest (SW) Indian Ocean for instance, the importance of marine mammal bycatch is spatially variable and it seems to primarily affect inshore species, including coastal delphinids (Kiszka et al 2008). Thus, delimiting biologically meaningful conservation units (i.e., based on population structure assessments) will be a crucial step toward preserving the marine megafauna of this region.

The spinner dolphin (*Stenella longirostris*) is one of the most abundant and widely distributed tropical delphinids (Perrin 2009). Four subspecies are currently recognised, based on morphological and ecological differences (Perrin and Gilpatrick 1994; Perrin et al. 1999). The Gray's spinner dolphin (*S. longirostris longirostris*), hereafter the spinner dolphin, is primarily an insular subspecies, and its distribution includes the Atlantic, Indian and Pacific Oceans (Perrin and Gilpatrick 1994). In French Polynesia, Hawaii and the Maldives, spinner dolphins enter atolls, sheltered bays and lagoons through reef channels in the morning and leave in the afternoon to feed offshore overnight (Würsig et al. 1994; Anderson 2005; Gannier and Petiau 2006), essentially on mesopelagic prey (Perrin et al. 1973; Dolar et al. 2003). Around the lagoon of Mayotte, in the Mozambique Channel (SW Indian Ocean), spinner dolphins primarily inhabit the outer slope of the barrier reef to rest and socialise, and rarely enter the lagoon (Kiszka et al. 2010a, 2011). During the past 20 years, extensive work has been conducted on the movements, behaviour, social, and genetic population structure of insular spinner dolphins, particularly in the Pacific Ocean (Norris et al. 1994; Karczmarski et al. 2005; Oremus et al. 2007; Andrews et al. 2010). These studies highlight that spinner dolphins may form “fission-fusion” societies, with groups forming and separating over short periods of times, such as around the big island of Hawaii (Karczmarski et al. 2005; Andrews et al. 2010). However, social structure may vary according to habitat characteristics and geographical isolation. Indeed, at the remote Midway atoll (Hawaii), spinner dolphins form stable groups with high level of site fidelity, limited emigration/immigration and strong inter-individual associations (Karczmarski et al. 2005). In this region, gene flow is more restricted among populations showing a fluid social structure (the Kona Coast of the island of Hawaii) than among populations with stable social groups (Midway and Kure Atolls) (Andrews et al. 2010).

In the southwestern Indian Ocean, the spinner dolphin is one of the most common small cetacean species in tropical and subtropical waters, particularly around islands and reef complexes off Zanzibar (Amir et al. 2002), Mayotte (Kiszka et al. 2010a, 2011) and the Comoros archipelago (Kiszka et al. 2010c), La Réunion (Dulau-Drouot et al. 2008, Condet and Dulau-Drouot, 2016), Madagascar (Rosenbaum 2003) and Mauritius (Webster et al. 2015). Spinner dolphins (particularly the *S. l. longirostris* form) are rarely observed in open ocean waters (> 2,000 m), but can occur for short periods of time between islands (particularly in island chains), mostly when undertaking overnight foraging trips (e.g. Kiszka et al. 2011, Mannocci et al. 2013, Thorne et al. 2012). As this species occurs in coastal and reef-associated waters, spinner dolphins are impacted by human activities, including past hunting and bycatch off Zanzibar (Stensland et al. 1998, Amir et al. 2012), direct hunting and bycatch in south-western Madagascar (Razafindrakoto et al. 2008) and disturbance from dolphin-watching activities such as on the west coast of Mauritius (Webster et al. 2015). However, the geographical extent of the influence of such direct and indirect effects on populations is unknown.

This study aims to characterise genetic diversity and population structure of spinner dolphins in the SW Indian Ocean, particularly from Zanzibar (Tanzania), Mayotte (Comoros archipelago) and La Réunion (Mascarene archipelago) (Fig. 1). These islands were selected because they are located along a west-east gradient, from continental waters of Africa (Zanzibar) to the most isolated and remote oceanic islands (La Réunion). We also characterise depth preferences of spinner dolphins at these sites to evaluate their reliance on shallow-water habitat and to estimate habitat size around the two oceanic islands. Resting habitat availability has been suggested to influence population size and dispersal in other island-associated spinner dolphin populations (Andrews et al. 2010). We evaluate the relationship between our estimations of genetic diversity

and habitat size: islands with more suitable habitats are expected to sustain greater population sizes, which would maintain greater genetic diversity. Geographic distances separating the coasts of the three sites are greater than 900 km. Considering previous knowledge gathered in the Pacific (Andrews et al. 2006; Oremus et al. 2007; Andrews et al. 2010), we hypothesised that geographic isolation is a major driver of the genetic population structure of spinner dolphins in this region, and that sampled islands should contain genetically distinct populations.

MATERIAL AND METHODS

Sample collection and DNA extraction

In total, tissue samples collected from 69 individual spinner dolphins off Zanzibar (n=21), Mayotte (n=29) and La Réunion (n=19) were available for this study (Fig. 1). For Zanzibar, muscle tissues were collected from bycaught animals in drift gillnets between 2000 and 2004 and were stored frozen at -20°C. Samples from Mayotte and La Réunion were collected during dedicated biopsy surveys undertaken in territorial waters from 2006 to 2011 from small boats. Biopsy attempts were made opportunistically, when groups and individuals were easily approachable and when conditions were optimal (Beaufort < 2, dolphins closely approaching the boat). Optimal weather conditions allowed stability of the research boat and better chances to sample the animals successfully and safely (Kiszka et al. 2010b). Blubber and skin biopsies were collected using a crossbow (BARNETT Veloci-Speed® Class, 68-kg draw weight and BARNETT Panzer V Class, 68-kg draw weight) with Finn Larsen (Ceta-Dart, Copenhagen, Denmark) bolts and tips (dart 20-mm long, 7-mm-diameter). Biopsy samples were preserved individually in 90% ethanol before shipping and subsequent analysis. Biopsy sampling was

conducted under French scientific permit #78/DAF/2004 (September 10, 2004) and permit #032/DAF/SEF/2008 (May 16, 2008) for Mayotte and MC/2009/336 for La Réunion. Genomic DNA was extracted from ~ 25 mg of tissue (muscle or skin) using a Nucleospin Tissue kit (Macherey-Nagel) following the manufacturer's protocol.

Microsatellite genotyping and mitochondrial DNA (mtDNA) sequencing
Twelve microsatellite loci previously optimized for *S. longirostris* were genotyped (Table 1). PCR reactions included ~20 ng of genomic DNA, 0.5 U Taq polymerase, 0.25 mM dNTPs, 1.5 mM MgCl₂, 1X PCR Buffer, 0.125 µM of each primer in a 20 µL final volume. PCR profiles were as follows: initial 5 min denaturation step at 94°C followed by 35 cycles of 30 s at 94°C, 30 s at a specific annealing temperature (see Table 1), 45 s at 72°C, and by a final 7-min extension step at 72°C. All PCRs were conducted in a Techne TC-5000 thermocycler. PCR products were visualized using polyacrylamide gels on the LICOR 4300 DNA Analyser. Allele sizes were determined by eye using a size standard, and by two different researchers to ensure consistency in scoring.

A portion of the mtDNA control region was amplified using primers Dlp-1.5 (5'-TCACCCAAAGCTGRARTTCTA-3') (Baker et al. 1998) and Dlp-8G (5'-GGAGTACTATGTCCTGTAACCA-3') (Dalebout et al. 2005). PCR reactions included 0.5 U Taq polymerase, 0.25 mM dNTPs, 1.5 mM MgCl₂, 1X PCR Buffer, 0.125 µM of each primer and ~50 ng of genomic DNA in a 50 µL reaction volume. PCRs were conducted in a Techne TC-5000 thermocycler using the following profile: initial 5 min denaturation step at 94°C followed by 35 cycles of 30 s at 94°C, 30 s at 54°C, 45 s at 72°C, and by a final 7-min extension step at

72°C. PCR products were purified and sequenced by Genoscreen (Lille, France). Sequences were edited in Chromas sequence viewer v. 2.33 (Chromas Technelysium) and were aligned using BioEdit v.5.0.6 (Hall 1999).

Microsatellite analyses

We tested for departures from Hardy-Weinberg or linkage equilibrium within each sampled site using Genepop v. 4.2 (Raymond and Rousset 1995) with 10,000 dememorizations, 1,000 batches and 10,000 iterations per batch. The sequential Bonferroni technique (Holm 1979) was applied to correct for multiple tests. The presence of null alleles and scoring errors was assessed using Micro-checker v. 2.2.3 (van Oosterhooft et al. 2004) within each site. We searched for potential duplicates within biopsied animals by comparing their multi-locus genotypes (i.e., searching for identical genotypes in the dataset, and for genotypes with less than three different alleles overall) using Genalex v. 6.41 (Peakall and Smouse 2006). The mitochondrial haplotypes of samples with matching microsatellite genotypes were compared to confirm they were duplicates of the same individual. We investigated whether related individuals were included in the dataset by calculating maximum-likelihood estimates of pairwise relatedness using ML-Relate (Kalinowski et al. 2006). To avoid biases in population inferences that could result from family structure (Anderson and Dunham 2008), we removed one individual from each pair of potential relatives, i.e., individuals showing a pairwise relatedness value greater than 0.45 (as in Viricel and Rosel 2014). Allele richness, observed and expected heterozygosity were calculated using FSTAT v. 2.9.3.2 (Goudet 1995) and Arlequin v. 3.5.1.2 (Excoffier and Lischer 2010), respectively. These molecular diversity indices were calculated for the whole dataset and for each site separately.

Population structure was assessed using a Bayesian approach implemented with Structure v. 2.3.4 (Pritchard et al. 2000), which infers the number of populations (K) present in a dataset based on assumptions of Hardy-Weinberg and linkage equilibria within populations. Analyses were conducted using the admixture and correlated allele frequencies models, with and without prior information on individual location (option “LOCPRIOR”). Including prior information on sample locations can improve population inferences, particularly when the level of population differentiation is weak or recent (Hubisz et al. 2009). To verify that using prior information did not artificially result in distinct clusters, we conducted additional Structure runs with the LOCPRIOR option after randomizing the sample location information in the input file. Three randomized input files were created. All Structure runs included 300,000 Markov chain Monte Carlo iterations and a 50,000 step burn in. Ten replicate runs were performed for K values between 1 and 5. Convergence was assessed by examining *alpha* and likelihood plots, and by comparing individual membership probabilities across replicate runs. The best K was chosen by comparing mean log probabilities among K values, and when $K = 1$ was ruled out, by applying Evanno’s method using ΔK (Evanno et al. 2005).

To assess genotypic variation among individuals and among the three locations, we applied a principal component analysis (PCA) to the microsatellite data using the package adegenet (Jombart 2008) in R v. 3.1.2 (R Core Team, 2015). In the PCA, allele frequencies were scaled using the centring option. An analysis of molecular variance (AMOVA, Excoffier et al. 1992) was conducted in Arlequin v. 3.5.1.2. (Excoffier and Lischer 2010) to estimate genetic differentiation among the three islands. Pairwise F_{ST} estimates were calculated and significance was assessed using 10,000 permutations. We tested for isolation by distance (IBD) by conducting

a Mantel test comparing pairwise genetic distances ($F_{ST}/(1-F_{ST})$, Rousset 1997) with log-transformed (base 10) geographic distances among sampling locations. Geographic distances between population pairs were calculated as the Euclidean distance between the approximate centres of the areas where samples were collected. The Mantel test was performed using IBDWS v. 3.23 (Jensen et al. 2005) with 10,000 randomizations. Finally, we investigated the occurrence of private alleles in each population identified using Genalex v. 6.41.

MtDNA sequence analyses

Diversity indices (haplotype and nucleotide diversities) were calculated for each site using Arlequin. We used JModeltest v. 0.1.1 (Guindon and Gascuel 2003; Posada 2008) and the Akaike Criterion to determine the most appropriate model of substitution given our sequence alignment. AMOVAs comparing the populations identified using Structure were performed in Arlequin. Genetic differentiation was measured using both F_{ST} and Φ_{ST} . For Φ_{ST} , distances between haplotypes were calculated using the model of substitution selected with JModeltest. Significance was assessed using 10,000 permutations. We evaluated IBD as described above for microsatellite data. A median-joining network was constructed using Network v.4.6.1.2 (Bandelt et al. 1999) with default parameters to represent relationships among haplotypes.

Habitat size

In order to relate patterns of genetic structure and availability of habitat used by spinner dolphins, we created spinner dolphin habitat maps based on their depth preferences around Zanzibar, Mayotte and La Réunion. From 2004 to 2008, sighting data were collected around Mayotte (n = 168 sightings in 224 days of survey) and La Réunion (n = 51 sightings in 278 days of survey)

during small-boat dedicated surveys (see Dulau-Drouot et al. 2008; Kiszka et al. 2011 for sampling protocols). For surveys undertaken around Mayotte and La Réunion, the sampling effort did not follow predefined transects and was not homogeneous. However, both surveys covered shallow inshore waters, the outer slope of the reef and deep oceanic habitats (depth > 500 m around Mayotte: Kiszka et al. 2011 and depth > 1,000 m around La Réunion: Dulau-Drouot et al. 2008). For Zanzibar, as sighting data were unavailable, we used geographic locations of bycatch events that were recorded during a bycatch monitoring program coordinated by the Institute for Marine Sciences, University of Dar es Salaam, based on Zanzibar between 2000 and 2007 (n = 27 records). These data were used to determine habitat preferences of spinner dolphins using depth as the main variable. We chose to focus habitat analyses on depth as it was previously identified as one of the main explanatory variables explaining spinner dolphin distribution patterns in this region (Kiszka et al. 2011, Condet & Dulau-Drouot 2016) and in other parts of the world (e.g. Thorne et al. 2012). Depth data were extracted from the GEBCO_2014 Gridded bathymetric dataset (30 arcsecond resolution) hosted on the British Oceanographic Data Center (<http://www.bodc.ac.uk>). We considered that the preferred habitat corresponded to the depth range where 95% of spinner dolphin observations were made. Thus, we excluded 2.5% of the deepest and 2.5% of the shallowest observations to determine i) the preferred depth range for Mayotte and La Réunion separately (we did not include Zanzibar in this comparison as observations from Zanzibar come from bycatch events and are thus not directly comparable to sighting data from the other two islands), ii) the overall preferred depth range for the three islands together. For the second calculation, group size information available for sightings around Mayotte was taken into account. Our reasoning was that using group size information (when available) better reflects the preferred habitat of these populations (i.e. observing large groups in an area carries more weight than observing a single individual). We

computed the total area available within the depth range limits obtained for the three islands together, and mapped these areas for Mayotte and La Réunion. It was not computed for Zanzibar since available habitat within this depth-range is virtually infinite along the East African continental shelf relative to the two islands. The retrieval of individual depths, the computation of projected surfaces and the mapping were all performed with the marmap package v0.9.2 (Pante and Simon-Bouhet 2013) in R v3.1.2.

RESULTS

Microsatellite data

Each sample was genotyped at 8 to 12 loci. No significant departure from Hardy-Weinberg or linkage equilibrium was observed (after sequential Bonferroni correction) in the three sampled sites. Furthermore, no scoring errors were identified using Micro-checker. Possible null alleles were detected for loci EV94 and 415/416 in Mayotte; however, since this issue was restricted to one site, we kept these two loci in all analyses. A duplicate sample was identified (100% matching genotypes and haplotype) in La Réunion, where the same individual was biopsied twice. Five pairs of potential relatives were detected using ML-Relate: two pairs in La Réunion, one pair in Zanzibar, and two pairs in Mayotte. Only one individual from each pair was kept in population analyses conducted with both types of markers (mtDNA and microsatellites) to avoid the potential bias of including relatives. The final microsatellite dataset included 63 individuals (Table 2). The number of alleles per locus ranged from six to 25 (Table 1). Observed heterozygosity and mean allele richness were similar among the three sites (Table 2).

No population subdivision was found when Structure was run without prior information (Fig. 2a). Conversely, genetic structure was detected when individual location was used as prior information. The most likely number of populations present in the dataset was three based on comparison of the mean log probabilities and ΔK (Fig. 2b), and corresponded to the three sampled sites (Fig. 2c): within each site, all individuals were assigned to the same cluster with high ancestry proportions ($q > 0.8$, mean $q = 0.95$). Randomizing sample locations in the Structure input file and applying the LOCPRIOR option resulted in no structure being detected by the program (most likely $K = 1$; see Appendix 1).

The first principal component of the PCA separated individuals from Mayotte from the other two locations, while the third principal component showed two separate clusters for individuals from La Réunion and Zanzibar (Appendix 2). Some overlap in the PC space was observed among the three clusters. Together, the first three principal components explained 12.7% of the total genotypic variation. The AMOVA conducted among the three populations revealed relatively weak, but significant genetic differentiation (Overall $F_{ST} = 0.020$ $p < 0.0001$; Table 3). The Mantel test conducted to examine IBD was not significant ($r = 0.21$, $R^2 = 0.04$, $p = 0.51$). Private alleles were detected within each of the three sites (Mayotte: 31; La Reunion: 9; Zanzibar: 11).

MtDNA data

The final control region sequence alignment was 720 bp-long and included 28 unique haplotypes (Genbank accession # XXXXXXXXX-XX). The sequence alignment included 42 substitutions and no indels. The model selected using JModeltest was Tamura-Nei (Tamura and Nei 1993) with a gamma correction ($\alpha = 0.726$). Analyses of mitochondrial sequences supported the population

structure detected using microsatellite loci: significant differences in mitochondrial haplotype frequencies (after sequential Bonferroni correction) were observed among all sites using F_{ST} (Overall $F_{ST} = 0.084$ $p < 0.0001$; Table 4). However, none of the pairwise comparisons were significant when distances among haplotypes were incorporated in the AMOVA (i.e., using Φ_{ST} , Table 4). The test for IBD was not significant ($r = 0.96$, $R^2 = 0.93$, $p = 0.17$). There was one shared haplotype between La Réunion and Zanzibar, and three shared haplotypes between Mayotte and Zanzibar. Mayotte and La Réunion had no haplotypes in common (Fig. 3). No obvious phylogeographic structure was observed on the haplotype network.

Habitat size

Comparing the depth distribution data of the two oceanic islands revealed that the preferred depth range of spinner dolphins around La Réunion (16 to 935 m) was larger than the one calculated for Mayotte (5 to 175 m). When all observation data are combined, the locations used by 95% of the individuals sighted around Zanzibar, Mayotte and La Réunion have a depth range of 9 to 162 m. The total surface available around Mayotte and La Réunion within this depth-range (9 to 162 m) was 1,036 and 327 km², respectively (Figure 4). The habitat surface corresponding to spinner dolphins' depth preferences is thus three times larger in Mayotte than in La Réunion.

DISCUSSION

Marine organisms with high dispersal capacities can show weak genetic structure across large geographic distances. For instance, the short-beaked common dolphin (*Delphinus delphis*) and

the Portuguese dogfish (*Centroscymnus coelolepis*) each form a single panmictic population across the eastern Atlantic (Veríssimo et al. 2011; Moura et al. 2013). However, even low levels of genetic differentiation can correspond to restricted levels of dispersal in a demographic sense and can be associated with adaptive divergence (e.g., Knutsen et al. 2011; Aykanat et al. 2015), and therefore represent important findings in terms of conservation and management.

In highly dispersive marine organisms, incorporating spatial information as prior information in genetic analyses can help reveal genetic differentiation (Selkoe et al. 2008), as illustrated in the present study. Indeed, we did not detect any structure when using a Bayesian approach to detect the number of genetically distinct populations based on microsatellite data alone. Including prior information about sample location in the Bayesian analysis allowed retrieving three populations, corresponding to the distinct islands that were sampled. Our results are in accordance with expectations of Latch et al. (2006) and Hubisz et al. (2009), as the F_{ST} estimates we obtained from analysis of microsatellite data fell right in the range where genetic structure can remain undetected without prior information ($0.01 < F_{ST} < 0.03$). This approach (Structure with LOCPRIOR: Hubisz et al. 2009) does not seem to falsely inflate genetic structure as shown in previous studies (e.g. Christie et al. 2010; Russello et al. 2012; Viricel 2012). Results from the three randomizations we conducted indicate the genetic clusters we observed using prior information are biologically significant (Appendix 1). Furthermore, the Mantel test comparing pairwise genetic and geographic distances was not significant, suggesting IBD did not confound our Structure results. The three distinct populations inferred from Structure and observed in the PCA based on nuclear data were supported by the significant genetic differentiation estimated with mitochondrial DNA sequences using F_{ST} . Possible explanations for the lack of significant differentiation observed using Φ_{ST} are that the observed genetic differentiation is recent and that

not enough time has passed for new mutations to accumulate within populations; or that migration rates among these populations are greater than the mutation rate of the mitochondrial DNA control region. In these cases, incorporating distances among haplotypes in the AMOVA can increase noise in the analysis (Bird et al. 2011), rendering Φ_{ST} less informative than F_{ST} .

The preferred depth range of spinner dolphins we inferred from sighting and bycatch data (i.e., for the three locations analysed together) confirmed that the species is associated with relatively shallow-water habitat at these locations, which corresponds to their resting grounds. The wider depth range observed for the population of La Réunion (also described in Dulau-Drouot et al. 2008) may reflect the very narrow continental shelf of the island, with depth increasing rapidly from the shore, compared to Mayotte. In fact, a recent habitat modelling study showed that despite the wide depth range of spinner dolphin observations around La Réunion, most sightings occur between 51 and 63 m of depth, within a “core habitat” also characterised by flat and light-colored seabeds (Condet & Dulau-Drouot 2016). Geographic isolation and the reliance of spinner dolphins on appropriate shallow-water resting habitat during daytime are likely factors causing and/or maintaining divergence between populations occupying Mayotte and La Réunion.

Although we cannot tease apart the relative role of each of these two factors, hypotheses can be made based on what has been observed in other small delphinid species. Indeed, large geographic distance from continental waters does not seem to be a sufficient driver to cause restricted gene flow in pelagic dolphins found around oceanic islands as illustrated by population structure studies on Atlantic spotted dolphin (*Stenella frontalis*) and common bottlenose dolphin (*T. truncatus*) populations from the Azores: the pelagic morphotypes within these two species form panmictic populations over large distances, from the Azores to offshore waters of the

northwestern Atlantic at least 4500 km away (Qurouil et al. 2007; Viricel & Rosel 2014). On the other hand, some pelagic island-associated delphinids show restricted gene flow across short distances within an archipelago such as common bottlenose dolphin populations around Hawaii (Martien et al. 2012), or the rough-toothed dolphin (*Steno bredanensis*), which displays long-term site fidelity around the Society Islands (French Polynesia), and fine-scale genetic differentiation between two islands only 170 km apart (Oremus et al. 2012). Thus, daily reliance on near-shore, insular habitats may be a predominant driver of population structure in pelagic delphinids, even more so than geographic isolation.

The genetic subdivision we observed is consistent with other studies conducted on *S. longirostris* in the Pacific Ocean, which indicated that insular populations are generally discrete. Levels of genetic differentiation were similar to those observed among Society (French Polynesia) and Hawaii archipelago (Oremus et al., 2007; Andrews et al. 2010). Spinner dolphins found around islands of the Society Archipelago (French Polynesia) form relatively closed communities showing strong island fidelity (Oremus et al. 2007). Gene flow among these communities is restricted, despite the short geographic distances separating some of these islands (i.e. tens of kilometers). Oremus et al. (2007) suggest that these communities are characterised by a metapopulation dynamics, which would explain the high genetic diversity and large island effective population sizes estimated from their molecular data. In the Hawaiian archipelago, fine-scale genetic structure is also observed, but patterns of gene flow vary according to social structure and habitat availability (Andrews et al. 2010). Overall, both the genetic and social structure of this species seems to be influenced by the availability and extent of resting areas (Karczmarski et al. 2005; Andrews et al. 2010). In Hawaiian populations, genetic differentiation increased with geographic distance among islands. In the present study, the tests for IBD we

conducted for both types of markers were not significant. We have to note, however, that these tests had low statistical power since only three populations were sampled. Thus, future studies analyzing samples from additional islands in the SW Indian Ocean would better allow testing for IBD. Additionally, photo-identification data would complement present findings, as site fidelity may constitute another factor driving population structure in spinner dolphins from the SW Indian Ocean.

The genetic diversity we measured (Table 2) was similar to what has been reported for this subspecies in French Polynesia (Oremus et al. 2007), and was greater than the diversity observed in Hawaii (Andrews et al. 2010). Despite the genetic differentiation of spinner dolphins from La Réunion and Mayotte, the genetic diversity of these two populations is similar to the diversity of spinner dolphins from the coast of Zanzibar. Factors influencing their population genetic diversity include effective population size (linked to drift) and immigration rates. Given the differences in habitat size among the three sites we compared, the respective local population size of the populations occupying these sites may differ. Thus, we hypothesize that the similar levels of genetic diversity we observe in these populations likely reflect low but recurrent gene flow, which may be sufficient to maintain genetic diversity within island communities. Spinner dolphins from the SW Indian Ocean could thus be under a metapopulation dynamics, similarly to what findings from Oremus et al. (2007) indicate for populations in the Society Archipelago. Alternatively, the genetic divergence we measured could be recent, and these populations may have so far retained ancestral polymorphisms.

The present study constitutes the first population structure assessment for the spinner dolphin in the SW Indian Ocean and our findings have important conservation implications. Indeed, the

habitat preferences and patterns of restricted gene flow we identified suggest spinner dolphin populations found off Mayotte and La Réunion are demographically independent from each other. Their differentiation makes them potentially vulnerable if directly impacted by human activities, and spinner dolphins found off these two islands should therefore be treated as two distinct conservation units at the national level. Analysing samples from other islands (e.g. Madagascar, Mauritius) within this region would allow further evaluating the genetic isolation of these populations. Bycatch levels of spinner dolphins in gillnets off Zanzibar (Amir et al. 2002) should be considered as a cause of concern, and this issue highlights the need of a population structure assessment along the east coast of Africa. Protecting important resting habitat is an important step toward the conservation of insular spinner dolphin populations. Dedicated surveys help assessing whether current marine protected areas encompass such habitats (e.g. in la Réunion: Dulau-Drouot et al. 2008) and habitat modelling studies (e.g. Thorne et al. 2012, Condet & Dulau-Drouot 2016) allow identifying key areas where new conservation efforts should be focused.

Acknowledgements

At Mayotte, samples and data from July 2004 to June 2006 were collected during a dolphin research project conducted by the *Office National de la Chasse et de la Faune Sauvage* (ONCFS, Game and Wildlife Service) and the Agriculture and Forestry Office (*Direction de l'Agriculture et de la Forêt*). From May 2007 to April 2009, data were collected during a joined programme of the University of La Rochelle, the *Office National de la Chasse et de la Faune Sauvage* and the *Collectivité Départementale de Mayotte*. DNA samples were prepared at the Molecular Core Facility at the University of La Rochelle. We would like to thank the Associate Editor and four anonymous reviewers for constructive comments on the manuscript.

471

472 **Compliance with ethical standards**

473

474 **Funding:** Funding was provided by the *Conseil Général de Mayotte*, the *Ministère de l'Energie*,
475 *l'Ecologie, le Développement Durable et de la Mer* (MEEDDM), the University of La Rochelle
476 and the *Office National de la Chasse et de la Faune Sauvage*. The samples and data collected in
477 Zanzibar was supported by funding grants from SIDA/Sarec. In Reunion, sampling was co-
478 funded by DEAL-Reunion, BNOI and IRD, as part of the Southwestern Indian Ocean Fisheries
479 Project (SWIOFP).

480 **Conflict of Interest:** The authors declare that they have no conflict of interest.

481 **Ethical approval:** All applicable international, national, and/or institutional guidelines for the
482 care and use of animals were followed.

483

References

- Amir OA, Berggren P, Jiddawi NS (2002) The incidental catch of dolphins in gillnet fisheries in Zanzibar, Tanzania. WIO J Mar Sci 1:155-162
- Amir OA, Berggren P, Jiddaw NS (2012) Recent records of marine mammals in Tanzanian waters. J. Cetacean Res. Manage. 12:249–253
- Amos B, Schlötterer C, Tautz D (1993) Social structure of pilot whales revealed by analytical DNA profiling. Science 260:670-672
- Anderson RC (2005) Observations of cetaceans in the Maldives, 1990-2002. J Cet Res Manag 7:119-135
- Anderson EC, Dunham KK (2008) The influence of family groups on inferences made with the program structure. Mol Ecol Resour 8:1219-1229
- Andrews KR et al (2006) Patterns of genetic diversity of the Hawaiian spinner dolphin (*Stenella longirostris*). Atoll Res Bull 543:65-73
- Andrews KR et al (2010) Rolling stones and stable homes: social structure, habitat diversity and population genetics of the Hawaiian spinner dolphin (*Stenella longirostris*). Mol Ecol 19:732-748
- Aykanat T, Johnston SE, Orell P, Niemelä E, Erkinaro J, Primmer CR. 2015. Low but significant genetic differentiation underlies biologically meaningful phenotypic divergence in a large Atlantic salmon population. Mol Ecol 24:5158-5174
- Baker C et al (1998) Mitochondrial DNA variation and maternal gene flow among humpback whales of the southern hemisphere. Mar Mamm Sci 14:721-737.
- Bandelt HJ, Forster P, Rohl A (1999) Median-joining networks for inferring intraspecific phylogenies. Mol Biol Evol 16:37-48.

508 Bérubé M, Jorgensen H, McEwing R, Palsbøll PJ (2000) Polymorphic di-nucleotide
 509 microsatellite loci isolated from the humpback whale, *Megaptera novaeangliae*. Mol Ecol
 510 9:2181-2182

511 Bird CE, Karl SA, Smouse PE, Toonen RJ (2011) Detecting and measuring genetic
 512 differentiation. In: Koenemann S, Held C, Schubart CD (ed) Crustacean Issues:
 513 Phylogeography and Population Genetics in Crustacea, 1st edn. CRC Press, Boca Raton, pp
 514 31-55

515 Caldwell M, Gaines MS, Hughes CR (2002) Eight polymorphic microsatellite loci for bottlenose
 516 dolphin and other cetacean species. Mol Ecol Notes 2:393-395

517 Christie MR, Johnson DW, Stallings CD, Hixon MA (2010) Self-recruitment and sweepstakes
 518 reproduction amid extensive gene flow in a coral-reef fish. Mol Ecol 19:1042-1057

519 Ciannelli L et al (2010) Small-scale genetic structure in a marine population in relation to water
 520 circulation and egg characteristics. Ecology 91(10):2918-2930

521 Condet M, Dulau-Drouot V (2016) Habitat selection of two island-associated dolphin species
 522 from the south-west Indian Ocean. Cont Shelf Res 125:18-27

523 Cowen RK, Lwiza MMK, Sponaugle S, Paris CB, Olson DB (2000) Connectivity of marine
 524 populations: Open or closed? Science 287:857-859

525 Cowen RK, Gawarkiewicz G, Pineda J, Thorrold SR, Werner FE (2007) Population connectivity
 526 in marine systems: An overview. Oceanography 20:14-21

527 Dalebout ML et al (2005) Worldwide structure of mtDNA diversity among Cuvier's beaked
 528 whales (*Ziphius cavirostris*): implications for threatened populations. Mol Ecol 14:3353-
 529 3371

530 Dolar MLL, Walker WA, Kooyman GL, Perrin WF (2003) Comparative feeding ecology of
 531 spinner dolphins (*Stenella longirostris*) and Fraser's dolphins (*Lagenodelphis hosei*) in the
 532 Sulu Sea. Mar Mamm Sci 19:1-1
 533 Dulau-Drouot V, Boucaud, V, Rota B (2008) Cetacean diversity off La Réunion Island (France).
 534 J M Biol Ass UK 88:1263-1272
 535 Evanno G, Regnaut S, Goudet J (2005) Detecting the number of clusters of individuals using the
 536 software STRUCTURE: a simulation study. Mol Ecol 14:2611-2620
 537 Excoffier L, Lischer HE (2010) Arlequin suite ver 3.5: a new series of programs to perform
 538 population genetics analyses under Linux and Windows. Mol Ecol Resour 10:564-567
 539 Excoffier L, Smouse P, Quattro J (1992) Analysis of molecular variance inferred from metric
 540 distances among DNA haplotypes: Application to human mitochondrial DNA restriction
 541 data. Genetics 131:479-491
 542 Fontaine MC et al (2007) Rise of oceanographic barriers in continuous populations of a cetacean:
 543 the genetic structure of harbour porpoises in Old World waters. BMC Biol 5:30
 544 Fullard KJ, Early G, Heide-Jørgensen MP, Bloch D, Rosing-Asvid A, Amos W (2000)
 545 Population structure of long-finned pilot whales in the North Atlantic: a correlation with sea
 546 surface temperature? Mol Ecol 9:949-958
 547 Gannier A, Petiau E (2006) Environmental variables affecting the residence of spinner dolphins
 548 (*Stenella longirostris*) in the Bay of Tahiti (French Polynesia). Aquat Mamm 32:202-211
 549 Goudet J (1995) FSTAT (version 1.2.): a computer program to calculate *F*-statistics. J Hered
 550 86:485-486
 551 Guindon S, Gascuel O (2003) A simple, fast, and accurate algorithm to estimate large
 552 phylogenies by maximum likelihood. Syst Biol 52:696-704

553 Hall T (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program
 554 for Windows 95/98/NT. Nucl Acid S 41:95-98
 555 Halpern BS et al (2008) A global map of human impact on marine ecosystems. Science 319:948-
 556 952
 557 Hoelzel AR (2009) Evolution of population genetic structure in marine mammal species. In:
 558 Bertorelle G, Bruford MW, Hauffe HC, Rizzoli A, Vernesi C (eds) Cambridge University
 559 Press, New York, pp 294-318
 560 Hoelzel AR, Dahlheim M, Stern SJ (1998) Low genetic variation among killer whales (*Orcinus*
 561 *orca*) in the eastern North Pacific and genetic differentiation between foraging specialists. J
 562 Hered 89:121-128
 563 Holm S (1979) A simple sequentially rejective multiple test procedure. Scand J Stat 6:65-70
 564 Hubisz MJ, Falush D, Stephens M, Pritchard J (2009) Inferring weak population structure with
 565 the assistance of sample group information. Mol Ecol Resour 9:1322-1332
 566 Jensen JL, Bohonak AJ, Kelley ST (2005) Isolation by distance, web service. BMC Genetics
 567 6:13. V.3.23 <http://ibdws.sdsu.edu/>
 568 Jombart T. (2008) adegenet: a R package for the multivariate analysis of genetic markers.
 569 Bioinformatics 24:1403-1405
 570 Kalinowski ST, Wagner AP, Taper ML (2006) ML-Relate: a computer program for maximum-
 571 likelihood estimation of relatedness and relationship. Mol Ecol Notes 6:576-579
 572 Karczmarski L, Würsig B, Gailey G, Larson KW, Vanderlip C (2005) Spinner dolphin in a
 573 remote Hawaiian atoll: social grouping and population structure. Behav Ecol 16:675-685
 574 Kiszka J et al (2008) Marine mammal bycatch in the Southwest Indian Ocean: Review and need
 575 for a comprehensive status assessment. Western Indian Ocean J Mar Sci 7:119-136

576 Kiszka J, Ersts PJ, Ridoux V (2010a) Structure of a toothed cetacean community around a
577 tropical island (Mayotte, Mozambique Channel). *Afr J Mar Sci* 32(3):543-551

578 Kiszka J, Simon-Bouhet B, Charlier F, Pusineri C, Ridoux V (2010b) Individual and group
579 behavioural reactions of small delphinids to remote biopsy sampling. *Anim Welfare* 19:411-
580 417

581 Kiszka J, Breysse O, Vely M (2010c) Preliminary account of cetacean diversity and humpback
582 whale (*Megaptera novaeangliae*) group characteristics around the Union of the Comoros
583 (Mozambique Channel). *Mammalia* 74:51-56

584 Kiszka J, Simon-Bouhet B, Martinez L, Pusineri C, Richard P, Ridoux V (2011) Ecological niche
585 segregation within a community of sympatric dolphins around a tropical island. *Mar Ecol*
586 *Prog Ser* 433:273-288

587 Knutsen H, Olsen EM, Jorde PE, Espeland SH, André C, Stenseth NC. 2011. Are low but
588 statistically significant levels of genetic differentiation in marine fishes ‘biologically
589 meaningful’? A case study of coastal Atlantic cod. *Mol Ecol* 20:768-783

590 Krützen M, Valsecchi E, Connor RC, Sherwin WB (2001) Characterization of microsatellite loci
591 in *Tursiops aduncus*. *Mol Ecol Notes* 1:170-172

592 Latch EK, Dharmarajan G, Glaubitz JC, Rhodes OE Jr (2006) Relative performance of Bayesian
593 clustering software for inferring population substructure and individual assignment at low
594 levels of population differentiation. *Conserv Genet* 7:295-302

595 Louis M et al (2014a) Ecological opportunities and specializations shaped genetic divergence in a
596 highly mobile marine top predator. *Proc R Soc B* 281:20141558

597 Louis M et al (2014b) Habitat-driven population structure of bottlenose dolphins, *Tursiops*
598 *truncatus*, in the North-East Atlantic. *Mol Ecol* 23:857-874

599 Mannocci L et al (2014) Predicting top predator habitats in the Southwest Indian Ocean.
600 Ecography 37(3):261-278

601 Martien KK et al (2012) Population structure of island-associated dolphins: evidence from
602 mitochondrial and microsatellite markers for common bottlenose dolphins (*Tursiops*
603 *truncatus*) around the main Hawaiian Islands. Mar Mamm Sci 28:E208-E232

604 Mendez M et al (2013). Integrating multiple lines of evidence to better understand the
605 evolutionary divergence of humpback dolphins along their entire distribution range: a new
606 dolphin species in Australian waters? Mol Ecol 22: 5936-5948.

607 Mendez M et al (2011) Molecular ecology meets remote sensing: environmental drivers to
608 population structure of humpback dolphins in the Western Indian Ocean. Heredity 107:349-
609 361.

610 Möller LM, Valdez FP, Allen S, Bilgmann K, Corrigan S, Beheregaray LB (2011) Fine-scale
611 genetic structure in short-beaked common dolphins (*Delphinus delphis*) along the East
612 Australian Current. Mar Biol 158:113-126

613 Möller LM, Wiszniewski J, Allen SJ, Beheregaray LB (2007) Habitat type promotes rapid and
614 extremely localised genetic differentiation in dolphins. Mar Freshwat Res 58:640-648

615 Moura AE, Natoli A, Rogan E, Hoelzel AR (2013) Atypical panmixia in a European dolphin
616 species (*Delphinus delphis*): implications for the evolution of diversity across oceanic
617 boundaries. J Evol Biol 26:63-75

618 Norris KS, Würsig B, Wells RS, Würsig M (1994) The Hawaiian Spinner Dolphin, University of
619 California Press, Berkeley, 408 pp.

620 van Oosterhooft C, Hutchinson W, Wills D, Shipley P (2004) MICRO-CHECKER: software for
621 identifying and correcting genotyping errors in microsatellite data. Mol Ecol Notes 4:535-
622 538

623 Oremus M, Poole MM, Steel D, Baker CS (2007) Isolation and interchange among insular
624 spinner dolphin communities in the South Pacific revealed by individual identification and
625 genetic diversity. *Mar Ecol Prog Ser* 336:275-289

626 Oremus M, Poole MM, Albertson GR, Baker CS (2012) Pelagic or insular? Genetic
627 differentiation of rough-toothed dolphins in the Society Islands, French Polynesia. *J Exp Mar*
628 *Biol Ecol* 423-433:37-46

629 Palsbøll PJ, Bérubé M, Larsen AH, Jørgensen H (1997) Primers for the amplification of tri- and
630 tetramer microsatellite loci in baleen whales. *Mol Ecol* 6:893-895

631 Pante E, Simon-Bouhet B (2013) marmap: A package for importing, plotting and analyzing
632 bathymetric and topographic data in R. *PLoS ONE* 8(9): e73051.
633 doi:10.1371/journal.pone.0073051

634 Peakall R, Smouse PE (2006) GENALEX 6 : genetic analysis in Excel. Population genetic
635 software for teaching and research. *Mol Ecol Notes* 6:288-295

636 Pelc RA, Warner RR, Gaines SD (2009) Geographical patterns of genetic structure in marine
637 species with contrasting life histories. *J Biogeogr* 36:1881-1890

638 Perrin WF (2009) Spinner dolphin *Stenella longirostris*. In: Perrin WF, Würzig B, Thewissen J
639 (ed) *Encyclopedia of Marine Mammals*, 2nd edn. Academic Press, San Diego, pp 1100-1103

640 Perrin WF, Dolar M, Louella L, Robineau D (1999). Spinner dolphins (*Stenella longirostris*) of
641 the western Pacific and Southeast Asia: pelagic and shallow-water forms. *Mar Mamm Sci*
642 15(4):1029-1053

643 Perrin WF, Gilpatrick JW Jr. (1994) Spinner dolphin *Stenella longirostris* (Gray, 1828). In:
644 Ridway SH and Harrison R (eds) *Handbook of marine mammals*, Academic Press, London,
645 pp 99-128.

646 Perrin WF, Warner RR, Fiscus CH, Holtz DB (1973) Stomach content of porpoise, *Stenella* spp.,
 647 and yellowfin tuna, *Thunnus albacares*, in mixed-species aggregations. Fish Bull 71:1077-
 648 1092
 649 Posada D (2008) jModelTest : Phylogenetic model averaging. Mol Biol Evol 25:1253-1256
 650 Pritchard JK, Stephens M, Donnelly P (2000) Inference of population structure using multilocus
 651 genotype data. Genetics 155:945-959
 652 Quérrouil S et al (2007) High gene flow in oceanic bottlenose dolphins (*Tursiops truncatus*) of the
 653 North Atlantic. Conserv Genet 8:1405-1419
 654 R Core Team (2015) R: A language and environment for statistical computing. R Foundation for
 655 Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>
 656 Raymont M, Rousset F (1995) Genepop (version 1.2), population genetics software for exact
 657 tests and ecumenicism. J Hered 86:248-249
 658 Razafindrakoto Y, Andrianarivelo N, Cerchio S, Rasoamananto I, Rosenbaum H (2008)
 659 Preliminary assessment of cetacean incidental mortality in artisanal fisheries in Anakao,
 660 southwestern region of Madagascar. WIO J Mar Sci 7(2):175-184
 661 Rosel PE, France SC, Wang JY, Kocher TD (1999) Genetic structure of harbour porpoise
 662 *Phocoena phocoena* populations in the northwest Atlantic based on mitochondrial and
 663 nuclear markers. Mol Ecol 8:541-554
 664 Rosenbaum HC (2003) Marine mammals of Madagascar. In: Godman S, Bengston J (eds), The
 665 natural history of Madagascar, University of Chicago Press, Chicago, pp 213-216
 666 Rousset F (1997) Genetic differentiation and estimation of gene flow from *F*-statistics under
 667 isolation by distance. Genetics 145:1219-1228
 668 Russello MA, Kirk SL, Frazer KK, Askey PJ (2012) Detection of outlier loci and their utility for
 669 fisheries management. Evol Appl 5:39-52

670 Selkoe KA, Henzler CM, Gaines SD (2008) Seascape genetics and the spatial ecology of marine
671 populations. *Fish Fish* 9:363-377

672 Stensland E, Berggren P, Johnstone R, Jiddawi N (1998) Marine mammals in Tanzanian waters:
673 urgent need for status assessment. *Ambio* 27: 771-774

674 Tamura K, Nei M (1993) Estimation of the number of nucleotide substitutions in the control
675 region of mitochondrial DNA in human and chimpanzees. *Mol Biol Evol* 10:512-526

676 Thorne LH et al (2012) Predictive modeling of spinner dolphin (*Stenella longirostris*) resting
677 habitat in the main Hawaiian Islands. *PLoS One* 7(8) :e43167

678 Valsecchi E, Amos W (1996) Microsatellite markers for the study of cetacean populations. *Mol*
679 *Ecol* 5:151-156

680 Veríssimo A, McDowell JR, Graves JE (2011) Population structure of a deep-water squaloid
681 shark, the Portuguese dogfish (*Centroscymus coelolepis*). *ICES J Mar Sci* 68:555-563

682 Viricel A (2012) Using genetics to assess population structure in three cetacean species and to
683 investigate the etiology of cardiomyopathy in *Kogia breviceps*. Dissertation, University of
684 Louisiana at Lafayette

685 Viricel A, Rosel PE (2014) Hierarchical population structure and habitat differences in a highly
686 mobile marine species: the Atlantic spotted dolphin. *Mol Ecol* 23:5018-5035

687 Webster I, Cockcroft VG, Cadinouche A (2015) Spinner dolphins *Stenella longirostris* off south-
688 west Mauritius: abundance and residency. *Afr J Mar Sci* 37(1):115-124

689 Würsig B, Wells RS, Norris KS, Würsig M (1994) A spinner dolphin's day. In: Norris KS,
690 Würsig B, Wells RS, Würsig M (eds) *The Hawaiian Spinner Dolphin* London, University of
691 California Press, pp 65-102

692

693

694

695

696

Tables

Table 1. Twelve microsatellite loci genotyped in this study. PCR annealing temperature (Ta), reference, number of alleles, observed (H_o) and expected (H_e) heterozygosity are given for each locus.

Locus	Ta (°C)	Reference	Number of alleles	H_o	H_e
415/416	45	Amos et al. 1993	12	0.772	0.889
GT575	60	Bérubé et al. 2000	10	0.679	0.778
GT6	60	Caldwell et al. 2002	6	0.635	0.676
AAT44	55	Caldwell et al. 2002	10	0.770	0.710
KWM12a	46	Hoelzel et al. 1998	10	0.889	0.844
MK5	50	Krützen et al. 2001	13	0.786	0.912
MK6	50	Krützen et al. 2001	18	0.879	0.931
GATA98	54	Palsbøll et al. 1997	9	0.655	0.804
PPHO142	50	Rosel et al. 1999	9	0.804	0.810
PPHO131	57	Rosel et al. 1999	12	0.755	0.834
EV1	47	Valsecchi and Amos 1996	16	0.868	0.902
EV94	54	Valsecchi and Amos 1996	25	0.772	0.925

Table 2. Mitochondrial DNA (mtDNA) and microsatellite diversity indices for *Stenella longirostris* from each site: N, sample size; No. haplo, number of haplotypes; π , nucleotide diversity; h , haplotype diversity; AR, allele richness; H_o , observed heterozygosity; H_e , expected heterozygosity

	mtDNA				Microsatellites			
	N	No. haplo	π	h	N	Mean AR	H_o	H_e
Zanzibar	20	9	0.013	0.826	20	7.1	0.747	0.790
Mayotte	19	14	0.015	0.965	27	8.2	0.773	0.835
La Réunion	16	9	0.012	0.900	16	7.5	0.798	0.818

Table 3. AMOVA results obtained from analysis of microsatellite data from 63 *S. longirostris* individuals. Pairwise F_{ST} values are shown below diagonal and corresponding p-values above diagonal.

	Zanzibar	Mayotte	La Réunion
Zanzibar	NA	<0.001*	<0.001*
Mayotte	0.024	NA	0.009*
La Réunion	0.025	0.013	NA

*significant p-value after sequential Bonferroni correction

Table 4. AMOVA results for mitochondrial DNA sequences. Pairwise F_{ST} and Φ_{ST} values are shown below and above diagonal, respectively.

	Zanzibar	Mayotte	La Réunion
Zanzibar	NA	0.050	0.014
Mayotte	0.058*	NA	0.066
La Réunion	0.130*	0.067*	NA

*significant p-value after sequential Bonferroni correction

Figure Legends

Figure 1. Study area and sample locations. Sample sizes are indicated for each site. The 200 m and 1,000 m isobaths are represented by darker lines.

Figure 2. Bayesian clustering analysis (Structure) results obtained from analysis of 12 microsatellite loci (a) without any prior information and (b) using prior information about sample location (“LOCPRIOR” option). The mean log probability ($\text{LnP}(K)$) is given for each K tested and the ΔK from Evanno’s method is shown between successive K values. (c) The barplot represents individual ancestry proportions for the three populations obtained using the “LOCPRIOR” option.

Figure 3. Median-joining network of 28 mitochondrial control region haplotypes observed in 63 *S. longirostris* individuals. Filled circles represent haplotypes and their size is proportional to their frequency in the dataset. Circles are shaded in colours proportionally to the number of individuals from each population (Mayotte: blue; La Reunion: green; Zanzibar: red). Unsamplred or extinct intermediate haplotypes are shown as black dots. Each line corresponds to one mutational step, except when a number of mutations is adjacent to it.

Figure 4. Maps showing the surface available for daytime resting for spinner dolphins around Mayotte (top) and La Réunion (bottom). These surfaces were estimated based on the preferred depth-range of spinner dolphins in the SW Indian Ocean. Maps include all sightings, including outlier observations that were not used to determine the preferred depth range (see Material and Methods).

739 **Figures**

740 Figure 1.

741

744 Figure 3

745

746 Figure 4

747

748