



Landscape heterogeneity related to organic and conventional farming systems: effects on natural enemies of crop pests and pest predation levels

Stéphanie Aviron, Alexandre Monteiro, Jacques Baudry, El Aziz Djoudi, Julien Petillon, Manuel Plantegenest, Camille Puech

► To cite this version:

Stéphanie Aviron, Alexandre Monteiro, Jacques Baudry, El Aziz Djoudi, Julien Petillon, et al.. Landscape heterogeneity related to organic and conventional farming systems: effects on natural enemies of crop pests and pest predation levels. International conference on Ecological Sciences (SFE Ecologie 2016), Oct 2016, Marseille, France. , 108 p., 2016, International Conference on Ecological Sciences - Sféologie 2016. hal-01458669

HAL Id: hal-01458669

<https://hal.science/hal-01458669>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SFE 2016 - International conference on ecological sciences

Proposition : oral communication - Theme : Agricultural Ecology

Landscape heterogeneity related to organic and conventional farming systems: effects on natural enemies of crop pests and pest predation levels

Aviron S.^a, Monteiro, A.^a, Baudry J.^a, Djoudi A.^b, Pétilion J.^c, Plantegenest M.^b, Poggi S.^b, Puech C.^{a,b}

^a INRA, UR 980, SAD-Paysage, F-35042 Rennes Cedex, France

^b INRA, UMR1349 IGEPP, F-35653 Le Rheu, France

^c Université de Rennes 1, EA7316 Biodiversité et Gestion des Territoires, 263 Avenue du Général Leclerc CS 74205, Campus de Beaulieu, F-35042 Rennes Cedex, France

Abstract

Organic farming is considered a promising type of production to enhance biological control of pests by their natural enemies, and therefore, to reduce pesticide use in modern agricultural systems. The effects of organic farming on natural enemies have been largely studied at the field scale, but the role of landscape heterogeneity related to the amount and spatial configuration of organic farming remains little explored. Moreover, little is known about local and landscape effects of farming systems on pest predation by natural enemies. The aim of our study was to investigate the effects of farming systems (organic vs. conventional) at the field and landscape scales, on the diversity of natural enemies and on prey predation levels in cereal crops. Natural enemies (predatory carabid beetles and ladybirds) and predation levels of sentinel preys (aphids and weeds) were sampled in 20 pairs of organic-conventional cereal crops located in 20 landscapes with varying percent cover of organic farming, in western France. Our results show that the diversity of natural enemies and pest predation levels were mainly affected by the type of farming system at the field scale, and were overall higher in organic crops. They were not influenced by the amount and spatial configuration of farming systems at the landscape scale, but were influenced by the amount of woody habitats (hedgerows). Our findings suggest that the promotion of biological control in hedgerow network landscapes mainly relies on the adoption of organic practices and, to a lesser extent, on the conservation of hedgerows.

Keywords : Organic farming, landscape context, carabid beetles, ladybirds, aphid and weed predation, hedgerows.