Oxidation of a PAH polluted soil using modified Fenton reaction in unsaturated condition affects biological and physico-chemical properties

Fabien Laurent, Aurélie Cébron, Christophe Schwartz, Corinne Leyval

► To cite this version:

Fabien Laurent, Aurélie Cébron, Christophe Schwartz, Corinne Leyval. Oxidation of a PAH polluted soil using modified Fenton reaction in unsaturated condition affects biological and physico-chemical properties. *Chemosphere*, 2012, 86, pp.659-664. 10.1016/j.chemosphere.2011.11.018 . hal-01486347

HAL Id: hal-01486347

<https://hal.science/hal-01486347>

Submitted on 27 Feb 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Oxidation of a PAH polluted soil using modified Fenton reaction in unsaturated
condition affects biological and physico-chemical properties**

Laurent F^{1,2}, Cébron A¹, Schwartz C², Leyval C^{1*}

¹ LIMOS UMR7137 Nancy Université CNRS, BP 70239, F-54506 Vandœuvre-lès-Nancy
cedex, France

² LSE UMR 1120 Nancy Université INRA, BP 172, F-54505 Vandœuvre-lès-Nancy cedex,
France

*Corresponding author: Corinne Leyval, LIMOS UMR7137 Nancy Université CNRS, Faculté
des Sciences et Technologies, BP 70239, 54506 Vandoeuvre-les-Nancy Cedex, France, tel: 33
(0)3 83 68 42 82, fax: 33 (0)3 83 68 42 84, e-mail: corinne.leyval@limos.uhp-nancy.fr

Abstract

A batch experiment was conducted to assess the impact of chemical oxidation using modified Fenton reaction on PAH content and on physico-chemical and biological parameters of an industrial PAH contaminated soil in unsaturated condition. Two levels of oxidant (H_2O_2 , 6 and 65 g kg^{-1}) and FeSO_4 were applied. Agronomic parameters, bacterial and fungal density, microbial activity, seed germination and ryegrass growth were assessed. Partial removal of PAHs (14 and 22%) was obtained with the addition of oxidant. The impact of chemical oxidation on PAH removal and soil physico-chemical and biological parameters differed depending on the level of reagent. The treatment with the highest concentration of oxidant decreased soil pH, cation exchange capacity and extractable phosphorus content. Bacterial, fungal, and PAH degrading bacteria densities were also lower in oxidized soil. However a rebound of microbial populations and an increased microbial activity in oxidized soil were measured after 5 weeks of incubation. Plant growth on soil treated by the highest level of oxidant was negatively affected.

Keywords: Fenton, oxidation, PAH, soil, fertility, biological activity

1. Introduction

The presence of pollutants in soils can affect soil functions (e.g. filter and exchange, support for plant growth and biodiversity) (Smith et al., 2006; Maxim and Spangenberg, 2009). Although remediation treatments reduce or eliminate pollutant concentration, their potential effect on soil physical, chemical and biological properties, and soil functions is rarely considered. Polycyclic aromatic hydrocarbons (PAHs) are widespread contaminants in soil.

41 Their effects on human health and their carcinogenic potential is largely studied (Bispo et al.,
42 1999; Joe et al., 2004; White and Claxton, 2004). Different remediation technologies were
43 developed in the last decades to remove PAHs from contaminated soils (Wang et al., 1990;
44 Cajthaml et al., 2002; Gan et al., 2009). *In situ* chemical oxidation (ISCO) is a technology
45 generally used to remediate organic contaminants in groundwater and saturated soil (Rivas,
46 2006; Huling and Pivetz, 2006). This remediation technique based on the injection of
47 different oxidant reagents does not require soil excavation and represents an alternative to
48 conventional treatment (*e.g.* thermal desorption, venting). Fenton reagents (hydrogen
49 peroxide and iron) are among the most commonly used and studied oxidizing agents (Rivas,
50 2006) for pollutants such as petroleum hydrocarbons, BTEX, chlorinated ethenes and PAHs
51 (Huling and Pivetz, 2006). In Fenton reaction, Fe (II) catalyzes the decomposition of
52 hydrogen peroxide to generate hydroxyl radicals ($\text{OH}^\bullet$) which are strong and non specific
53 oxidants. However, some authors showed that $\text{OH}^\bullet$ radicals preferentially attack the aromatic
54 compounds (Westerhoff et al., 1999; Mikutta et al., 2005) and can also react with sorbed and
55 aqueous PAHs. Fenton oxidation generally includes acid addition since a low pH (2-3) is
56 required for optimum reaction, but modified Fenton reaction at near neutral pH using ferric
57 ions and chelating agents was also performed (Nam et al., 2001).

58 Fenton reaction was mainly studied in saturated conditions (Jonsson et al., 2006; Sirguey et
59 al., 2008; Sun and Yan, 2008; Valderrama et al., 2009) with large liquid/solid ratios. Further,
60 the amounts of oxidant used in many studies (Nam et al., 2001; Palmroth et al., 2006a;
61 Valderrama et al., 2009) were very high for field treatment. Chemical oxidation in unsaturated
62 conditions was also applied, although fewer studies were reported (Palmroth et al. 2006a,
63 Huling and Pivetz, 2006). The addition of chemical oxidants could impact other soil
64 characteristics than organic pollutant concentration and for surface soils this could lead to a
65 global loss of fertility. Chemical oxidation of PAHs was shown to induce a decrease of

organic carbon, total nitrogen and phosphorus contents in soils (Leifeld and Kögel-Knabner, 2001; Mikutta et al., 2005; Sirguey et al., 2008). Microorganism populations were negatively affected following ISCO but the impact seemed to be temporary with a rebound of cell numbers after a few weeks (Sahl and Munakata-Marr, 2006). After oxidation treatment, PAH biodegradation was enhanced, suggesting the possible combination of both chemical oxidation and biodegradation of organic pollutants (Lee and Hosomi, 2001; Nam et al., 2001; Palmroth et al., 2006a; 2006b; Valderrama et al., 2009)

The present study was conducted to assess the impact of chemical oxidation to surface soil using modified Fenton reaction on PAH removal and on soil physico-chemical and biological parameters. Since surface soils are mostly unsaturated, we performed the oxidation in unsaturated condition. An industrial PAH contaminated soil was treated with a volume of reagents calculated to keep the soil at water holding capacity. We used a modified Fenton reaction without acid addition to the soil to avoid a strong pH decrease that may drastically affect biological activity. Two concentrations of reagents were tested. Analyses of PAH concentration, soil characteristics (pH, cation exchange capacity, organic carbon, nitrogen, phosphorus), microbial density and activity and seed germination and plant growth using ryegrass (*Lolium perenne*) were performed with oxidized and control soil.

2. Materials and methods

2.1. Soil sampling and analysis

An industrial soil sample was collected from a former coking plant site (Neuves-Maisons, France) (Biache et al., 2008; Monsérié et al., 2009). The soil is a sandy loamy soil (11.9% clay, 21.5% silt and 66.6% sand). The soil presents a high content of organic carbon (50g kg⁻¹), a neutral pH (7.3) and the contamination by the 16 US-EPA priority PAHs reaches

1200 mg kg⁻¹. Large amount of soil was excavated and homogenized by quartering. Sub-samples were sieved at 2 mm and air dried at 25°C. Soil analyses were performed by INRA laboratory (Arras, France) according to international standard methods: water holding capacity (WHC) (ISO 11464), pH, total organic carbon (ISO 10694), total nitrogen (ISO 13878), extractable phosphorus (ISO 11263), cation exchange capacity (CEC, NF X 31-130).

2.2. Oxidation experiment

The modified Fenton reaction was performed with two levels of reagents corresponding to 6 g kg⁻¹ (low, L) and 65 g kg⁻¹ (high, H) of H₂O₂. Three glass containers were filled with 1500 g of 2 mm sieved soil at 5 % humidity. In two of them, the soil sample was mixed with 7 (L) or 82 (H) g of solid ferrous sulfate (FeSO₄, 7H₂O, 99.5% purity) to catalyse the decomposition of H₂O₂ without increasing the volume of solution added. Then, 450 mL of a solution of H₂O₂ at either 0.54 (L) or 6.3 M (H) were added gradually to reach water holding capacity. The third one was a control with addition of 450 mL of water and no reagent addition. After 48 h, the soil sample in each container was homogenized and sub-sampled for further analysis. The soil was then air dried and sieved to 2 mm again. For microbial analysis, sub-samples were directly used for quantification of culturable heterotroph bacteria and incubation. For DNA extraction sub-samples were stored at -20°C.

2.3. Measurement of microbial activity (carbon mineralization) in soil after oxidation

Fenton oxidized and control soil samples (20 g), with five replicates each, were incubated at 100% WHC in 150 mL serum bottles with a Teflon septum for 5 weeks at 24°C to measure microbial activity by CO₂ production. To avoid anaerobic conditions during incubation, bottles were aired out every week. Carbon dioxide (CO₂) release was measured every day by

infrared spectrophotometer (Bios, absorption at 2325.6 cm^{-1}) (Quantin et al., 2005). After 5 weeks, five replicates were taken for microbial and PAH analysis as described hereafter. An abiotic control of each modality (autoclaved twice at an interval of 48 h) was performed for a week in order to evaluate the abiotic release of CO_2 .

2.4. MPN counts of heterotroph microorganisms

Soil (1 g fresh) was suspended in 10 mL sterile ultra-pure water with 6 g sterile glass-beads and soil suspension was shaken during 1 h. Serial dilution of soil suspension (10^{-2} to 10^{-7}) was prepared in NaCl (0.85%). The most-probable-number (MPN) was counted in 96-well microplates. A volume (25 μL) of dilution was added with 200 μL nutrient broth liquid medium (1 g L^{-1}) in each well with 40 replicates by dilution. Microplates were incubated for 48 h at 28°C then absorbance was measured at 620 nm and MPN counts calculated by a computer program using McCrady table (Binet et al., 2000).

2.5. Quantification of bacteria, fungi and PAH-degrading bacteria

DNA extractions were performed independently for each biological replicate. DNA was extracted using the FastDNA[®] Spin kit for soil according to manufacturers recommendations (MP Biomedicals) from 1 g of soil. Bacteria, fungi and PAH-degrading bacteria were quantified by targeting 16S rDNA, 18S rDNA and PAH-RHD _{α} genes using real-time PCR quantification of gene copy number, with the primer sets 968F/1401R (Felske et al., 1998), FF390R/Fung5F (Lueders et al., 2004), PAH-RHD _{α} GN F/R and PAH-RHD _{α} GP F/R (Cébron et al., 2008), respectively. Real-Time PCR experiments were conducted in triplicate with an iCycler iQ (Bio-Rad), associated with iCycler Optical System Interface software (version 2.3). The final volume used (20 μL) contained 10 μL of iQ SYBR GREEN SuperMix (Biorad), 0.4 μM each primer, 0.06% bovine serum albumine (BSA), 0.2 μL DMSO, 0.08 μL

T4gp32 (MP Biomedicals) and 1 μL of DNA as described in Cébron et al. (2008). Quantitative calibrations of Real-Time PCR assays were performed with 10 times dilution series (from 10^8 to 10^3 target gene copies μL^{-1}) of standard linearized plasmids (pCR2.1 vector from Invitrogen) with target gene fragment inserted (Cébron et al., 2008).

2.6. Germination and plant growth test

Fenton oxidized and control soil samples (100 g) with five replicates were placed in 200 mL pots. Soil was watered with deionised water to 70% WHC and soil moisture was daily adjusted. Twenty seeds were sown on each pot, thinned to 5 plants after germination. The pots were placed in a growth chamber at $24^\circ\text{C}/20^\circ\text{C}$, 16 h day light, $250 \mu\text{mol m}^{-2} \text{s}^{-1}$ intensity. The germination rate was recorded every day and conducted over a period of 2 weeks. After 4 weeks, plants were harvested, and roots gently rinsed in water. Shoot and root dry weights were measured after drying at 105°C for 24 h.

2.7. PAH analysis

PAHs were extracted from soil samples in triplicates using ASE 200 (Accelerated Solvent Extraction Dionex). Anhydrous sodium sulfate (1 g) and 1 g of Florisil® was added to each sample (1 g) in the extraction cell to purify the extract. The extraction solvent was composed of 50% acetone and 50% methylene chloride and extraction was run at 100°C and 130 bars for 5 min (Ouvrard et al., 2011). After evaporation of most of the solvent, samples were diluted in 5 mL of acetonitrile and evaporated to obtain exclusive acetonitrile extracts. Analysis of 16 US-EPA priority PAHs concentrations was performed by high-performance liquid chromatography (Varian) (Monsérié et al., 2009). Results were reported for dry soil for comparison between treatments.

2.8. Statistical analysis

Statistical analyses were performed using XLSTAT© Software. Comparisons were made by ANOVA after verification of applicability conditions. A multiple post hoc comparison (Tukey) test was applied to determine the difference per pair. Gene quantification data were log transformed for statistical comparisons. To test time effect of bacteria recolonization a Student t-test was applied. For seed germination a Fisher test was applied to calculate LSD (least significant difference).

3. Results

3.1. Effect of Fenton oxidation on soil PAH concentration and physico-chemical characteristics

Total PAH concentration was significantly lower in Fenton oxidized than in control soil with 22% and 14% decrease with H and L oxidant concentration, respectively (Table 1). The same result was found for the individual PAHs except for the three low molecular weight PAHs naphthalene, acenaphtylene and acenaphthene where the concentration decrease was not significant (Table 1). No clear difference was observed for the degradation rate between the 3 (18 to 36% decrease with H oxidant concentration), 4 (20.3 to 24.3%), 5 (16 to 22.1%) and 6 (24.3 to 24.6 %) ring PAHs. No further significant PAH concentration decrease occurred during the 5 weeks incubation (data not shown). Although no replicate analysis was performed for organic carbon and nitrogen content, the carbon content tended to be lower in oxidized than control soil, while nitrogen content was not affected. After Fenton H treatment, several parameters were strongly affected: a CEC decrease of 2 units, a pH decrease of 2.5 units and a decrease of available phosphorus (77% of the initial content) were measured.

3.2. *Effect of Fenton reaction on microbial density and activity*

No production of CO₂ was observed in the abiotic controls (data not shown). The Fenton H treatment significantly and rapidly increased carbon mineralization rate, while mineralization rate was not significantly different with Fenton L treated and control soil (Figure 1). After oxidation (T0), the number of culturable heterotrophs, the 16S, 18S rDNA and PAH degrading (PAH-RHD_a) gene copy numbers were significantly lower in Fenton H treated than in control soil (Figure 2). After 5 weeks of incubation (T5), these numbers were significantly higher than at T0 in Fenton H treated soil (Figure 2), while they did not significantly differ between T0 and T5 for control and Fenton L treated soils. After 5 weeks the number of culturable heterotrophs and of 18S rDNA gene copies were even significantly higher in Fenton H treated than in control soil. However, in Fenton L treated soil there was no significant difference in culturable heterotrophs, 16S, 18S and PAH-RHD gene copy number with the control soil at T0 and five weeks after incubation (T5).

3.3. *Effect of treatment on germination and growth*

Ryegrass germination rate reached 75-80% in Fenton treated and control soil after 14 days. However, a 3-days significant delay in seed germination was observed for the Fenton H treated soil (Figure 3). Root and shoot biomass was significantly lower in Fenton H treated soil (35% and 43% decrease) than in the control soil, but it was not affected in Fenton L treated soil (Figure 4).

4. Discussion

214 The modified Fenton reaction applied in unsaturated conditions significantly reduced the
215 amount of PAHs in soil. The degradation concerned the different PAHs except the lower
216 molecular weight ones. Similar degradation rates of PAHs regardless of the number of
217 aromatic rings or lower degradation rates of higher molecular weight PAHs were previously
218 reported depending on the $\text{H}_2\text{O}_2/\text{Fe}^{2+}$ ratio (Tran et al., 2010). However, our degradation rate,
219 obtained with 5-56 g oxidant added for 1 g PAHs, was lower than observed in other studies
220 where higher concentrations of oxidants were applied in saturated conditions (100 to 4000 g
221 oxidant for 1 g PAHs) for spiked (Nam et al., 2001; Kulik et al., 2006) or aged contaminated
222 soils (Nam et al., 2001; Jonsson et al., 2006, 2007). In soil, the non specific action of hydroxyl
223 radicals could oxidize the targeted molecules as well as the organic matter, which increases
224 the quantity of oxidant needed to degrade PAHs. The soil contains more than 50 g kg^{-1} of
225 organic carbon and PAHs represent only 2% of it (Monsérié et al., 2009). The apparent lower
226 C_{org} content in both oxidized treatments than in the control soil, could be due to the action of
227 hydroxyl radicals, but this should be confirmed by further investigations. Moreover, the soil
228 used in this study had a high carbonate content which can limit the efficiency of Fenton
229 reaction. At the same time, this reaction lead to a decrease of pH, probably due to the
230 injection of an acidic H_2O_2 solution and to the reaction between Fe (III) and H_2O_2 (Huling and
231 Pivetz, 2006). Carbonates can also buffer the environment of hydroxyl radicals which present
232 a higher efficiency in acidic conditions (Huling and Pivetz, 2006). Another explanation could
233 be the low accessibility of aged pollutants in this historically contaminated soil (Katayama et
234 al., 2010; Ouvrard et al., 2011). Contrary to other studies, the contact between PAHs and the
235 reagent was not enhanced by the use of a high volume of solution. The fact that hydroxyl
236 radicals have a short half life (Sies, 1993) added to the difficulty to access PAHs located in
237 newly formed aggregates specific to the industrial soil (Monsérié et al., 2009) can be another
238 limitation for the degradation by such chemical processes.

Soil fertility was affected by the Fenton oxidation with the higher concentration of oxidant added. The main effect was a decrease of soil pH, a negative effect on plant growth, as previously observed by Sirguey et al. (2008) and a decrease of CEC and available P. The decrease in CEC could be explained by the consequence of the pH drop on the charge of the organic matter carboxyl groups, which partly turned to COOH instead of COO⁻ (Sirguey et al., 2008). Moreover, it can result from the ionisation in acidic conditions of mineral OH groups. Cation exchange capacity reflects the mobility and the availability of ions which are important components of soil fertility. The decrease of available phosphorus content could result from the formation of ferric phosphate in acid condition which is an insoluble form of phosphorus. The pH decrease, the low available phosphorus content and the decrease of cation exchange capacity could explain the decrease of biomass and root elongation. Soil fertility and plant growth could possibly be restored by addition of ameliorants (lime, fertilizer) (Séré et al., 2008).

Besides decreasing soil fertility, the oxidation treatment affected microbial density and activity. The decrease in bacterial, fungal, and PAH degrading bacteria population densities could be explained by the toxicity of hydroxyl radicals strongly reacting with biological membranes (Slater, 1984). Moreover, the heat produced by the exothermic Fenton reaction and the decrease of pH could contribute to the decreased microbial number and activity just after chemical oxidation. The decrease of the number of microorganisms after Fenton reaction as well as the rebound have been previously observed (Ferguson et al., 2004; Palmroth et al., 2006b; Sahl and Munakata-Marr, 2006). The increased microbial activity, estimated by the increased CO₂ release, and the increased number of cultivable and total bacteria and fungi during incubation after oxidation indicated that recolonization occurred. Cultivable bacteria which are heterotrophic preferentially use the most available organic carbon. The dead bacterial cells (Palmroth et al., 2006b) as well as the organic carbon affected by the oxidation

treatment (Leifeld and Kögel-Knabner, 2001; Mikutta et al., 2005) could potentially provide a supplementary available amount of organic carbon. Contrary to other studies, no further degradation of PAHs occurred in our experiment following oxidation, and no variations in PAH-degrading populations were observed. However, it was previously shown that PAH availability in this historically contaminated soil was low and limited natural attenuation (Ouvrard et al. 2011).

5. Conclusion

Partial removal of PAHs (14 and 22%) in a historically contaminated soil was possible in unsaturated conditions using Fenton reagents with the low and high doses of H₂O₂ (6 and 65 g kg⁻¹). The impact of chemical oxidation on PAH concentration and soil physico-chemical and biological parameters differed with the level of reagent used. PAH degradation was not proportional to the quantity of reagent added. With the highest oxidant level applied, soil parameters were highly impacted, especially pH value and biological parameters such as the quantity of microorganisms, seed germination and plant growth. The impact on microbiological parameters was temporary and disappeared rapidly. Specific microbial functions could be investigated to further check whether soil microbial functioning was affected. Reduction of soil pH was a major limiting factor for plant growth at high level of oxidant. Further investigations could be performed with different concentrations of reagents to optimize PAH degradation without impact on vegetation. Otherwise, chemical oxidation could be combined with soil restoration strategies.

Acknowledgments

This work was funded by the ANR-Precodd Oxysol project, the French Government and the Lorraine Region (GISFI project, www.gisfi.fr). The authors thank Prof. Marie-Odile Simonnot and Jean-Christophe Renat (Valterra Company) for fruitful discussions and advices.

References

- Biache, C., Mansuy-Huault, L., Faure, P., Munier-Lamy, C., Leyval, C., 2008. Effects of thermal desorption on the composition of two coking plant soils: Impact on solvent extractable organic compounds and metal bioavailability. *Environ. Poll.* 156, 671-677.
- Binet, P., Portal, J.M., Leyval, C., 2000. Dissipation of 3-6-ring polycyclic aromatic hydrocarbons in the rhizosphere of ryegrass. *Soil Biol. Biochem.* 32, 2011-2017.
- Bispo, A., Jourdain, M.J., Jauzein, M., 1999. Toxicity and genotoxicity of industrial soils polluted by polycyclic aromatic hydrocarbons (PAHs). *Org. Geochem.* 30, 947-952.
- Cajthaml, T., Bhatt, M., Šašek, V., Matějů, V., 2002. Bioremediation of PAH-contaminated soil by composting: A case study. *Folia Microbiol.* 47, 696-700.
- Cébron, A., Norini, M.-P., Beguiristain, T., Leyval, C., 2008. Real-Time PCR quantification of PAH-ring hydroxylating dioxygenase (PAH-RHD[alpha]) genes from Gram positive and Gram negative bacteria in soil and sediment samples. *J. Microbiol. Methods.* 73, 148-159.
- Felske, A., Akkermans, A.D.L., De Vos, W.M., 1998. Quantification of 16S rRNAs in complex bacterial communities by multiple competitive reverse transcription-PCR in temperature gradient gel electrophoresis fingerprints. *Appl. Environ. Microbiol.* 64, 4581-4587.

311 Ferguson, S.H., Woinarski, A.Z., Snape, I., Morris, C.E., Revill, A.T., 2004. A field trial of in
 312 situ chemical oxidation to remediate long-term diesel contaminated Antarctic soil. *Cold*
 313 *Regions Sci. Tech.* 40, 47-60.

314 Fismes, J., Schwartz, C., Perrin-Ganier, C., Morel, J.-L., Charissou, A.-M., Jourdain, 2004.
 315 Risk of contamination for edible vegetables growing on soils polluted by polycyclic
 316 aromatic hydrocarbon. *Polycyclic Aromat. Compd.* 24, 827-836.

317 Gan, S., Lau, E.V., Ng, H.K., 2009. Remediation of soils contaminated with polycyclic
 318 aromatic hydrocarbons (PAHs). *J. Hazard. Mater.* 172, 532-549.

319 Huling, S.G., Pivetz, B.E., 2006. In-Situ Chemical Oxidation - Engineering Issue.
 320 (EPA/600/R-06/072), pp. 1-60.

321 Jonsson, S., Persson, Y., Frankki, S., Lundstedt, S., Bavel, B.v., Haglund, P., Tysklind, M.,
 322 2006. Comparison of Fenton's reagent and ozone oxidation of polycyclic aromatic
 323 hydrocarbons in aged contaminated soils. *J. Soil. Sediment.* 6, 208-214.

324 Jonsson, S., Persson, Y., Frankki, S., van Bavel, B., Lundstedt, S., Haglund, P., Tysklind, M.,
 325 2007. Degradation of polycyclic aromatic hydrocarbons (PAHs) in contaminated soils by
 326 Fenton's reagent: A multivariate evaluation of the importance of soil characteristics and PAH
 327 properties. *J. Hazard. Mater.* 149, 86-96.

328 Katayama, A., Bhula, R., Burns, G.R., Carazo, E., Felsot, A., Hamilton, D., Harris, C., Kim,
 329 Y.-H., Kleter, G., Koerdel, W., Linders, J., Peijnenburg, J.G.M.W., Sabljic, A., Stephenson,
 330 R.G., Racke, D.K., Rubin, B., Tanaka, K., Unsworth, J., Wauchope, R.D., 2010.
 331 Bioavailability of xenobiotics in the soil environment. *Rev. Environ. Contam. T.* 203, 1-86.

332 Kulik, N., Goi, A., Trapido, M., Tuhkanen, T., 2006. Degradation of polycyclic aromatic
 333 hydrocarbons by combined chemical pre-oxidation and bioremediation in creosote
 334 contaminated soil. *J. Environ. Manage.* 78, 382-391.

335 Lee, B.-D., Hosomi, M., 2001. A hybrid Fenton oxidation-microbial treatment for soil highly
 336 contaminated with benz(a)anthracene. *Chemosphere* 43, 1127-1132.

337 Leifeld, J., Kögel-Knabner, I., 2001. Organic carbon and nitrogen in fine soil fractions after
 338 treatment with hydrogen peroxide. *Soil. Biol. Biochem.* 33, 2155-2158.

339 Lueders, T., Wagner, B., Claus, P., Friedrich, M.W., 2004. Stable isotope probing of rRNA
 340 and DNA reveals a dynamic methylotroph community and trophic interactions with fungi and
 341 protozoa in oxic rice field soil. *Environ. Microbiol.* 6, 60-72.

342 Maxim, L., Spangenberg, J.H., 2009. Driving forces of chemical risks for the European
 343 biodiversity. *Ecolog. Econ.* 69, 43-54.

344 Mikutta, R., Kleber, M., Kaiser, K., Jahn, R., 2005. Review: organic matter removal from
 345 soils using hydrogen peroxide, sodium hypochlorite, and disodium peroxodisulfate. *Soil Sci.*
 346 *Soc. Am. J.* pp. 120-135.

347 Monsérié, M.-F., Watteau, F., Villemin, G., Ouvrard, S., Morel, J.-L., 2009. Technosol
 348 genesis: identification of organo-mineral associations in a young Technosol derived from
 349 coking plant waste materials. *J. Soil. Sediment.* 9, 537-546.

350 Nam, K., Rodriguez, W., Kukor, J.J., 2001. Enhanced degradation of polycyclic aromatic
 351 hydrocarbons by biodegradation combined with a modified Fenton reaction. *Chemosphere* 45,
 352 11-20.

353 Ouvrard S., Barnier C., Bauda P., Beguiristain T., Biache C., Bonnard M., Caupert C., Cébron
 354 A., Cortet J., Cotellet S., Dazy M., Faure P., Masfaraud J. F., Nahmani J., Palais F., Poupin P.,
 355 Raoult N., Vasseur P., Morel J.L., Leyval C., 2011. *In situ* assessment of phytotechnologies
 356 for multicontaminated soil management. *Int. J. Phytoremediat.* , 13:sup1, 245-263.

357 Palmroth, M., Langwaldt, J., Aunola, T., Goi, A., Puhakka, J., Tuhkanen, T., 2006a.
 358 Treatment of PAH-contaminated soil by combination of Fenton's reaction and biodegradation.
 359 *J. Chem. Technol. Biotechnol.* 81, 598-607.

360 Palmroth, M., Langwaldt, J., Aunola, T., Goi, A., Münster, U., Puhakka, J., Tuhkanen, T.,
 361 2006b. Effect of modified Fenton's reaction on microbial activity and removal of PAHs in
 362 creosote oil contaminated soil. *Biodegradation* 17, 29-39.
 363 Quantin, C., Joner, E.J., Portal, J.M., Berthelin, J., 2005. PAH dissipation in a contaminated
 364 river sediment under oxic and anoxic conditions. *Environ. Poll.* 134, 315-322.
 365 Rivas, F.J., 2006. Polycyclic aromatic hydrocarbons sorbed on soils: A short review of
 366 chemical oxidation based treatments. *J. Hazard. Mater.* 138, 234-251.
 367 Sahl, J., Munakata-Marr, J., 2006. The effects of in situ chemical oxidation on
 368 microbiological processes: A review. *Remed. J.* 16, 57-70.
 369 Séré, G., Schwartz, C., Ouvrard, S., Sauvage, C., Renat, J.-C., Morel, J., 2008. Soil
 370 construction: A step for ecological reclamation of derelict lands. *J. Soil. Sediment.* 8, 130-
 371 136.
 372 Sies, H., 1993. Strategies of antioxidant defense. *Eur. J. Biochem.* 215, 213-219.
 373 Sirguey, C., De Souza e Silva, P.T., Schwartz, C., Simonnot, M.-O., 2008. Impact of chemical
 374 oxidation on soil quality. *Chemosphere* 72, 282-289.
 375 Slater, T.F., 1984. Free-radical mechanisms in tissue injury. *Biochem. J.* 222, 1-15.
 376 Smith, M.J., Flowers, T.H., Duncan, H.J., Alder, J., 2006. Effects of polycyclic aromatic
 377 hydrocarbons on germination and subsequent growth of grasses and legumes in freshly
 378 contaminated soil and soil with aged PAHs residues. *Environ. Poll.* 141, 519-525.
 379 Sun, H.-W., Yan, Q.-s., 2008. Influence of pyrene combination state in soils on its treatment
 380 efficiency by Fenton oxidation. *J. Environ. Manage.* 88, 556-563.
 381 Tran, L.-H., Drogui, P., Mercier, G., Blais, J.-F., 2010. Comparison between Fenton oxidation
 382 process and electrochemical oxidation for PAH removal from an amphoteric surfactant
 383 solution. *J. Appl. Electrochem.* 40, 1493-1510.
 384

385 Valderrama, C., Alessandri, R., Aunola, T., Cortina, J.L., Gamisans, X., Tuhkanen, T., 2009.
386 Oxidation by Fenton's reagent combined with biological treatment applied to a creosote-
387 contaminated soil. *J. Hazard. Mater.* 166, 594-602.

388 Wang, X., Yu, X., Bartha, R., 1990. Effect of bioremediation on polycyclic aromatic
389 hydrocarbon residues in soil. *Environ. Sci. Technol.* 24, 1086-1089.

390 Westerhoff, P., Aiken, G., Amy, G., Debroux, J., 1999. Relationships between the structure of
391 natural organic matter and its reactivity towards molecular ozone and hydroxyl radicals.
392 *Water Res.* 33, 2265-2276.

393 White, P.A., Claxton, L.D., 2004. Mutagens in contaminated soil: a review. *Mutat. Res., Rev.*
394 *Mutat. Res.* 567, 227-345.

395
396
397
398
399
400
401
402
403
404
405
406
407
408
409

Figure captions

Figure 1: Cumulated production of CO₂ over time of incubation with oxidized (Fenton L and H) and control soil (Control). Mean $\pm$ SE (n=5). Arrows represent days of oxigenation.

Figure 2: Density of cultivable bacteria, 16S rDNA, PAH-RDH α and 18s rDNA gene copy numbers in oxidized (Fenton L and H) and control soil (Control) at the start (T0) and after 5 weeks (T5) of incubation. As the evolution of PAH-RHD α GN and GP populations was not significantly different, the two populations were added to represent the global population of degraders. Mean $\pm$ SE (n=3). LD: detection limit. Different letters within category data indicate significant differences between treatments; stars indicate significant differences between T₀ and T₆ by Mann and Whitney test.

Figure 3: Germination rate (%) of *Lolium perenne* seeds in oxidized (Fenton L and H) and control soil. Mean $\pm$ SD (n=5). Stars indicate a significant difference between Fenton H treated and Control soil

Figure 4: Shoot and root dry weight of *Lolium perenne* after 4 weeks on oxidized (Fenton L and H) and control soil. (Mean $\pm$ SE. n=5). Values followed by the same letter are not significantly different at the 5% level (two-way ANOVA, Tukey test)

Table 1: Physico-chemical characteristics of the soil after Fenton treatment at low (L) and high (H) concentration of oxidant. Concentrations of PAHs, given as mean, followed by the same letter are not significantly different at the 5% level (one-way ANOVA, Tukey test)

		Control	Fenton L	Fenton H
	Units			
Org C	(g kg ⁻¹)	61,7	57.7	57.1
N	(g kg ⁻¹)	2.4	2.4	2.3
Olsen P	(g kg ⁻¹)	0.049	0.068	0.011
CEC	(cmol kg ⁻¹)	10.3	10.5	8.7
pH		7.3	7.4	4.9
Naphthalene		24.8 ^a	26.4 ^a	22.0 ^a
Acenaphthylene		0.7 ^a	0.6 ^a	0.8 ^a
Acenaphthene		57.9 ^a	54.8 ^a	52.7 ^a
Fluorene		25.9 ^a	22.9 ^b	20.4 ^b
Phenanthrene		89.1 ^a	80.9 ^{ab}	72.9 ^b
Anthracene		36.8 ^a	26.2 ^b	23.6 ^b
Fluoranthene		235.5 ^a	206.1 ^b	186.9 ^c
Pyrene		128.8 ^a	111.4 ^b	102.6 ^c
Benzo(a)Anthracene	(μg kg ⁻¹)	101.2 ^a	87.0 ^b	78.4 ^c
Chrysene		56.7 ^a	47.4 ^b	42.9 ^c
Benzo(b)Fluoranthene		96.7 ^a	83.5 ^b	77.3 ^b
Benzo(k)Fluoranthene		30.8 ^a	26.6 ^b	24.5 ^b
Benzo(a)Pyrene		108.8 ^a	91.5 ^b	84.7 ^b
Dibenzo(a,h)Anthracene		13.1 ^a	11.3 ^b	11.0 ^b
Benzo(g,h,i)Perylene		64.2 ^a	52.7 ^b	48.4 ^b
Indeno(1,2,3-cd)Pyrene		90.9 ^a	75.4 ^b	68.8 ^b
Total PAHs		1161.9 ^a	1004.9 ^b	917.9 ^c

Figure 1

Figure 2

Figure 3

Figure 4