

# Galerkin approximations of nonlinear optimal control problems in Hilbert spaces

Mickaël D. Chekroun, Axel Kröner, Honghu Liu

► **To cite this version:**

Mickaël D. Chekroun, Axel Kröner, Honghu Liu. Galerkin approximations of nonlinear optimal control problems in Hilbert spaces . *Electronic Journal of Differential Equations*, Texas State University, Department of Mathematics, 2017, 2017 (189), pp.1-40. hal-01501178

**HAL Id: hal-01501178**

**<https://hal.archives-ouvertes.fr/hal-01501178>**

Submitted on 3 Apr 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# GALERKIN APPROXIMATIONS OF NONLINEAR OPTIMAL CONTROL PROBLEMS IN HILBERT SPACES

MICKAËL D. CHEKROUN, AXEL KRÖNER, AND HONGHU LIU

ABSTRACT. Nonlinear optimal control problems in Hilbert spaces are considered for which we derive approximation theorems for Galerkin approximations. Approximation theorems are available in the literature. The originality of our approach relies on the identification of a set of natural assumptions that allows us to deal with a broad class of nonlinear evolution equations and cost functionals for which we derive convergence of the value functions associated with the optimal control problem of the Galerkin approximations. This convergence result holds for a broad class of nonlinear control strategies as well. In particular, we show that the framework applies to the optimal control of semilinear heat equations posed on a general compact manifold without boundary. The framework is then shown to apply to geoenvironment and mitigation of greenhouse gas emissions formulated for the first time in terms of optimal control of energy balance climate models posed on the sphere  $S^2$ .

## CONTENTS

1. Introduction	1
2. Galerkin approximations for optimal control problems: Convergence results	4
2.1. Preliminaries	4
2.2. Convergence of Galerkin approximations: Trajectory-wise result	5
2.3. Local-in- $u$ approximation result	8
2.4. Convergence of Galerkin approximations: Uniform-in- $u$ result	11
2.5. Galerkin approximations of optimal control and value functions: Convergence results	15
2.6. Galerkin approximations of optimal control and value functions: Error estimates	18
2.7. Examples that satisfy Assumption (A7)	21
3. Application to optimal control of energy balance climate models	22
3.1. Preliminary from differential geometry	22
3.2. Galerkin approximations of controlled semilinear heat equations on $S^n$	24
3.3. Energy balance models (EBMs)	26
3.4. Optimal control of climate?	28
3.5. Optimal control of EBMs: Convergence results of value functions	29
4. Concluding remarks	31
Acknowledgments	32
Appendix A. Existence of optimal controls	32
References	33

## 1. INTRODUCTION

Optimal control problems of infinite dimensional systems play an important role in a broad range of applications in engineering and various scientific disciplines [Lio71, Fat99, Fur00, BDPDM07, HPUU09,

Trö10]. Various methods for solving numerically the related optimization problems are available; see e.g. [HPUU09, MTZ08]. The case of linear evolution equations has benefited from a long tradition, and an abundant literature exists about finite element techniques or Galerkin methods for the design of approximate optimal controls; see e.g. [MBJ73, Gib79, Las80, Mal82, Kno82, BK84, LT87, AM89, LT00]. The case of Galerkin approximations of optimal control problems for nonlinear evolutions seems to have been much less addressed. Semidiscrete Ritz-Galerkin approximations of nonlinear parabolic boundary control problems have been considered for which convergence of the approximate controls have been obtained; see [Trö93, Trö94]. We refer also to [MV07, NV12] for error estimates concerned with space-time finite element approximations of the state and control to optimal control problems governed by semilinear parabolic equations, and to [DH04] for finite element approximations of optimal control problems associated with the Navier-Stokes equations.

In this article, we study Galerkin approximations for (possibly non-quadratic) optimal control problems over a finite horizon  $[0, T]$ , of nonlinear evolution equations in Hilbert space. Our framework covers not only a broad class of semilinear parabolic equations but also includes systems of nonlinear delay differential equations (DDEs) [CGLW16] and allows in each case for a broad class of nonlinear control strategies. The main contribution of this article is to identify for such equations a set of easily checkable conditions in practice, from which we prove the pointwise convergence of the value functions associated with the optimal control problem of the Galerkin approximations, and for a broad class of cost functionals; see Theorem 2.2, our main result. This convergence at the level of value functions results essentially from a double uniform convergence—with respect to time and the set of admissible controllers—of the controlled Galerkin states; see Theorem 2.1 and Corollary 2.1.

The treatment adopted here is based on the classical Trotter-Kato approximation approach from the  $C_0$ -semigroup theory [Gol85, Paz83], which can be viewed as the functional analysis operator version of the Lax equivalence principle.<sup>1</sup> Within this approach, we generalize, in particular, the convergence results about value functions obtained in the earlier work [Fer97] concerned with the Galerkin approximations to optimal control problems governed by linear evolution equations in Hilbert space. Given a Hilbert state space  $\mathcal{H}$ , denoting by  $\Pi_N$  the orthogonal projector associated with the  $N$ -dimensional Galerkin subspace, and by  $y_N(\cdot; \Pi_N x, u)$  the controlled Galerkin state (driven by  $u$ ) and emanating from  $\Pi_N x$ , a key property to ensure convergence of the value functions for such optimal control problems is the following double uniform convergence

$$(1.1) \quad \lim_{N \rightarrow \infty} \sup_{u \in \mathcal{U}_{ad}} \sup_{t \in [0, T]} \|y_N(t; \Pi_N x, u) - y(t; x, u)\|_{\mathcal{H}} = 0, \quad \forall x \in \mathcal{H},$$

where  $\mathcal{U}_{ad}$  denotes a set of admissible controls; see [Fer97, Theorem 4.2].

When the evolution equation involves state- or control-dependent nonlinear terms, the conditions provided in [Fer97, Proposition 2.1] to ensure (1.1) needs to be amended. Whether the nonlinear terms involve the controls or the system's state, our working assumption regarding the linear terms of the original equation and of its Galerkin approximations, is (as in [Fer97]) to satisfy respectively the “stability” and “consistency” conditions required in the Trotter-Kato theorem; see Assumption (A1) and Assumption (A2) in Sect. 2.1. In the case of a linear equation with nonlinear control terms, a simple compactness assumption about the set of admissible controls (see Assumption (A5) in Sect. 2.4) is sufficient to ensure (1.1); see Remark 2.2.

In the case of an evolution equation depending nonlinearly on the system's state and on the controls, a key assumption is introduced to ensure (1.1) that adds up to standard local Lipschitz conditions on the state-dependent nonlinear terms (Assumption (A3)) and the nonlinear control operator  $\mathcal{C} : V \rightarrow \mathcal{H}$ , where  $V$  denotes an auxiliary Hilbert space in which the controls take values. Introducing  $\Pi_N^\perp := \text{Id}_{\mathcal{H}} - \Pi_N$ , this assumption concerns a double uniform convergence about the residual

<sup>1</sup>i.e., if “consistency” and “stability” are satisfied, then “convergence” holds, and reciprocally.

energy  $\|\Pi_N^\perp y(t; x, w)\|_{\mathcal{H}}$  (see Assumption **(A7)** in Sect. 2.4), namely

$$(1.2) \quad \lim_{N \rightarrow \infty} \sup_{u \in \mathcal{U}_{ad}} \sup_{t \in [0, T]} \|\Pi_N^\perp y(t; x, u)\|_{\mathcal{H}} = 0.$$

With this assumption at hand, and the rest of our working assumptions, standard *a priori* bounds (Assumption **(A6)**) allow us to ensure (1.1) for a broad class of nonlinear evolution equations in Hilbert spaces. The pointwise convergence of the value functions associated with the optimal control problem of the corresponding Galerkin approximations is then easily derived for a broad class of cost functionals; see Theorem 2.2.

The relevance of assumption (1.2) for applications is addressed through various angles. First, from the proof of Corollary 2.1 (and thus Theorem 2.1) in which Theorem 2.2 relies. In that respect, a sort of pedagogical detour is made in Sect. 2.3 in which we show essentially that a weaker (than (1.1)) local-in- $u$  approximation result (Lemma 2.3) follows from the rest of our working assumptions (except Assumptions **(A6)** and **(A7)**<sup>2</sup>) and from a local-in- $u$  estimate about the residual energy (Lemma 2.1); the latter resulting itself from the continuity of the mapping  $u \mapsto y(t; x, u)$ . Condition (1.2) constitutes thus a natural strengthening of inherent properties to the approximation problem.

From a more applied perspective, sufficient conditions concerning the spectrum of the linear part—such as self-adjointness and compact resolvent—are pointed out in Sect. 2.7 to ensure (1.2); see Lemma 2.6 and Remark 2.6. Finally, Sect. 2.6 provides error estimates concerning the value function and the optimal control that complete the picture and emphasize from another perspective the relevance of the residual energy in the analysis of the approximation problem; see Theorem 2.3 and Corollary 2.2.

With this preamble in mind, we provide now the more formal organization of this article. First, we present in Sect. 2.1 the type of state equation and its corresponding Galerkin approximations that we will be working on. A trajectory-wise convergence result for each fixed control  $u$  is then derived in Sect. 2.2. As mentioned earlier, it relies essentially on the theory of  $C_0$ -semigroups and the Trotter-Kato theorem [Paz83, Thm. 4.5, p.88]; see Lemma 2.1. In a second step, we derive a “local-in- $u$ ” approximation result in Sect. 2.3 for controls that lie within a neighborhood of a given control  $u$ ; see Lemma 2.3. As discussed above, a key approximation property about the residual energy of solutions (see (2.33)) is then amended into an assumption (see Assumption **(A7)**) to ensure a uniform-in- $u$  convergence result; see Theorem 2.1 of Sect. 2.4. As shown in Sect. 2.5, this uniform convergence result helps us derive—in the spirit of dynamic programming (see Corollary 2.1)—the convergence of the value functions associated with optimal control problems based on Galerkin approximations; see Theorem 2.2. For this purpose, some standard sufficient conditions for the existence of optimal controls are also recalled in Appendix A. Simple and useful error estimates about the value function and the optimal control are then provided in Sect. 2.6. In Sect. 2.7 we point out a broad class of evolution equations for which Assumption **(A7)** is satisfied.

As applications of the theoretical results derived in Sect. 2, we show in Sect. 3 that our framework allows to provide rigorous Galerkin approximations to the optimal control of a broad class of semilinear heat problems, posed on a compact (smooth) manifold without boundary. As a concrete example, the framework is shown to apply to geoengineering and the mitigation of greenhouse gas (GHG) emissions formulated for the first time here in terms of optimal control of energy balance models (EBMs) arising in climate modeling; see [Bud69, Sel69, Ghi76, NCC81] for an introduction on EBMs, and [DH98, BCDT09] for a mathematical analysis. After recalling some fundamentals of differential geometry in Sect. 3.1 to prepare the analysis, a general convergence result of Galerkin approximations to controlled semilinear heat problems posed on an  $n$ -dimensional sphere  $S^n$  is formulated

---

<sup>2</sup>More precisely, by assuming a weaker version of Assumption **(A6)**, namely Assumption **(A4)**, and without assuming **(A7)**.

in Sect. 3.2; see Corollary 3.1. The application to the optimal control of EBMs is then presented in Sect. 3.3 in the context of geoenvironment and GHG emissions for which approximation of the value function and error estimates about the optimal control are obtained. Finally, Sect. 4 outlines several possible directions for future research the framework introduced in this article opens up.

## 2. GALERKIN APPROXIMATIONS FOR OPTIMAL CONTROL PROBLEMS: CONVERGENCE RESULTS

We present in this section, rigorous convergence results for semi-discretization of optimal control problems based on Galerkin approximations. In particular, we derive the pointwise convergence of the value functions associated with optimal control problems based on Galerkin approximations in Sect. 2.5.

**2.1. Preliminaries.** We consider in this section finite-dimensional approximations of the following initial-value problem (IVP):

$$(2.1) \quad \begin{aligned} \frac{dy}{dt} &= Ly + F(y) + \mathfrak{C}(u(t)), \quad t \in (0, T], \\ y(0) &= x, \end{aligned}$$

where  $x$  lies in  $\mathcal{H}$ , and  $\mathcal{H}$  denotes a separable Hilbert space. The time-dependent forcing  $u$  lives in a separable Hilbert space  $V$  (possibly different than  $\mathcal{H}$ ); the (possibly nonlinear) mapping  $\mathfrak{C} : V \rightarrow \mathcal{H}$  is assumed to be such that  $\mathfrak{C}(0) = 0$ . Other assumptions regarding  $\mathfrak{C}$  will be made precise when needed.

We assume that the linear operator  $L : D(L) \subset \mathcal{H} \rightarrow \mathcal{H}$  is the infinitesimal generator of a  $C_0$ -semigroup of bounded linear operators  $T(t)$  on  $\mathcal{H}$ . Recall that in this case the domain  $D(L)$  of  $L$  is dense in  $\mathcal{H}$  and that  $L$  is a closed operator; see [Paz83, Cor. 2.5, p. 5].

Under the above assumptions on the operator  $L$ , recall that there exists  $M \geq 1$  and  $\omega \geq 0$  [Paz83, Thm. 2.2, p. 4] such that

$$(2.2) \quad \|T(t)\| \leq Me^{\omega t}, \quad t \geq 0,$$

where  $\|\cdot\|$  denotes the operator norm subordinated to  $\|\cdot\|_{\mathcal{H}}$ .

For the moment, we take the set of admissible controls to be

$$(2.3) \quad \mathcal{U} := L^q(0, T; V),$$

with  $q \geq 1$ . In the later subsections, further assumptions on the admissible controls will be specified when needed.

Let  $u$  be in  $\mathcal{U}$  given by (2.3), a *mild solution* to (2.1) over  $[0, T]$  is a function  $y$  in  $C([0, T], \mathcal{H})$  such that

$$(2.4) \quad y(t) = T(t)x + \int_0^t T(t-s)F(y(s))ds + \int_0^t T(t-s)\mathfrak{C}(u(s))ds, \quad t \in [0, T].$$

In what follows we will often denote by  $y(s; x, u)$  a mild solution to (2.1).

Let  $\{\mathcal{H}_N : N \in \mathbb{Z}_+^*\}$  be a sequence of finite-dimensional subspaces of  $\mathcal{H}$  associated with *orthogonal projectors*

$$(2.5) \quad \Pi_N : \mathcal{H} \rightarrow \mathcal{H}_N,$$

such that

$$(2.6) \quad \|(\Pi_N - \text{Id})x\| \xrightarrow{N \rightarrow \infty} 0, \quad \forall x \in \mathcal{H},$$

and

$$(2.7) \quad \mathcal{H}_N \subset D(L), \quad \forall N \geq 1.$$

The corresponding Galerkin approximation of (2.1) associated with  $\mathcal{H}_N$  is then given by:

$$(2.8) \quad \begin{aligned} \frac{dy_N}{dt} &= L_N y_N + \Pi_N F(y_N) + \Pi_N \mathfrak{C}(u(t)), \quad t \in [0, T], \\ y_N(0) &= \Pi_N x, \quad x \in \mathcal{H}, \end{aligned}$$

where

$$(2.9) \quad L_N := \Pi_N L \Pi_N : \mathcal{H} \rightarrow \mathcal{H}_N.$$

In particular, the domain  $D(L_N)$  of  $L_N$  is  $\mathcal{H}$ , because of (2.7).

Throughout this section, we assume the following set of assumptions:

- (A0) The linear operator  $L : D(L) \subset \mathcal{H} \rightarrow \mathcal{H}$  is the infinitesimal generator of a  $C_0$ -semigroup of bounded linear operators  $T(t)$  on  $\mathcal{H}$ .
- (A1) For each positive integer  $N$ , the linear flow  $e^{L_N t} : \mathcal{H}_N \rightarrow \mathcal{H}_N$  extends to a  $C_0$ -semigroup  $T_N(t)$  on  $\mathcal{H}$ . Furthermore the following uniform bound is satisfied by the family  $\{T_N(t)\}_{N \geq 1, t \geq 0}$

$$(2.10) \quad \|T_N(t)\| \leq M e^{\omega t}, \quad N \geq 1, \quad t \geq 0,$$

where  $\|T_N(t)\| := \sup\{\|T_N(t)x\|_{\mathcal{H}}, \|x\|_{\mathcal{H}} \leq 1, x \in \mathcal{H}\}$  and the constants  $M$  and  $\omega$  are the same as given in (2.2).

- (A2) The following convergence holds

$$(2.11) \quad \lim_{N \rightarrow \infty} \|L_N \phi - L \phi\|_{\mathcal{H}} = 0, \quad \forall \phi \in D(L).$$

- (A3) The nonlinearity  $F$  is locally Lipschitz in the sense given in (2.12) below.

Following the presentation commonly adopted for the Trotter-Kato approach, the assumptions (A0)-(A2) are concerned with the linear parts of the original system (2.1) and of its Galerkin approximation (2.8). Assumption (A3) is concerned with the nonlinearity in (2.1). Other assumptions regarding the latter will be made in the sequel. Throughout this article, a mapping  $f : \mathcal{W}_1 \rightarrow \mathcal{W}_2$  between two Banach spaces,  $\mathcal{W}_1$  and  $\mathcal{W}_2$ , is said to be locally Lipschitz if for any ball  $\mathfrak{B}_r \subset \mathcal{W}_1$  with radius  $r > 0$  centered at the origin, there exists a constant  $\text{Lip}(f|_{\mathfrak{B}_r}) > 0$  such that

$$(2.12) \quad \|f(y_1) - f(y_2)\|_{\mathcal{W}_2} \leq \text{Lip}(f|_{\mathfrak{B}_r}) \|y_1 - y_2\|_{\mathcal{W}_1}, \quad \forall y_1, y_2 \in \mathfrak{B}_r.$$

**2.2. Convergence of Galerkin approximations: Trajectory-wise result.** As a preparation for the main result given in Sect. 2.5, we derive hereafter a trajectory-wise convergence result for the solutions to the Galerkin approximations (2.8); see Lemma 2.1 below.

With this purpose in mind, besides (A0)-(A3), we will also make use of the following assumption.

- (A4) For each  $T > 0$ ,  $x$  in  $\mathcal{H}$  and each  $u$  in  $\mathcal{U}$  with  $\mathcal{U}$  defined in (2.3), the problem (2.1) admits a unique mild solution  $y(\cdot; x, u)$  in  $C([0, T], \mathcal{H})$ , and its Galerkin approximation (2.8) admits a unique solution  $y_N(\cdot; \Pi_N x, u)$  in  $C([0, T], \mathcal{H}_N)$  for each  $N$  in  $\mathbb{Z}_+^*$ . Moreover, there exists a constant  $C := C(T, x, u)$  such that

$$(2.13) \quad \|y_N(t; \Pi_N x, u)\|_{\mathcal{H}} \leq C, \quad \forall t \in [0, T], \quad N \in \mathbb{Z}_+^*.$$

Note that in applications, (A4) is typically satisfied via a priori estimates; see Remark 2.1-(ii) below.

**Lemma 2.1.** *Let  $\mathcal{U}$  be the set of admissible controls given by (2.3), with  $q$  defined therein. Consider the IVP (2.1) and the associated Galerkin approximation (2.8). Assume that the nonlinear operator  $\mathfrak{C} : V \rightarrow \mathcal{H}$  satisfies, for  $1 \leq p \leq q$ , the following growth condition,*

$$(2.14) \quad \|\mathfrak{C}(w)\|_{\mathcal{H}} \leq \gamma_1 \|w\|_V^p + \gamma_2, \quad \forall w \in V,$$

where  $\gamma_1 > 0$  and  $\gamma_2 \geq 0$ .

Assume also that **(A0)**–**(A4)** hold. Then for each fixed  $T > 0$ ,  $x$  in  $\mathcal{H}$  and  $u$  in  $\mathcal{U}$ , the following convergence result is satisfied:

$$(2.15) \quad \lim_{N \rightarrow \infty} \sup_{t \in [0, T]} \|y_N(t; \Pi_N x, u) - y(t; x, u)\|_{\mathcal{H}} = 0.$$

*Proof.* To simplify the notations, only the  $x$ -dependency that matters to the estimates derived hereafter will be made explicit. Let  $u$  be given in  $\mathcal{U}$  and  $x$  in  $\mathcal{H}$ , then by the variation-of-constants formula applied to Eq. (2.8) we have, for  $0 \leq t \leq T$  and  $N$  in  $\mathbb{Z}_+^*$ , that

$$(2.16) \quad y_N(t; u) = e^{L_N t} \Pi_N x + \int_0^t e^{L_N(t-s)} \Pi_N F(y_N(s; u)) ds + \int_0^t e^{L_N(t-s)} \Pi_N \mathfrak{C}(u(s)) ds.$$

Then, it follows from (2.4) and (2.16) that the difference  $w_N(t; u) := y(t; u) - y_N(t; u)$  satisfies

$$(2.17) \quad \begin{aligned} w_N(t; u) &= T(t)x - e^{L_N t} \Pi_N x + \int_0^t T(t-s)F(y(s; u)) ds - \int_0^t e^{L_N(t-s)} \Pi_N F(y_N(s; u)) ds \\ &\quad + \int_0^t T(t-s)\mathfrak{C}(u(s)) ds - \int_0^t e^{L_N(t-s)} \Pi_N \mathfrak{C}(u(s)) ds \end{aligned}$$

and hence, we have

$$(2.18) \quad \begin{aligned} w_N(t; u) &= T(t)x - e^{L_N t} \Pi_N x + \int_0^t (T(t-s) - e^{L_N(t-s)} \Pi_N) F(y(s; u)) ds \\ &\quad + \int_0^t e^{L_N(t-s)} \Pi_N (F(y(s; u)) - F(y_N(s; u))) ds + \int_0^t (T(t-s) - e^{L_N(t-s)} \Pi_N) \mathfrak{C}(u(s)) ds. \end{aligned}$$

Let us introduce for every  $s$  in  $[0, T]$ ,  $x$  in  $\mathcal{H}$ , and  $u$  in  $\mathcal{U}$ ,

$$(2.19a) \quad r_N(s; u) := \|y(s; u) - y_N(s; u)\|_{\mathcal{H}},$$

$$(2.19b) \quad \zeta_N(x) := \sup_{t \in [0, T]} \|T(t)x - e^{L_N t} \Pi_N x\|_{\mathcal{H}},$$

$$(2.19c) \quad d_N(s; u) := \sup_{t \in [s, T]} \|(T(t-s) - e^{L_N(t-s)} \Pi_N) F(y(s; u))\|_{\mathcal{H}},$$

and for almost every  $s$  in  $[0, T]$ ,

$$(2.20) \quad \tilde{d}_N(s; u) := \sup_{t \in [s, T]} \|(T(t-s) - e^{L_N(t-s)} \Pi_N) \mathfrak{C}(u(s))\|_{\mathcal{H}}.$$

For  $x$  in  $\mathcal{H}$  and  $u$  in  $\mathcal{U}$ , let us denote by  $\mathfrak{B}$  the closed ball in  $\mathcal{H}$  with radius  $C$  centered at the origin where  $C$  is the upper bound in estimate (2.13) for the Galerkin solutions (Assumption **(A4)**).

Since by assumption,  $t \mapsto y(t; x, u)$  lies in  $C([0, T], \mathcal{H})$  for any  $x$  in  $\mathcal{H}$  and  $u$  in  $\mathcal{U}$  given by (2.3), one can assume without loss of generality that  $y(t; x, u)$  stays in  $\mathfrak{B}$  for all  $t$  in  $[0, T]$ , by possibly redefining  $C$ .

We obtain then from (2.18) that

$$(2.21) \quad \begin{aligned} r_N(t; u) &\leq \zeta_N(x) + \int_0^t d_N(s; u) ds + \int_0^t \|e^{L_N(t-s)} \Pi_N (F(y(s; u)) - F(y_N(s; u)))\|_{\mathcal{H}} ds + \int_0^t \tilde{d}_N(s; u) ds \\ &\leq \zeta_N(x) + \int_0^t d_N(s; u) ds + M \text{Lip}(F|_{\mathfrak{B}}) \int_0^t e^{\omega(t-s)} r_N(s; u) ds + \int_0^t \tilde{d}_N(s; u) ds \\ &\leq \zeta_N(x) + \int_0^t d_N(s; u) ds + M \text{Lip}(F|_{\mathfrak{B}}) e^{\omega T} \int_0^t r_N(s; u) ds + \int_0^t \tilde{d}_N(s; u) ds, \end{aligned}$$

where we have used that  $F$  is locally Lipschitz (Assumption **(A3)**) and the uniform bound (2.10) of Assumption **(A1)**.

It follows then from Gronwall's inequality that

$$(2.22) \quad r_N(t; u) \leq \left( \zeta_N(x) + \int_0^T d_N(s; u) ds + \int_0^T \tilde{d}_N(s; u) ds \right) \exp(M\text{Lip}(F|_{\mathfrak{B}})e^{\omega T}T), \quad \forall t \in [0, T].$$

We are thus left with the estimation of  $\zeta_N(x)$ ,  $\int_0^T d_N(s; u) ds$  and  $\int_0^T \tilde{d}_N(s; u) ds$  as  $N \rightarrow \infty$ . Note that the assumptions **(A1)** and **(A2)** allow us to use a version of the Trotter-Kato theorem [Paz83, Thm. 4.5, p.88] which implies together with (2.6) that

$$(2.23) \quad \lim_{N \rightarrow \infty} e^{LNt} \Pi_N \phi = T(t)\phi, \quad \forall \phi \in \mathcal{H},$$

uniformly in  $t$  lying in bounded intervals; see also [CGLW16, footnote 6].

It follows that

$$(2.24) \quad \lim_{N \rightarrow \infty} \zeta_N(x) = 0,$$

and that  $d_N(\cdot; u)$  and  $\tilde{d}_N(\cdot; u)$  converge point-wisely to zero on  $[0, T]$ , i.e.,

$$(2.25) \quad \lim_{N \rightarrow \infty} d_N(s; u) = 0, \quad \forall s \in [0, T],$$

and

$$(2.26) \quad \lim_{N \rightarrow \infty} \tilde{d}_N(s; u) = 0, \quad \forall s \in [0, T].$$

On the other hand, by using (2.2) and (2.10) and from the local Lipschitz assumption on  $F$ , we get

$$(2.27) \quad \begin{aligned} & \| (T(t-s) - e^{LN(t-s)} \Pi_N) F(y(s; u)) \|_{\mathcal{H}} \\ & \leq 2Me^{\omega(t-s)} \| F(y(s; u)) \|_{\mathcal{H}} \\ & \leq 2Me^{\omega(t-s)} \left( \| F(y(s; u)) - F(0) \|_{\mathcal{H}} + \| F(0) \|_{\mathcal{H}} \right) \\ & \leq 2Me^{\omega(t-s)} (\text{Lip}(F|_{\mathfrak{B}}) \| y(s; u) \|_{\mathcal{H}} + \| F(0) \|_{\mathcal{H}}), \quad 0 \leq s \leq t \leq T, \end{aligned}$$

which implies

$$(2.28) \quad d_N(s, u) \leq 2Me^{\omega T} (\text{Lip}(F|_{\mathfrak{B}}) \| y(s; u) \|_{\mathcal{H}} + \| F(0) \|_{\mathcal{H}}), \quad \forall s \in [0, T].$$

Since  $y(\cdot; u)$  lies in  $C([0, T]; \mathcal{H})$ , the mapping  $s \mapsto \| y(s, u) \|_{\mathcal{H}}$  is in particular integrable on  $[0, T]$ , and the Lebesgue dominated convergence theorem allows us to conclude from (2.25) and (2.28) that

$$(2.29) \quad \lim_{N \rightarrow \infty} \int_0^T d_N(s; u) ds = 0.$$

Let us estimate  $\int_0^T \tilde{d}_N(s; u) ds$  as  $N \rightarrow \infty$ . By using the growth condition (2.14), we get

$$(2.30) \quad \| (T(t-s) - e^{LN(t-s)} \Pi_N) \mathfrak{C}(u(s)) \|_{\mathcal{H}} \leq 2Me^{\omega T} \left( \gamma_1 \| u(s) \|_V^p + \gamma_2 \right), \text{ for a.e. } s \in [0, T] \text{ and all } t \in [s, T].$$

Since  $u$  lies in  $L^q([0, T]; V)$  with  $q \geq p$ , then  $u$  lies in  $L^p([0, T]; V)$  and the right hand side (RHS) of (2.30) is integrable on  $[0, T]$ . The Lebesgue dominated convergence theorem allows us then to conclude from (2.26) and (2.30) that

$$(2.31) \quad \lim_{N \rightarrow \infty} \int_0^T \tilde{d}_N(s; u) ds = 0.$$

The desired convergence result (2.15) follows now from (2.22) by using (2.24), (2.29) and (2.31).  $\square$



**2.3. Local-in- $u$  approximation result.** In this section, we consider for  $q \geq 1$ ,  $\mathcal{U}_{ad}$ , to be the subset of  $L^q(0, T; V)$ , constituted by measurable functions that take values in  $U$ , a bounded subset of the Hilbert space  $V$ . In other words,

$$(2.32) \quad \mathcal{U}_{ad} := \{f \in L^q(0, T; V) : f(s) \in U \text{ for a.e. } s \in [0, T]\}, \quad q \geq 1.$$

The set  $\mathcal{U}_{ad}$  will be endowed with the induced topology from that of  $L^q(0, T; V)$ .

We present hereafter a natural property that is derived from our working assumptions, namely that given an finite-dimensional approximation  $\mathcal{H}_N$  of  $\mathcal{H}$ , the residual energy of solutions to the IVP (2.1), i.e.

$$\|(\text{Id}_{\mathcal{H}} - \Pi_N)y(t; x, w)\|_{\mathcal{H}},$$

can be made small as  $N \rightarrow \infty$  and uniformly in  $w$ , provided that  $w$  lies within a sufficiently small open set of  $\mathcal{U}_{ad}$  given in (2.32).

This is the purpose of Lemma 2.2 which boils down to proving the continuity of the mapping  $u \mapsto y(t; x, u)$ ; see (2.45) below. As a consequence a local-in- $u$  approximation result is naturally inferred; see Lemma 2.3. However, as pointed out and amended in Sect. 2.4 below, this is insufficient to guarantee convergence results for the value functions associated with Galerkin approximations of optimal control problems subordinated to (2.1).

The merit of Lemma 2.2 below is nevertheless not only to identify a symptom, but also to help us propose a cure. Indeed, by requiring the residual energy of the solution to the IVP (2.1) to vanish uniformly (in  $u$ ) as  $N \rightarrow \infty$ , we are able to conclude about the desired convergence results for the value functions. The latter uniform property (i.e. the ‘‘cure’’) is shown below to hold for a broad class of IVPs; see Sect. 2.7. For the moment, let us present the ‘‘symptom,’’ i.e. the local-in- $u$  approximation results. For that purpose, we start with a local-in- $u$  estimate about the residual energy,

$$\|(\text{Id}_{\mathcal{H}} - \Pi_N)y(t; x, w)\|_{\mathcal{H}},$$

where  $w$  lives in some neighborhood of  $u$ .

**Lemma 2.2.** *Assume that (A0)–(A4) hold. Assume furthermore that  $\mathfrak{C} : V \rightarrow \mathcal{H}$  is locally Lipschitz and that the admissible controls lie in  $\mathcal{U}_{ad}$  given by (2.32) and endowed with the induced  $L^q(0, T; V)$ -topology, for  $q > 1$ .*

*Then, for any  $(x, u)$  in  $\mathcal{H} \times \mathcal{U}_{ad}$  and any  $\epsilon > 0$  there exists a neighborhood  $\mathcal{O}_u \subset \mathcal{U}_{ad}$  of  $u$  and  $N_0 \geq 0$  such that the mild solution  $y(t; x, u)$  to (2.1) satisfies*

$$(2.33) \quad \sup_{w \in \mathcal{O}_u} \sup_{t \in [0, T]} \|(Id_{\mathcal{H}} - \Pi_N)y(t; x, w)\|_{\mathcal{H}} \leq \epsilon, \quad \forall N \geq N_0.$$

*Proof.* As we will see, we mainly need to show that the solution  $y(t; x, u)$  to the IVP (2.1) depends continuously on the control  $u$  in  $\mathcal{U}_{ad}$ , endowed with the  $L^q(0, T; V)$ -topology. The above estimate (2.33) follows then directly from this continuous dependence as explained at the end of the proof.

Given  $u$  in  $\mathcal{U}_{ad}$  and  $r > 0$ , we denote by  $\mathfrak{B}_{\mathcal{U}_{ad}}(u, r)$  the closed ball centered at  $u$  with radius  $r$ , for the induced  $L^q(0, T; V)$ -topology on  $\mathcal{U}_{ad}$ . For any  $w$  in  $\mathfrak{B}_{\mathcal{U}_{ad}}(u, r)$ ,  $x$  in  $\mathcal{H}$  and  $t$  in  $[0, T]$ , consider  $\phi(t) := y(t; x, u) - y(t; x, w)$  and note that  $\phi(0) = 0$ . It follows from (2.4) that

$$(2.34) \quad \phi(t) = \int_0^t T(t-s)(F(y(s; x, u)) - F(y(s; x, w)))ds + \int_0^t T(t-s)(\mathfrak{C}(u(s)) - \mathfrak{C}(w(s)))ds.$$

We have then

$$(2.35) \quad \|\phi(t)\|_{\mathcal{H}} \leq \int_0^t Me^{\omega(t-s)} \|F(y(s; x, u)) - F(y(s; x, w))\|_{\mathcal{H}} ds + \int_0^t Me^{\omega(t-s)} \|\mathfrak{C}(u(s)) - \mathfrak{C}(w(s))\|_{\mathcal{H}} ds.$$

Let  $C > 0$  be chosen such that  $\|y(t; x, u)\|_{\mathcal{H}} \leq C$  for all  $t \in [0, T]$ , and let  $w$  be in  $\mathfrak{B}_{\mathcal{U}_{ad}}(u, r)$ , we define then

$$(2.36) \quad t^* := \max\{t \in [0, T] : \|y(t; x, w)\|_{\mathcal{H}} < 2C\}.$$

First let us note that  $t^* > 0$ . Indeed recalling that  $\|y(t; x, u)\|_{\mathcal{H}} \leq C$  by assumption, we have in particular that  $\|x\|_{\mathcal{H}} \leq C$ . Now due to the continuity for any  $(x, w)$  in  $\mathcal{H} \times \mathcal{U}_{ad}$  of the mapping

$$\begin{aligned} [0, T] &\longrightarrow \mathcal{H} \\ t &\longmapsto y(t; x, w), \end{aligned}$$

(since  $y(t; x, w)$  is a mild solution), we infer, since  $\|x\|_{\mathcal{H}} \leq C$ , for each  $w$  in  $\mathfrak{B}_{\mathcal{U}_{ad}}(u, r)$  the existence of  $t'(w) > 0$  such that

$$\|y(t; x, w)\|_{\mathcal{H}} < 2C, \quad \forall t \in [0, t'(w)],$$

and therefore  $t^* > 0$ .

Denote also  $\mathfrak{B}_{\mathcal{H}} \subset \mathcal{H}$  the closed ball centered at the origin with radius  $2C$ . Let  $\mathfrak{B}_V$  be the smallest closed ball in  $V$  containing the bounded set  $U$ . By using the local Lipschitz property of  $F$  and  $\mathfrak{C}$ , we obtain from (2.35) that

$$(2.37) \quad \begin{aligned} \|\phi(t)\|_{\mathcal{H}} &\leq \int_0^t Me^{\omega(t-s)} \text{Lip}(F|_{\mathfrak{B}_{\mathcal{H}}}) \|\phi(s)\|_{\mathcal{H}} ds + \int_0^t Me^{\omega(t-s)} \text{Lip}(\mathfrak{C}|_{\mathfrak{B}_V}) \|u(s) - w(s)\|_V ds \\ &\leq M \text{Lip}(F|_{\mathfrak{B}_{\mathcal{H}}}) \int_0^t e^{\omega(t-s)} \|\phi(s)\|_{\mathcal{H}} ds + e^{\omega t^*} M \text{Lip}(\mathfrak{C}|_{\mathfrak{B}_V}) \int_0^t \|u(s) - w(s)\|_V ds, \quad \forall t \in [0, t^*]. \end{aligned}$$

By Hölder's inequality, we have

$$(2.38) \quad \int_0^t \|u(s) - w(s)\|_V ds \leq t^{\frac{p-1}{p}} \|u - w\|_{L^p(0, T; V)} \leq t^{\frac{p-1}{p}} r,$$

which leads to

$$(2.39) \quad \|\phi(t)\|_{\mathcal{H}} \leq M \text{Lip}(F|_{\mathfrak{B}_{\mathcal{H}}}) \int_0^t e^{\omega(t-s)} \|\phi(s)\|_{\mathcal{H}} ds + (t^*)^{\frac{p-1}{p}} r e^{\omega t^*} M \text{Lip}(\mathfrak{C}|_{\mathfrak{B}_V}), \quad \forall t \in [0, t^*].$$

It follows then from Gronwall's inequality that

$$(2.40) \quad \begin{aligned} \|\phi(t)\|_{\mathcal{H}} &\leq (t^*)^{\frac{p-1}{p}} r M \text{Lip}(\mathfrak{C}|_{\mathfrak{B}_V}) \exp(2\omega t^* + t^* M \text{Lip}(F|_{\mathfrak{B}_{\mathcal{H}}})) \\ &\leq T^{\frac{p-1}{p}} r M \text{Lip}(\mathfrak{C}|_{\mathfrak{B}_V}) \exp(2\omega T + T M \text{Lip}(F|_{\mathfrak{B}_{\mathcal{H}}})) , \quad \forall t \in [0, t^*]. \end{aligned}$$

Now, let  $C_* := T^{\frac{p-1}{p}} M \text{Lip}(\mathfrak{C}|_{\mathfrak{B}_V}) \exp(2\omega T + T M \text{Lip}(F|_{\mathfrak{B}_{\mathcal{H}}}))$  and

$$r_1 := \frac{C}{2C_*}.$$

We claim that  $t^* = T$  if  $r \leq r_1$ . Otherwise, if  $t^* < T$ , applying (2.40) at  $t = t^*$ , we get

$$(2.41) \quad \|\phi(t^*)\|_{\mathcal{H}} \leq \frac{C}{2},$$

which leads then to

$$(2.42) \quad \|y(t^*; x, w)\|_{\mathcal{H}} \leq \|y(t^*; x, u)\|_{\mathcal{H}} + \|\phi(t^*)\|_{\mathcal{H}} \leq \frac{3}{2}C.$$

This last inequality contradicts with the definition of  $t^*$  given by (2.36).

We obtain thus for each  $r \in (0, r_1]$  that

$$(2.43) \quad \|\phi(t)\|_{\mathcal{H}} \leq r C_* , \quad \forall t \in [0, T].$$

Now, it follows from (2.43) that for any fixed  $\epsilon > 0$  there exists  $r_\epsilon > 0$  sufficiently small such that

$$(2.44) \quad \|\phi(t)\|_{\mathcal{H}} \leq \frac{1}{2}\epsilon, \quad \forall t \in [0, T].$$

Recalling the definition of  $\phi$ , we have thus proved that

$$(2.45) \quad \sup_{w \in \mathfrak{B}_{\mathcal{U}_{ad}}(u, r_\epsilon)} \sup_{t \in [0, T]} \|y(t; x, u) - y(t; x, w)\|_{\mathcal{H}} \leq \frac{1}{2}\epsilon.$$

We turn now to the last arguments needed to prove (2.33). It consists first to note that the convergence property (2.6) and the continuity of  $t \mapsto y(t; x, u)$  imply, for the given  $\epsilon > 0$ , the existence of a positive integer  $N_0$  for which

$$(2.46) \quad \sup_{t \in [0, T]} \|(\text{Id}_{\mathcal{H}} - \Pi_N)y(t; x, u)\|_{\mathcal{H}} \leq \frac{1}{2}\epsilon, \quad \forall N \geq N_0.$$

Now by defining  $\Pi_N^\perp := \text{Id}_{\mathcal{H}} - \Pi_N$ , and noting that

$$(2.47) \quad \begin{aligned} \|\Pi_N^\perp y(t; x, w)\|_{\mathcal{H}} &\leq \|\Pi_N^\perp(y(t; x, w) - y(t; x, u))\|_{\mathcal{H}} + \|\Pi_N^\perp y(t; x, u)\|_{\mathcal{H}} \\ &\leq \|y(t; x, w) - y(t; x, u)\|_{\mathcal{H}} + \|\Pi_N^\perp y(t; x, u)\|_{\mathcal{H}}, \end{aligned}$$

we conclude—from (2.45) and (2.46)—to the desired estimate (2.33) with  $\mathcal{O}_u$  taken to be  $\mathfrak{B}_{\mathcal{U}_{ad}}(u, r_\epsilon)$ , the open ball in  $\mathcal{U}_{ad}$  of radius  $r_\epsilon$ . □

We conclude this section with a local-in- $u$  approximation result.

**Lemma 2.3.** *Assume the assumptions of Lemma 2.2 hold. Then, for any  $(x, u)$  in  $\mathcal{H} \times \mathcal{U}_{ad}$  and any  $\epsilon > 0$  there exists a neighborhood  $\mathcal{O}_u \subset \mathcal{U}_{ad}$  of  $u$  and  $N_0 \geq 1$  such that the mild solution  $y(t; x, u)$  to (2.1) satisfies*

$$(2.48) \quad \sup_{w \in \mathcal{O}_u} \sup_{t \in [0, T]} \|y_N(t; \Pi_N x, w) - y(t; x, w)\|_{\mathcal{H}} \leq \epsilon, \quad \forall N \geq N_0,$$

where  $y_N$  denotes the solution to the Galerkin approximation (2.8).

*Proof.* First, let us remark that even if here  $\mathfrak{C}$  does not satisfy the growth condition (2.14), one can still derive the trajectory-wise convergence result (2.15) by exploiting the fact that  $u$  lies in  $\mathcal{U}_{ad}$  and  $\mathfrak{C}$  is locally Lipschitz. The only change in the proof consists indeed of replacing the estimate (2.30) by the following:

$$(2.49) \quad \|(T(t-s) - e^{L_N(t-s)}\Pi_N)\mathfrak{C}(u(s))\|_{\mathcal{H}} \leq 2Me^{\omega T} \text{Lip}(\mathfrak{C}|_{\mathfrak{B}_V}) \|u(s)\|_V, \text{ for a.e. } s \in [0, T] \text{ and all } t \in [s, T],$$

where  $\mathfrak{B}_V$  denotes the smallest closed ball in  $V$  containing the bounded set  $U$ . Since  $u$  lies in  $L^q([0, T]; V)$ , then  $u$  lies in  $L^1([0, T]; V)$  and the RHS of (2.49) is integrable on  $[0, T]$ . We can then still use the Lebesgue dominated convergence theorem to obtain (2.31) as before.

We explain now how to derive (2.48) from (2.15). Recall that from (2.45) derived in the proof of Lemma 2.2, the mild solutions to the IVP (2.1) depend continuously on the control  $u$  in  $\mathcal{U}_{ad}$ , endowed with the  $L^q([0, T]; V)$ -topology. An estimate similar to (2.45) can be derived for the solutions to Galerkin approximation (2.8), ensuring thus also their continuous dependence on  $u$ .

As a consequence, for any given  $\epsilon > 0$  and  $u$  in  $\mathcal{U}_{ad}$ , there exists a neighborhood  $\mathcal{O}_u \subset \mathcal{U}_{ad}$  containing  $u$  for which

$$(2.50) \quad \begin{aligned} \sup_{w \in \mathcal{O}_u} \sup_{t \in [0, T]} \|y(t; x, u) - y(t; x, w)\|_{\mathcal{H}} &\leq \frac{1}{3}\epsilon, \\ \sup_{w \in \mathcal{O}_u} \sup_{t \in [0, T]} \|y_N(t; \Pi_N x, u) - y_N(t; \Pi_N x, w)\|_{\mathcal{H}} &\leq \frac{1}{3}\epsilon, \quad \forall N \in \mathbb{Z}_+^*. \end{aligned}$$

On the other hand, the trajectory-wise convergence (2.15) ensures the existence of a positive integer  $N_0$ , such that

$$(2.51) \quad \sup_{t \in [0, T]} \|y_N(t; \Pi_N x, u) - y(t; x, u)\|_{\mathcal{H}} \leq \frac{1}{3}\epsilon, \quad \forall N \geq N_0.$$

Since

$$(2.52) \quad \begin{aligned} \|y_N(t; \Pi_N x, w) - y(t; x, w)\|_{\mathcal{H}} &\leq \|y_N(t; \Pi_N x, w) - y_N(t; \Pi_N x, u)\|_{\mathcal{H}} \\ &\quad + \|y_N(t; \Pi_N x, u) - y(t; x, u)\|_{\mathcal{H}} \\ &\quad + \|y(t; x, u) - y(t; x, w)\|_{\mathcal{H}}, \end{aligned}$$

the desired estimate (2.48) follows from (2.50) and (2.51). □

**2.4. Convergence of Galerkin approximations: Uniform-in- $u$  result.** In the previous section, the local-in- $u$  approximation result has been derived under a boundedness assumption on  $U$  arising in the definition of  $\mathcal{U}_{ad}$ . Here compactness will substitute the boundedness to derive convergence results that are uniform in  $u$ . More precisely, we will assume for that purpose

**(A5)** *The set of admissible controls  $\mathcal{U}_{ad}$  is given by (2.32) with  $U$  being a compact subset of the Hilbert space  $V$ .*

We will make also use of the following assumptions.

**(A6)** *Let  $\mathcal{U}_{ad}$  be given by (2.32). For each  $T > 0$ ,  $(x, u)$  in  $\mathcal{H} \times \mathcal{U}_{ad}$ , the problem (2.1) admits a unique mild solution  $y(\cdot; x, u)$  in  $C([0, T], \mathcal{H})$ , and for each  $N \geq 0$ , its Galerkin approximation (2.8) admits a unique solution  $y_N(\cdot; \Pi_N x, u)$  in  $C([0, T], \mathcal{H})$ . Moreover, there exists a constant  $\mathcal{C} := \mathcal{C}(T, x)$  such that*

$$(2.53) \quad \begin{aligned} \|y(t; x, u)\|_{\mathcal{H}} &\leq \mathcal{C}, \quad \forall t \in [0, T], u \in \mathcal{U}_{ad}, \\ \|y_N(t; \Pi_N x, u)\|_{\mathcal{H}} &\leq \mathcal{C}, \quad \forall t \in [0, T], N \in \mathbb{Z}_+^*, u \in \mathcal{U}_{ad}. \end{aligned}$$

**(A7)** *Let  $\mathcal{U}_{ad}$  be given by (2.32). For each fixed  $T > 0$ , and any mild solution  $y(\cdot; x, u)$  to (2.1) with  $(x, u)$  in  $\mathcal{H} \times \mathcal{U}_{ad}$ , it holds that*

$$(2.54) \quad \lim_{N \rightarrow \infty} \sup_{u \in \mathcal{U}_{ad}} \sup_{t \in [0, T]} \|(\text{Id}_{\mathcal{H}} - \Pi_N)y(t; x, u)\|_{\mathcal{H}} = 0.$$

**Remark 2.1.**

- (i) Note that **(A6)** differs from **(A4)** by the inequality  $\|y(t; x, u)\|_{\mathcal{H}} \leq \mathcal{C}$ , and that the constant in (2.53) is independent of the control  $u$ .
- (ii) Let  $u$  be in  $\mathcal{U}_{ad}$  given by (2.32). Then uniform bounds such as in (2.53) are guaranteed if e.g. an a priori estimate of the following type holds for the IVPs (2.1) and (2.8):

$$(2.55) \quad \sup_{t \in [0, T]} \|y(t; x, u)\|_{\mathcal{H}} \leq \alpha(\|x\|_{\mathcal{H}} + \|u\|_{L^q(0, T; V)}) + \beta, \quad \alpha > 0, \beta \geq 0.$$

- (iii) We refer to Sect. 2.7 below for a broad class of IVPs for which Assumption **(A7)** holds.

As a preparatory lemma to Theorem 2.1, we first prove that given  $x$  in  $\mathcal{H}$ ,

$$(\text{Id}_{\mathcal{H}} - \Pi_N)F(y(t; x, u)) \xrightarrow{N \rightarrow \infty} 0,$$

uniformly in  $u$  lying in  $\mathcal{U}_{ad}$  and  $t$  in  $[0, T]$ . For that only assumptions **(A3)**, **(A6)** and **(A7)** are used, and  $U$  involved in the definition (2.32) of  $\mathcal{U}_{ad}$  is assumed to be bounded (not necessarily compact). We have

**Lemma 2.4.** *Assume that **(A3)**, **(A6)** and **(A7)** hold. Then,*

$$(2.56) \quad \lim_{N \rightarrow \infty} \sup_{u \in \mathcal{U}_{ad}} \sup_{t \in [0, T]} \|(\text{Id}_{\mathcal{H}} - \Pi_N)F(y(t; x, u))\|_{\mathcal{H}} = 0.$$

*Proof.* For any given  $\epsilon > 0$ , by **(A7)**, there exists  $N_0$  in  $\mathbb{Z}_+^*$  such that

$$(2.57) \quad \sup_{u \in \mathcal{U}_{ad}} \sup_{t \in [0, T]} \|(\text{Id}_{\mathcal{H}} - \Pi_N)y(t; x, u)\|_{\mathcal{H}} \leq \epsilon, \quad \forall N \geq N_0.$$

Note also that for all  $x$  in  $\mathcal{H}$ , and  $u$  in  $\mathcal{U}_{ad}$  given by (2.32)

$$(2.58) \quad \begin{aligned} \|(\text{Id}_{\mathcal{H}} - \Pi_N)F(y(t; x, u))\|_{\mathcal{H}} &\leq \|(\text{Id}_{\mathcal{H}} - \Pi_N)F(\Pi_{N_0}y(t; x, u))\|_{\mathcal{H}} \\ &\quad + \|(\text{Id}_{\mathcal{H}} - \Pi_N)(F(y(t; x, u)) - F(\Pi_{N_0}y(t; x, u)))\|_{\mathcal{H}}. \end{aligned}$$

For the second term on the RHS, we have

$$(2.59) \quad \begin{aligned} &\|(\text{Id}_{\mathcal{H}} - \Pi_N)(F(y(t; x, u)) - F(\Pi_{N_0}y(t; x, u)))\|_{\mathcal{H}} \\ &\leq \|F(y(t; x, u)) - F(\Pi_{N_0}y(t; x, u))\|_{\mathcal{H}} \\ &\leq \text{Lip}(F|_{\mathfrak{B}}) \|(\text{Id}_{\mathcal{H}} - \Pi_{N_0})y(t; x, u)\|_{\mathcal{H}}, \quad \forall t \in [0, T], u \in \mathcal{U}_{ad}, \end{aligned}$$

where  $\mathfrak{B}$  denotes the ball in  $\mathcal{H}$  centered at the origin with radius  $\mathcal{C}$  given by (2.53).

It follows then from (2.57) that

$$(2.60) \quad \sup_{u \in \mathcal{U}_{ad}} \sup_{t \in [0, T]} \|(\text{Id}_{\mathcal{H}} - \Pi_N)(F(y(t; x, u)) - F(\Pi_{N_0}y(t; x, u)))\|_{\mathcal{H}} \leq \text{Lip}(F|_{\mathfrak{B}}) \epsilon.$$

Due to **(A6)**, for each  $x$  in  $\mathcal{H}$ ,  $\Pi_{N_0}y(s; x, u)$  lies in a uniformly (in  $s$  and  $u$ ) bounded subset of  $\mathcal{H}_{N_0}$ . The finite dimensionality of  $\mathcal{H}_{N_0}$  (as a Galerkin subspace) ensures the compactness of the set

$$\mathcal{E}_x := \overline{\{\Pi_{N_0}y(s; x, u), u \in \mathcal{U}_{ad}, s \in [0, T]\}}^{\mathcal{H}}.$$

For each  $z$  in  $\mathcal{E}_x$  and  $\epsilon > 0$ , due to (2.6) there exists an integer  $N_1(z)$  such that

$$\|(\text{Id}_{\mathcal{H}} - \Pi_N)F(z)\| \leq \frac{\epsilon}{2}, \quad \forall N \geq N_1(z).$$

Since  $F$  is continuous, there exists a neighborhood  $\mathcal{N}_z$  of  $z$  in  $\mathcal{H}_{N_0}$  such that

$$(2.61) \quad \|(\text{Id}_{\mathcal{H}} - \Pi_N)F(w)\| \leq \epsilon, \quad \forall N \geq N_1(z), w \in \mathcal{N}_z.$$

From the compactness of  $\mathcal{E}_x$  we can extract a finite cover of  $\mathcal{E}_x$  by such neighborhoods  $\mathcal{N}_z$  for which (2.61) holds, and thus one can ensure the existence of an integer  $N_1$  for which

$$(2.62) \quad \sup_{z \in \mathcal{E}_x} \|(\text{Id}_{\mathcal{H}} - \Pi_N)F(z)\|_{\mathcal{H}} \leq \epsilon, \quad \forall N \geq N_1.$$

This last inequality ensures for each  $x$  in  $\mathcal{H}$ , the existence an integer  $N_1$  for which

$$(2.63) \quad \sup_{u \in \mathcal{U}_{ad}} \sup_{t \in [0, T]} \|(\text{Id}_{\mathcal{H}} - \Pi_N)F(\Pi_{N_0}y(t; x, u))\|_{\mathcal{H}} \leq \epsilon, \quad \forall N \geq N_1.$$

Then, (2.56) follows from (2.58), (2.60) and (2.63). □

We are now in position to formulate a uniform (in  $u$ ) version of Lemma 2.1 in which the growth condition (2.14) is no longer required.

**Theorem 2.1.** *Assume that (A0)–(A3) and (A5)–(A7) hold. Assume also that  $\mathfrak{C} : V \rightarrow \mathcal{H}$  is continuous. Then, for any  $(x, u)$  in  $\mathcal{H} \times \mathcal{U}_{ad}$ , the mild solution  $y(t; x, u)$  to (2.1) satisfies the following uniform convergence result*

$$(2.64) \quad \lim_{N \rightarrow \infty} \sup_{u \in \mathcal{U}_{ad}} \sup_{t \in [0, T]} \|y_N(t; \Pi_N x, u) - y(t; x, u)\|_{\mathcal{H}} = 0,$$

where  $y_N$  denotes the solution to the Galerkin approximation (2.8).

*Proof.* Compared to Lemma 2.1, we have replaced (A4) by the stronger assumption (A6) and the set of admissible controls  $\mathcal{U}_{ad}$  is as given in (A5). By following the proof of Lemma 2.1, in order to obtain the uniform convergence result (2.64), it suffices to show that the two terms  $\int_0^T d_N(s; u) ds$  and  $\int_0^T \tilde{d}_N(s; u) ds$  involved in the RHS of (2.22) converge to zero as  $N \rightarrow \infty$  uniformly with respect to  $u$  in  $\mathcal{U}_{ad}$ .

Recall from (2.19c) that  $d_N(s; u) = \sup_{t \in [s, T]} \|(T(t-s) - e^{L_N(t-s)} \Pi_N) F(y(s; x, u))\|_{\mathcal{H}}$ , which is defined for every  $s$  in  $[0, T]$  and  $u$  in  $\mathcal{U}_{ad}$ . Thanks to Lemma 2.4, for any fixed  $\epsilon > 0$ , there exists  $N_0$  in  $\mathbb{Z}_+^*$ , such that

$$(2.65) \quad \|(\text{Id}_{\mathcal{H}} - \Pi_{N_0}) F(y(s; x, u))\|_{\mathcal{H}} < \epsilon, \quad s \in [0, T], u \in \mathcal{U}_{ad}.$$

Now, for  $N_0$  chosen above, we have

$$(2.66) \quad \begin{aligned} \|(T(t-s) - e^{L_N(t-s)} \Pi_N) F(y(s; x, u))\|_{\mathcal{H}} &\leq \|(T(t-s) - e^{L_N(t-s)} \Pi_N) \Pi_{N_0} F(y(s; x, u))\|_{\mathcal{H}} \\ &\quad + \|(T(t-s) - e^{L_N(t-s)} \Pi_N) (\text{Id}_{\mathcal{H}} - \Pi_{N_0}) F(y(s; x, u))\|_{\mathcal{H}}. \end{aligned}$$

By using (2.2) and Assumption (A1), we obtain:

$$(2.67) \quad \begin{aligned} &\sup_{t \in [s, T]} \|(T(t-s) - e^{L_N(t-s)} \Pi_N) (\text{Id}_{\mathcal{H}} - \Pi_{N_0}) F(y(s; x, u))\|_{\mathcal{H}} \\ &\leq 2Me^{\omega T} \|(\text{Id}_{\mathcal{H}} - \Pi_{N_0}) F(y(s; x, u))\|_{\mathcal{H}} \\ &\leq 2Me^{\omega T} \epsilon, \quad \forall s \in [0, T], u \in \mathcal{U}_{ad}, \end{aligned}$$

where the last inequality follows from (2.65).

Due to (A6), for each  $x$  in  $\mathcal{H}$ ,  $y(s; x, u)$  lies in a uniformly (in  $s$  and  $u$ ) bounded subset of  $\mathcal{H}$ . This together with the locally Lipschitz property of  $F$  implies that  $\Pi_{N_0} F(y(s; x, u))$  lies in a bounded subset of  $\mathcal{H}_{N_0}$  for all  $u$  in  $\mathcal{U}_{ad}$  and for all  $s$  in  $[0, T]$ . The finite dimensionality of  $\mathcal{H}_{N_0}$  ensures then the compactness of the set

$$\mathcal{E}'_x = \overline{\{\Pi_{N_0} F(y(s; x, u)), u \in \mathcal{U}_{ad}, s \in [0, T]\}}^{\mathcal{H}}.$$

Now, for each  $z$  in  $\mathcal{E}'_x$  and  $\epsilon > 0$ , the convergence property (2.23) valid uniformly over bounded time-intervals allows us to ensure the existence of an integer  $N_1(z)$ , for which

$$\sup_{t \in [0, T]} \|(T(t) - e^{L_N t} \Pi_N) z\|_{\mathcal{H}} \leq \frac{\epsilon}{2}, \quad \forall N \geq N_1(z).$$

Then, by using (2.2) and Assumption (A1), there exists a neighborhood  $\mathcal{N}_z$  of  $z$  in  $\mathcal{H}_{N_0}$  such that

$$(2.68) \quad \sup_{t \in [0, T]} \|(T(t) - e^{L_N t} \Pi_N) w\|_{\mathcal{H}} \leq \epsilon, \quad \forall N \geq N_1(z), w \in \mathcal{N}_z.$$

From the compactness of  $\mathcal{E}'_x$  we can extract a finite cover of  $\mathcal{E}'_x$  by such neighborhoods in which (2.68) holds, and thus one can ensure the existence of an integer  $N_1$  for which

$$(2.69) \quad \sup_{t \in [0, T]} \|(T(t) - e^{L_N(t)} \Pi_N) w\|_{\mathcal{H}} \leq \epsilon, \quad \forall N \geq N_1, w \in \mathcal{E}'_x.$$

Now, for each fixed  $s$  in  $[0, T]$  and  $u$  in  $\mathcal{U}_{ad}$ , by taking  $w = \Pi_{N_0} F(y(s; x, u))$ , we get from (2.69) that

$$\sup_{t \in [0, T]} \|(T(t) - e^{L_N(t)} \Pi_N) \Pi_{N_0} F(y(s; x, u))\|_{\mathcal{H}} \leq \epsilon, \quad \forall N \geq N_1.$$

It follows then,

$$(2.70) \quad \sup_{t \in [s, T]} \|(T(t-s) - e^{L_N(t-s)} \Pi_N) \Pi_{N_0} F(y(s; x, u))\|_{\mathcal{H}} \leq \epsilon, \quad \forall N \geq N_1, s \in [0, T], u \in \mathcal{U}_{ad}.$$

By using (2.67) and (2.70) in (2.66), we get

$$d_N(s; u) \leq (1 + 2Me^{\omega T})\epsilon, \quad \forall N \geq N_1, s \in [0, T], u \in \mathcal{U}_{ad},$$

which leads to

$$(2.71) \quad \sup_{u \in \mathcal{U}_{ad}} \int_0^T d_N(s; u) ds \leq (1 + 2Me^{\omega T})T\epsilon, \quad \forall N \geq N_1.$$

We consider now the term  $\sup_{u \in \mathcal{U}_{ad}} \int_0^T \tilde{d}_N(s; u) ds$  with

$$\tilde{d}_N(s; u) = \sup_{t \in [s, T]} \|(T(t-s) - e^{L_N(t-s)} \Pi_N) \mathfrak{C}(u(s))\|_{\mathcal{H}}$$

as defined in (2.20) for almost every  $s$  in  $[0, T]$  and every  $u$  in  $\mathcal{U}_{ad}$  here.

Since the set  $U \subset V$  is compact (cf. Assumption **(A5)**) and  $\mathfrak{C} : V \rightarrow \mathcal{H}$  is continuous, then  $\mathfrak{C}(U)$  is a compact set of  $\mathcal{H}$ . Following a compactness argument similar to that used to derive (2.69), we can ensure the existence of an integer  $N_2$  such that

$$(2.72) \quad \sup_{t \in [0, T]} \|(T(t) - e^{L_N(t)} \Pi_N) \mathfrak{C}(w)\|_{\mathcal{H}} \leq \epsilon, \quad \forall N \geq N_2, w \in U.$$

Now, for each  $u$  in  $\mathcal{U}_{ad}$ , since  $u(s)$  takes value in  $U$  for almost every  $s$  in  $[0, T]$ , we obtain from (2.72) that

$$(2.73) \quad \sup_{t \in [s, T]} \|(T(t-s) - e^{L_N(t-s)} \Pi_N) \mathfrak{C}(u(s))\|_{\mathcal{H}} \leq \epsilon, \quad \text{for all } N \geq N_2 \text{ and a.e. } s \in [0, T].$$

It follows then

$$(2.74) \quad \int_0^T \tilde{d}_N(s; u) ds = \int_0^T \sup_{t \in [s, T]} \|(T(t-s) - e^{L_N(t-s)} \Pi_N) \mathfrak{C}(u(s))\|_{\mathcal{H}} ds \leq T\epsilon, \quad \forall N \geq N_2, u \in \mathcal{U}_{ad}.$$

Namely,

$$(2.75) \quad \sup_{u \in \mathcal{U}_{ad}} \int_0^T \tilde{d}_N(s; u) ds \leq T\epsilon, \quad \forall N \geq N_2.$$

The desired uniform convergence of  $\int_0^T d_N(s; u) ds$  and  $\int_0^T \tilde{d}_N(s; u) ds$  follows from (2.71) and (2.75). The proof is now complete.  $\square$

**Remark 2.2.** From the proof given above, it is clear that Assumption **(A5)** is made to ensure the uniform convergence of  $\int_0^T \tilde{d}_N(s; u) ds$ , while Assumptions **(A6)** and **(A7)** are made to ensure the uniform convergence of  $\int_0^T d_N(s; u) ds$ . The last two assumptions are thus not needed if the nonlinear term  $F$  is identically zero.

Note that one can readily check that the convergence result stated in Theorem 2.1 also holds when (2.1) is initialized at any other time instance  $t$  in  $[0, T]$ . This will be needed in the next subsection to derive approximation results for value functions associated with optimal control problems for the IVP (2.1).

More precisely, for each  $(t, x)$  in  $[0, T] \times \mathcal{H}$ , we consider the following evolution problem

$$(2.76) \quad \begin{aligned} \frac{dy}{ds} &= Ly + F(y) + \mathfrak{C}(u(s)), \quad s \in (t, T], u \in \mathcal{U}_{ad}[t, T], \\ y(t) &= x \in \mathcal{H}, \end{aligned}$$

with

$$(2.77) \quad \mathcal{U}_{ad}[t, T] := \{u|_{[t, T]} : u \in \mathcal{U}_{ad}\},$$

and the corresponding Galerkin approximation:

$$(2.78) \quad \begin{aligned} \frac{dy_N}{ds} &= L_N y_N + \Pi_N F(y_N) + \Pi_N \mathfrak{C}(u(s)), \quad s \in (t, T], u \in \mathcal{U}_{ad}[t, T], \\ y_N(t) &= x_N, \quad x_N := \Pi_N x \in \mathcal{H}_N. \end{aligned}$$

Hereafter, we denote by  $y_{t,x}(\cdot; u)$  the solution to (2.76) emanating from  $x$  at time  $t$ , and by  $y_{t,x_N}^N(\cdot; u)$  the solution to (2.78) emanating from  $x_N := \Pi_N x$  at time  $t$ . We have then the following corollary of Theorem 2.1:

**Corollary 2.1.** Assume that the conditions of Theorem 2.1 hold. Then, for any mild solution  $y_{t,x}(\cdot; u)$  to (2.76) over  $[0, T]$ , with  $x$  in  $\mathcal{H}$  and  $u$  in  $\mathcal{U}_{ad}[t, T]$  given in (2.77), the following convergence result is satisfied:

$$(2.79) \quad \lim_{N \rightarrow \infty} \sup_{t \in [0, T]} \sup_{u \in \mathcal{U}_{ad}[t, T]} \sup_{s \in [t, T]} \|y_{t,x_N}^N(s; u) - y_{t,x}(s; u)\|_{\mathcal{H}} = 0,$$

where  $y_{t,x_N}^N$  denotes the solution to the Galerkin approximation (2.78).

*Proof.* Since both the linear operator  $L$  and the nonlinearity  $F$  in (2.76) are time independent, then the supremum of  $\sup_{u \in \mathcal{U}_{ad}[t, T]} \sup_{s \in [t, T]} \|y_{t,x_N}^N(s; u) - y_{t,x}(s; u)\|_{\mathcal{H}}$  as  $t$  varies in  $[0, T]$ , is achieved at  $t = 0$ , i.e.,

$$(2.80) \quad \begin{aligned} \sup_{t \in [0, T]} \sup_{u \in \mathcal{U}[t, T]} \sup_{s \in [t, T]} \|y_{t,x_N}^N(s; u) - y_{t,x}(s; u)\|_{\mathcal{H}} &= \sup_{u \in \mathcal{U}[0, T]} \sup_{s \in [0, T]} \|y_{0,x_N}^N(s; u) - y_{0,x}(s; u)\|_{\mathcal{H}} \\ &= \sup_{u \in \mathcal{U}} \sup_{s \in [0, T]} \|y_N(s; x_N, u) - y(s; x, u)\|_{\mathcal{H}}. \end{aligned}$$

The desired estimate (2.79) follows then from (2.64). □

**2.5. Galerkin approximations of optimal control and value functions: Convergence results.** We assume in this section that  $U$  is a compact and convex subset of the Hilbert space  $V$ . In particular this ensures that  $\mathcal{U}_{ad}$  defined in (2.32) is a bounded, closed and convex set. For such an admissible set of controls, conditions of existence to optimal control problems associated with the IVPs (2.1) and (2.8) are recalled in Appendix A.

We introduce next the cost functional,  $J: \mathcal{H} \times \mathcal{U}_{ad} \rightarrow \mathbb{R}^+$ , associated with the IVP (2.1):

$$(2.81) \quad J(x, u) := \int_0^T [\mathcal{G}(y(s; x, u)) + \mathcal{E}(u(s))] ds, \quad x \in \mathcal{H},$$



where  $\mathcal{G} : \mathcal{H} \rightarrow \mathbb{R}^+$  and  $\mathcal{E} : V \rightarrow \mathbb{R}^+$  are assumed to be *continuous*, and  $\mathcal{G}$  is assumed to satisfy furthermore the condition:

(C1)  $\mathcal{G}$  is locally Lipschitz in the sense given of (2.12).

In particular non-smooth cost functionals are allowed such as  $L^1$  (on  $u$ ) arising in fuel-optimal control problems [AF13, Chaps. 6.13].

The associated optimal control problem then writes

$$(P) \quad \min J(x, u) \quad \text{s.t.} \quad (y, u) \in L^2(0, T; \mathcal{H}) \times \mathcal{U}_{ad} \text{ solves (2.1) with } y(0) = x \in \mathcal{H}.$$

The cost functional,  $J_N : \mathcal{H}_N \times \mathcal{U}_{ad} \rightarrow \mathbb{R}^+$ , associated with the Galerkin approximation (2.8) is given by

$$(2.82) \quad J_N(\Pi_N x, u) := \int_0^T [\mathcal{G}(y_N(s; \Pi_N x, u)) + \mathcal{E}(u(s))] ds, \quad x \in \mathcal{H},$$

and the corresponding optimal control problem reads:

$$(P_N) \quad \min J_N(\Pi_N x, u) \quad \text{s.t.} \quad (y_N, u) \in L^2(0, T; \mathcal{H}_N) \times \mathcal{U}_{ad} \text{ solves (2.8) with } y_N(0) = \Pi_N x \in \mathcal{H}_N.$$

We assume hereafter that both problems, (P) and (P<sub>N</sub>), possess each a solution.

We analyze the convergence of the corresponding value functions by adopting a dynamic programming approach. For that purpose, we consider for each  $(t, x)$  in  $[0, T] \times \mathcal{H}$  a family of optimal control problems associated with (2.76) and the following cost functional  $J_{t,x}$ :

$$(2.83) \quad J_{t,x}(u) := \int_t^T [\mathcal{G}(y_{t,x}(s; u)) + \mathcal{E}(u(s))] ds, \quad t \in [0, T], \quad u \in \mathcal{U}_{ad}[t, T],$$

The cost functional associated with the corresponding Galerkin approximation (2.78) is given by

$$(2.84) \quad J_{t,x_N}^N(u) := \int_t^T [\mathcal{G}(y_{t,x_N}^N(s; u)) + \mathcal{E}(u(s))] ds, \quad t \in [0, T], \quad u \in \mathcal{U}_{ad}[t, T],$$

in which we have denoted  $\Pi_N x$  by  $x_N$ .

The value functions corresponding to the optimal control problems associated respectively with (2.76) and with (2.78), are then defined as follows:

$$(2.85a) \quad v(t, x) := \inf_{u \in \mathcal{U}_{ad}[t, T]} J_{t,x}(u), \quad \forall (t, x) \in [0, T] \times \mathcal{H} \quad \text{and} \quad v(T, x) := 0,$$

$$(2.85b) \quad v_N(t, x_N) := \inf_{u \in \mathcal{U}_{ad}[t, T]} J_{t,x_N}^N(u), \quad \forall (t, x_N) \in [0, T] \times \mathcal{H}_N \quad \text{and} \quad v_N(T, x_N) := 0.$$

We have then the following result.

**Theorem 2.2.** *Assume that the conditions in Theorem 2.1 together with (C1) hold. Furthermore, let there exists for each pair  $(t, x)$  a minimizer  $u_{t,x}^*$  (resp.  $u_{t,x}^{N,*}$ ) in  $\mathcal{U}_{ad}[t, T]$  of the minimization problem in (2.85a) (resp. in (2.85b)).*

*Then for any  $x$  in  $\mathcal{H}$ , it holds that*

$$(2.86) \quad \lim_{N \rightarrow \infty} \sup_{t \in [0, T]} |v_N(t, \Pi_N x) - v(t, x)| = 0.$$

*Proof.* By the definition of the value functions in (2.85), we have

$$(2.87) \quad v(t, x) = J_{t,x}(u_{t,x}^*) \leq J_{t,x}(u_{t,x}^{N,*}),$$

and

$$(2.88) \quad v_N(t, \Pi_N x) = J_{t,x_N}^N(u_{t,x}^{N,*}),$$

with  $x_N := \Pi_N x$ .

The inequality (2.87) and the definition of  $J$  give then

$$(2.89) \quad v(t, x) \leq \int_t^T [\mathcal{G}(y_{t,x}(s; u_{t,x}^{N,*})) + \mathcal{E}(u_{t,x}^{N,*}(s))] ds.$$

By subtracting  $J_{t,x_N}^N(u_N^*)$  on both sides of the above inequality, we get

$$(2.90) \quad \begin{aligned} v(t, x) - v_N(t, \Pi_N x) &\leq \int_t^T [\mathcal{G}(y_{t,x}(s; u_{t,x}^{N,*})) + \mathcal{E}(u_{t,x}^{N,*}(s))] ds - \int_t^T [\mathcal{G}(y_{t,x_N}^N(s; u_{t,x}^{N,*})) + \mathcal{E}(u_{t,x}^{N,*}(s))] ds \\ &= \int_t^T [\mathcal{G}(y_{t,x}(s; u_{t,x}^{N,*})) - \mathcal{G}(y_{t,x_N}^N(s; u_{t,x}^{N,*}))] ds. \end{aligned}$$

Besides, since both  $L$  and  $F$  are time-independent, it follows from **(A6)** that there exists a positive constant  $\mathcal{C}$  such that

$$(2.91) \quad \begin{aligned} \|y_{t,x}(s; u_{t,x}^{N,*})\| &\leq \mathcal{C}, \quad \forall t \in [0, T], s \in [t, T], n \in \mathbb{Z}_+^*, \\ \|y_{t,x_N}^N(s; u_{t,x}^{N,*})\| &\leq \mathcal{C}, \quad \forall t \in [0, T], s \in [t, T], n \in \mathbb{Z}_+^*. \end{aligned}$$

Now by denoting by  $\mathfrak{B}$  the ball in  $\mathcal{H}$  with radius  $\mathcal{C}$  centered at the origin, we have

$$(2.92) \quad \begin{aligned} &\int_t^T [\mathcal{G}(y_{t,x}(s; u_{t,x}^{N,*})) - \mathcal{G}(y_{t,x_N}^N(s; u_{t,x}^{N,*}))] ds \\ &\leq \text{Lip}(\mathcal{G}|_{\mathfrak{B}}) \int_t^T \|y_{t,x}(s; u_{t,x}^{N,*}) - y_{t,x_N}^N(s; u_{t,x}^{N,*})\|_{\mathcal{H}} ds \\ &\leq T \text{Lip}(\mathcal{G}|_{\mathfrak{B}}) \sup_{s \in [t, T]} \|y_{t,x}(s; u_{t,x}^{N,*}) - y_{t,x_N}^N(s; u_{t,x}^{N,*})\|_{\mathcal{H}}, \end{aligned}$$

which together with (2.90) leads to

$$(2.93) \quad v(t, x) - v_N(t, \Pi_N x) \leq T \text{Lip}(\mathcal{G}|_{\mathfrak{B}}) \sup_{s \in [t, T]} \|y(s; u_{t,x}^{N,*}) - y_{t,x_N}^N(s; u_{t,x}^{N,*})\|_{\mathcal{H}}.$$

Similarly, we have

$$(2.94) \quad v_N(t, \Pi_N x) - v(t, x) \leq T \text{Lip}(\mathcal{G}|_{\mathfrak{B}}) \sup_{s \in [t, T]} \|y(s; u_{t,x}^*) - y_{t,x_N}^N(s; u_{t,x}^*)\|_{\mathcal{H}}.$$

The convergence result (2.86) follows then from (2.93), (2.94), and Corollary 2.1. □

**Remark 2.3.** If the nonlinearity  $F$  in the IVP (2.1) depends also on time, by modifying Assumption **(A3)** accordingly, all the results of Section 2 still hold literally except that of Lemma 2.4 that needs to be amended. For instance, by replacing **(A3)** by the following assumption

**(A3')** The nonlinearity  $F : [0, T] \times \mathcal{H} \rightarrow \mathcal{H}$  satisfies that  $F(\cdot, y) \in L^\infty(0, T; \mathcal{H})$  for every  $y \in \mathcal{H}$ ,  $F(t, \cdot)$  is locally Lipschitz for almost every  $t \in [0, T]$ , and for any given bounded set  $\mathfrak{B} \subset \mathcal{H}$ , the mapping  $t \mapsto \text{Lip}(F(t, \cdot)|_{\mathfrak{B}})$  is in  $L^\infty(0, T)$ , where  $\text{Lip}(F(t, \cdot)|_{\mathfrak{B}})$  denotes the Lipschitz constant of  $F(t, \cdot)$  on the set  $\mathfrak{B}$ .

Under this new assumption, by replacing in Lemma 2.4 the supremum for  $t$  over  $[0, T]$  in (2.56) with the essential supremum, we have

$$(2.95) \quad \lim_{N \rightarrow \infty} \sup_{u \in \mathcal{U}_{ad}} \text{ess sup}_{t \in [0, T]} \|(Id_{\mathcal{H}} - \Pi_N)F(t, y(t; x, u))\|_{\mathcal{H}} = 0.$$

**2.6. Galerkin approximations of optimal control and value functions: Error estimates.** We provide in this section some simple and useful error estimates in terms of their interpretations.

For that purpose, we assume throughout this subsection the following set of assumptions collected as follows

- (E) – The linear operator  $L : D(L) \subset \mathcal{H} \rightarrow \mathcal{H}$  is self-adjoint.
- The Galerkin approximations (2.8) are constructed based on the eigen-subspaces  $\mathcal{H}_N := \text{span}\{e_k : k = 1, \dots, N\}$ ,  $N \in \mathbb{Z}_+^*$ , where the  $e_k$ 's are the eigenfunctions of  $L$ .
- Assumption (A6).
- Assumption (C1).

We take the set of admissible controls,  $\mathcal{U}_{ad}$ , to be given by (2.32). But in contrast to Sect. 2.5, the set  $U$  in the definition of  $\mathcal{U}_{ad}$  is not assumed to be compact in  $V$ .

Hereafter within this subsection,  $\mathfrak{B}$  denotes the ball in  $\mathcal{H}$  centered at the origin with radius  $\mathcal{C}$ , where  $\mathcal{C}$  is the same as given in Assumption (A6). We first starts with a basic pointwise estimate between the cost functional  $J_{t,x}$  given by (2.83) and its approximation  $J_{t,x_N}^N$  given by (2.84).

**Lemma 2.5.** *Under the set of assumptions given by (E), for any  $(t, x) \in [0, T] \times \mathcal{H}$ , and  $u \in \mathcal{U}_{ad}[t, T]$ , there exists  $\gamma > 0$ , independent of  $N$ , such that*

$$(2.96) \quad |J_{t,x}(u) - J_{t,x_N}^N(u)| \leq \text{Lip}(\mathcal{G}|_{\mathfrak{B}}) \left[ \sqrt{T-t} + \gamma(T-t) \right] \|\Pi_N^\perp y_{t,x}(\cdot; u)\|_{L^2(t,T;\mathcal{H})}, \quad \forall u \in \mathcal{U}_{ad}[t, T].$$

*Proof.* Note that by the definitions of  $J_{t,x}$  and  $J_{t,x_N}^N$  and the locally Lipschitz condition (C1) on  $\mathcal{G}$ , we have

$$(2.97) \quad |J_{t,x}(u) - J_{t,x_N}^N(u)| \leq \text{Lip}(\mathcal{G}|_{\mathfrak{B}}) \int_t^T \|y_{t,x}(s; u) - y_{t,x_N}^N(s; u)\|_{\mathcal{H}} ds,$$

where  $\mathfrak{B} \subset \mathcal{H}$  denotes a ball centered at the origin that contains  $y_{t,x}(s; u)$  and  $y_{t,x_N}^N(s; u)$  for all  $s$  in  $[t, T]$ . By rewriting  $y_{t,x}(s; u)$  as  $\Pi_N y_{t,x}(s; u) + \Pi_N^\perp y_{t,x}(s; u)$ , we have thus

$$(2.98) \quad |J_{t,x}(u) - J_{t,x_N}^N(u)| \leq \text{Lip}(\mathcal{G}|_{\mathfrak{B}}) \int_t^T \left( \|\Pi_N y_{t,x}(s; u) - y_{t,x_N}^N(s; u)\|_{\mathcal{H}} + \|\Pi_N^\perp y_{t,x}(s; u)\|_{\mathcal{H}} \right) ds.$$

The estimate (2.96) follows then immediately if one proves that there exists  $\gamma^2 > 0$  independent of  $N$  such that for any  $(t, x) \in [0, T] \times \mathcal{H}$ , and  $u \in \mathcal{U}_{ad}[t, T]$ ,

$$(2.99) \quad \|\Pi_N y_{t,x}(s; u) - y_{t,x_N}^N(s; u)\|_{\mathcal{H}}^2 \leq \gamma^2 \int_t^s \|\Pi_N^\perp y_{t,x}(s'; u)\|_{\mathcal{H}}^2 ds', \quad s \in [t, T].$$

Indeed, in such a case we get

$$(2.100) \quad \int_t^T \|\Pi_N y_{t,x}(s; u) - y_{t,x_N}^N(s; u)\|_{\mathcal{H}} ds \leq \gamma \int_t^T \left( \int_t^s \|\Pi_N^\perp y_{t,x}(s'; u)\|_{\mathcal{H}}^2 ds' \right)^{\frac{1}{2}} ds \\ \leq \gamma(T-t) \|\Pi_N^\perp y_{t,x}(\cdot; u)\|_{L^2(t,T;\mathcal{H})},$$

and by noting from Hölder's inequality that

$$\int_t^T \|\Pi_N^\perp y_{t,x}(s; u)\|_{\mathcal{H}} ds \leq \sqrt{T-t} \|\Pi_N^\perp y_{t,x}(\cdot; u)\|_{L^2(t,T;\mathcal{H})},$$

we arrive at (2.96).

We are thus left with the proof of (2.99) which is easily derived as follows. Let us introduce

$$(2.101) \quad w(s) := \Pi_N y_{t,x}(s; u) - y_{t,x_N}^N(s; u).$$

By applying  $\Pi_N$  to both sides of Eq. (2.76), we obtain that  $\Pi_N y_{t,x}(\cdot; u)$  satisfies the following IVP:

$$\begin{aligned} \frac{d\Pi_N y}{ds} &= L_N \Pi_N y + \Pi_N F(\Pi_N y + \Pi_N^\perp y_{t,x}(s; u)) + \Pi_N \mathfrak{C}(u(s)), \quad s \in (t, T], \\ \Pi_N y(t) &= \Pi_N x \in \mathcal{H}_N. \end{aligned}$$

This together with (2.78) implies that  $w$  satisfies the following problem:

$$(2.102) \quad \begin{aligned} \frac{dw}{ds} &= L_N w + \Pi_N F(\Pi_N y + \Pi_N^\perp y_{t,x}(s; u)) - \Pi_N F(y_N), \quad s \in (t, T], \\ w(t) &= 0. \end{aligned}$$

By taking the  $\mathcal{H}$ -inner product on both sides of (2.102) with  $w$ , we obtain:

$$(2.103) \quad \frac{1}{2} \frac{d\|w\|_{\mathcal{H}}^2}{ds} = \langle L_N w, w \rangle + \langle \Pi_N (F(\Pi_N y + \Pi_N^\perp y_{t,x}(s; u)) - F(y_N)), w \rangle.$$

The local Lipschitz property of  $F$  implies then that

$$(2.104) \quad \begin{aligned} \langle \Pi_N (F(\Pi_N y + \Pi_N^\perp y_{t,x}(s; u)) - F(y_N)), w \rangle &\leq \text{Lip}(F|_{\mathfrak{B}}) \|\Pi_N y + \Pi_N^\perp y_{t,x}(s; u) - y_N\|_{\mathcal{H}} \|w\|_{\mathcal{H}} \\ &\leq \text{Lip}(F|_{\mathfrak{B}}) (\|w\| + \|y_N\|_{\mathcal{H}}) \|w\|_{\mathcal{H}} \\ &\leq \text{Lip}(F|_{\mathfrak{B}}) \left( \frac{3}{2} \|w\|_{\mathcal{H}}^2 + \frac{1}{2} \|y_N\|_{\mathcal{H}}^2 \right). \end{aligned}$$

Since  $L$  is self-adjoint, we have

$$(2.105) \quad \langle L_N w(s), w(s) \rangle = \sum_{i=1}^N \beta_i \|w_i(s)\|_{\mathcal{H}}^2 \leq \beta_1 \|w(s)\|_{\mathcal{H}}^2,$$

where  $\beta_i$  is the eigenvalue associated with its  $i^{\text{th}}$  eigenmode  $e_i$ .

Using (2.104) and (2.105) in (2.103), we finally arrive at

$$(2.106) \quad \frac{1}{2} \frac{d\|w(s)\|_{\mathcal{H}}^2}{ds} \leq \left( \beta_1 + \frac{3}{2} \text{Lip}(F|_{\mathfrak{B}}) \right) \|w(s)\|_{\mathcal{H}}^2 + \frac{1}{2} \text{Lip}(F|_{\mathfrak{B}}) \|\Pi_N^\perp y_{t,x}(s; u)\|_{\mathcal{H}}^2,$$

which for all  $s$  in  $[t, T]$ , by a standard application of Gronwall's inequality, leads to

$$(2.107) \quad \begin{aligned} \|\Pi_N y_{t,x}(s; u) - y_{t,x_N}^N(s; u)\|_{\mathcal{H}}^2 &= \|w(s)\|_{\mathcal{H}}^2 \\ &\leq \text{Lip}(F|_{\mathfrak{B}}) \int_t^s e^{2[\beta_1 + \frac{3}{2} \text{Lip}(F|_{\mathfrak{B}})](s-s')} \|\Pi_N^\perp y_{t,x}(s'; u)\|_{\mathcal{H}}^2 ds' \\ &\leq e^{2[\beta_1 + \frac{3}{2} \text{Lip}(F|_{\mathfrak{B}})]T} \text{Lip}(F|_{\mathfrak{B}}) \int_t^s \|\Pi_N^\perp y_{t,x}(s'; u)\|_{\mathcal{H}}^2 ds', \end{aligned}$$

taking into account that  $w(t) = 0$  due to (2.101). The estimate (2.99) is thus verified, and the proof is complete.  $\square$

We have then

**Theorem 2.3.** *Assume the set of assumptions given by (E). Assume also that for each  $(t, x) \in [0, T] \times \mathcal{H}$ , there exists a minimizer  $u_{t,x}^*$  (resp.  $u_{t,x_N}^{N,*}$ ) for the value function  $v$  (resp.  $v_N$ ) defined in (2.85).*

*Then for any  $(t, x) \in [0, T] \times \mathcal{H}$  it holds that*

$$(2.108) \quad \begin{aligned} &|v(t, x) - v_N(t, x_N)| \\ &\leq \text{Lip}(\mathcal{G}|_{\mathfrak{B}}) \left[ \sqrt{T-t} + \gamma(T-t) \right] \left( \|\Pi_N^\perp y_{t,x}(\cdot; u_{t,x}^*)\|_{L^2(t,T;\mathcal{H})} + \|\Pi_N^\perp y_{t,x}(\cdot; u_{t,x_N}^{N,*})\|_{L^2(t,T;\mathcal{H})} \right), \end{aligned}$$

where the constant  $\gamma$  is the same as in Lemma 2.5.

*Proof.* The result is a direct consequence of Lemma 2.5. Indeed, since

$$v(t, x) = J_{t,x}(u_{t,x}^*) \leq J_{t,x}(u_{t,x_N}^{N,*}) \quad \text{and} \quad v_N(t, x_N) = J_{t,x_N}^N(u_{t,x_N}^{N,*}),$$

we get

$$v(t, x) - v_N(t, x_N) \leq J_{t,x}(u_{t,x}^{N,*}) - J_{t,x_N}^N(u_{t,x_N}^{N,*}).$$

It follows then from (2.96) that

$$(2.109) \quad v(t, x) - v_N(t, x_N) \leq \text{Lip}(\mathcal{G}|_{\mathfrak{B}}) \left[ \sqrt{T-t} + \gamma(T-t) \right] \|\Pi_N^\perp y_{t,x}(\cdot; u_{t,x_N}^{N,*})\|_{L^2(t,T;\mathcal{H})}$$

Similarly,

$$(2.110) \quad v_N(t, x_N) - v(t, x) \leq \text{Lip}(\mathcal{G}|_{\mathfrak{B}}) \left[ \sqrt{T-t} + \gamma(T-t) \right] \|\Pi_N^\perp y_{t,x}(\cdot; u_{t,x}^*)\|_{L^2(t,T;\mathcal{H})}$$

The estimate (2.108) results then from (2.109) and (2.110).  $\square$

**Remark 2.4.** Note that in the RHS of (2.108), it is not clear a priori that

$$(2.111) \quad \lim_{N \rightarrow \infty} \|\Pi_N^\perp y_{t,x}(\cdot; u_{t,x_N}^{N,*})\|_{L^2(t,T;\mathcal{H})} = 0.$$

The reason relies on the dependence on  $u_{t,x_N}^{N,*}$  of  $\|\Pi_N^\perp y_{t,x}(\cdot; u_{t,x_N}^{N,*})\|_{L^2(t,T;\mathcal{H})}$ , where  $u_{t,x_N}^{N,*}$  denotes the control synthesized from the  $N$ -dimensional Galerkin approximation.

Estimate (2.108) provides thus another perspective regarding the usage of Assumption (A7) to ensure the convergence of the value functions associated with the Galerkin approximations presented in Sect. 2.5. Note that Assumption (A7) is just a slightly strengthened form of (2.111). See Sect. 2.7 for a broad class of IVPs for which Assumption (A7) is satisfied.

**Corollary 2.2.** Assume that the conditions given in Theorem 2.3 hold. Let us also denote  $u^* := u_{0,x}^*$  and  $u_N^* := u_{0,x_N}^{N,*}$ . Assume furthermore that there exists  $\sigma > 0$  such that the following local growth condition is satisfied for the cost functional  $J$  defined in (2.81):

$$(2.112) \quad \sigma \|u^* - v\|_{L^q(0,T;V)}^q \leq J(x, v) - J(x, u^*),$$

for all  $v$  in some neighborhood  $\mathcal{W} \subset \mathcal{U}_{ad}$  of  $u^*$ , with  $\mathcal{U}_{ad}$  given by (2.32). Assume finally that  $u_N^*$  lies in  $\mathcal{W}$ . Then,

$$(2.113) \quad \|u^* - u_N^*\|_{L^q(0,T;V)}^q \leq \frac{1}{\sigma} \text{Lip}(\mathcal{G}|_{\mathfrak{B}}) \left[ \sqrt{T} + \gamma T \right] \left( \|\Pi_N^\perp y^*(\cdot; u^*)\|_{L^2(0,T;\mathcal{H})} + 2\|\Pi_N^\perp y(\cdot; u_N^*)\|_{L^2(0,T;\mathcal{H})} \right),$$

where the constant  $\gamma$  is the same as in Lemma 2.5.

*Proof.* By the assumptions, we have

$$(2.114) \quad \|u^* - u_N^*\|_{L^q(0,T;V)}^q \leq \frac{1}{\sigma} (J(x, u_N^*) - J(x, u^*)).$$

Note also that

$$(2.115) \quad \begin{aligned} J(x, u_N^*) - J(x, u^*) &= J(x, u_N^*) - J_N(x_N, u_N^*) + J_N(x_N, u_N^*) - J(x, u^*) \\ &= J_{0,x}(u_N^*) - J_{0,x_N}^N(u_N^*) + v_N(0, x_N) - v(0, x), \end{aligned}$$

where we used the fact that

$$J(x, u_N^*) = J_{0,x}(u_N^*), \quad J_N(x_N, u_N^*) = J_{0,x_N}^N(u_N^*), \quad J_N(x_N, u_N^*) = v_N(0, x_N) \quad \text{and} \quad J(x, u^*) = v(0, x).$$

The result follows by applying the estimate (2.96) to  $J_{0,x}(u_N^*) - J_{0,x_N}^N(u_N^*)$  and the estimate (2.108) to  $v_N(0, x_N) - v(0, x)$ .

□

**Remark 2.5.** Note that (2.112) ensures uniqueness of the local minimizer  $u^*$  in  $\mathcal{W}$ .

**2.7. Examples that satisfy Assumption (A7).** We consider in this subsection a special but important case of Galerkin approximations (2.8) built from the eigenfunctions<sup>3</sup>  $\{e_k\}_{k \geq 1}$  of  $L$ , and for which we assume the following properties:

- (i) The set of admissible controls  $\mathcal{U}_{ad}$  is given as in (2.32) with  $q > 1$ .
- (ii) The linear operator  $L : D(L) \subset \mathcal{H} \rightarrow \mathcal{H}$  is self-adjoint with compact resolvent and satisfies Assumption (A0).
- (iii) The mapping  $F : \mathcal{H} \rightarrow \mathcal{H}$  and  $\mathfrak{C} : V \rightarrow \mathcal{H}$  are locally Lipschitz, and  $\mathfrak{C}(0) = 0$ .
- (iv) Assumption (A6) is satisfied.

We have then the following useful Lemma for applications; see Sect. 3.

**Lemma 2.6.** The convergence property (2.54) in (A7) holds under assumptions (i)-(iv) above.

*Proof.* Since  $L$  is assumed to be self-adjoint with compact resolvent, it follows from spectral theory of self-adjoint compact operator [Bré10, Thm. 6.8, Prop. 6.9, and Thm. 6.11] that the eigenfunctions of  $L$  form an orthonormal basis of  $\mathcal{H}$ , and the eigenvalues  $\{\beta_k\}_{k \geq 1}$  of  $L$  approach either  $\infty$  or  $-\infty$  as  $k$  approaches  $\infty$ . Since  $L$  is also assumed to be the infinitesimal generator of a  $C_0$ -semigroup (Assumption (A0)),  $\beta_k$  is bounded above [Paz83, Thm. 5.3]. It follows then that

$$(2.116) \quad \beta_N \rightarrow -\infty \quad \text{as} \quad N \rightarrow \infty.$$

As before  $\Pi_N : \mathcal{H} \rightarrow \mathcal{H}_N$  denotes the orthogonal projector associated with  $\mathcal{H}_N$  spanned by the first  $N$  eigenfunctions. We also denote  $\Pi_N^\perp$  to be the orthogonal projector associated with the orthogonal complement of  $\mathcal{H}_N$  in  $\mathcal{H}$ , namely

$$(2.117) \quad \Pi_N^\perp := \text{Id} - \Pi_N.$$

Now, by applying  $\Pi_N^\perp$  to both sides of (2.4) and introducing the notation,  $y_{N>} := \Pi_N^\perp y$ , we get:

$$(2.118) \quad y_{N>}(t) = T(t)y_{N>}(0) + \int_0^t T(t-s)\Pi_N^\perp(F(y(s;x,u)) + \mathfrak{C}(u(s)))ds, \quad t \in [0, T],$$

where we also used the fact that  $\Pi_N^\perp$  commutes with the semigroup  $\{T(t)\}_{t \geq 0}$ .

Let  $\mathcal{C}$  be the constant arising in the upper bounds of (2.53) in Assumption (A6), and denote by  $\mathfrak{B}_{\mathcal{H}}$  the closed ball in  $\mathcal{H}$  centered at the origin with radius  $\mathcal{C}$ . Let also  $\mathfrak{B}_V$  be the smallest closed ball in  $V$  containing the compact set  $U$  given in (2.32). We get then for  $t \in [0, T]$ :

$$(2.119) \quad \begin{aligned} |y_{N>}(t)| &\leq e^{\beta_{N+1}t}|y_{N>}(0)| + \text{Lip}(F|_{\mathfrak{B}_{\mathcal{H}}})(\mathcal{C} + \|F(0)\|_{\mathcal{H}}) \int_0^t e^{\beta_{N+1}(t-s)}ds + \text{Lip}(\mathfrak{C}|_{\mathfrak{B}_V}) \int_0^t e^{\beta_{N+1}(t-s)}\|u(s)\|_V ds \\ &\leq e^{\beta_{N+1}t}|y_{N>}(0)| + \frac{1}{|\beta_{N+1}|} \text{Lip}(F|_{\mathfrak{B}_{\mathcal{H}}})(\mathcal{C} + \|F(0)\|_{\mathcal{H}}) + \frac{1}{(|\beta_{N+1}|q')^{\frac{1}{q'}}} \text{Lip}(\mathfrak{C}|_{\mathfrak{B}_V})\|u\|_{L^q([0,T;V])}, \end{aligned}$$

where we have used the Hölder inequality to estimate  $\int_0^t e^{\beta_{N+1}(t-s)}\|u(s)\|_V ds$  with  $q' = \frac{q}{q-1}$ .

Now, let  $C_U := \sup_{w \in U} \|w\|_V$ . We have

$$(2.120) \quad \|u\|_{L^q(0,T;V)} \leq C_U T^{\frac{1}{q}}, \quad \forall u \in \mathcal{U}_{ad}.$$

<sup>3</sup>i.e. with  $\mathcal{H}_N := \text{span}\{e_1, \dots, e_N\}$  and  $L_N := \Pi_N L \Pi_N$ , for which Assumptions (A0)-(A2) are satisfied.

Using the above bound in (2.119), we obtain

$$(2.121) \quad \begin{aligned} |y_{N>}(t)| &\leq e^{\beta_{N+1}t} |y_{N>}(0)| + \frac{1}{|\beta_{N+1}|} \text{Lip}(F|_{\mathfrak{B}_{\mathcal{H}}})(\mathcal{C} + \|F(0)\|_{\mathcal{H}}) \\ &\quad + \frac{1}{(|\beta_{N+1}|q')^{\frac{1}{q'}}} C_U T^{\frac{1}{q'}} \text{Lip}(\mathfrak{C}|_{\mathfrak{B}_V}), \quad \forall t \in [0, T], u \in \mathcal{U}_{ad}. \end{aligned}$$

Since  $\beta_N$  approaches  $-\infty$  as  $N$  approaches  $\infty$  by assumption, we get that for each  $\epsilon > 0$ ,  $T > 0$  and  $x$  in  $\mathcal{H}$ , there exists a positive integer  $N_0$  such that

$$(2.122) \quad \sup_{u \in \mathcal{U}_{ad}} \sup_{0 \leq t \leq T} \|(\text{Id}_{\mathcal{H}} - \Pi_N)y(t; x, u)\|_{\mathcal{H}} \leq \epsilon, \quad \forall N \geq N_0.$$

Thus Assumption (A7) is satisfied.  $\square$

**Remark 2.6.** From the proof given above, it is clear that Assumption (ii) in Lemma 2.6 can be relaxed to e.g.

(ii') The eigenfunctions of the linear operator  $L : D(L) \subset \mathcal{H} \rightarrow \mathcal{H}$  forms an orthonormal basis of  $\mathcal{H}$  and the eigenvalues of  $L$  approaches  $-\infty$ .

Note also that Lemma 2.6 still holds if  $L$  has complex eigenvalues. We just need to work with a complexification of the operator  $L$  and the underlying state space  $\mathcal{H}$  (cf. e.g. [CLW15b, footnote 17 on p. 55]), and make the corresponding changes in Assumption (ii') above.

### 3. APPLICATION TO OPTIMAL CONTROL OF ENERGY BALANCE CLIMATE MODELS

We show in this section that our framework allows us to provide rigorous Galerkin approximations to the optimal control of broad class of semilinear heat problems, posed on a compact (smooth) manifold without boundary. As an application, we show in Sect. 3.3 that the optimal control of energy balance models (EBMs) arising in the context of geoengineering and climate change, can be thus approximated by optimal control problems of ODEs, more tractable numerically. We first recall some fundamentals of differential geometry to prepare the analysis.

**3.1. Preliminary from differential geometry.** To properly write an EBM on the sphere, we recall how differential operators are defined on an abstract compact smooth manifold  $\mathfrak{M}$  without boundary, of dimension  $n$  and endowed with a Riemannian metric<sup>4</sup>  $\mathfrak{g}$ . First, given a smooth function  $u$  on  $\mathfrak{M}$ , the gradient  $\nabla_{\mathfrak{g}}u$  is a vector field on  $\mathfrak{M}$ , that takes in local coordinates the form

$$(3.1) \quad \nabla_{\mathfrak{g}}u := \sum_{i=1}^n \left( \sum_{j=1}^n \mathfrak{g}^{ij} \partial_j u \right) \partial_i.$$

The divergence of a vector field  $X = \sum_{j=1}^n X^j \partial_j$  takes the following form in local coordinates

$$(3.2) \quad \text{div}_{\mathfrak{g}} X = \frac{1}{\sqrt{\det \mathfrak{g}}} \sum_{i=1}^n \partial_i \left( X^i \sqrt{\det \mathfrak{g}} \right).$$

The Laplacian on  $(\mathfrak{M}, \mathfrak{g})$  takes then the form

<sup>4</sup>Recall that a Riemannian metric  $\mathfrak{g}$  is a smooth family of inner products on the tangent spaces  $T_p\mathfrak{M}$ . Namely,  $\mathfrak{g}$  associates (smoothly) to each  $p$  in  $\mathfrak{M}$  a positive definite symmetric bilinear form  $\varphi_p$  on  $T_p\mathfrak{M} \times T_p\mathfrak{M}$ . In local coordinates,

$$\mathfrak{g}_{ij}(p) = \varphi_p(\partial_i, \partial_j),$$

and thus  $\det \mathfrak{g}(p) > 0$ .

$$(3.3) \quad \begin{aligned} \Delta_{\mathfrak{g}} &:= \operatorname{div}_{\mathfrak{g}} \circ \nabla_{\mathfrak{g}} \\ &= \frac{1}{\sqrt{\det \mathfrak{g}}} \sum_{i=1}^n \partial_i \left( \sum_{j=1}^n \mathfrak{g}^{ij} \sqrt{\det \mathfrak{g}} \partial_j \right). \end{aligned}$$

Set

$$(3.4) \quad \mathbb{S}^n := \{x \in \mathbb{R}^{n+1} : |x| = 1\}.$$

Let  $y = (y_1, \dots, y_{n+1})$  be any point of  $\mathbb{S}^n$  and  $(x_1, \dots, x_n)$  be its image under the stereographic projection from the “north pole”  $N = (0, \dots, 0, 1)$  onto the space

$$(3.5) \quad \mathbb{R}^n \equiv \{(\xi_1, \dots, \xi_{n+1}) \in \mathbb{R}^{n+1} \mid \xi_{n+1} = 0\}.$$

The canonical Riemannian metric on  $\mathbb{S}^n$  takes then the form

$$(3.6) \quad \mathfrak{g}_{ij} = \frac{4}{(1 + |x|^2)^2} \delta_{ij}, \quad 1 \leq i, j \leq n.$$

Note that  $\mathfrak{g}^{ij}$  introduced above has here the explicit expression:

$$(3.7) \quad \mathfrak{g}^{ij} = (\mathfrak{g}_{ii})^{-1} \delta_{ij}, \quad 1 \leq i, j \leq n.$$

In what follows we denote by  $\langle \cdot, \cdot \rangle_{\mathfrak{g}}$  the inner Riemannian product on  $(\mathbb{S}^n, \mathfrak{g})$  and by  $L^2(\mathbb{S}^n)$  the space of square-integrable real-valued functions for the norm induced by this inner product. The space  $\vec{L}^2(\mathbb{S}^n)$  is then defined as

$$\vec{L}^2(\mathbb{S}^n) := \underbrace{L^2(\mathbb{S}^n) \times \dots \times L^2(\mathbb{S}^n)}_n.$$

Introduce now the polar coordinates on  $\mathbb{S}^n$ . Consider the “south pole”  $S = (0, \dots, 0, -1)$ . For any  $y \in \mathbb{S}^n \setminus \{N, S\}$ , define  $\rho \in (0, \pi)$  and  $\theta \in \mathbb{S}^{n-1}$  by

$$(3.8) \quad \cos \rho = y^{n+1}, \quad \theta = \frac{y'}{|y'|},$$

where  $y' = (y_1, y_2, \dots, y_n, 0)$ .

These variables have the following interpretation: the polar radius  $\rho$  represents the angle between the position vectors of  $y$  and  $N$ ; it can be also regarded as the latitude of the point  $y$  measured from  $N$ . The polar angle  $\theta$  can be regarded as the longitude of the point  $y$ .

The canonical spherical Riemannian metric has the following expression in the polar coordinates:

$$(3.9) \quad ds^2 = \sum_{i,j=1}^n \mathfrak{g}_{ij} dx_i dx_j = d\rho^2 + (\sin \rho)^2 d\theta^2.$$

The operator  $\Delta_{\mathfrak{g}}$  then takes the form

$$(3.10) \quad \Delta_{\mathfrak{g}} = \frac{\partial}{\partial \rho^2} + (n-1) \cot \rho \frac{\partial}{\partial \rho} + \frac{1}{(\sin \rho)^2} \Delta_{\theta},$$

where  $\Delta_{\theta}$  denotes the Laplace-Beltrami operator on  $\mathbb{S}^{n-1}$ .

Let us introduce the linear operator  $\mathcal{L}_{\mathfrak{g}} := -\Delta_{\mathfrak{g}}$  with domain

$$(3.11) \quad \mathcal{D}(\mathcal{L}_{\mathfrak{g}}) := \{u \in L^2(\mathbb{S}^n) : \nabla_{\mathfrak{g}} u \in \vec{L}^2(\mathbb{S}^n), \Delta_{\mathfrak{g}} u \in L^2(\mathbb{S}^n)\}.$$

It is known that  $\mathcal{L}_{\mathfrak{g}}$  is self-adjoint [Gri09, Sect. 4.2]. Moreover, the following results hold:



**Theorem 3.1.** *The spectrum of  $\mathcal{L}_g$  with domain  $\mathcal{D}(\mathcal{L}_g)$  defined in (3.11) is discrete and consists of an increasing sequence  $\{\lambda_k\}_{k=1}^\infty$  of non-negative eigenvalues (counted according to multiplicity) such that*

$$\lim_{k \rightarrow \infty} \lambda_k = +\infty.$$

*There is an orthonormal basis  $\{e_k\}_{k=1}^\infty$  in  $L^2(\mathbb{S}^n)$  such that each function  $e_k$  is an eigenfunction of  $\mathcal{L}_g$  with the eigenvalue  $\lambda_k$ .*

Let  $\{E_\lambda\}_{\lambda \geq 0}$  denote the spectral resolution of  $\mathcal{L}_g$ . We can define then for each  $t \geq 0$ ,

$$(3.12) \quad P(t) := \int_0^\infty e^{-t\lambda} dE_\lambda,$$

that constitutes a bounded linear operator acting on  $L^2(\mathbb{S}^n)$  satisfying the properties summarized below.

**Theorem 3.2.** *Let  $P(t)$  defined by (3.12), then*

(i) *For any  $t \geq 0$ ,  $P(t)$  is a bounded self-adjoint operator on  $L^2(\mathbb{S}^n)$ , and*

$$(3.13) \quad \|P(t)\| \leq 1.$$

(ii) *The family  $\{P(t)\}_{t \geq 0}$  satisfies the semigroup identity:*

$$(3.14) \quad P(t)P(s) = P(t+s),$$

*for all  $t, s \geq 0$ .*

(iii) *The mapping  $t \mapsto P(t)$  is strongly continuous on  $[0, \infty)$ . That is, for any  $t \geq 0$  and  $f \in L^2(\mathbb{S}^n)$ ,*

$$(3.15) \quad \lim_{s \rightarrow t} P(s)f = P(t)f,$$

*where the limit is understood in the norm of  $L^2(\mathbb{S}^n)$ . In particular, for any  $f \in L^2(\mathbb{S}^n)$ ,*

$$(3.16) \quad \lim_{t \rightarrow 0^+} P(t)f = f.$$

(iv) *For all  $f \in L^2(\mathbb{S}^n)$  and  $t > 0$ , we have that  $P(t)f$  lies in  $\mathcal{D}(\mathcal{L}_g)$  and*

$$(3.17) \quad \frac{d}{dt}(P(t)f) = -\mathcal{L}_g(P(t)f).$$

The above theorems are particular cases of results presented in [Gri09] for the Laplace operator defined on general *weighted smooth manifolds* [Gri09, Def. 3.17]. See [Gri09, Thm. 10.13] for Theorem 3.1 and [Gri09, Thm. 4.9] for Theorem 3.2.

**3.2. Galerkin approximations of controlled semilinear heat equations on  $\mathbb{S}^n$ .** Given  $\mathcal{L}_g = -\Delta_g$  with domain  $\mathcal{D}(\mathcal{L}_g)$  given in (3.11), we consider the following abstract controlled semilinear heat problem of the form (2.1) posed in  $L^2(\mathbb{S}^n)$ :

$$(3.18) \quad \begin{aligned} \frac{dy}{dt} &= -\mathcal{L}_g y + F(y) + \mathfrak{C}(u(t)), \quad t \in (0, T], \\ y(0) &= x \in L^2(\mathbb{S}^n). \end{aligned}$$

In what follows we denote by  $\mathcal{H}$  the space  $L^2(\mathbb{S}^n)$ . Based on Theorem 2.1 of Section 2, we show the uniform convergence of Galerkin approximations to (3.18) associated with the reduced state space  $\mathcal{H}_N \subset L^2(\mathbb{S}^n)$  defined by

$$(3.19) \quad \mathcal{H}_N := \text{span}\{e_k : k = 1, \dots, N\}, \quad N \in \mathbb{Z}_+^*,$$

in which the  $e_k$ 's denote the eigenfunctions of  $-\mathcal{L}_g$  lying in  $\mathcal{D}(\mathcal{L}_g)$ ; see Theorem 3.1.

The linear approximations  $L_N$  of the operator  $-\mathcal{L}_g$  are then naturally defined as

$$(3.20) \quad L_N := \Pi_N \Delta_g \Pi_N : \mathcal{H} \rightarrow \mathcal{H}_N,$$

where  $\Pi_N$  denotes the orthogonal projector associated with  $\mathcal{H}_N$ .

Theorem 2.1 leads then to the following corollary about uniform convergence of Galerkin approximations of (3.18).

**Corollary 3.1.** *Let  $V$  be a Hilbert space. Assume that  $F : L^2(\mathbb{S}^n) \rightarrow L^2(\mathbb{S}^n)$  and  $\mathfrak{C} : V \rightarrow L^2(\mathbb{S}^n)$  are both globally Lipschitz, and  $\mathfrak{C}(0) = 0$ . Assume the set of admissible controls  $\mathcal{U}_{ad}$  is given by (2.32) with  $U$  therein being a compact subset of the Hilbert space  $V$  and with  $q > 1$ .*

*Then, for any  $T > 0$  and any  $(x, u)$  in  $L^2(\mathbb{S}^n) \times \mathcal{U}_{ad}$ , the problem (3.18) admits a unique mild solution  $y(\cdot; x, u)$  in  $C([0, T], \mathcal{H})$ , and its Galerkin approximation (2.8) associated with the eigen-subspaces (3.19) admits a unique solution  $y_N(\cdot; \Pi_N x, u)$  in  $C([0, T], \mathcal{H}_N)$  for each  $N$  in  $\mathbb{Z}_+^*$ . Moreover, the following uniform convergence result holds:*

$$(3.21) \quad \lim_{N \rightarrow \infty} \sup_{u \in \mathcal{U}_{ad}} \sup_{t \in [0, T]} \|y_N(t; \Pi_N x, u) - y(t; x, u)\|_{\mathcal{H}} = 0.$$

*Proof.* We only need to check the conditions (A0)–(A2), (A6), and (A7) assumed in Theorem 2.1. This is done below in four steps.

STEP 1: Checking (A0)–(A1). From Theorem 3.2, we know that  $-\mathcal{L}$  is the infinitesimal generator of the  $C_0$ -semigroup of contractions,  $\{P(t)\}_{t \geq 0}$ , on  $L^2(\mathbb{S}^n)$ .

Since  $-\mathcal{L}_g$  is self-adjoint, the operator  $L_N$  is a finite-rank diagonal operator acting on  $\mathcal{H}$  and we have for each  $\phi$  in  $\mathcal{H}$ ,

$$(3.22) \quad e^{tL_N} \phi := \sum_{k=1}^N e^{-t\lambda_k} \langle \phi, e_k \rangle_g e_k.$$

It follows that each operator  $L_N$  generates a  $C_0$ -semigroup of contractions on  $\mathcal{H}$ , which will be denoted by  $\{P_N(t)\}_{t \geq 0}$ . We have thus checked Assumptions (A0)–(A1) given in Section 2 with  $M = 1$  and  $\omega = 0$ , namely

$$(3.23) \quad \|P(t)\| \leq 1 \quad \text{and} \quad \|P_N(t)\| \leq 1, \quad N \geq 0, \quad t \geq 0.$$

STEP 2: Checking (A2). This condition results from the self-adjointness of  $-\mathcal{L}_g$ . Indeed, for any given  $\phi \in \mathcal{D}(\mathcal{L}_g)$ , since both  $\phi$  and  $\Delta_g \phi$  belong to  $L^2(\mathbb{S}^n)$ , the following expansions against the eigenbasis hold:

$$(3.24) \quad \phi = \sum_{i=1}^{\infty} a_i e_i, \quad \Delta_g \phi = \sum_{i=1}^{\infty} b_i e_i,$$

where

$$(3.25) \quad a_i = \langle \phi, e_i \rangle_g, \quad b_i = \langle \Delta_g \phi, e_i \rangle_g, \quad i \in \mathbb{Z}_+^*.$$

Note that

$$(3.26) \quad b_i = \langle \Delta_g(\phi), e_i \rangle_g = \langle \phi, \Delta_g e_i \rangle_g = \lambda_i \langle \phi, e_i \rangle_g = \lambda_i a_i.$$

We get then that

$$(3.27) \quad \begin{aligned} \|L_N \phi - \Delta_g \phi\|_{L^2(\mathbb{S}^n)} &= \left\| \Pi_N \Delta_g \left( \sum_{i=1}^N a_i e_i \right) - \Delta_g \phi \right\|_{L^2(\mathbb{S}^n)} \\ &= \left\| \sum_{i=1}^N \lambda_i a_i e_i - \sum_{i=1}^{\infty} b_i e_i \right\|_{L^2(\mathbb{S}^n)} = \left\| \sum_{i=N+1}^{\infty} b_i e_i \right\|_{L^2(\mathbb{S}^n)} \end{aligned}$$

and Assumption **(A2)** follows.

**STEP 3: Checking (A6).** Since  $F$  and  $\mathfrak{C}$  are globally Lipschitz, the existence of a unique mild solution to (3.18) and to its Galerkin approximation (2.8) follows directly from a classical fixed point argument and standard Gronwall's estimates; see e.g. [CH98, Prop. 4.3.3 and Thm. 4.3.4]. In particular, by (3.23), we obtain the following a priori estimates for all  $t$  in  $[0, T]$  and  $N$  in  $\mathbb{Z}_+^*$ :

$$(3.28) \quad \begin{aligned} \|y(t; x, u)\|_{L^2(\mathbb{S}^n)} &\leq e^{\text{Lip}(F)t} \|x\|_{L^2(\mathbb{S}^n)} + \int_0^t g(s) ds + \text{Lip}(F) \int_0^t g(s) e^{\text{Lip}(F)(t-s)} ds, \\ \|y_N(t; \Pi_N x, u)\|_{L^2(\mathbb{S}^n)} &\leq e^{\text{Lip}(F)t} \|x\|_{L^2(\mathbb{S}^n)} + \int_0^t g(s) ds + \text{Lip}(F) \int_0^t g(s) e^{\text{Lip}(F)(t-s)} ds, \end{aligned}$$

where

$$g(s) = \|F(0)\|_{L^2(\mathbb{S}^n)} + \text{Lip}(\mathfrak{C}) \|u(s)\|_{L^2(\mathbb{S}^n)}, \quad \text{for a.e. } s \in [0, T].$$

Moreover, since by assumption  $u(t)$  takes value in a compact thus bounded set  $U$  for each  $u \in \mathcal{U}_{ad}$ , the a priori estimates (3.28) also ensure the required uniform boundedness estimates (2.53) stated in Assumption **(A6)**.

**STEP 4: Checking (A7).** Due to our assumptions and from what precedes, the conditions of Lemma 2.6 are satisfied and thus Assumption **(A7)** is satisfied. The proof is complete.  $\square$

**Remark 3.1.** Corollary 3.1 has been formulated in the case where  $F$  and  $\mathfrak{C}$  are globally Lipschitz, but actually the conclusions of this corollary still hold if these conditions are relaxed to be locally Lipschitz as long as Assumption **(A6)** is satisfied with the relevant a priori estimates as a consequence of Theorem 2.1.

**Remark 3.2.** Similar to Remark 2.3, we note that Corollary 3.1 still holds when the nonlinearity  $F$  depends also on time, i.e.,  $F : [0, T] \times L^2(\mathbb{S}^n) \rightarrow L^2(\mathbb{S}^n)$ , and satisfies for instance that  $F(\cdot, y) \in L^\infty(0, T; L^2(\mathbb{S}^n))$  for each  $y \in L^2(\mathbb{S}^n)$ ,  $F(t, \cdot)$  is globally Lipschitz for almost every  $t \in [0, T]$  and the mapping  $t \mapsto \text{Lip}(F(t, \cdot))$  is in  $L^\infty(0, T)$ .

**Remark 3.3.** Note also that Corollary 3.1 still holds when the semilinear heat problem (3.18) is posed on a general  $n$ -dimensional Riemannian smooth and compact manifold  $(\mathfrak{M}, \mathfrak{g})$  without boundary, with the Riemannian metric  $\mathfrak{g}$  not limited thus to  $\mathfrak{g}$  defined in (3.6). Similarly the case of semilinear heat problem posed on a non-empty relatively compact subset  $\Omega$  of  $(\mathfrak{M}, \mathfrak{g})$  with homogenous Dirichlet boundary conditions can also be dealt with. This is because Theorems 3.1 and 3.2 still hold for such cases; see again [Gri09, Thm. 4.9 and Thm. 10.13]. In particular, Corollary 3.1 holds when the Riemannian metric  $\mathfrak{g}$  on  $\mathbb{S}^n$  is replaced by another Riemannian metric  $\tilde{\mathfrak{g}}$  and the Laplacian  $\Delta_{\mathfrak{g}}$  in (3.18) is replaced by  $\Delta_{\tilde{\mathfrak{g}}}$  accordingly. This is remark about the change of Riemannian metric is used in Sect. 3.3 that follows.

**3.3. Energy balance models (EBMs).** Energy balance models (EBMs) are among the simplest climate models developed for the study of climate sensitivity. They are formulated based on the energy balance on the Earth surface [NCC81, SLW<sup>+</sup>12] and have the Earth surface temperature as the only dependent variable. First made popular by the works [Bud69, Sel69], these models have been extensively studied since both analytically and numerically; see e.g. [BCDT09, NCC81, RCC<sup>+</sup>14] and references therein.

With suitable tuning of their parameters, EBMs that resolve the Earth's land-sea geography and are forced by the seasonal insolation cycle have been shown to mimic, to a certain extent, the observed zonal temperatures for the observed present climate [NSM83, Cro00]. Once EBMs are fitted to the observations [Sel69, Ghi76, GLN93] or to the simulated climate of general circulation models (GCMs) [HCKN89, Cro00], they can be used to estimate the temporal response patterns to various forcing

scenarios; such a methodology is of particular interest in the detection and attribution of climate change [SAS<sup>+</sup>07].

Depending on whether zonal or meridional averages are taken, the modeled surface temperature can either depend on the latitude only, or depend on both the latitude and the longitude, resulting respectively in 1D, or 2D models. In the 2D case, the model is posed on the two-dimensional unit sphere  $S^2$ , and takes typically the following form [NSM83]:

$$(3.29) \quad \frac{\partial T(\xi, t)}{\partial t} = \operatorname{div}_{\mathfrak{g}}(D(x)\nabla_{\mathfrak{g}}T(\xi, t)) + f(t, x, T(\xi, t)) - g(T(\xi, t)) + E(\xi, t), \quad \xi \in S^2, t > 0.$$

Here the gradient  $\nabla_{\mathfrak{g}}$  and the divergence  $\operatorname{div}_{\mathfrak{g}}$  on the Riemannian manifold  $(S^2, \mathfrak{g})$  are given respectively by (3.1) and (3.2), and the Riemannian metric  $\mathfrak{g}$  is given by (3.6). The diffusion term  $\operatorname{div}_{\mathfrak{g}}(D(x)\nabla_{\mathfrak{g}}T(\xi, t))$  describes the redistribution of heat on the surface of the Earth by conduction and convection, the reaction terms  $f(t, x, T(\xi, t)) - g(T(\xi, t))$  express the balance between incoming and outgoing radiations, and  $E(\xi, t)$  denotes an anthropogenic forcing. See Table 1 for the precise meaning of the symbols involved in (3.29). We refer to [BCDT09] for the rigorous approximation of (3.29) via finite elements on manifolds.

TABLE 1. Glossary of model's parameters & variables

Symbol	Interpretation
$\xi := (\rho, \theta)$	$\rho \in (0, \pi)$ denotes the latitude and $\theta \in (-\pi, \pi)$ denotes the longitude
$T(\xi, t)$	sea surface temperature at $\xi, t$
$x$	sine of the latitude, i.e. $x = \sin(\rho)$
$D(x)$	diffusion coefficient for heat transport, assumed to be zonally averaged and hence does not depend on the longitude $\theta$
$f(t, x, T(\xi, t))$	incoming solar radiation
$g(T(\xi, t))$	outgoing infrared radiation
$E(\xi, t)$	forcing representing greenhouse gas emissions

The functions  $f(t, x, T(\xi, t))$  and  $g(T(\xi, t))$  are typically of the following form [Bud69]:

$$(3.30) \quad \begin{aligned} f(t, x, T(\xi, t)) &= QS(x, t)(1 - \alpha(x, T)), \\ g(T(\xi, t)) &= a + bT(\xi, t). \end{aligned}$$

Here  $Q$  is the so-called solar constant,  $S(x, t)$  denotes a solar insolation distribution function,  $\alpha(x, T)$  denotes the albedo, and  $a$  and  $b$  are empirical constants typically estimated from satellite observations; see e.g. [GLN93]. We refer to [ALB05] for the calibration of other parameters including  $Q$ , or coefficients such as  $S(x, t)$  or contained in  $\alpha(x, T)$ ; see also [RCC<sup>+</sup>14].

We also note that the LHS of (3.29) should be multiplied by a factor  $\kappa(\xi)$ , which measures the effective heat capacity per unit area. Here, we have assumed that  $\kappa(\xi)$  is a constant which is taken to be 1 after a scaling in the time variable.

As a preparation to cast a controlled version of (3.29) into the form of (3.18), we will make use of a new Riemannian metric so that the diffusion term in (3.29) becomes simply the Laplacian under this new metric. For this purpose, we assume that

**(H1)** the diffusion coefficient  $D(x)$  is  $C^1$ -smooth and is strictly positive.

By introducing the new Riemannian metric

$$(3.31) \quad \tilde{\mathfrak{g}} = \frac{1}{D(x)}\mathfrak{g},$$

it is known that the Laplace operator under this new Riemannian metric is given by

$$(3.32) \quad \Delta_{\tilde{g}} = \operatorname{div}_{\tilde{g}}(D(x)\nabla_{\tilde{g}}).$$

Therefore, Eq. (3.29) can be rewritten into the following form on  $(\mathbb{S}^2, \tilde{g})$ :

$$(3.33) \quad \frac{\partial T(\xi, t)}{\partial t} = \Delta_{\tilde{g}}T(\xi, t) + f(t, x, T(\xi, t)) - g(T(\xi, t)) + E(\xi, t).$$

With this rewriting, we consider the operator  $\mathcal{L}_{\tilde{g}} = -\Delta_{\tilde{g}}$  with domain  $\mathcal{D}(\mathcal{L}_{\tilde{g}})$  given by (3.11) in which  $\tilde{g}$  replaces  $g$ . We are now in position to apply the general results of Sect. 3.2 in particular Corollary 3.1 to a contemporary problem related to geoengineering that we address here in the framework of optimal control of EBMs such as (3.33).

**3.4. Optimal control of climate?** In 1955, John von Neumann envisioned that “probably intervention in atmospheric and climate matters will come in a few decades, and will unfold on a scale difficult to imagine at present;” see [VN55]. As our planet enters a period of changing climate never before experienced in recorded human history, primarily caused by the rapid buildup of carbon dioxide in the atmosphere from the burning of fossil fuels, interest is growing in the potential for deliberate large-scale intervention in the climate system to counter climate change; see e.g. [She09, Cou15b, Cou15a]. Although we are still far away from large-scale implementation of what John von Neumann envisioned decades ago, the consideration of climate engineering—also known as geoengineering—is raising in the scientific community with a literature that became more abundant on the topic over the recent years; see e.g. [LV09, She09, VL11, Cou15b, Cou15a].

At the simplest level, the surface temperature of the Earth results from the net balance of incoming solar (shortwave) radiation and outgoing terrestrial (longwave) radiation [KT97]. Proposed geoengineering methods attempt to rectify the current and potential future radiative imbalance and they are usually divided into two basic categories: (i) carbon dioxide removal techniques which remove  $\text{CO}_2$  from the atmosphere to increase the amount of longwave radiation emitted by the Earth; and (ii) solar radiation management techniques that reduce the amount of solar (shortwave) radiation absorbed by the Earth by reflecting a small percentage of the sun’s light and heat back into space.

While a lot of efforts have been devoted to describing different geoengineering options in detail and discussing their advantages, effectiveness, potential side effects and drawbacks, still more understanding is required before any method could even be seriously considered for deployment on the requisite international scale [She09]. On the other hand, policies to reduce global greenhouse gas (GHG) emissions is a pressing topic on any political agenda, and uncertainties to climate change [MSB<sup>+</sup>04, MMH<sup>+</sup>09] add to the difficulty in quantifying unambiguously the effects of forcing variations on the climate system.

From a mathematical perspective, since any geoengineering methods or GHG mitigation policies can be expressed as controls acting on the climate system, it is natural and important to investigate whether a given type of controls, corresponding e.g. to one or a combination of several geoengineering methods, can drive the climate system from a given “current” state to a desired state over a targeted finite time horizon. This controllability aspect has indeed been investigated within the context of climatology based on some types of EBMs; see e.g. [Día94a, Día94b].

Given that any large-scale decision for addressing climate change have economic [Hal09], societal or physical constraints, it also seems natural to frame the problem as an optimal control of the climate system to seek for controls within a chosen set of geoengineering strategies that lead to the minimization of a relevant cost functional. To our knowledge, this optimal control perspective has not yet been investigated much from a fundamental viewpoint. In the following, we aim to provide a sufficiently general formulation for this purpose, based on the class of EBMs encompassed by Eq. (3.29). The latter equation will serve as our underlying state equation in what follows.

Since EBMs are known to provide reliable models of the mean annual global temperature distribution around the globe [HCKN89, Cro00, SAS<sup>+</sup>07], they constitute a natural laboratory for such an investigation before one moves onto more sophisticated and detailed climate models such as GCMs [GCS08]. In that respect, it is also worth mentioning that EBMs can actually be derived from the thermodynamics equation of the atmosphere primitive equations via an averaging procedure [Kie92]. See also [BEGX13] for the design of optimal economic mitigation policies based on EBMs coupled with an economic growth model.

**3.5. Optimal control of EBMs: Convergence results of value functions.** We formulate the optimal control of EBMs within the general setting of Sect. 2, by relying on the properties of the heat semi-group on the sphere recalled in Sect. 3.2, here generated by  $-\mathcal{L}_{\bar{g}}$ .

First, in order to allow for geoengineering strategies of different nature in different geographic regions, we consider a collection of open subsets  $\{\Omega_i \subset \mathbb{S}^2 : i = 1, \dots, M\}$ , where  $M$  denotes the number of such strategies, one for each region  $\Omega_i$ , with possible overlapping.

The set of admissible controls is defined as follows. For region  $\Omega_i$  we define the Hilbert space of functions

$$(3.34) \quad V_i = L^2(\Omega_i, \mathbb{R}^{n_i}),$$

with  $n_i$  some positive integer, and introduce

$$(3.35) \quad V := V_1 \times \dots \times V_M.$$

Consider for each  $i$ ,  $U_i$  to be a compact subset of  $V_i$  and let us introduce the set

$$(3.36) \quad U := \{v = (v_1, \dots, v_M) \in V : v_i \in U_i, i = 1, \dots, M\}.$$

Let the set of admissible controls be given by:

$$(3.37) \quad \mathcal{U}_{ad} := \{u \in L^q(0, t_f; V) : u(s) \in U \text{ a.e. } \}.$$

where  $q > 1$  is fixed. Note that since each  $V_i$  is a space of functions defined over a region  $\Omega_i$ , an admissible control  $u$  in  $\mathcal{U}_{ad}$  is actually a locally distributed control. Note that given  $u$  in  $\mathcal{U}_{ad}$  and  $t$  in  $(0, t_f)$ , its  $i^{\text{th}}$ -component  $u_i(t)$  is a spatial function that lies in  $V_i$ . We will denote hereafter by  $u_i(t)[\bar{\zeta}]$  its value taken at  $\bar{\zeta} \in \mathbb{S}^2$ .

Finally, we assume that the combined effects of the geoengineering strategies on the global temperature field  $T(\bar{\zeta}, t)$ , is represented via a nonlinear function

$$(3.38) \quad \begin{aligned} G : \mathbb{R}^{n_1} \times \dots \times \mathbb{R}^{n_M} &\rightarrow \mathbb{R} \\ (\zeta_1, \dots, \zeta_M) &\longmapsto G(\zeta_1, \dots, \zeta_M), \end{aligned}$$

that forces Eq. (3.33). In practice, the modeler has to specify the function  $G$  (and  $V$ ), depending on the geoengineering strategy or the GHG mitigation policy adopted as well as the EBM retained. As explained below, our framework ensures that a global Lipschitz assumption<sup>5</sup> on  $G$  allows for convergence of Galerkin approximations and thus provide a rigorous basis for a numerical investigation of various control scenarios.

We consider thus, for each  $u$  in  $\mathcal{U}_{ad}$ , the following controlled version of the EBM (3.33) which writes for each  $\bar{\zeta} \in \mathbb{S}^2$  and  $t \in [0, t_f]$  as,

$$(3.39) \quad \frac{\partial T(\bar{\zeta}, t)}{\partial t} = \Delta_{\bar{g}} T(\bar{\zeta}, t) + f(t, x, T(\bar{\zeta}, t)) - g(T(\bar{\zeta}, t)) + E(\bar{\zeta}, t) + G(\tilde{u}_1(t, \bar{\zeta}), \dots, \tilde{u}_M(t, \bar{\zeta})),$$

<sup>5</sup>That can be relaxed to a local Lipschitz assumption as long as a priori error estimates are available to ensure Assumption (A6); see Remark 3.1.

supplemented with an initial condition  $T_0$  in  $L^2(\mathbb{S}^2)$ . Here

$$(3.40) \quad \tilde{u}_i(t, \xi) = \begin{cases} u_i(t)[\xi], & \text{if } \xi \in \Omega_i, \\ 0, & \text{otherwise.} \end{cases}$$

In order to recast the IVP associated with (3.39) into the abstract form (3.18), we introduce the following function spaces

$$(3.41) \quad \mathcal{H} := L^2(\mathbb{S}^2), \quad \mathcal{H}_1 := \mathcal{D}(\mathcal{L}_{\tilde{g}}),$$

where  $\mathcal{D}(\mathcal{L}_{\tilde{g}})$  is defined in (3.11).

We make the following assumption.

**(H2)** The functions  $f$  and  $g$  take the forms given by (3.30), where the solar insolation distribution function  $S$  therein lives in the space  $L^\infty((0, \infty), L^\infty(\mathbb{S}^2))$  and the albedo  $\alpha(x, T)$  is a continuous, piecewise-linear ramp function such as given in [Ghi76, Eq. (2a)] or [RCC<sup>+</sup>14, Eq. (2.2)]. The GHG emission term  $E$  belongs to  $L^1_{\text{loc}}((0, \infty), L^2(\mathbb{S}^2))$ .

We define now the nonlinearity  $F : [0, t_f] \times \mathcal{H} \rightarrow \mathcal{H}$  to be:

$$(3.42) \quad F(t, v)[\xi] := f(t, x, v(\xi)) - g(v(\xi)) + E(\xi, t), \quad \text{for all } v \in \mathcal{H}, \text{ and a.e. } \xi \in \mathbb{S}^2, t \in [0, t_f].$$

Under Assumption **(H2)**, for each  $v$  in  $\mathcal{H}$  and almost every  $t \in [0, t_f]$ ,  $F(t, v)$  belongs to  $\mathcal{H}$ .

Finally, we define the nonlinear operator  $\mathfrak{C} : V \rightarrow \mathcal{H}$  associated with the control to be:

$$(3.43) \quad \mathfrak{C}(v)[\xi] := G(\tilde{v}_1(\xi), \dots, \tilde{v}_M(\xi)), \quad \text{for all } v \in V, \text{ and a.e. } \xi \in \mathbb{S}^2,$$

with

$$(3.44) \quad \tilde{v}_i(\xi) = \begin{cases} v_i(\xi), & \text{if } \xi \in \Omega_i, \\ 0, & \text{otherwise.} \end{cases}$$

Then, Eq. (3.39) can be rewritten as Eq. (3.18) of Sect. 3.2, with the (nonlinear) operators  $F$  and  $\mathfrak{C}$  defined above. Having the purpose in mind of driving the temperature field  $T(\xi, t)$  to a state sufficiently close to a specified profile at the final time  $t_f$ , while keeping the control cost “low”, we consider the cost functional:

$$(3.45) \quad J(T_0, u) = \int_0^{t_f} \left( \frac{1}{2} \|T(\cdot, t; T_0, u) - T_d\|_{\mathcal{H}}^2 + \frac{\mu}{2} \|u(\cdot, t)\|_V^2 \right) dt, \quad T_0 \in \mathcal{H}, \mu \geq 0.$$

Here  $T_d$  denotes the targeted temperature field over the globe (that lies in  $\mathcal{H}$ ) and  $T(\cdot, t; T_0, u)$  denotes the mild solution to (3.39) that emanates from  $T_0$ .

The associated optimal control problem reads then:

$$(3.46) \quad \min J(T_0, u) \quad \text{s.t.} \quad (T, u) \in L^2(0, T; \mathcal{H}) \times \mathcal{U}_{ad} \text{ solves Eq. (3.39)} \\ \text{subject to the initial condition } T(\cdot, 0) = T_0 \in \mathcal{H}.$$

**Approximation of the value function and error estimates about the optimal control.** Note that thanks to Assumption **(H2)**, the nonlinearity  $F$  defined in (3.42) satisfies the conditions required in Remark 3.2. If we assume furthermore that the nonlinear operator  $\mathfrak{C}$  defined by (3.43) to be globally Lipschitz as a mapping from  $V$  to  $\mathcal{H}$  (and  $\mathfrak{C}(0) = 0$ ) then by Corollary 3.1 and Remarks 3.2–3.3, the uniform convergence result given by (3.21) in Corollary 3.1 holds for the IVP associated with (3.39). Namely,

$$(3.47) \quad \lim_{N \rightarrow \infty} \sup_{u \in \mathcal{U}_{ad}} \sup_{t \in [0, T]} \|y_N(t; \Pi_N x, u) - y(t; x, u)\|_{\mathcal{H}} = 0,$$

where  $y_N(\cdot; \Pi_N x, u)$  denotes the solution to the Galerkin approximation of (3.39) associated with the eigen-subspace spanned by the first  $N$  eigenfunctions of the Laplacian  $\Delta_{\mathbb{g}}$ .

Note also that Condition (C1) on the cost functional  $J$  defined in (3.45) is clearly satisfied. Then, Theorem 2.2 applies if we assume furthermore that there exists for each pair  $(t, x)$  a minimizer  $u_{t,x}^*$  (resp.  $u_{t,x}^{N,*}$ ) in  $\mathcal{U}_{ad}[t, T]$  of the minimization problem in (2.85a) (resp. in (2.85b)) associated here with the optimal control problem (3.46). Thus, for any  $x$  in  $\mathcal{H}$ , it holds that

$$(3.48) \quad \lim_{N \rightarrow \infty} \sup_{t \in [0, T]} |v_N(t, \Pi_N x) - v(t, x)| = 0.$$

Observe that since  $\mathfrak{C}$  is globally Lipschitz, Assumption (H2) leads to simple a priori error estimates such as (3.28), showing thus that Assumption (A6) is satisfied; see also Remark 3.2. The other conditions in Assumption (E) of Sect. 2.6 are trivially satisfied here and thus the error estimates of Sect. 2.6 hold. In particular Corollary 2.2 applies with  $q = 2$  and  $\sigma = \mu/2$  which, using the notations of this corollary, leads to

$$(3.49) \quad \|u^* - u_N^*\|_{L^2(0, T; V)}^2 \leq \frac{4\mathcal{C} + 4\|T_d\|_{\mathcal{H}}}{\mu} \left[ \sqrt{T} + \gamma T \right] \left( \|\Pi_N^\perp y^*(\cdot; u^*)\|_{L^2(0, T; \mathcal{H})} + 2\|\Pi_N^\perp y(\cdot; u_N^*)\|_{L^2(0, T; \mathcal{H})} \right),$$

after a simple estimate of  $\text{Lip}(\mathcal{G}|_{\mathfrak{B}})$  where  $\mathcal{G}(y) = \|y - T_d\|_{\mathcal{H}}^2$ ,  $y \in \mathcal{H}$ .

With the rigorous convergence results (3.47) and (3.48) and error estimate (3.49), the numerical approximation of solutions to (3.46) becomes affordable via e.g. a Pontryagin-Maximum-Principle approach applied to Galerkin approximations [CL15] of Eq. (3.29) built here from spherical harmonics. Indeed, a relatively realistic EBM such as given by Eq. (3.29) is known to be simulated accurately out of few spherical harmonics (typically  $20 \leq N \leq 30$ ); see e.g. [NSM83, HCKN89].

#### 4. CONCLUDING REMARKS

Thus, by means of rigorous Galerkin approximations of nonlinear evolution equations in Hilbert spaces, this article provides a natural framework for the synthesis of approximate optimal controls, along with approximations of the value functions. The framework opens up several possible directions for future research. We outline some of these issues below.

1. The usage of spectral methods for solving more complex climate models than EBMs considered in Sect. 3 is standard. By its natural assumptions to verify in practice, the framework presented above makes possible to address the problem of geoengineering strategies or GHG mitigation policies in terms of optimal control of Galerkin approximations of such models, enabling thus, at least theoretically, to reduce the dimension of the problem. However, very often these models include e.g. nonlinear advection terms that require to deal with a loss of regularity. Our framework needs thus to be amended to deal with such a case. The use of interpolated spaces to deal with the loss of regularity and formulations of the Trotter-Kato theorem exploiting Gelfand triple are natural tools to cope with this difficulty. Analogues of Theorems 2.1 and 2.2, as well as Corollary 2.2, seem thus to be reasonably accessible within this approach.

2. Another promising direction is the synthesis of (approximate) optimal controls in a feedback form from the Hamilton-Jacobi-Equation (HJB) associated with Galerkin approximations [KKZ15, GK16]. This is particularly relevant for the optimal control of systems near the first criticality in which only very few modes<sup>6</sup> have lost their stability and where Galerkin approximations are very

<sup>6</sup>Such as a pair of modes in the case of a Hopf bifurcation [MW05].



often useful to approximate the dynamics near the bifurcated states [DTV<sup>+</sup>16] although center manifold reduction techniques lead often to further reducing the number of resolved modes needed to approximate accurately the dynamics [MW05, MW14, CLW15a].

3. However, far from the first criticality or when the nonlinear effects get amplified a larger number of modes is required to dispose of good Galerkin approximations of, already, the uncontrolled dynamics; see [CL15, CLW15b]. The numerical burden of the synthesis of controls at a nearly optimal cost—by solving the HJB equation corresponding to the Galerkin approximation—becomes then quickly prohibitive, especially for the case of locally distributed controls; see [CL15, Sect. 7]. One avenue to do deal with reduced state space of further reduced dimension, is to search for high-mode parameterizations that help reduce the residual energy contained in the unresolved modes, i.e. to reduce the RHS of (2.113) involving the terms  $\|\Pi_N^\perp y^*(\cdot; u^*)\|_{L^2(0, T; \mathcal{H})}$  and  $\|\Pi_N^\perp y(\cdot; u_N^*)\|_{L^2(0, T; \mathcal{H})}$  in Corollary 2.2. The theory of parameterizing manifolds (PM) [CL15, CL16, CLM16] allows for such a reduction leading typically to approximate controls coming with error estimates that introduce multiplying factors  $0 \leq Q < 1$  in the “RHS-like” of (2.113); see [CL15, Theorem 1 & Corollary 2]. The combination of the Galerkin framework introduced here with the PM reduction techniques of [CL15] constitutes thus an idea that is worth pursuing.

4. Finally, we emphasize that the framework introduced here is not limited to Galerkin approximations built from eigenfunctions of the linear part. This is particularly useful for evolution equations for which such eigenfunctions are not the best choice to build Galerkin approximations. As explained in [CGLW16], systems of nonlinear delay differential equations (DDEs) constitute such a type of evolution systems. The optimal control of systems of nonlinear DDEs may thus benefit from the framework introduced here and will be pursued elsewhere. Such an approach may be also relevant to study the possible impact of geoengineering strategies or GHG mitigation policies on large-scale climatic phenomena such as the El Niño–Southern Oscillation (ENSO) and for which DDEs are known to provide good models able to capture some of the essential features of ENSO’s irregularity; see [TSCJ94, NBH<sup>+</sup>98, CNK<sup>+</sup>14, CGN17].

#### ACKNOWLEDGMENTS

This work has been partially supported by the Office of Naval Research (ONR) Multidisciplinary University Research Initiative (MURI) grant N00014-12-1-0911 and N00014-16-1-2073 (MDC), and by the National Science Foundation grants DMS-1616981 (MDC) and DMS-1616450 (HL). MDC and AK were supported by the project “Optimal control of partial differential equations using parameterizing manifolds, model reduction, and dynamic programming” funded by the *Foundation Hadamard/Gaspard Monge Program for Optimization and Operations Research (PGMO)*.

#### APPENDIX A. EXISTENCE OF OPTIMAL CONTROLS

We recall hereafter some standard sufficient conditions for the existence of optimal controls. Our approach follows [HPUU09] that we adapt to our framework. Let  $\mathcal{U} := L^q(0, T; U)$  with  $q \geq 1$ , with here  $U$  denoting a bounded, closed and convex subset of a separable Hilbert space  $V$ . In particular,  $\mathcal{U}$  is also bounded, closed and convex.

Let  $Y$  denotes a separable Hilbert space. Let us assume that we can write the state equation in the form

$$(A.1) \quad e(y, u) = 0,$$

with  $e: Y \times \mathcal{U} \rightarrow Z$  a continuous mapping taking values in a Hilbert space  $Z$ .

The optimal control problem reads as

$$(A.2) \quad \min_{(y,u) \in Y \times \mathcal{U}} J(y, u), \quad \text{subject to } e(y, u) = 0.$$

Existence to this optimal control problem can be shown under the following assumptions.

*Assumption A.1.*

- (i) The state equation (A.1) has a bounded solution operator  $\mathcal{U} \rightarrow Y, u \mapsto y(u)$ .
- (ii) The mapping  $Y \times \mathcal{U} \rightarrow Z, (y, u) \mapsto e(y, u)$  is continuous for the weak topology.
- (iii) The cost function  $J: Y \times \mathcal{U} \rightarrow \mathbb{R}^+$  is weakly lower semicontinuous.

**Remark A.1.** *Assumption A.1 (iii) is satisfied if  $J$  is convex and continuous.*

We introduce the feasible set

$$(A.3) \quad F_{\text{ad}} := \{(y, u) \in Y \times \mathcal{U} : e(y, u) = 0\}.$$

**Definition A.1.** *The pair  $(\bar{y}, \bar{u}) \in Y \times \mathcal{U}$  is called a solution of (A.2) if*

$$(A.4) \quad J(\bar{u}, \bar{y}) \leq J(y, u) \quad \text{for all } (y, u) \in F_{\text{ad}}.$$

**Theorem A.1.** *Under Assumption A.1 problem (A.2) has a non-empty set of solutions.*

*Proof.* The proof is classical. We present only a brief sketch; e.g. [HPUU09, Section 1.5.2]. Since  $J \geq 0$  and  $F_{\text{ad}}$  is nonempty, the infimum exists and there is a minimizing sequence  $(y_k, u_k) \subset F_{\text{ad}}$  with  $J^* := \lim_{k \rightarrow \infty} J(y_k, u_k)$ . Furthermore,  $(u_k)$  is bounded as a sequence in  $\mathcal{U}$ , and by Assumption A.1 (i) the sequence  $(y_k)$  is bounded. Thus, by reflexivity of  $L^2(0, T; \mathcal{H}) \times Y$  we can select a subsequence that converges weakly to some limit  $(\bar{u}, \bar{y})$ . Since  $\mathcal{U}$  is convex and closed, we have that  $\bar{u}$  lies in  $\mathcal{U}$ , and together with Assumption A.1 (ii) we have that the set  $F_{\text{ad}}$  is sequentially weakly closed; hence, we have  $(\bar{y}, \bar{u}) \in F_{\text{ad}}$ . With Assumption A.1 (iii) we obtain the assertion by a classical argument.  $\square$

## REFERENCES

- [AF13] M. Athans and P.L. Falb, *Optimal Control: An Introduction to the Theory and its Applications*, Courier Corporation, 2013.
- [ALB05] V.A. Alexeev, P.L. Langen, and J.R. Bates, *Polar amplification of surface warming on an aquaplanet in “ghost forcing” experiments without sea ice feedbacks*, *Climate Dynamics* **24** (2005), no. 7-8, 655–666.
- [AM89] W. Alt and U. Mackenroth, *Convergence of finite element approximations to state constrained convex parabolic boundary control problems*, *SIAM J. Control Optim.* **27** (1989), 718–736.
- [BCDT09] R. Bermejo, J. Carpio, J.I. Díaz, and L. Tello, *Mathematical and numerical analysis of a nonlinear diffusive climate energy balance model*, *Mathematical and Computer Modelling* **49** (2009), no. 5, 1180–1210.
- [BDPDM07] A. Bensoussan, G. Da Prato, M.C. Delfour, and S. Mitter, *Representation and control of infinite dimensional systems*, Springer Science & Business Media, 2007.
- [BEGX13] W. A. Brock, G. Engström, D. Grass, and A. Xepapadeas, *Energy balance climate models and general equilibrium optimal mitigation policies*, *J. Economic Dynamics and Control* **37** (2013), 2371–2396.
- [BK84] H. T. Banks and K. Kunisch, *The linear regulator problem for parabolic systems*, *SIAM J. Control Optim.* **22** (1984), 684–698.
- [Bré10] H. Brézis, *Functional Analysis, Sobolev Spaces and Partial Differential Equations*, Springer, 2010.
- [Bud69] M. I. Budyko, *The effect of solar radiation variations on the climate of the Earth*, *Tellus* **21** (1969), 611–619.
- [CGLW16] M. D. Chekroun, M. Ghil, H. Liu, and S. Wang, *Low-dimensional Galerkin approximations of nonlinear delay differential equations*, *Disc. Cont. Dyn. Sys. A* **36** (2016), no. 8, 4133–4177.
- [CGN17] M.D. Chekroun, M. Ghil, and J. D. Neelin, *Pullback attractor crisis in a delay differential ENSO model*, *Advances in Nonlinear Geosciences*, to appear (A. Tsonis, ed.), Springer, 2017.
- [CH98] T. Cazenave and A. Haraux, *An Introduction to Semilinear Evolution Equations*, Oxford Lecture Series in Mathematics and its Applications, vol. 13, The Clarendon Press, Oxford, 1998.
- [CL15] M. D. Chekroun and H. Liu, *Finite-horizon parameterizing manifolds, and applications to suboptimal control of nonlinear parabolic PDEs*, *Acta Applicandae Mathematicae* **135** (2015), no. 1, 81–144.

- [CL16] ———, *Post-processing finite-horizon parameterizing manifolds for optimal control of nonlinear parabolic PDEs*, 2016 IEEE 55th Conference on Decision and Control (CDC), IEEE, 2016, pp. 1411–1416.
- [CLM16] M.D. Chekroun, H. Liu, and J.C. McWilliams, *The emergence of fast oscillations in a reduced primitive equation model and its implications for closure theories*, *Computers & Fluids* (2016).
- [CLW15a] M. D. Chekroun, H. Liu, and S. Wang, *Approximation of Stochastic Invariant Manifolds: Stochastic Manifolds for Nonlinear SPDEs I*, Springer Briefs in Mathematics, Springer, New York, 2015.
- [CLW15b] ———, *Stochastic Parameterizing Manifolds and Non-Markovian Reduced Equations: Stochastic Manifolds for Nonlinear SPDEs II*, Springer Briefs in Mathematics, Springer, New York, 2015.
- [CNK<sup>+</sup>14] M. D. Chekroun, J. D. Neelin, D. Kondrashov, J. C. McWilliams, and M. Ghil, *Rough parameter dependence in climate models: The role of Ruelle–Pollicott resonances*, *Proc. Natl. Acad. Sci. USA* **111** (2014), no. 5, 1684–1690.
- [Cou15a] National Research Council, *Climate intervention: Carbon dioxide removal and reliable sequestration*, National Academies Press, 2015.
- [Cou15b] ———, *Climate intervention: Reflecting sunlight to cool earth*, National Academies Press, 2015.
- [Cro00] T.J. Crowley, *Causes of climate change over the past 1000 years*, *Science* **289** (2000), no. 5477, 270–277.
- [DH98] J.I. Díaz and G. Hetzer, *A quasilinear functional reaction-diffusion equation arising in climatology*, *Equations aux dérivées partielles et applications (Articles dédiés à J.-L. Lions)*, Gauthier-Villars (Éditions Scientifique et Médicales Elsevier), 1998, pp. 461–480.
- [DH04] K. Deckelnick and M. Hinze, *Semidiscretization and error estimates for distributed control of the instationary Navier-Stokes equations*, *Numerische Mathematik* **97** (2004), no. 2, 297–320.
- [Día94a] J. I. Díaz, *Approximate controllability for some simple climate models*, *Environment, economic and their mathematical models* (J. I. Díaz and J. L. Lions, eds), *Research Notes in Applied Mathematics*, no. 35, Masson, Paris, France, 1994, pp. 29–43.
- [Día94b] ———, *Controllability and obstruction for some non linear parabolic problems in climatology*, *Modelado de Sistemas en Oceanografía, Climatología y Ciencias Medio Ambientales* (C. Pares and A. Valle, eds), Universidad de Málaga (1994), 43–58.
- [DTV<sup>+</sup>16] H.A. Dijkstra, A. Tantet, J. Viebahn, E. Mulder, M. Hebbink, D. Castellana, H. van den Pol, J. Frank, S. Baars, F. Wubs, M.D. Chekroun, and Ch. Kuehn, *A numerical framework to understand transitions in high-dimensional stochastic dynamical systems*, *Dynamics and Statistics of the Climate System* **1** (2016), no. 1, dzw003.
- [Fat99] H. O. Fattorini, *Infinite Dimensional Optimization and Control Theory*, *Encyclopedia of Mathematics and its Applications*, vol. 62, Cambridge University Press, 1999.
- [Fer97] R. Ferretti, *Internal approximation schemes for optimal control problems in Hilbert spaces*, *J. Mathematical Systems Estimation and Control* **7** (1997), 1–25.
- [Fur00] A. V. Fursikov, *Optimal Control of Distributed Systems: Theory and Applications*, *Translations of Mathematical Monographs*, vol. 187, American Mathematical Society, 2000.
- [GCS08] M. Ghil, M. D. Chekroun, and E. Simonnet, *Climate dynamics and fluid mechanics: Natural variability and related uncertainties*, *Physica D: Nonlinear Phenomena* **237** (2008), no. 14, 2111–2126.
- [Ghi76] M. Ghil, *Climate stability for a Sellers-type model*, *J. Atmos. Sci.* **33** (1976), 3–20.
- [Gib79] J. S. Gibson, *The Riccati integral equations for optimal control problems on Hilbert spaces*, *SIAM J. Control Optim.* **17** (1979), 537–565.
- [GK16] J. Garcke and A. Kröner, *Suboptimal feedback control of PDEs by solving HJB equations on adaptive sparse grids*, *Journal of Scientific Computing* (2016), 1–28.
- [GLN93] C.E. Graves, W.-H. Lee, and G.R. North, *New parameterizations and sensitivities for simple climate models*, *Journal of geophysical research* **98** (1993), no. D3, 5025–5036.
- [Gol85] J. A. Goldstein, *Semigroups of Linear Operators and Applications*, Oxford University Press, 1985.
- [Gri09] A. Grigor’yan, *Heat Kernel and Analysis on Manifolds*, American Mathematical Soc., 2009.
- [Hal09] S. Hallegatte, *Strategies to adapt to an uncertain climate change*, *Global environmental change* **19** (2009), no. 2, 240–247.
- [HCKN89] W.T. Hyde, T.J. Crowley, K.-Y. Kim, and G.R. North, *Comparison of GCM and energy balance model simulations of seasonal temperature changes over the past 18 000 years*, *Journal of Climate* **2** (1989), no. 8, 864–887.
- [HPUU09] M. Hinze, R. Pinnau, M. Ulbrich, and S. Ulbrich, *Optimization with PDE Constraints*, *Mathematical Modeling: Theory and Applications*, vol. 23, Springer, 2009.
- [Kie92] J. T. Kiehl, *Atmospheric general circulation modeling*, *Climate system modeling* (K. E. Trenberth, ed.), Cambridge University Press, 1992, pp. 317–370.
- [KKZ15] A. Kröner, K. Kunisch, and H. Zidani, *Optimal feedback control for undamped wave equations by solving a HJB equation*, *ESAIM Control Optim. Calc. Var.* **21** (2015), no. 2, 442–464. MR 3348407

- [Kno82] G. Knowles, *Finite element approximation of parabolic time optimal control problems*, SIAM Journal on Control and Optimization **20** (1982), no. 3, 414–427.
- [KT97] J. T. Kiehl and K. E. Trenberth, *Earth's annual global mean energy budget*, Bull. Am. Meteorol. Soc. **78** (1997), 197–208.
- [Las80] I. Lasiecka, *Unified theory for abstract parabolic boundary problems—a semigroup approach*, Applied mathematics & optimization **6** (1980), no. 1, 287–333.
- [Lio71] J.-L. Lions, *Optimal Control of Systems Governed by Partial Differential Equations*, Translated from the French by S. K. Mitter. Die Grundlehren der mathematischen Wissenschaften, Band 170, Springer-Verlag, New York-Berlin, 1971.
- [LT87] I. Lasiecka and R. Triggiani, *The regulator problem for parabolic equations with Dirichlet boundary control*, Applied Mathematics and Optimization **16** (1987), no. 1, 147–168.
- [LT00] ———, *Control Theory for Partial Differential Equations: Continuous and Approximation Theories. I*, Encyclopedia of Mathematics and its Applications, vol. 74, Cambridge University Press, Cambridge, 2000.
- [LV09] T. M. Lenton and N. E. Vaughan, *The radiative forcing potential of different climate geoengineering options*, Atmos. Chem. Phys. **9** (2009), 5539–5561.
- [Mal82] K. Malanowski, *Convergence of approximations vs regularity of solutions for convex, control-constrained optimal-control problems*, Applied Mathematics & Optimization **8** (1982), no. 1, 69–95.
- [MBJ73] R. S. McKnight and W. E. Bosarge JR., *The Ritz-Galerkin procedure for parabolic control problems*, SIAM J. Control Optim. **11** (1973), 510–524.
- [MMH<sup>+</sup>09] M. Meinshausen, N. Meinshausen, W. Hare, S.C.B. Raper, K. Frieler, R. Knutti, D.J. Frame, and M.R. Allen, *Greenhouse-gas emission targets for limiting global warming to 2 °C*, Nature **458** (2009), no. 7242, 1158–1162.
- [MSB<sup>+</sup>04] J.M. Murphy, D.M.H. Sexton, D.N. Barnett, G.S. Jones, M.J. Webb, M. Collins, and D.A. Stainforth, *Quantification of modeling uncertainties in a large ensemble of climate change simulations*, Nature **430** (2004), no. 7001, 768–772.
- [MTZ08] T.T. Medjo, R. Temam, and M. Ziane, *Optimal and robust control of fluid flows: some theoretical and computational aspects*, Applied Mechanics Reviews **61** (2008), no. 1, 010802.
- [MV07] D. Meidner and B. Vexler, *Adaptive space-time finite element methods for parabolic optimization problems*, SIAM Journal on Control and Optimization **46** (2007), no. 1, 116–142.
- [MW05] T. Ma and S. Wang, *Bifurcation Theory and Applications*, vol. 53, World Scientific, 2005.
- [MW14] ———, *Phase Transition Dynamics*, Springer, New York, 2014. MR 3154868
- [NBH<sup>+</sup>98] J. D. Neelin, D. S. Battisti, A. C. Hirst, F.-F. Jin, Y. Wakata, T. Yamagata, and S. E. Zebiak, *ENSO theory*, Journal of Geophysical Research: Oceans (1978–2012) **103** (1998), no. C7, 14261–14290.
- [NCC81] G. R. North, R. F. Cahalan, and J. A. Coakley, *Energy balance climate models*, Reviews of Geophysics **19** (1981), 91–121.
- [NSM83] G. R. North, D. A. Short, and J. G. Mengel, *Simple energy balance model resolving the seasons and the continents: application to the astronomical theory of the ice ages*, J. Geophys. Res. **88** (1983), 6576–6586.
- [NV12] I. Neitzel and B. Vexler, *A priori error estimates for space–time finite element discretization of semilinear parabolic optimal control problems*, Numerische Mathematik **120** (2012), no. 2, 345–386.
- [Paz83] A. Pazy, *Semigroups of Linear Operators and Applications to Partial Differential Equations*, Applied Mathematical Sciences, vol. 44, Springer-Verlag, New York, 1983.
- [RCC<sup>+</sup>14] L. Roques, M. Chekroun, M. Cristofol, S. Soubeyrand, and M. Ghil, *Parameter estimation for energy balance models with memory*, Proc. R. Soc. A **470** (2014), 20140349.
- [SAS<sup>+</sup>07] D. Stone, M. R. Allen, F. Selten, M. Kliphuis, and P.A. Stott, *The detection and attribution of climate change using an ensemble of opportunity*, Journal of Climate **20** (2007), no. 3, 504–516.
- [Sel69] W. D. Sellers, *A global climatic model based on the energy balance of the Earth-Atmosphere system*, J. Appl. Meteorol. **8** (1969), 392–400.
- [She09] J. G. Shepherd, *Geoengineering the climate: Science, governance and uncertainty*, Royal Society, 2009.
- [SLW<sup>+</sup>12] G.L. Stephens, J. Li, M. Wild, C. A. Clayson, N. Loeb, S. Kato, T. Lecuyer, P.W. Stackhouse Jr., M. Lebsock, and T. Andrews, *An update on Earth's energy balance in light of the latest global observations*, Nature Geoscience **5** (2012), no. 10, 691–696.
- [Trö93] F. Tröltzsch, *Semidiscrete Ritz-Galerkin approximation of nonlinear parabolic boundary control problems*, Optimal control (Freiburg, 1991), Internat. Ser. Numer. Math., vol. 111, Birkhäuser, Basel, 1993, pp. 57–68.
- [Trö94] ———, *Semidiscrete Ritz-Galerkin approximation of nonlinear parabolic boundary control problems—strong convergence of optimal controls*, Appl. Math. Optim. **29** (1994), no. 3, 309–329.
- [Trö10] F. Tröltzsch, *Optimal Control of Partial Differential Equations: Theory, Methods and Applications*, Graduate Studies in Mathematics, vol. 112, American Mathematical Society, 2010.

- [TSCJ94] E. Tziperman, L. Stone, M.A. Cane, and H. Jarosh, *El Niño chaos: Overlapping of resonances between the seasonal cycle and the Pacific ocean-atmosphere oscillator*, *Science* **264** (1994), no. 5155, 72–74.
- [VL11] N. E. Vaughan and T. M. Lenton, *A review of climate geoengineering proposals*, *Climatic change* **109** (2011), 745–790.
- [VN55] J. Von Neumann, *Can we survive technology?*, *Fortune* (June, 1955).

DEPARTMENT OF ATMOSPHERIC & OCEANIC SCIENCES, UNIVERSITY OF CALIFORNIA, LOS ANGELES, CA 90095-1565, USA

*E-mail address:* mchekroun@atmos.ucla.edu

INRIA AND CMAP, ÉCOLE POLYTECHNIQUE, CNRS, UNIVERSITÉ PARIS SACLAY, 91128 PALAISEAU, FRANCE

*E-mail address:* axel.kroener@inria.fr

DEPARTMENT OF MATHEMATICS, VIRGINIA POLYTECHNIC INSTITUTE AND STATE UNIVERSITY, BLACKSBURG, VIRGINIA 24061, USA

*E-mail address:* hhliu@vt.edu