Assessing Stabilization Through π - π Interactions in Aromatic Oligoamide β -Sheet Foldamers

Laure Sebaoun,^{†,¶} Brice Kauffmann,^{‡,§} Thomas Delclos,^{†,¶} Victor Maurizot,^{†,¶} and Ivan Huc*,^{†,¶}

[†] Univ. Bordeaux, CBMN (UMR 5248), Institut Européen de Chimie Biologie, 2 rue Escarpite 33600 Pessac, France

[¶] CNRS, CBMN (UMR 5248), France

[‡] Univ. Bordeaux, IECB (UMS 3033/US 001), 2 rue Escarpite, 33600 Pessac, France

[#] CNRS, IECB (UMS 3033), France

[§] INSERM, IECB (US 001), France

Supporting Information Placeholder

ABSTRACT: We have recently introduced aromatic oligoamide β -sheet foldamers based on rigid turn units and short linear strands that undergo intramolecular π - π stacking.¹ We now report that conformational stability in these structures can be reached using less rigid turn units and more extensive π - π interactions between longer linear strands. For this study, two-stranded sheets of variable length were prepared. Their conformation was assessed in solution by ¹H NMR and in the solid state by x-ray crystallography.

In comparison with helical synthetic foldamers, sheet-like structures have much less frequently been observed.² A possible reason for this may be that, when removed from the context of a tertiary structure as found in proteins, sheets tend to extensively aggregate and precipitate which makes them difficult to isolate and characterize. In contrast, helices generally are well-behaved discrete and soluble objects that fulfill their potential for non-covalent interactions intramolecularly. Indeed, synthetic foldamer sheets based on hydrogen-bonding are often designed to be dimeric³ in order to keep aggregation and solubility under control. Thus, the design of a soluble multistranded β -sheet foldamer would entail the orchestration of non covalent interactions so that they operate preferentially in an intramolecular fashion, thereby preventing extensive aggregation. Following this principle, we recently introduced aromatic oligoamide β -sheet foldamers stabilized by π - π interactions between short linear segments and rigid turn units such as **5** (Chart 1) that hold them in a face-to-face orientation.¹ A key feature of **5** lies in its four methyl groups *ortho* to the aromatic amine functions holding the two xylyl rings parallel to each other and perpendicular to the dinitrobenzene unit, as reflected by strong ring current effects which cause a large upfield shift of the NMR signal of the proton *meta* to both nitro groups pointing towards the interior of the structure (*H*_{int}).¹ As an extension to this initial design, we hereby demonstrate that rigidity at the turn units can be loosened and compensated by enhanced π - π interactions between extended linear aromatic oligoamide segments, without causing aggregation.

Chart 1. Two stranded aromatic oligoamide β -sheets possessing linear segments of increasing length.

Turn unit **1**, which lacks the methyl groups of **5**, was prepared from 4,6-dinitro-1,3-difluorobenzene using previously described procedures.¹ Compound **1** possesses two Boc-protected amine functions meant to attach linear strands and form sheet structures. Two isobutoxy chains were introduced to enhance its solubility in non protic organic solvents and to favor a coplanar orientation of each *para*-phenylenediamine units and its appended linear aromatic oligoamide segment, imparted by an intramolecular hydrogen bond between the terminal NH and adjacent isobutoxy oxygen atom, as found in other aromatic oligoamide foldamers.⁴

Trimeric turn **1** was first elongated using terephthalic acid monomers to form pentamer **2**. In order to substantially enhance face-to-face intramolecular π - π stacking, it was anticipated that larger units than simple benzene derivatives would be necessary. For this purpose, we introduced a new monomer, 6-amino-4-isobutoxy-7-methoxy-2-quinolinecarboxylic acid. As described in the supporting, its synthesis follows routes developed for other amino-2-quinolinecarboxylic acid monomers.⁵ The main feature of the new monomer is that the relative orientation of its 6-amino and 2-acid functions will confer a linear structure to oligoamide segments in which it is inserted.⁶ Elongation of pentamer **2** proceeded *via* the saponification of its terminal esters, and the subsequent coupling of two quinoline monomers or two quinoline dimers to yield **3** and **4**, respectively.

The chemical shift values of the H_{int} protons was monitored in CDCl_3 as a probe of the conformation of the new oligomers in this solvent (Fig 1) as they were found to undergo significant variations while $\delta_{H_{\text{ext}}}$ remained essentially unchanged. In rigid turn **5**, this resonance is strongly upfield shifted ($\delta_{H_{\text{int}}} = 4.91$ ppm), while the absence of methyl group in **1** makes its structure be more flexible and much reduce intramolecular ring current effects ($\delta_{H_{\text{int}}} = 6.68$ ppm). Upon elongation of trimer **1** into pentamer **2** and heptamer **3**, no major change is observed; H_{int} resonances actually undergo minor downfield shifts. In contrast, a strong upfield shift is observed between heptamer **3** ($\delta_{H_{\text{int}}} = 6.80$ ppm) and nonamer **4** ($\delta_{H_{\text{int}}} = 6.08$ ppm, $\Delta\delta = 0.72$ ppm). The effect on $\delta_{H_{\text{int}}}$ of the additional quinoline monomers at the end of the linear strands from **3** to **4** is very remote and suggests a better defined conformation of **4** leading to stronger ring current effects upon H_{int} . The non linear trend of $\delta_{H_{\text{int}}}$ values from **1** to **4** is indicative of cooperative effects.⁷

A series of multidimensional NMR experiments (HSQC, HMBC, TOCSY and ROESY) allowed us to fully assign the ^1H NMR spectrum of **4**. As shown in Fig. 2, ROESY experiments revealed contacts that unambiguously establish a folded two-stranded hairpin structure of **4** in solution. In particular inter-strand (as opposed to intra-strand) correlations are demonstrated when they occur between protons that are too distant to establish a contact if they would belong to the same aromatic ring such as, for example, 7-methoxy protons and the proton in position 5 of equivalent quinoline rings (OMe/ H_7 and OMe/ H_{10} correlations in Fig. 2). In contrast, no such correlations were observed in the spectra of heptamer **3**. Thus, the greater flexibility of turn **1** as compared to turn **5** can be compensated by extensive intramolecular π - π interactions between long linear strands. All observed correlations, and in particular OMe/ H_7 and OMe/ H_{10} (Fig. 2), are consistent with an antiparallel orientation of the two strands of **4**, with each ring head-to-tail with respect to the ring on which it stacks, as observed previously with multistranded sheets having shorter linear segments.¹ Yet the existence of conformers with a parallel orientation of the two strands of **4** cannot be completely ruled out from solution studies. Indeed, the interstrand NOE correlations expected in a parallel arrangement may not be distinguished from intrastrand correlations, unless the two strands are strongly offset in which case interstrand correlations may match with those expected in an antiparallel arrangement of the two strands. Spectra recorded over a wide temperature range (-50°C to 40°C) did not reveal any major change (Figure S2), suggesting that the same structure prevails over this range. Even at low temperature, the diastereotopicity of side chain isobutoxy protons does not appear

in the spectrum, indicating fast rotational dynamics of each linear strand with respect to the other. As a last point worth noting, it is unclear whether the H_{int} resonance in the folded conformation of **4** remains at lower field than in **5** because the latter is still better organized, because the former adopts a conformation at the turn in which ring current effects are weaker, or because ring currents are simply less intense in the former due to the different substitution patterns of its *para*-phenylenediamine rings.

Figure 1. Part of the 300 MHz ^1H NMR spectra in CDCl_3 at 298K of: a) hairpin turn trimer **1**; b) hairpin turn pentamer **2**; c) hairpin turn heptadecamer **3**; d) hairpin turn nonamer **4**; and e) hairpin turn trimer **5**. Signals of H_{ext} are marked with white circles. Signals of H_{int} are marked with full black circles. Stars indicate signals belonging to an impurity.

Figure 2. Schematic representation of ^1H - ^1H correlations observed in 400 MHz ROESY-2D spectra of nonamer **4** at 25°C in CDCl_3 . Blue arrows represent correlations within the hairpin turn; pink arrows represent interstrand correlations; others correlations are represented by grey arrows.

Solid state investigations were undertaken and fully confirmed solution data. Pentamer **2** was crystallized with two independent molecules in the asymmetric unit both showing divergent linear segments with no apparent stacking interactions between the *para*-phenylenediamine units or the terminal terephthalic acid units (Fig. 3a, 3b). On the contrary, the crystal structure of **4** (Fig. 3c, 3d) revealed a folded structure with extensive face-to-face π - π overlap between its two linear segments. The dinitrobenzene turn is perpendicular to the linear strands, which are found to be in an antiparallel orientation, presumably favored by local dipole-dipole interactions. Consistent with solution data, quinoline 7-methoxy protons are found much closer ($3.7 \text{ \AA}$) to protons in position 5 of the quinoline stacked above them than to the protons in position 5 of the quinoline to which they belong ($5.7 \text{ \AA}$). It is apparent in the structure of **4** that the interstrand distance imparted by the turn unit (4.7

Å) is too large for tight aromatic stacking to occur in short sequences such as **1** and **2**. Only the long strands of **4** compensate for this impediment, allowing its quinoline rings to stack at a distance of 3.4 Å.

The packing of **4** in the crystal shows extended head-to-tail stacks suggestive of a multi-stranded sheet-like aggregation mode (Fig. 3e). Nevertheless, this organization is restricted to the solid state, and compound **4** shows good solubility (> 13 mM in CDCl₃) and concentration independent chemical shift values up to 13 mM indicating limited aggregation in solution. Thus, the face-to-face π - π interactions that stabilize the two-stranded hairpin-turn structure of **4** are strong enough to be effective intramolecularly and weak enough not to prevail intermolecularly.

In summary, we have shown that π - π stacking may direct the folding of extended two-stranded β -hairpin structures in an organic solvent even when a relatively flexible turn unit is used. The balance between intra- and intermolecular interactions allows folding to occur and aggregation to be prevented. These objects are reminiscent or related foldamer structures including some crescent-like macrocycles,⁸ zippers based on oligoanthranilamides,⁹ as well as dimeric, trimeric and tetrameric aromatic oligoamide β -helices.¹⁰ It is anticipated that folding and aggregation of such β -hairpins will operate differently in protic media due to the very strong solvophobic component that makes π - π stacking be much stronger in these solvents.¹¹ Research along this line is in progress and will be reported in due course.

ASSOCIATED CONTENT

Supporting Information

Synthetic schemes, experimental procedures, full characterization of new compounds, crystallographic data, detailed NMR investigations including complete ¹H NMR assignment and solution structure elucidation of **4**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

* E-mail: i.huc@iecb.u-bordeaux.

Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

Notes

The authors declare no competing financial interest.

Figure 3. Side views of the crystal structures of: a) form A of hairpin turn pentamer **2**; b) form B of hairpin turn pentamer **2**; c) hairpin turn nonamer **4** and d) its top view. Packing of **4** in the crystal, e) side and f) front views. Protons and included solvent molecules have been removed for clarity.

ACKNOWLEDGMENT

This work was supported by the French Ministry of Research (pre-doctoral fellowship to L.S.) and by the European Research Council under the European Union's Seventh Framework Programme (Grant Agreement No. ERC-2012-AdG-320892).

REFERENCES

Graphic entry for the Table of Contents (TOC)

REFERENCES

- ¹ Sebaoun, L.; Maurizot, V.; Granier, T.; Kauffmann, B.; Huc, I. *J. Am. Chem. Soc.* **2014**, *136*, 2168.
- ² Guichard, G.; Huc, I. *Chem. Commun.* **2011**, 47, 5933.
- ³ (a) Nowick, J. S.; Chung, D. M.; Maitra, K.; Maitra, S.; Stigers, K. D.; Sun, Y. *J. Am. Chem. Soc.* **2000**, *122*, 7654. (b) Nowick, J. S.; Brower, J. O. *J. Am. Chem. Soc.* **2003**, *125*, 876. (c) Khakshoor, O.; Demeler, B.; Nowick, J. S. *J. Am. Chem. Soc.* **2007**, *129*, 5558. (d) Khakshoor, O.; Lin, A. J.; Korman, T. P.; Sawaya, M. R.; Tsai, S. C.; Eisenberg, D.; Nowick, J. S. *J. Am. Chem. Soc.* **2010**, *132*, 11622. (e) Cheng, P.-N.; Pham, J. D.; Nowick, J. S. *J. Am. Chem. Soc.* **2013**, *135*, 5477. (f) Cheng, P. N.; Pham, J. D.; Nowick, J. S. *J. Am. Chem. Soc.* **2013**, *135*, 5477. (g) Zhu, J.; Lin, J.-B.; Xu, Y.-X.; Shao, X.-B.; Jiang, X.-K.; Li, Z.-T. *J. Am. Chem. Soc.*, **2006**, *128*, 12307. (h) Gong, B.; Yan, Y.; Zeng, H.; Skrzypczak-Jankunn, E.; Kim, Y. W.; Zhu, J.; Ickes, H. *J. Am. Chem. Soc.* **1999**, *121*, 5607. (i) Yang, X.; Martinovic, S.; Smith, R. D.; Gong, B. *J. Am. Chem. Soc.* **2003**, *125*, 9932. (j) Gong, B. *Acc. Chem. Res.* **2012**, *45*, 2077. (k) Archer, E. A.; Sochia, A. E.; Krische, M. J. *Chem. Eur. J.* **2001**, *7*, 2059. (l) Archer, E. A.; Krische, M. J. *J. Am. Chem. Soc.*, **2002**, *124*, 5074 (m) Gong, H.; Krische, M. J. *J. Am. Chem. Soc.* **2005**, *127*, 1719.
- ⁴ For reviews about aromatic oligoamides foldamers, see: (a) Huc, I. *Eur. J. Org. Chem.* **2004**, 17. (b) Zhang, D.-W.; Zhao, X.; Hou, J. L.; Li, Z. *T. Chem. Rev.* **2012**, *112*, 5271.
- ⁵ (a) Qi, T.; Deschrijver, T.; Huc, I. *Nat. Protoc.* **2013**, *8*, 693. (b) Ferrand, Y.; Kendhale, A. M.; Garric, J.; Kauffmann, B.; Huc, I. *Angew. Chem. Int. Ed.* **2010**, *49*, 1778.
- ⁶ (a) Hamuro, Y.; Geib, S. J.; Hamilton, A. D. *J. Am. Chem. Soc.* **1996**, *118*, 7529. (b) Wu, Z.-Q.; Jiang, X.-K.; Zhu, S.-Z.; Li, Z.-T. *Org. Lett.* **2004**, *6*, 229. (c) Odriozola, I.; Kyrtsakas, N.; Lehn, J.-M. *Chem. Commun.* **2004**, 62.
- ⁷ Stone, M. T.; Heemstra, J. M.; Moore, J. S. *Acc. Chem. Res.* **2006**, *39*, 11-20.
- ⁸ Lin, L.; Zhang, J.; Wu, X.; Liang, G.; He, L.; Gong, B. *Chem. Commun.* **2010**, 46, 7361.
- ⁹ Nair, R. V.; Kheria, S.; Rayavarapu, S.; Kotmale, A. S.; Jagadeesh, B.; Gonnade, R. G.; Puranik, V. G.; Rajamohanam, P. R.; Sanjayan, G. J. *J. Am. Chem. Soc.* **2013**, *135*, 11477.
- ¹⁰ (a) Berl, V.; Huc, I.; Khoury, R.; Krische, M. J.; Lehn, J.-M. *Nature* **2000**, *407*, 720. (b) Berni, E.; Kauffmann, B.; Bao, C.; Lefeuvre, J.; Bassani, D. M.; Huc, I. *Chem. Eur. J.* **2007**, *13*, 8463. (c) Baptiste, B.; Zhu, J.; Haldar, D.; Kauffmann, B.; Léger, J.-M.; Huc, I. *Chem. Asian J.* **2010**, *5*, 1364. (d) Ferrand, Y.; Kendhale, A. M.; Garric, J.; Kauffmann, B.; Huc, I. *Angew. Chem. Int. Ed.* **2010**, *49*, 1778. (e) Gan, Q.; Bao, C.; Kauffmann, B.; Grélard, A.; Xiang, J.; Liu, S.; Huc, I.; Jiang, H. *Angew. Chem. Int. Ed.* **2008**, *47*, 1715.
- ¹¹ (a) Qi, T.; Maurizot, V.; Noguchi, H.; Charoenraks, T.; Kauffmann, B.; Takafuji, M.; Ihara, H.; Huc, I. *Chem. Commun.* **2012**, 48, 6337. (b) Bradford, V. J.; Iverson, B. L. *J. Am. Chem. Soc.* **2008**, *130*, 1517. (c) Zych, A. J.; Iverson, B. L. *J. Am. Chem. Soc.* **2000**, *122*, 8898. (d) Gabriel, G. J.; Sorey, S.; Iverson, B. L. *J. Am. Chem. Soc.* **2005**, *127*, 2637. (e) B. L. Lokey, R. S.; Iverson, B. L. *Nature* **1995**, 375, 303.