

Labeled embedding of $(n, n - 2)$ -graphs in their complements

Mohammed Amin Tahraoui, Eric Duchene, Hamamache Kheddouci, Mariusz Wozniak

► **To cite this version:**

Mohammed Amin Tahraoui, Eric Duchene, Hamamache Kheddouci, Mariusz Wozniak. Labeled embedding of $(n, n - 2)$ -graphs in their complements. *Discussiones Mathematicae Graph Theory*, University of Zielona Góra, 2017, <10.7151/dmgt.1977>. <hal-01518715>

HAL Id: hal-01518715

<https://hal.archives-ouvertes.fr/hal-01518715>

Submitted on 29 Nov 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

4 **LABELED EMBEDDING OF $(N, N - 2)$ -GRAPHS IN THEIR**
5 **COMPLEMENTS**

6 M-A. TAHRAOUI

7 *Université de Lyon, CNRS, Université Lyon 1, LIRIS,*
8 *UMR5205, F-69622, France*

9 E. DUCHÊNE

10 *Université de Lyon, CNRS, Université Lyon 1, LIRIS,*
11 *UMR5205, F-69622, France*

12 **e-mail:** eric.duchene@univ-lyon1.fr

13 H. KHEDDOUCI

14 *Université de Lyon, CNRS, Université Lyon 1, LIRIS,*
15 *UMR5205, F-69622, France*

17 AND

18 M. WOŹNIAK

19 *Department of Discrete Mathematics, Faculty of Applied Mathematics,*
20 *AGH-University of Science and Technology, Krakow, Poland.*

21 **Abstract**

22 Graph packing generally deals with unlabeled graphs. In [4], the authors
23 have introduced a new variant of the graph packing problem, called the
24 *labeled packing of a graph*. This problem has recently been studied on trees
25 [6] and cycles [4]. In this note, we present a lower bound on the labeled
26 packing number of any $(n, n - 2)$ -graph into K_n . This result improves the
27 bound given by Woźniak in [7].

28 **Keywords:** Packing of graphs, Labeled packing, Permutation.

29 **2010 Mathematics Subject Classification:** 05C70.

30

1. CONTEXT AND DEFINITIONS

31 **Graph theoretical definitions**

32 All graphs considered in this paper are finite, undirected, without loops or
 33 multiple edges. If T is a rooted tree of order n , we define an *end vertex* as a
 34 vertex which does not have any son, and a *leaf-parent* as a vertex whose all of its
 35 sons are end vertices.

36 Given a positive integer n , the graphs K_n , P_n and C_n will denote respectively the
 37 complete graph, the path and the cycle on n vertices. For a graph G , we will use
 38 $V(G)$ and $E(G)$ to denote its vertex and edge sets respectively. Given $V' \subset V$,
 39 the subgraph $G[V']$ denotes the subgraph of G induced by V' , i.e., $E(G[V'])$
 40 contains all the edges of E which have both extremities in V' . If a graph G has
 41 order n and size m , we say that G is an (n, m) -graph.

42 An independent set of G is a subset of vertices $X \subseteq V$, such that no two vertices
 43 in X are adjacent. An independent set is said to be maximal if no independent
 44 set properly contains it. An independent set of maximum cardinality is called a
 45 maximum independent set. For undefined terms, we refer the reader to [2]. A
 46 permutation σ is a one-to-one mapping of $\{1, \dots, n\}$ into itself. We say that a
 47 permutation σ is *fixed-point-free* if $\sigma(x) \neq x$ for all x of $\{1, \dots, n\}$.

48 **The graph packing problem**

49 The graph packing problem was introduced by Bollobás and Eldridge [1] and
 50 Sauer and Spencer [5] in the late 1970s. Let G_1, \dots, G_k be k graphs of order
 51 n . We say that there is a packing of G_1, \dots, G_k (into the complete graph K_n
 52) if there exist bijections $\sigma_i : V(G_i) \rightarrow V(K_n)$, where $1 \leq i \leq k$, such that
 53 $\sigma_i^*(E(G_i)) \cap \sigma_j^*(E(G_j)) = \emptyset$ for $i \neq j$, and here the map $\sigma_i^* : E(G_i) \rightarrow E(K_n)$
 54 is the one induced by σ_i . A packing of k copies of a graph G will be called a
 55 k -placement of G . A packing of two copies of G (i.e., a 2-placement) is also called
 56 an embedding of G (into its complement \overline{G}). In other words, an embedding of
 57 a graph G is a permutation σ on $V(G)$ such that for each edge uv belonging to
 58 $E(G)$, its image $\sigma(u)\sigma(v)$ does not belong to $E(G)$.

59

60 In the literature, the question of the existence of an embedding of a given
 61 graph received a great attention (see the survey papers [8, 9]). In [3], full char-
 62 acterizations of all the $(n, n-1)$ and (n, n) embeddable graphs are given. The
 63 case of $(n, n-2)$ -graphs was also solved independently in [1, 3, 5]. In particular,
 64 it is proved in [5] that any pair of $(n, n-2)$ -graphs can be packed into K_n .

65

66 In [4], Duchêne *et al.* introduced and studied the graph packing problem for
 67 a vertex labeled graph. Roughly speaking, it consists of a graph packing which

68 preserves the labels of the vertices. We give below the formal definition of this
 69 problem.

70 **Definition** [4]. Given a positive integer p , let G be a graph of order n and f be
 71 a mapping from $V(G)$ to the set $\{1, \dots, p\}$. The mapping f is called a p -labeled-
 72 packing of k copies of G into K_n if there exist bijections $\sigma_i : V(G) \rightarrow V(K_n)$
 73 for $1 \leq i \leq k$, such that:

- 74 1. $\sigma_i^*(E(G)) \cap \sigma_j^*(E(G)) = \emptyset$ for all $i \neq j$.
- 75 2. For every vertex v of G , we have $f(v) = f(\sigma_1(v)) = f(\sigma_2(v)) = \dots = f(\sigma_k(v))$.

76 The maximum positive integer p for which G admits a p -labeled-packing of k
 77 copies of G is called the *labeled packing number* of k copies of G and is denoted
 78 by $\lambda^k(G)$. Throughout this paper, a labeled packing of two copies of G will be
 79 called a labeled embedding of G . It will be denoted by a pair (f, σ) .

80 Remark that the existence of a packing of k copies of a graph G is a neces-
 81 sary condition for the existence of p -labeled-packing of k copies of G . Indeed, it
 82 suffices to choose $p = 1$. Therefore, the result of Sauer and Spencer [5] ensures
 83 the existence of a p -labeled packing for $(n, n - 2)$ -graphs. An estimation of the
 84 labeled packing number of such graphs is the main issue of the current paper.

85

86 The following result was proved in [4]. It gives an upper bound for the labeled
 87 packing number of two copies of a general graph.

Lemma 1 (Duchêne et al., 11). *Let G be a graph of order n and let I be a maximum independent set of G . If there exists an embedding of G into K_n , then*

$$\lambda^2(G) \leq |I| + \lfloor \frac{n - |I|}{2} \rfloor$$

In [4], exact values of $\lambda^2(G)$ are given when G is a cycle or a path. In almost all cases, the upper bound of the above lemma is reached. More precisely, it is shown that for all $n \geq 6$,

$$\lambda^2(P_n) \in \{ \lfloor \frac{3n}{4} \rfloor, \lfloor \frac{3n}{4} \rfloor + 1 \}$$

$$\lambda^2(C_n) = \lfloor \frac{3n}{4} \rfloor$$

88 The case of trees is also considered [6], but only a lower bound is proposed.

89

2. LABELED EMBEDDING OF GRAPHS AND PERMUTATIONS

90 In this section, we give a strong relationship between a labeled embedding and
91 its permutation structure.

92 A permutation σ of a finite set can be written as the disjoint union of cycles
93 (two cycles being disjoint if they do not have any common element). Here, a
94 cycle (a_1, \dots, a_n) is a permutation sending a_i to a_{i+1} for $1 \leq i \leq n-1$ and a_n to
95 a_1 . This representation is called the *cyclic decomposition of σ* and is denoted by
96 $C(\sigma)$. According to this definition, the cycles of length one correspond to fixed
97 points of σ . For example, the cyclic decomposition of the permutation induced
98 by the labeled embedding of T (in Figure 1) is: $\{(v_1), (v_2), (v_3), (v_4), (v_5), (v_6),$
99 $(v_7), (v_8, v_{10}), (v_{11}, v_{13}), (v_9, v_{12})\}$.

100 We now recall a fundamental property of labeled embeddings (see [4]). For
101 any labeled embedding (f, σ) of a graph G , one can remark that the vertices of
102 every cycle of $C(\sigma)$ share the same label. In other words, the labeled embedding
103 number of G exactly corresponds to the maximum number of cycles induced by
104 an embedding of G . It means that if G admits an embedding with k cycles, then
105 $\lambda^2(G) \geq k$.

106

107 Although this correlation between labeled embeddings and the permutation's
108 number of cycles was recently stated, several studies can be found about the
109 permutation structure of an embedding. In particular, the permutation structure
110 of embeddings of $(n, n-2)$ -graphs was investigated by Woźniak in [7]:

111 **Theorem 2** (Woźniak, 94). *Let G be a graph of order n , different from $K_3 \cup 2K_1$
112 and $K_4 \cup 4K_1$. If $|E(G)| \leq n-2$, then there exists a permutation σ on $V(G)$
113 such that $\sigma_1, \sigma_2, \sigma_3$ define a 3-placement of G . Moreover, σ has all its cycles
114 of length 3, except for one of length one if $n \equiv 1 \pmod{3}$ or two of length one if
115 $n \equiv 2 \pmod{3}$.*

116 According to our previous remarks, the above theorem induces the following
117 result in the context of labeled embeddings.

Corollary 3. *Let G be a graph of order n , different from $K_3 \cup 2K_1$ and $K_4 \cup 4K_1$.
If $|E(G)| \leq n-2$, then*

$$\lambda^2(G) \geq \lfloor \frac{n}{3} \rfloor + n \pmod{3}$$

118 In the next section, we will show that the lower bound of Corollary 3 can be
119 improved (including for the excluded graphs).

120

3. MAIN RESULT

121 We first define the notion of *good permutation* for a graph.

122 **Definition.** Given a graph G , a permutation σ on $V(G)$ is said to be *good* if

- 123 • σ is an embedding of G ,
- 124 • σ has at least $\lfloor \frac{2n}{3} \rfloor$ cycles,
- 125 • every cycle of σ is of order at most 2, *i.e.*, for every pair of distinct vertices
- 126 u, v of G , if $\sigma(u) = v$, then $\sigma(v) = u$.

127 The following lemma will be useful in a special case of our main result.

128 **Lemma 4.** For $k > 0$, the graph $kC_3 \cup 2K_1$ admits a good permutation.

129 **Proof.** According to the diagram below (Figure 1), first remark that $3C_3$ admits
 130 a good permutation. Indeed, the numbers inside the vertices correspond to a
 131 labeled embedding with 6 labels, with at most two vertices sharing the same
 label.




Figure 1. Good permutation for $3C_3$

132

133 Now let k be a positive integer and G be the graph $kC_3 \cup 2K_1$. Let u and t
 134 be the two isolated vertices of G . For $1 \leq i \leq k$, let $\{v_{i1}, v_{i2}, v_{i3}\}$ be the vertices
 135 of the i^{th} triangle C_3 . For $k = 1$, consider the permutation σ where v_{11} is a
 136 fixed point, v_{12} and u are mutual images, as well as v_{13} and t . One can easily
 137 check that σ is good for G . For $k = 2$, Figure 2 shows a good permutation (more
 precisely, the corresponding labeled embedding).




Figure 2. Good permutation for $2C_3 \cup 2K_1$

138

139 For $k = 3$, consider the permutation σ corresponding to the labeled embed-
 140 ding of Figure 1, and extend it to G by setting $\sigma(u) = u$ and $\sigma(t) = t$. Then σ
 141 remains good for G . For $k > 3$, we can now conclude to the existence of a good
 142 permutation for G by pairing good permutations of $3C_3$ with a good permutation
 143 of $rC_3 \cup 2K_1$ where r is in $\{1, 2, 3\}$. ■

144 We now present a lower bound for the labeled embedding number of any
145 $(n, n - 2)$ -graph.

Theorem 5. *Let $n \geq 2$ and G be an (n, m) -graph with $m \leq n - 2$. The following inequality holds:*

$$\lambda^2(G) \geq \lfloor \frac{2n}{3} \rfloor$$

146 **Proof.** Let $n \geq 2$ and G be an (n, m) -graph with $m \leq n - 2$. Without loss
147 of generality, we can assume $|E(G)| = n - 2$. We will show that G admits a
148 good permutation by induction on n . If $n = 2, 3, 4$, then $G \in \{2K_1, 3K_1, K_1 \cup$
149 $K_2, 2K_2, K_{1,2} \cup K_1\}$. In each case, one can quickly check that there exist good
150 permutations with at least two cycles. The property still holds for $n = 5$, where
151 $G \in \{K_3 \cup 2K_1, K_1 \cup K_{1,3}, K_2 \cup K_{1,2}, P_4 \cup K_1\}$. Good permutations with at least
152 three cycles can be found.

153 Now let $n \geq 6$ and assume there exists a good permutation for every $(n', n' -$
154 $2)$ -graph of order $n' < n$ with $n' \geq 3$. Since G is an $(n, n - 2)$ -graph, at least
155 two of its connected components are trees. Denote by T and H two trees of G of
156 highest orders such that $|V(T)| \geq |V(H)|$. In what follows, we choose to consider
157 T and H as rooted trees. We consider the following four cases:

158

159 **Case 1:** $|V(T)| \geq 3$ and $|V(H)| \geq 3$. Two subcases are considered as follows:

160 **Subcase 1.1:** T or H admits a leaf-parent of degree 2. By symmetry, we may
161 assume that T admits a leaf-parent say x_1 , of degree 2. Let x_0 and x_2 be the
162 two vertices of T such that (x_0, x_1, x_2) is an induced path of T and x_2 is an end
163 vertex. Let y_1 be an end vertex of H and y_0 its parent. Now consider the graph
164 $G' = G - \{x_1, x_2, y_1\}$. Clearly, G' is an $(n - 3, n - 5)$ -graph with $n - 3 \geq 3$. Hence
165 the induction hypothesis guarantees the existence of a good permutation σ' for
166 G' . This permutation can be extended to a good permutation σ for G as follows:

167

$$\sigma(x_1) = \begin{cases} y_1 & \text{if } \sigma'(x_0) = x_0, \\ x_1 & \text{otherwise.} \end{cases} \quad \sigma(x_2) = \begin{cases} x_2 & \text{if } \sigma'(x_0) = x_0, \\ y_1 & \text{otherwise.} \end{cases}$$

168

$$\sigma(y_1) = \begin{cases} x_1 & \text{if } \sigma'(x_0) = x_0, \\ x_2 & \text{otherwise.} \end{cases} \quad \sigma(v) = \sigma'(v) \text{ if } v \in V(G')$$

169

170 Since the number of cycles of $\sigma|_{G-G'}$ equals two, and they all are of length
171 at most 2, it ensures that σ is a good permutation for G .

172 **Subcase 1.2:** T and H do not have any leaf-parent of degree 2 and T is a
173 star. Let ℓ be a leaf of H and v be its neighbour (of degree at least 3). Let u
174 be the unique vertex of degree at least 3 in T . Now consider the graph $G' =$
175 $G - T - \{\ell, v\}$. Clearly, G' is an (k, j) -graph with $k - 2 \geq j$ and $k \geq 2$. Hence
176 the induction hypothesis guarantees the existence of a good permutation σ' for

177 G' . This permutation can be extended to a good permutation σ for G by setting
 178 $\sigma(u) = v$, $\sigma(v) = u$, $\sigma(\ell) = \ell$ and $\sigma(l_i) = l_i$ for all leaf l_i of T .

179 **Subcase 1.3:** T and H do not have any leaf-parent of degree 2 and T is not
 180 a star. Hence T admits two leaf parents, say u and t . Let u_1, \dots, u_k (resp.
 181 $t_1, \dots, t_{k'}$) be the leaves adjacent to u (resp. t), with $k, k' \geq 2$. Let ℓ be a leaf of
 182 H and v be its neighbour (of degree at least 3). Let x be the non-leaf neighbour
 183 of u (it exists since T is not a star and $x = t$ if T is a bistar). See Figure 3 for a
 184 better view of these notations. Without loss of generality, we will assume $k \geq k'$.




Figure 3. Subcase (1.3)

185 If $k \geq 3$, consider the graph $G' = G - \{u, u_1, \dots, u_k, t_1, \ell\}$. Hence G' is a
 186 $(n - k - 3, n - k - 5)$ -graph satisfying $|V(G')| \geq 3$ (since $|V(H)| \geq 4$), and the
 187 induction hypothesis can be applied. Let σ' be a good permutation for G' . We
 188 now define a permutation σ for G as follows:

$$\sigma(u) = \begin{cases} t_1 & \text{if } \sigma'(x) = v, \\ \ell & \text{otherwise.} \end{cases} \quad \sigma(t_1) = \begin{cases} u & \text{if } \sigma'(x) = v, \\ u_1 & \text{otherwise.} \end{cases}$$

$$189 \quad \sigma(u_1) = \begin{cases} \ell & \text{if } \sigma'(x) = v, \\ t_1 & \text{otherwise.} \end{cases} \quad \sigma(\ell) = \begin{cases} u_1 & \text{if } \sigma'(x) = v, \\ u & \text{otherwise.} \end{cases}$$

$$190 \quad \sigma(u_i)_{i>1} = u_i \quad \sigma(v) = \sigma'(v) \text{ if } v \in V(G')$$

191 This permutation remains good as $(k + 1)$ cycles of lengths 1 or 2 and $(k + 3)$
 192 vertices have been added to σ' at G' respectively.

193
 194 If $k = k' = 2$, we first consider the case where $x = t$ (i.e., T is a bistar).
 195 Consider the graph $G' = G - \{u_1, t_1, \ell\}$ and proceed by induction as above. Since
 196 ut is an edge of T , we have $\sigma'(u) \neq \sigma'(t)$ and u and t cannot be fixed points
 197 simultaneously. Therefore, at least u_1 or t_1 can be set as a fixed point in σ
 198 and the two remaining vertices in $\{u_1, t_1, \ell\}$ can be images of each other (thus
 199 implying that σ is good).

200 In the case where $x \neq t$, consider the graph $G' = G - \{u, u_1, u_2, t_1, t_2, \ell\}$, and
 201 proceed as previously with the following permutation:

$$\sigma(u) = \begin{cases} t_1 & \text{if } \sigma'(t) = t, \\ t_1 & \text{if } \sigma'(t) \neq t \text{ and } \sigma'(x) = v, \\ \ell & \text{otherwise.} \end{cases} \quad \sigma(t_1) = \begin{cases} u & \text{if } \sigma'(t) = t, \\ u & \text{if } \sigma'(t) \neq t \text{ and } \sigma'(x) = v, \\ t_1 & \text{otherwise.} \end{cases}$$

$$202 \quad \sigma(t_2) = \begin{cases} \ell & \text{if } \sigma'(t) = t, \\ t_2 & \text{if } \sigma'(t) \neq t \text{ and } \sigma'(x) = v, \\ t_2 & \text{otherwise.} \end{cases} \quad \sigma(\ell) = \begin{cases} t_2 & \text{if } \sigma'(t) = t, \\ \ell & \text{if } \sigma'(t) \neq t \text{ and } \sigma'(x) = v, \\ u & \text{otherwise.} \end{cases}$$

$$203 \quad \sigma(u_i)_{i=1,2} = u_i \qquad \sigma(v) = \sigma'(v) \text{ if } v \in V(G')$$

204 One can easily check that σ remains good since 6 vertices and at least 4
 205 cycles have been added to σ' .

206 **Case 2:** $|V(T)| \geq 3$ and $H = K_1$.

207 **Subcase 2.1:** There exists a leaf-parent, say x , of degree at least 3. Let ℓ be
 208 one of its leaves, and y be the unique vertex of H . Now consider the graph
 209 $G' = G - \{x, \ell, y\}$, which satisfies $|E(G')| \leq |V(G')| - 2$. Hence it admits a good
 210 permutation σ' by induction hypothesis. A good permutation σ of G can thus be
 211 extended from G' by setting $\sigma(x) = y$, $\sigma(y) = x$ and $\sigma(\ell) = \ell$.

212 **Subcase 2.2:** All the vertices of T which are adjacent to leaves are of degree 2.
 213 Let x_0 be such a vertex (it exists since $|V(T)| \geq 3$), let ℓ_1 be its adjacent leaf,
 214 and x_1 its second neighbor. Now let ℓ_2 be a distinct leaf from ℓ_1 in T , and x_2 be
 215 its neighbor. If $x_1 \neq \ell_2$ and $x_1 \neq x_2$, then consider $G' = G - \{x_0, \ell_1, \ell_2\}$, which
 217 admits a good permutation σ' by induction hypothesis. Then set $\sigma|_{G'} = \sigma'$ and

218 • If $\sigma'(x_1) = x_1$:

$$219 \quad \sigma(x_0) = \ell_2, \sigma(\ell_2) = x_0, \text{ and } \sigma(\ell_1) = \ell_1.$$

220 • If $\sigma'(x_1) \neq x_1$:

$$221 \quad \sigma(x_0) = x_0, \sigma(\ell_2) = \ell_1, \text{ and } \sigma(\ell_1) = \ell_2.$$

222 One can now easily check that σ is good for G . If $x_1 = \ell_2$ or $x_1 = x_2$, then T
 223 is either a P_3 or a P_4 . Since $n \geq 6$, it implies that G admits at least another
 224 connected component which is an $(n, n-1)$ or an (n, n) connected graph with
 225 $n \geq 2$ (since $|E(G)| \geq 4$, there exists at least one edge in $G - T$). In other words,
 226 this component is either a tree T' , or a tree with an edge $T' \cup \{e\}$. Let ℓ_3 be
 227 a leaf in T' . Note that we do not care whether ℓ_3 is adjacent to e or not. By
 228 considering $G' = G - \{x_0, \ell_1, \ell_3\}$ together with the above permutation where ℓ_2

229 is replaced by ℓ_3 , we find a good permutation for G .

230

231 **Case 3:** $|V(T)| = 2$. Let $T = (x_0, x_1)$ and let y be a vertex of degree 2 of G .
 232 Such a vertex exists since $n \geq 6$. Consider the graph $G' = G - \{x_0, x_1, y\}$. By
 233 induction hypothesis, there exists a good permutation for G' , say σ' . We set
 234 $\sigma(x_0) = y, \sigma(y) = x_0, \sigma(x_1) = x_1$ and for every vertex $v \in V(G')$, $\sigma(v) = \sigma'(v)$,
 235 which defines a good permutation for G .

236

237 **Case 4:** $|V(T)| = 1$. In this case, G contains isolated vertices (at least two) and
 238 non-tree connected components. Two subcases are considered as follows:

239 **Subcase 4.1:** G has a vertex, say x , of degree at least 3. Let y and z be two
 240 isolated vertices of G . Consider the graph $G' = G - \{x, y, z\}$. The induction
 241 hypothesis guarantees the existence of a good permutation σ' for G' . By putting
 242 $\sigma(x) = y, \sigma(y) = x, \sigma(z) = z$ and for every vertex $v \in V(G')$, $\sigma(v) = \sigma'(v)$, we
 243 get a good permutation for G .

244 **Subcase 4.2:** *The complementary subcase to (4.1), i.e., G is the sum of two*
 245 *isolated vertices and an union of cycles.* This case is solved as follows:

246 (a) $G = kC_3 \cup 2K_1$ for some $k \geq 1$. Lemma 4 allows us to conclude.

247 (b) G has at least one cycle, say H , of order at least 4, and one cycle, say Q ,
 248 of order at least 3: let (x_1, x_2, x_3) be an induced path of H , let x_4 be a vertex of
 249 Q and z, t be the two isolated vertices of G . Denote by x (resp. y) the neighbor
 250 of x_1 (resp. x_3) different from x_2 . Note that we may have $x = y$ in the case
 251 $H = C_4$. See Figure 4 for a graphical depiction of these notations.




Figure 4. Case (4.2.b)

252 Consider the graph $G' = G - \{x_1, x_2, x_3, x_4, z, t\}$. Since $|V(G)| \geq 9$, we
 253 have $|V(G')| \geq 3$ and the induction hypothesis guarantees the existence of a
 254 good permutation σ' for G' . The permutation σ' can be extended to a good
 255 permutation σ of G by setting $\sigma(t) = t$, and


256 Hence $\sigma|_{G-G'}$ has four cycles of size at most 2, and σ is thus good for G .

257 (c) G is the sum of C_m (for some $m \geq 4$) and two isolated vertices. If $m < 8$,
 258 then Figure 5 shows labeled embeddings corresponding to good permutations.

$$\sigma(x_1) = \begin{cases} x_1 & \text{if } \sigma'(x) \neq x \text{ and } \sigma'(y) \neq y, \\ x_4 & \text{if } \sigma'(x) = x, \\ z & \text{otherwise.} \end{cases} \quad \sigma(x_2) = \begin{cases} x_4 & \text{if } \sigma'(x) \neq x \text{ and } \sigma'(y) \neq y, \\ x_2 & \text{otherwise.} \end{cases}$$

$$\sigma(x_3) = \begin{cases} x_3 & \text{if } \sigma'(x) \neq x \text{ and } \sigma'(y) \neq y, \\ z & \text{if } \sigma'(x) = x, \\ x_4 & \text{otherwise.} \end{cases} \quad \sigma(x_4) = \begin{cases} x_2 & \text{if } \sigma'(x) \neq x \text{ and } \sigma'(y) \neq y, \\ x_1 & \text{if } \sigma'(x) = x, \\ x_3 & \text{otherwise.} \end{cases}$$

$$\sigma(z) = \begin{cases} z & \text{if } \sigma'(x) \neq x \text{ and } \sigma'(y) \neq y, \\ x_3 & \text{if } \sigma'(x) = x, \\ x_1 & \text{otherwise.} \end{cases}$$

Figure 5. Case $C_m \cup 2K_1$ for $m = 4, \dots, 7$

259 If $m \geq 8$, let (x_1, \dots, x_8) be a path of C_m . Let z, t be the two isolated vertices
260 of G . We consider the graph $G' = G - \{x_2, x_3, x_6, x_7, z, t\}$ which admits a good
261 permutation σ' by induction hypothesis. Since v_4 and v_5 are adjacent, at least
262 one of them is not a fixed point under σ' . Without loss of generality, assume
263 $\sigma'(x_4) \neq x_4$. The permutation σ' can be extended to a good permutation σ for
264 G as follows: set x_3 and t as fixed points. If $\sigma'(x_5) \neq x_1$, we set $\sigma(x_2) = x_6$,

265 $\sigma(x_6) = x_2$, $\sigma(x_7) = z$, and $\sigma(z) = x_7$. Otherwise, we set $\sigma(x_2) = x_7$, $\sigma(x_7) =$
 266 x_2 , $\sigma(x_6) = z$, and $\sigma(z) = x_6$. For the same reasons as in case (4.2.b), this
 267 permutation is good for G .

268

■

269 CONCLUSION

270 Theorem 5 gives a first lower bound about the labeled embedding number of
 271 $(n, n - 2)$ -graphs. Yet, the computation of the exact value remains an open
 272 question, as this bound is not exact for many families of $(n, n - 2)$ -graphs. As an
 273 example, consider a cycle C_n without two edges. Its labeled packing number is
 274 at least the one of C_n , (i.e., $\lfloor 3n/4 \rfloor$). Yet, for any large value of n , we can find an
 275 $(n, n - 2)$ -graph for which the bound is tight. Indeed, consider G as an union of k
 276 disjoint triangles with $K_2 \cup K_1$. The size of a maximum independent set for this
 277 graph equals $k+2$. According to Lemma 1, we have that $\lambda_2(G) = 2k+2 = \lfloor 2n/3 \rfloor$.

278 In addition, we mention that this result can be used to study the labeled
 279 embedding of $(n, n - 1)$ -graphs. One can show for example that the same bound
 280 is valid for the union of cycles with a single tree.

281 REFERENCES

282 [1] B. Bollobás and S. E. Eldridge, *Packing of graphs and applications to com-*
 283 *putational complexity*, J. Comb. Theory (B) **25** (105–124) 1978.

284 [2] J. A. Bondy and U. S. R. Murty, *Graph Theory with Applications* (McMil-
 285 lan, London; Elsevier, New York, 1976).

286 [3] D. Burns and S. Schuster, *Every $(p, p - 2)$ graph is contained in its comple-*
 287 *ment*, J. Graph Theory **1** (277–279) 1977.

288 [4] E. Duchêne, H. Kheddouci, R.J. Nowakowski and M.A. Tahraoui, *Labeled*
 289 *packing of graphs*, Australasian Journal of Combinatorics **57** (109–126) 2013.

290 [5] N. Sauer and J. Spencer, *Edge disjoint placement of graphs*, J. Combin.
 291 Theory Ser. B **25** (295–302) 1978.

292 [6] M. A. Tahraoui, E. Duchêne and H. Kheddouci, *Labeled 2-packings of trees*,
 293 Discrete Mathematics **338(5-6)** (816–824) 2015.

294 [7] M. Woźniak, *Embedding Graphs of Small Size*, Discrete Applied Mathemat-
 295 ics **51(1-2)** (233–241) 1994.

296 [8] M. Woźniak, *Packing of graphs and permutations—a survey* Discrete Mathe-
 297 matics **276** (1-3) 379–391 2004

- ²⁹⁸ [9] H. P. Yap Packing of graphs—a survey *Discrete Mathematics* 72:395–404 (1988)