



Correlated equilibria and communication in games

Françoise Forges

► To cite this version:

Françoise Forges. Correlated equilibria and communication in games. Computational Complexity. Theory, Techniques, and Applications, pp.295-704, 2012, 10.1007/978-1-4614-1800-9_45. hal-01519895

HAL Id: hal-01519895

<https://hal.science/hal-01519895>

Submitted on 9 May 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Correlated Equilibrium and Communication in Games

Françoise Forges, CEREMADE, Université Paris-Dauphine

Article Outline

Glossary

I. Definition of the Subject and its Importance

II. Introduction

III. Correlated Equilibrium: Definition and Basic Properties

IV. Correlated Equilibrium and Communication

V. Correlated Equilibrium in Bayesian Games

VI. Related Topics and Future Directions

VII. Bibliography

Acknowledgements

The author wishes to thank Elchanan Ben-Porath, Frédéric Koessler, R. Vijay Krishna, Ehud Lehrer, Bob Nau, Indra Ray, Jérôme Renault, Eilon Solan, Sylvain Sorin, Bernhard von Stengel, Tristan Tomala, Amparo Urbano, Yannick Viossat and, especially, Olivier Gossner and Péter Vida, for useful comments and suggestions.

Glossary

Bayesian game: an interactive decision problem consisting of a set of n players, a set of types for every player, a probability distribution which accounts for the players' beliefs over each others' types, a set of actions for every player and a *von Neumann-Morgenstern utility function* defined over n -tuples of types and actions for every player.

Nash equilibrium: in an n -person strategic form game, a strategy n -tuple from which unilateral deviations are not profitable.

von Neumann-Morgenstern utility function: a *utility function* which reflects the individual's preferences over lotteries. Such a utility function is defined over outcomes and can be extended to any lottery λ by taking expectation with respect to λ .

Pure strategy (or simply strategy): a mapping which, in an interactive decision problem, associates an action with the information of a player whenever this player can make a choice.

Sequential equilibrium: a refinement of the *Nash equilibrium* for n -person multistage interactive decision problems, which can be loosely defined as a strategy n -tuple together with beliefs over past information for every player, such that every player maximizes his expected utility given his beliefs and the others' strategies, with the additional condition that the beliefs satisfy (possibly sophisticated) Bayes updating given the strategies.

Strategic (or normal) form game: an interactive decision problem consisting of a set of n players, a set of strategies for every player and a (typically, *von Neumann-Morgenstern*) *utility function* defined over n -tuples of strategies for every player.

Utility function: a real valued mapping over a set of outcomes which reflects the preferences of an individual by associating a utility level (a “pay-off”) with every outcome.

I. Definition of the Subject and its Importance

The correlated equilibrium is a game theoretic solution concept. It was proposed by Aumann (1974, 1987) in order to capture the strategic correlation opportunities that the players face when they take into account the extraneous environment in which they interact. The notion is illustrated in Section II. A formal definition is given in Section III. The correlated equilibrium also appears as the appropriate solution concept if preplay communication is allowed between the players. As shown in Section IV, this property can be given several precise statements according to the constraints imposed on the players' communication, which can go from plain conversation to exchange of messages through noisy channels. Originally designed for static games with complete information, the correlated equilibrium applies to any strategic form game. It is geometrically and computationally more tractable than the better known Nash equilibrium. The solution concept has been extended to dynamic games, possibly with incomplete information. As an illustration, we define in details the communication equilibrium for Bayesian games in Section V.

II. Introduction

Example

Consider the two-person game known as “chicken”, in which each player i can take a “pacific” action (denoted as p^i) or an “aggressive” action (denoted as a^i):

	p^2	a^2
p^1	(8, 8)	(3, 10)
a^1	(10, 3)	(0, 0)

The interpretation is that player 1 and player 2 simultaneously choose an action and then get a payoff, which is determined by the pair of chosen actions according to the previous matrix. If both players are pacific, they both get 8. If both are aggressive, they both get 0. If one player is aggressive and the other is pacific, the aggressive player gets 10 and the pacific one gets 3. This game has two pure Nash equilibria (p^1, a^2) , (a^1, p^2) and one mixed Nash equilibrium in which both players choose the pacific action with probability $\frac{3}{5}$, resulting in the expected payoff 6 for both players. A possible justification for the latter solution is that the players make their choices as a function of independent extraneous random signals. The assumption of independence is strong. Indeed, there may be no way to prevent the players’ signals from being correlated.

Consider a random signal which has no effect on the players’ payoffs and takes three possible values: low, medium or high, occurring each with probability $\frac{1}{3}$. Assume that, before the beginning of the game, player 1 distinguishes whether the signal is high or not, while player 2 distinguishes whether the signal is low or not. The relevant interactive decision problem is then the extended game in which the players can base their action on the private information they get on the random signal, while the payoffs only depend on the players’ actions. In this game, suppose that player 1 chooses the aggressive action when the signal is high and the pacific action otherwise. Similarly, suppose that player 2 chooses the aggressive action when the signal is low and the pacific action otherwise. We show that these strategies form an equilibrium in the extended game. Given player 2’s strategy, assume that player 1 observes a high signal. Player 1 deduces that the signal cannot be low so that player 2 chooses the pacific action; hence player 1’s best response is to play aggressively. Assume now that player 1 is informed that the signal is not high; he deduces that with probability $\frac{1}{2}$, the signal is medium (*i.e.*,

not low) so that player 2 plays pacific and with probability $\frac{1}{2}$, the signal is low so that player 2 plays aggressive. The expected payoff of player 1 is 5.5 if he plays pacific and 5 if he plays aggressive; hence, the pacific action is a best response. The equilibrium conditions for player 2 are symmetric. To sum up, the strategies based on the players' private information form a Nash equilibrium in the extended game in which an extraneous signal is first selected. We shall say that these strategies form a "correlated equilibrium". The corresponding probability distribution over the players' actions is

$$\begin{array}{ccc} & p^2 & a^2 \\ p^1 & \frac{1}{3} & \frac{1}{3} \\ a^1 & \frac{1}{3} & 0 \end{array} \quad (1)$$

and the expected payoff of every player is 7. This probability distribution can be used directly to make private recommendations to the players before the beginning of the game (see the canonical representation below).

III. Correlated Equilibrium: Definition and Basic Properties

Definition

A game in strategic form $G = (N, (\Sigma^i)_{i \in N}, (u^i)_{i \in N})$ consists of a set of players N together with, for every player $i \in N$, a set of strategies (for instance, a set of actions) Σ^i and a (von Neumann-Morgenstern) utility function $u^i : \Sigma \rightarrow \mathbb{R}$, where $\Sigma = \prod_{j \in N} \Sigma^j$ is the set of all strategy profiles. We assume that the sets N and Σ^i , $i \in N$, are finite.

A correlation device $d = (\Omega, q, (\mathcal{P}^i)_{i \in N})$ is described by a finite set of signals Ω , a probability distribution q over Ω and a partition \mathcal{P}^i of Ω for every player $i \in N$. Since Ω is finite, the probability distribution q is just a real vector $q = (q(\omega))_{\omega \in \Omega}$ such that $q(\omega) \geq 0$ and $\sum_{\omega \in \Omega} q(\omega) = 1$.

From G and d , we define the extended game G_d as follows:

- ω is chosen in Ω according to q
- every player i is informed of the element $P^i(\omega)$ of \mathcal{P}^i which contains ω
- G is played: every player i chooses a strategy σ^i in Σ^i and gets the utility $u^i(\sigma)$, $\sigma = (\sigma^j)_{j \in N}$.

A (pure) strategy for player i in G_d is a mapping $\alpha^i : \Omega \rightarrow \Sigma^i$ which is \mathcal{P}^i -measurable, namely, such that $\alpha^i(\omega') = \alpha^i(\omega)$ if $\omega' \in P^i(\omega)$. The interpretation is that, in G_d , every player i chooses his strategy σ^i as a function of his private information on the random signal ω which is selected before the beginning of G .

According to Aumann (1974), a *correlated equilibrium* of G is a pair (d, α) , which consists of a correlation device $d = (\Omega, q, (\mathcal{P}^i)_{i \in N})$ and a Nash equilibrium $\alpha = (\alpha^i)_{i \in N}$ of G_d . The equilibrium conditions of every player i , conditionally on his private information, can be written as:

$$\sum_{\omega' \in P^i(\omega)} q(\omega' | P^i(\omega)) u^i(\alpha(\omega')) \geq \sum_{\omega' \in P^i(\omega)} q(\omega' | P^i(\omega)) u^i(\tau^i, \alpha^{-i}(\omega')) \quad (2)$$

$$\forall i \in N, \forall \tau^i \in \Sigma^i, \forall \omega \in \Omega : q(\omega) > 0$$

where $\alpha^{-i} = (\alpha^j)_{j \neq i}$.

A mixed Nash equilibrium $\rho = (\rho^i)_{i \in N}$ of G can be viewed as a correlated equilibrium of G . By definition, every ρ^i is a probability distribution over Σ^i , the finite set of pure strategies of player i . Let us consider the correlation device $d = (\Omega, q, (\mathcal{P}^i)_{i \in N})$ in which $\Omega = \Sigma = \prod_{j \in N} \Sigma^j$, q is the product probability distribution induced by the mixed strategies (*i.e.*, $q((\sigma^j)_{j \in N}) = \prod_{j \in N} \rho^j(\sigma^j)$) and for each i , \mathcal{P}^i is the partition of Ω generated by Σ^i (*i.e.*, for $\omega, \nu \in \Omega$, $\nu \in P^i(\omega) \Leftrightarrow \nu^i = \omega^i$). Let $\alpha^i : \Sigma \rightarrow \Sigma^i$ be the projection over Σ^i (*i.e.*, $\alpha^i(\sigma) = \sigma^i$). The correlation device d and the strategies α^i defined in this way form a correlated equilibrium. As we shall see below, this correlated equilibrium is “canonical”.

Canonical representation

A *canonical* correlated equilibrium of G is a correlated equilibrium in which $\Omega = \Sigma = \prod_{j \in N} \Sigma^j$ while for every player i , the partition \mathcal{P}^i of Σ is generated by Σ^i and $\alpha^i : \Sigma \rightarrow \Sigma^i$ is the projection over Σ^i . A canonical correlated equilibrium is thus fully specified by a probability distribution q over Σ . A natural interpretation is that a mediator selects $\sigma = (\sigma^j)_{j \in N}$ according to q and privately recommends σ^i to player i , for every $i \in N$. The players are not forced to obey the mediator, but σ is selected in such a way that player i cannot benefit from deviating unilaterally from the recommendation σ^i , *i.e.*, $\tau^i = \sigma^i$ maximizes the conditional expectation of player i 's payoff $u^i(\tau^i, \sigma^{-i})$ given the recommendation σ^i . A probability distribution q over Σ

thus defines a canonical correlated equilibrium if and only if it satisfies the following linear inequalities:

$$\sum_{\sigma^{-i} \in \Sigma^{-i}} q(\sigma^{-i} | \sigma^i) u^i(\sigma^i, \sigma^{-i}) \geq \sum_{\sigma^{-i} \in \Sigma^{-i}} q(\sigma^{-i} | \sigma^i) u^i(\tau^i, \sigma^{-i})$$

$$\forall i \in N, \forall \sigma^i \in \Sigma^i : q(\sigma^i) > 0, \forall \tau^i \in \Sigma^i$$

or, equivalently,

$$\sum_{\sigma^{-i} \in \Sigma^{-i}} q(\sigma^i, \sigma^{-i}) u^i(\sigma^i, \sigma^{-i}) \geq \sum_{\sigma^{-i} \in \Sigma^{-i}} q(\sigma^i, \sigma^{-i}) u^i(\tau^i, \sigma^{-i})$$

$$\forall i \in N, \forall \sigma^i, \tau^i \in \Sigma^i \quad (3)$$

The equilibrium conditions can also be formulated *ex ante*:

$$\sum_{\sigma \in \Sigma} q(\sigma) u^i(\sigma) \geq \sum_{\sigma \in \Sigma} q(\sigma) u^i(\alpha^i(\sigma^i), \sigma^{-i})$$

$$\forall i \in N, \forall \alpha^i : \Sigma^i \rightarrow \Sigma^i$$

The following result is an analog of the “revelation principle” in mechanism design (see, e.g., Myerson (1982)): *let (α, d) be a correlated equilibrium associated with an arbitrary correlation device $d = (\Omega, q, (\mathcal{P}^i)_{i \in N})$. The corresponding “correlated equilibrium distribution”, namely, the probability distribution induced over Σ by q and α , defines a canonical correlated equilibrium.* For instance, in the introduction, (1) describes a canonical correlated equilibrium.

Duality and existence

From the linearity of (3), duality theory can be used to study the properties of correlated equilibria, in particular to prove their existence without relying on Nash (1951)’s theorem and its fixed point argument (recall that every mixed Nash equilibrium is a correlated equilibrium). Hart and Schmeidler (1989) establish the existence of a correlated equilibrium by constructing an auxiliary two person zero-sum game and applying the minimax theorem. Nau and McCardle (1990) derive another elementary proof of existence from an extension of the “no arbitrage opportunities” axiom that underlies subjective probability theory. They introduce jointly coherent strategy profiles, which do not expose the players as a group to arbitrage from an outside

observer. They show that a strategy profile is jointly coherent if and only if it occurs with positive probability in some correlated equilibrium. From a technical point of view, both proofs turn out to be similar. Myerson (1997) makes further use of the linear structure of correlated equilibria by introducing dual reduction, a technique to replace a finite game with a game with fewer strategies, in such a way that any correlated equilibrium of the reduced game induces a correlated equilibrium of the original game.

Geometric properties

As (3) is a system of linear inequalities, the set of all correlated equilibrium distributions is a convex polytope. Nau et al. (2004) show that if it has “full” dimension (namely, dimension $|\Sigma| - 1$), then all Nash equilibria lie on its relative boundary. Viossat (2006) characterizes in addition the class of games whose correlated equilibrium polytope contains a Nash equilibrium in its relative interior. Interestingly, this class of games includes two person zero-sum games but is not defined by “strict competition” properties. In two person games, all extreme Nash equilibria are also extreme correlated equilibria (Evangelista and Raghavan (1996), Gomez-Canovas et al. (1999)); this result does not hold with more than two players. Finally, Viossat (2005) proves that having a unique correlated equilibrium is a robust property, in the sense that the set of n person games with a unique correlated equilibrium is open. The same is not true for Nash equilibrium (unless $n = 2$).

Complexity

From (3), correlated equilibria can be computed by linear programming methods. Gilboa and Zemel (1989) show more precisely that the complexity of standard computational problems is “NP-hard” for the Nash equilibrium and *polynomial* for the correlated equilibrium. Examples of such problems are: “Does the game G have a Nash (resp., correlated) equilibrium which yields a payoff greater than r to every player (for some given number r)?” and “Does the game G have a unique Nash (resp., correlated) equilibrium?”. Papadimitriou (2005) develops a polynomial-time algorithm for finding correlated equilibria, which is based on a variant of the existence proof of Hart and Schmeidler (1989).

Foundations

By re-interpreting the previous canonical representation, Aumann (1987) proposes a decision theoretic foundation for the correlated equilibrium in

games with complete information, in which Σ^i for $i \in N$, stands merely for a set of *actions* of player i . Let Ω be the space of all states of the world; an element ω of Ω thus specifies all the parameters which may be relevant to the players' choices. In particular, the action profile in the underlying game G is part of the state of the world. A partition \mathcal{P}^i describes player i 's information on Ω . In addition, every player i has a prior belief, *i.e.*, a probability distribution, q^i over Ω . Formally, the framework is similar as above except that the players possibly hold different beliefs over Ω . Let $\alpha^i(\omega)$ denote player i 's action at ω ; a natural assumption is that player i knows the action he chooses, namely that α^i is \mathcal{P}^i -measurable. According to Aumann (1987), player i is *Bayes-rational* at ω if his action $\alpha^i(\omega)$ maximizes his expected payoff (with respect to q^i) given his information $P^i(\omega)$. Note that this is a separate rationality condition for every player, not an equilibrium condition. Aumann (1987) proves the following result: *Under the common prior assumption (namely, $q^i = q$, $i \in N$), if every player is Bayes-rational at every state of the world, the distribution of the corresponding action profile α is a correlated equilibrium distribution.* The key to this decision theoretic foundation of the correlated equilibrium is that, under the common prior assumption, Bayesian rationality amounts to (2).

If the common prior assumption is relaxed, the previous result still holds, with *subjective* prior probability distributions, for the *subjective* correlated equilibrium which was also introduced by Aumann (1974). The latter solution concept is defined in the same way as above, by considering a device $(\Omega, (q^i)_{i \in N}, (\mathcal{P}^i)_{i \in N})$, with a probability distribution q^i for every player i , and by writing (2) in terms of q^i instead of q . Brandenburger and Dekel (1987) show that (a refinement of) the subjective correlated equilibrium is equivalent to (correlated) rationalizability, another well-established solution concept which captures players' minimal rationality. Rationalizable strategies reflect that the players commonly know that each of them makes an optimal choice given some belief. Nau and McCardle (1991) reconcile objective and subjective correlated equilibrium by proposing the no arbitrage principle as a unified approach to individual and interactive decision problems. They argue that the objective correlated equilibrium concept applies to a game that is revealed by the players' choices, while the subjective correlated equilibrium concept applies to the "true game"; both lead to the same set of jointly coherent outcomes.

IV. Correlated Equilibrium and Communication

As seen in the previous section, correlated equilibria can be achieved in practice with the help of a mediator and emerge in a Bayesian framework embedding the game in a full description of the world. Both approaches require to extend the game by taking into account information which is not generated by the players themselves. Can the players reach a correlated equilibrium without relying on any extraneous correlation device, by just communicating with each other before the beginning of the game?

Consider the game of “chicken” presented in the introduction. The probability distribution

$$\begin{array}{cc} & \begin{matrix} p^2 & a^2 \end{matrix} \\ \begin{matrix} p^1 \\ a^1 \end{matrix} & \begin{pmatrix} 0 & \frac{1}{2} \\ \frac{1}{2} & 0 \end{pmatrix} \end{array} \quad (4)$$

describes a correlated equilibrium, which amounts to choosing one of the two pure Nash equilibria, with equal probability. Both players get an expected payoff of 6.5. Can they safely achieve this probability distribution if no mediator tosses a fair coin for them? The answer is positive, as shown by Aumann et al. (1968). Assume that before playing “chicken”, the players independently toss a coin and simultaneously reveal to each other whether heads or tails obtains. Player 1 tells player 2 “ h^1 ” or “ t^1 ” and, at the same time, player 2 tells player 1 “ h^2 ” or “ t^2 ”. If both players use a fair coin, reveal correctly the result of the toss and play (p^1, a^2) if both coins fell on the same side (*i.e.*, if (h^1, h^2) or (t^1, t^2) is announced) and (a^1, p^2) otherwise (*i.e.*, if (h^1, t^2) or (t^1, h^2) is announced), they get the same effect as a mediator using (4). Furthermore, none of them can gain by unilaterally deviating from the described strategies, even at the randomizing stage: the two relevant outcomes, $[(h^1, h^2) \text{ or } (t^1, t^2)]$ and $[(h^1, t^2) \text{ or } (t^1, h^2)]$, happen with probability $\frac{1}{2}$ provided that one of the players reveals the toss of a fair coin. This procedure is known as a “jointly controlled lottery”. An important feature of the previous example is that, in the correlated equilibrium described by (4), the players know each other’s recommendation. Hence, they can easily reproduce (4) by exchanging messages that they have selected independently. In the correlated equilibrium described by the probability distribution (1), the private character of recommendations is crucial to guarantee that (p^1, p^2) be played with positive probability. Hence one cannot hope that a simple procedure of direct preplay communication be sufficient to generate (1). However, the

fact that direct communication is necessarily public is typical of two-person games.

Given the game $G = (N, (\Sigma^i)_{i \in N}, (u^i)_{i \in N})$, let us define a (bounded) “cheap talk” extension $ext(G)$ of G as a game in which T stages of costless, unmediated preplay communication are allowed before G is played. More precisely, let M_t^i be a finite set of messages for player i , $i \in N$, at stage t , $t = 1, 2, \dots, T$; at every stage t of $ext(G)$, every player i selects a message in M_t^i ; these choices are made simultaneously before being revealed to a subset of players at the end of stage t . The rules of $ext(G)$ thus determine a set of “senders” for every stage t (those players i for whom M_t^i contains more than one message) and a set of “receivers” for every stage t . The players perfectly recall their past messages. After the communication phase, they choose their strategies (e.g., their actions) as in G ; they are also rewarded as in G , independently of the preplay phase, which is thus “cheap”. Communication has an indirect effect on the final outcome in G , since the players make their decisions as a function of the messages that they have exchanged. Specific additional assumptions are often made on $ext(G)$, as we will see below.

Let us fix a cheap talk extension $ext(G)$ of G and a Nash equilibrium of $ext(G)$. As a consequence of the previous definitions, the distribution induced by this Nash equilibrium over Σ defines a correlated equilibrium of G (this can be proved in the same way as the canonical representation of correlated equilibria stated in Section III). The question raised in this section is whether the reverse holds.

If the number of players is two, the Nash equilibrium distributions of cheap talk extensions of G form a subset of the correlated equilibrium distributions: the convex hull of Nash equilibrium distributions. Indeed, the players have both the same information after any direct exchange of messages. Conversely, by performing repeated jointly controlled lotteries like in the example above, the players can achieve any convex combination (with rational weights) of Nash equilibria of G as a Nash equilibrium of a cheap talk extension of G . The restriction on probability distributions whose components are rational numbers is only needed as far as we focus on *bounded* cheap talk extensions.

Bárány (1992) establishes that, if the number of players of G is at least four, every (rational) correlated equilibrium distribution of G can be realized as a Nash equilibrium of a cheap talk extension $ext(G)$, provided that $ext(G)$ allows the players to publicly check the record of communication under some circumstances. The equilibria of $ext(G)$ constructed by Bárány involve that

a receiver gets the same message from two different senders; the message is nevertheless not public thanks to the assumption on the number of players. At every stage of $ext(G)$, every player can ask for the revelation of all past messages, which are assumed to be recorded. Typically, a receiver can claim that the two senders' messages differ. In this case, the record of communication surely reveals that either one of the senders or the receiver himself has cheated; the deviator can be punished (at his minmax level in G) by the other players.

The punishments in Bárány (1992)'s Nash equilibria of $ext(G)$ need not be credible threats. Instead of using double senders in the communication protocols, Ben-Porath (1998, 2003) proposes a procedure of random monitoring, which prescribes a given behavior to every player in such a way that unilateral deviations can be detected with probability arbitrarily close to 1. This procedure applies if there are at least three players, which yields an analog of Bárány's result already in this case. If the number of players is exactly three, Ben-Porath (2003) needs to assume, as Bárány (1992), that public verification of the record of communication is possible in $ext(G)$ (see Ben-Porath (2006)). However, Ben-Porath concentrates on (rational) correlated equilibrium distributions which allow for strict punishment on a Nash equilibrium of G ; he constructs *sequential* equilibria which generate these distributions in $ext(G)$, thus dispensing with incredible threats. At the price of raising the number of players to five or more, Gerardi (2004) proves that every (rational) correlated equilibrium distribution of G can be realized as a sequential equilibrium of a cheap talk extension of G which does not require any message recording. For this, he builds protocols of communication in which the players base their decisions on majority rule, so that no punishment is necessary.

We have concentrated on two extreme forms of communication: mediated communication, in which a mediator performs lotteries and sends private messages to the players and cheap talk, in which the players just exchange messages. Many intermediate schemes of communication are obviously conceivable. For instance, Lehrer (1996) introduces (possibly multistage) "mediated talk": the players send private messages to a mediator, but the latter can only make deterministic public announcements. Mediated talk captures real-life communication procedures, like elections, especially if it lasts only for a few stages. Lehrer and Sorin (1997) establish that whatever the number of players of G , every (rational) correlated equilibrium distribution of G can be realized as a Nash equilibrium of a single stage mediated talk extension of

G. Ben-Porath (1998) proposes a variant of cheap talk in which the players do not only exchange verbal messages but also “hard” devices such as urns containing balls. This extension is particularly useful in two-person games to circumvent the equivalence between the equilibria achieved by cheap talk and the convex hull of Nash equilibria. More precisely, the result of Ben-Porath (1998) stated above holds for two-person games if the players first check together the content of different urns, and then each player draws a ball from an urn that was chosen by the other player, so as to guarantee that one player only knows the outcome of a lottery while the other one only knows the probabilities of this lottery.

The various extensions of the basic game G considered up to now, with or without a mediator, implicitly assume that the players are fully rational. In particular, they have unlimited computational abilities. By relaxing that assumption, Urbano and Vila (2002) and Dodis et al. (2000) build on earlier results from cryptography so as to implement any (rational) correlated equilibrium distribution through unmediated communication, including in two-person games.

As the previous paragraphs illustrate, the players can modify their initial distribution of information by means of many different communication protocols. Gossner (1998) proposes a general criterion to classify them: a protocol is “secure” if under all circumstances, the players cannot mislead each other nor spy on each other. For instance, given a cheap talk extension $ext(G)$, a protocol P describes, for every player, a strategy in $ext(G)$ and a way to interpret his information after the communication phase of $ext(G)$. P induces a correlation device $d(P)$ (in the sense of Section III). P is secure if, *for every game G* and every Nash equilibrium α of $G_{d(P)}$, the following procedure is a Nash equilibrium of $ext(G)$: communicate according to the strategies described by P in order to generate $d(P)$ and make the final choice, in G , according to α . Gossner (1998) gives a tractable characterization of secure protocols.

V. Correlated Equilibrium in Bayesian Games

A Bayesian game $\Gamma = (N, (T^i)_{i \in N}, p, (A^i)_{i \in N}, (v^i)_{i \in N})$ consists of: a set of players N ; for every player $i \in N$, a set of types T^i , a probability distribution p^i over $T = \prod_{j \in N} T^j$, a set of actions A^i and a (von Neumann-Morgenstern) utility function $v^i : T \times A \rightarrow \mathbb{R}$, where $A = \prod_{j \in N} A^j$. For simplicity, we make the common prior assumption: $p^i = p$ for every $i \in N$. All sets are assumed

finite. The interpretation is that a virtual move of nature chooses $t = (t^j)_{j \in N}$ according to p ; player i is only informed of his own type t^i ; the players then choose simultaneously an action. We will focus on two possible extensions of Aumann (1974)'s solution concept to Bayesian games: the strategic form correlated equilibrium and the communication equilibrium. Without loss of generality, the definitions below are given in “canonical form” (see Section III).

Strategic form correlated equilibrium

A (pure) strategy of player i in Γ is a mapping $\sigma^i : T^i \rightarrow A^i$, $i \in N$. The strategic form of Γ is a game $G(\Gamma)$, like the game G considered in Section III, with sets of pure strategies $\Sigma_i = A_i^{T_i}$ and utility functions u^i over $\Sigma = \prod_{j \in N} \Sigma^j$ computed as expectations with respect to p : $u^i(\sigma) = E[v^i(t, \sigma(t))]$, with $\sigma(t) = (\sigma^i(t^i))_{i \in N}$. A strategic form correlated equilibrium, or simply, a correlated equilibrium, of a Bayesian game Γ is a correlated equilibrium, in the sense of Section III, of $G(\Gamma)$. A canonical correlated equilibrium of Γ is thus described by a probability distribution Q over Σ , which selects an N -tuple of pure strategies $(\sigma^i)_{i \in N}$. This lottery can be thought of as being performed by a mediator who privately recommends σ^i to player i , $i \in N$, before the beginning of Γ , *i.e.*, before (or in any case, independently of) the chance move choosing the N -tuple of types. The equilibrium conditions express that, once he knows his type t^i , player i cannot gain in unilaterally deviating from $\sigma^i(t^i)$.

Communication equilibrium

Myerson (1982) transforms the Bayesian game Γ into a mechanism design problem by allowing the mediator to collect information from the players before making them recommendations. Following Forges (1986) and Myerson (1986a), a canonical communication device for Γ consists of a system q of probability distributions $q = (q(\cdot|t))_{t \in T}$ over A . The interpretation is that a mediator invites every player i , $i \in N$, to report his type t^i , then selects an N -tuple of actions a according to $q(\cdot|t)$ and privately recommends a^i to player i . The system q defines a communication equilibrium if none of the players can gain by unilaterally lying on his type or by deviating from the recommended action, namely if

$$\sum_{t^{-i} \in T^{-i}} p(t^{-i}|t^i) \sum_{a \in A} q(a|t) v^i(t, a) \geq \sum_{t^{-i} \in T^{-i}} p(t^{-i}|t^i) \sum_{a \in A} q(a|s^i, t^{-i}) v^i(t, \alpha^i(a^i), a^{-i})$$

$$\forall i \in N, \forall t^i, s^i \in T^i, \forall \alpha^i : A^i \rightarrow A^i$$

Correlated equilibrium, communication equilibrium and cheap talk

Every correlated equilibrium of the Bayesian game Γ induces a communication equilibrium of Γ , but the converse is not true, as the following example shows.

Consider the two-person Bayesian game in which $T^1 = \{s^1, t^1\}$, $T^2 = \{t^2\}$, $A^1 = \{a^1, b^1\}$, $A^2 = \{a^2, b^2\}$, $p(s^1) = p(t^1) = \frac{1}{2}$ and payoffs are described by

		a^2	b^2
s^1	a^1	(1, 1)	(-1, -1)
	b^1	(0, 0)	(0, 0)
		a^2	b^2
t^1	a^1	(0, 0)	(0, 0)
	b^1	(-1, -1)	(1, 1)

In this game, the communication equilibrium $q(a^1, a^2|s^1) = q(b^1, b^2|t^1) = 1$ yields the expected payoff of 1 to both players. However the maximal expected payoff of every player in a correlated equilibrium is $\frac{1}{2}$. In order to see this, one can derive the strategic form of the game (in which player 1 has four strategies and player 2 has two strategies). Let us turn to the game in which player 1 can cheaply talk to player 2 just after having learned his type. In this new game, the following strategies form a Nash equilibrium: player 1 truthfully reveals his type to player 2 and plays a^1 if s^1 , b^1 if t^1 ; player 2 chooses a^2 if s^1 , b^2 if t^1 . These strategies achieve the same expected payoffs as the communication equilibrium.

As in Section IV, one can define cheap talk extensions $ext(\Gamma)$ of Γ . A wide definition of $ext(\Gamma)$ involves an *ex ante* preplay phase, before the players learn their types, and an *interim* preplay phase, after the players learn their types but before they choose their actions. Every Nash equilibrium of $ext(\Gamma)$ induces a communication equilibrium of Γ . In order to investigate the converse, namely whether cheap talk can simulate mediated communication in a Bayesian game, two approaches have been developed. The first one

(Forges (1990), Gerardi (2000, 2004), Vida (2007)) proceeds in two steps, by reducing communication equilibria to correlated equilibria before applying the results obtained for strategic form games (see Section IV). The second approach (Ben-Porath (2003), Krishna (2007)) directly addresses the question in a Bayesian game.

By developing a construction introduced for particular two person games (Forges (1985)), Forges (1990) shows that every communication equilibrium outcome of a Bayesian game Γ with at least four players can be achieved as a correlated equilibrium outcome of a two stage *interim* cheap talk extension $ext_{int}(\Gamma)$ of Γ . No punishment is necessary in $ext_{int}(\Gamma)$: at the second stage, every player gets a message from three senders and uses majority rule if the messages are not identical. Thanks to the underlying correlation device, each receiver is able to privately decode his message. Vida (2007) extends Forges (1990) to Bayesian games with three or even two players. In the proof, he constructs a correlated equilibrium of a long, but almost surely finite, *interim* cheap talk extension of Γ , whose length depends both on the signals selected by the correlation device and the messages exchanged by the players. No recording of messages is necessary to detect and punish a cheating player. If there are at least four players in Γ , once a communication equilibrium of Γ has been converted into a correlated equilibrium of $ext_{int}(\Gamma)$, one can apply Bárány (1992)'s result to $ext_{int}(\Gamma)$ in order to transform the correlated equilibrium into a Nash equilibrium of a further, *ex ante*, cheap talk preplay extension of Γ . Gerardi (2000) modifies this *ex ante* preplay phase so as to postpone it at the *interim* stage. This result is especially useful if the initial move of nature in Γ is just a modelling convenience. Gerardi (2004) also extends his result for at least five person games with complete information (see Section IV) to any Bayesian game with full support (*i.e.*, in which all type profiles have positive probability: $p(t) > 0$ for every $t \in T$) by proving that every (rational) communication equilibrium of Γ can be achieved as a *sequential* equilibrium of a cheap talk extension of Γ .

Ben-Porath (2003) establishes that if Γ is a three (or more) person game with full support, every (rational) communication equilibrium of Γ which strictly dominates a Nash equilibrium of Γ for every type t^i of every player i , $i \in N$, can be implemented as a Nash equilibrium of an *interim* cheap talk extension of Γ in which public verification of past record is possible (see also Ben-Porath (2006)). Krishna (2007) extends Ben-Porath (1998)'s result on two person games (see Section IV) to the incomplete information framework. The other results mentioned at the end of Section IV have also

been generalized to Bayesian games (see Gossner (1998), Lehrer and Sorin (1997) and Urbano and Vila (2004a)).

VI. Related Topics and Future Directions

In this brief article, we concentrated on two solution concepts: the strategic form correlated equilibrium, which is applicable to any game, and the communication equilibrium, which we defined for Bayesian games. Other extensions of Aumann (1974)'s solution concept have been proposed for Bayesian games, as the agent normal form correlated equilibrium and the (possibly belief invariant) Bayesian solution (see Forges (1993, 2006) for definitions and references). The Bayesian solution is intended to capture the players' rationality in games with incomplete information in the spirit of Aumann (1987) (see Nau (1992) and Forges (1993)). Lehrer et al. (2006) open a new perspective in the understanding of the Bayesian solution and other equilibrium concepts for Bayesian games by characterizing the classes of equivalent information structures with respect to each of them. Comparison of information structures, which goes back to Blackwell (1951, 1953) for individual decision problems, was introduced by Gossner (2000) in the context of games, both with complete and incomplete information. In the latter model, information structures basically describe how extraneous signals are selected as a function of the players' types; two information structures are equivalent with respect to an equilibrium concept if, in every game, they generate the same equilibrium distributions over outcomes.

Correlated equilibria, communication equilibria and related solution concepts have been studied in many other classes of games, like multistage games (see, e.g., Forges (1986), Myerson (1986a)), repeated games with incomplete information (see, e.g., Forges (1985, 1988)) and stochastic games (see, e.g., Solan (2001), Solan and Vieille (2002)). The study of correlated equilibrium in repeated games with imperfect monitoring, initiated by Lehrer (1991, 1992), proved to be particularly useful and is still undergoing. Lehrer (1991) showed that if players are either fully informed of past actions or get no information ("standard-trivial" information structure), correlated equilibria are equivalent to Nash equilibria. In other words, all correlations can be generated internally, namely by the past histories, on which players have differential information. The schemes of internal correlation introduced to establish this result are widely applicable and inspired those of Lehrer (1996) (see Section IV). In general repeated games with imperfect monitoring, Renault and Tomala (2004) characterize communication equilibria but the amount of

correlation that the players can achieve in a Nash equilibrium is still an open problem (see, e.g., Gossner and Tomala (2007) and Urbano and Vila (2004b) for recent advances).

Throughout this article, we defined a correlated equilibrium as a *Nash* equilibrium of an extension of the game under consideration. The solution concept can be strengthened by imposing some refinement, *i.e.*, further rationality conditions, to the Nash equilibrium in this definition (see, e.g., Dhillon and Mertens (1996), Myerson (1986b)). Refinements of communication equilibria have also been proposed (see, e.g., Myerson (1986a), Gerardi (2004), Gerardi and Myerson (2007)). Some authors (see, e.g., Milgrom and Roberts (1996), Moreno and Wooders (1996), Ray (1996)) have also developed notions of *coalition proof* correlated equilibria, which resist not only to unilateral deviations, as in this article, but even to multilateral ones. A recurrent difficulty is that, for many of these stronger solution concepts, a useful canonical representation (as derived in Section III) is not available.

Except for two or three references, we deliberately concentrated on the results published in the game theory and mathematical economics literature, while substantial achievements in computer science would fit in this survey. Both streams of research pursue similar goals but rely on different formalisms and techniques. For instance, computer scientists often make use of cryptographic tools which are not familiar in game theory. Halpern (2007) gives an idea of recent developments at the interface of computer science and game theory (see in particular the section “implementing mediators”) and contains a number of references.

Finally, the assumption of full rationality of the players can also be relaxed. Evolutionary game theory has developed models of learning in order to study the long term behavior of players with bounded rationality. Many possible dynamics are conceivable to represent more or less myopic attitudes with respect to optimization. Under appropriate learning procedures, which express for instance that agents want to minimize the regret of their strategic choices, the empirical distribution of actions converge to correlated equilibrium distributions (see, e.g., Foster and Vohra (1997), Hart and Mas Colell (2000) and Hart (2005) for a survey). However, standard procedures, as the “replicator dynamics”, may even eliminate all the strategies which have positive probability in a correlated equilibrium (see Viossat (2007)).

VII. Bibliography

Primary literature

1. Aumann, R. J. (1974), Subjectivity and correlation in randomized strategies, *Journal of Mathematical Economics* 1, 67-96.
2. Aumann, R. J. (1987), Correlated equilibrium as an expression of Bayesian rationality, *Econometrica* 55, 1-18.
3. Aumann, R. J., Maschler, M. and Stearns, R. (1968), Repeated games with incomplete information: an approach to the nonzero sum case, Reports to the U.S. Arms Control and Disarmament Agency, ST-143, Chapter IV, 117-216 (reprinted in Aumann, R. J. and Maschler M. (1995), *Repeated Games of Incomplete Information*, Cambridge: M.I.T. Press)
4. Bárány, I. (1992), Fair distribution protocols or how players replace fortune, *Mathematics of Operations Research* 17, 327-340.
5. Ben-Porath, E. (1998), Correlation without mediation: expanding the set of equilibrium outcomes by cheap pre-play procedures, *Journal of Economic Theory* 80, 108-122.
6. Ben-Porath, E. (2003), Cheap talk in games with incomplete information, *Journal of Economic Theory* 108, 45-71.
7. Ben-Porath, E. (2006), A correction to “Cheap talk in games with incomplete information”, mimeo, Hebrew University of Jerusalem.
8. Blackwell, D. (1951), Comparison of experiments, *Proceedings of the Second Berkeley Symposium on Mathematical Statistics and Probability*, University of California Press, 93-102.
9. Blackwell, D. (1953), Equivalent comparison of experiments, *Annals of Mathematical Statistics* 24, 265-272.
10. Brandenburger, A. and Dekel, E. (1987), Rationalizability and correlated equilibria, *Econometrica* 55, 1391-1402.
11. Dhillon, A. and Mertens, J. F. (1996), Perfect correlated equilibria, *Journal of Economic Theory* 68, 279-302.

12. Dodis, Y., Halevi, S. and Rabin, T. (2000), A cryptographic solution to a game theoretic problem, CRYPTO 2000: 20th International Cryptology Conference, Springer-Verlag, 112-130.
13. Evangelista, F. and Raghavan, T.E.S. (1996), A note on correlated equilibrium, International Journal of Game Theory 25, 35-41.
14. Forges, F. (1985), Correlated equilibria in a class of repeated games with incomplete information, International Journal of Game Theory 14, 129-150.
15. Forges, F. (1986), An approach to communication equilibrium, Econometrica 54, 1375-1385.
16. Forges, F. (1988), Communication equilibria in repeated games with incomplete information, Mathematics of Operations Research, 13, 191-231.
17. Forges, F. (1990), Universal mechanisms, Econometrica 58, 1341-1364.
18. Forges, F. (1993), Five legitimate definitions of correlated equilibrium in games with incomplete information, Theory and Decision 35, 277-310.
19. Forges, F. (2006), Correlated equilibrium in games with incomplete information revisited, Theory and Decision, 61, 329-344.
20. Foster, D. and Vohra, R. (1997), Calibrated learning and correlated equilibrium, Games and Economic Behavior 21, 40-55.
21. Gerardi, D. (2000), Interim pre-play communication, mimeo, Yale University.
22. Gerardi, D. (2004), Unmediated communication in games with complete and incomplete information, Journal of Economic Theory 114, 104-131.
23. Gerardi, D. and Myerson, R. (2007), Sequential equilibria in Bayesian games with communication, Games and Economic Behavior 60, 104-134.

24. Gilboa, I. and Zemel, E. (1989), Nash and correlated equilibria: some complexity considerations, *Games and Economic Behavior* 1, 80-93.
25. Gomez-Canovas S., Hansen P. and Jaumard B. (1999), Nash Equilibria from the correlated equilibria viewpoint, *International Game Theory Review* 1, 33-44.
26. Gossner, O. (1998), Secure protocols or how communication generates correlation, *Journal of Economic Theory* 83, 69-89.
27. Gossner, O. (2000), Comparison of information structures, *Games and Economic Behavior* 30, 44-63.
28. Gossner, O. and Tomala, T. (2007), Secret correlation in repeated games with signals, *Mathematics of Operations Research* 32, 413-424.
29. Hart, S. and Schmeidler, D. (1989), Existence of correlated equilibria, *Mathematics of Operations Research* 14, 18-25.
30. Hart, S. and Mas-Colell, A. (2000), A simple adaptive procedure leading to correlated equilibrium, *Econometrica* 68, 1127-1150.
31. Krishna, R. V. (2007), Communication in games of incomplete information: two players, *Journal of Economic Theory* 132, 584-592.
32. Lehrer, E. (1991), Internal correlation in repeated games, *International Journal of Game Theory* 19, 431-456.
33. Lehrer, E. (1992), Correlated equilibria in two-player repeated games with non-observable actions, *Mathematics of Operations Research* 17, 175-199.
34. Lehrer, E. (1996), Mediated talk, *International Journal of Game Theory* 25, 177-188.
35. Lehrer, E. and Sorin, S. (1997), One-shot public mediated talk, *Games and Economic Behavior* 20, 131-148.
36. Lehrer, E., Rosenberg, D. and Shmaya, E. (2006), Signaling and mediation in Bayesian games, mimeo, Tel Aviv University.

37. Milgrom, P., and Roberts, J. (1996), Coalition-proofness and correlation with arbitrary communication possibilities, *Games and Economic Behavior* 17, 113-128.
38. Moreno, D., and Wooders, J. (1996), Coalition-proof equilibrium, *Games and Economic Behavior* 17, 80-112.
39. Myerson, R. (1982), Optimal coordination mechanisms in generalized principal-agent problems, *Journal of Mathematical Economics*, 10, 67-81.
40. Myerson, R. (1986a), Multistage games with communication, *Econometrica* 54, 323-358.
41. Myerson, R. (1986b), Acceptable and predominant correlated equilibria, *International Journal of Game Theory* 15, 133-154.
42. Myerson, R. (1997), Dual reduction and elementary games, *Games and Economic Behavior* 21, 183-202.
43. Nash, J. (1951), Non-cooperative games, *Annals of Mathematics* 54, 286-295.
44. Nau, R.F. (1992), Joint coherence in games with incomplete information, *Management Science* 38, 374-387.
45. Nau, R.F. and McCardle, K.F. (1990), Coherent behavior in noncooperative games, *Journal of Economic Theory* 50, 2, 424-444.
46. Nau, R.F. and McCardle, K.F. (1991), Arbitrage, rationality and equilibrium, *Theory and Decision* 31, 199-240.
47. Nau, R. F., Gomez-Canovas, S. and Hansen, P. (2004), On the geometry of Nash equilibria and correlated equilibria, *International Journal of Game Theory* 32, 443-453.
48. Papadimitriou, C. H. (2005), Computing correlated equilibria in multiplayer games, *Proceedings of the 37th ACM Symposium on Theory of Computing*, 49-56.
49. Ray, I. (1996), Coalition-proof correlated equilibrium: a definition, *Games and Economic Behavior* 17, 56-79.

50. Renault, J. and Tomala, T. (2004), Communication equilibrium payoffs in repeated games with imperfect monitoring, *Games and Economic Behavior* 49, 313-344.
51. Solan, E. (2001), Characterization of correlated equilibrium in stochastic games, *International Journal of Game Theory*, 30, 259-277.
52. Solan, E. and N. Vieille (2002), Correlated equilibrium in stochastic games, *Games and Economic Behavior*, 38, 362-399.
53. Urbano, A. and Vila, J. (2002), Computational complexity and communication: coordination in two-player games, *Econometrica* 70, 1893-1927.
54. Urbano, A. and Vila, J. (2004a), Computationally restricted unmediated talk under incomplete information, *Economic Theory*, 23, 283-320.
55. Urbano, A. and Vila, J. (2004b), Unmediated communication in repeated games with imperfect monitoring, *Games and Economic Behavior*, 46, 143-173.
56. Vida, P. (2007), From communication equilibria to correlated equilibria, mimeo, University of Vienna.
57. Viossat, Y. (2005), Is having a unique equilibrium robust? to appear in *Journal of Mathematical Economics*.
58. Viossat, Y. (2006), The geometry of Nash equilibria and correlated equilibria and a generalization of zero-sum games, mimeo, S-WoPEc working paper 641, Stockholm School of Economics.
59. Viossat, Y. (2007), The replicator dynamics does not lead to correlated equilibria, *Games and Economic Behavior* 59, 397-407.

Reviews

1. Forges, F. (1994), Non-zero sum repeated games and information transmission, in: Megiddo, N., ed., *Essays in Game Theory in Honor of Michael Maschler*, Berlin: Springer, 65-95.

2. Halpern, J. (2007), Computer science and game theory: a brief survey, mimeo, Cornell University, to appear in the New palgrave Dictionary of Economics.
3. Hart, S. (2005), Adaptative heuristics, *Econometrica* 73, 1401-1430.
4. Mertens, J.-F. (1994), Correlated- and communication equilibria, in Mertens, J.-F and Sorin, S., eds., *Game Theoretic Methods in General Equilibrium Analysis*, Dordrecht: Kluwer, 243-248.
5. Myerson, R. (1985), Bayesian equilibrium and incentive compatibility, in Hurwicz, L. Schmeidler D. and Sonnenschein, H., eds., *Social Goals and Social Organization*, Cambridge: Cambridge University Press, 229-259.
6. Myerson, R. (1994), Communication, correlated equilibria and incentive compatibility, in Aumann, R. and Hart S., eds., *Handbook of Game Theory*, Volume 2, Amsterdam: Elsevier, 827-847.
7. Sorin, S. (1997), Communication, correlation and cooperation, in Mas Colell, A. and Hart, S., eds., *Cooperation: Game Theoretic Approaches*, Berlin: Springer, 198-218.