A simple allometric model for estimating blueberry fruit weight from diameter measurements

Emilio Jorquera-Fontena, Michel Génard, Alejandra Ribera-Fonseca, Nicolas Franck

► To cite this version:

Emilio Jorquera-Fontena, Michel Génard, Alejandra Ribera-Fonseca, Nicolas Franck. A simple allometric model for estimating blueberry fruit weight from diameter measurements. *Scientia Horticulturae*, 2017, 219, pp.131 - 134. 10.1016/j.scienta.2017.03.009 . hal-01521538

HAL Id: hal-01521538

<https://hal.science/hal-01521538>

Submitted on 26 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution - ShareAlike| 4.0 International License

Short Communication

A simple allometric model for estimating blueberry fruit weight from diameter measurements

Emilio Jorquera-Fontena¹, Michel Génard², Alejandra Ribera-Fonseca^{3,4*}, Nicolás Franck⁵

¹Facultad de Ciencias Agropecuarias y Forestales, Universidad de La Frontera, Avenida Francisco Salazar 01145, Casilla 54-D, Temuco, Chile. emilio.jorquera@ufrontera.cl

²UR 1115 Plantes et Systèmes de culture Horticoles, Institut National de la Recherche Agronomique, Avignon, France. michel.genard@avignon.inra.fr

³Departamento de Producción Agropecuaria, Facultad de Ciencias Agropecuarias y Forestales, Universidad de La Frontera, Casilla 54-D, Temuco, Chile. alejandra.ribera@ufrontera.cl

⁴Center of Plant-Soil Interaction and Natural Resources Biotechnology, Scientific and Technological Bioresources Nucleus (BIOREN-UFRO), Universidad de La Frontera, Casilla 54-D, Temuco, Chile.

⁵Centro de Estudios de Zonas Áridas, Facultad de Ciencias Agronómicas, Universidad de Chile, Casilla 129, Coquimbo, Chile. nfranck@uchile.cl

*Corresponding author:

Alejandra Ribera Fonseca

Phone: +56 45 2734123/2325669/2325458

E-mail address: alejandra.ribera@ufrontera.cl

Abstract. Monitoring of fruit growth is a measurement widely used in physiology and agronomy studies. This is normally done from detached fruits, which can lead to erroneous results when fruits grow asynchronously in clusters as occurs in blueberry plants (*Vaccinium spp.*). The aim of this communication is to develop a simple allometric model for estimating blueberry weight from diameter measurements. In three growth seasons, weight and diameter of a total of 416 fruits were measured from four field-grown blueberry cultivars. The obtained data set was then randomly divided resulting in 200 data for calibrating model and 216 for validating. Prior to calibration procedure, the use of cultivar-specific models or a common model for four cultivars was evaluated by Analysis of Covariance between logarithm transformed diameter and cultivar/season variables. From analysis, no interactive effect was observed, so a common allometric model was constructed, giving a power functional relationship between berry diameter and weight. The allometric model was validated with the independent data, giving errors ranging from 4.7% to 6.9%. Residuals showed a random dispersion pattern. Our results showed that a common model was adequate for providing a reliable means of estimating blueberry fruit weight of different cultivars.

Key words: Regression, validation, non-destructive methods

1. Introduction

Monitoring of fruit growth is a measurement widely used in studies of physiology and evaluation of management practices (e.g. Godoy et al., 2008; Minchin et al., 2010). Thus, its knowledge has been an important parameter in understanding source-sink relationships, water and nutrient use efficiency, crop quality and yield potential (Génard and Lescourret, 2004, Grossman and DeJong, 1995, Spreer et al., 2009). Habitually, fruit growth curves are developed from detached samples; however, this method can increase variance and lead to erroneous results for species in which fruits grow in clusters (Coombe, 1976). This is because several development stages of the fruit can be observed on a plant at a given time, as occur typically in several fruits species including blueberry (e.g. Tuccio et al., 2011; Godoy et al., 2008). A common method for non-destructively estimating biomass of plant organs is to fit simple equations to data relating biomass to some dimensional characteristic of the plant such as diameter, height, length and width (e.g. Antunes et al., 2008; Roxburgh et al., 2015). In this line, estimating fruit weight from equations using simple measurements of fruit diameter can represent an inexpensive and rapid alternative for accurately assessing fruit weight. In blueberry industry, fruit segregation on the basis of weight and size, is crucial to obtain higher export market value. The aim of this communication is to develop a simple allometric model for estimating blueberry weight from diameter measurements.

2. Materials and methods

2.1. Fruit data

In three growth seasons, a total of 416 healthy berries from three northern highbush blueberry cultivars (*V. corymbosum* cultivars 'Bluegold', 'Brigitta' and 'Elliot') and one southern blueberry cultivar (*V. corymbosum* x *Vaccinium darrowi* Camp. cultivar 'Legacy') were collected from three commercial orchards located in the Región de La Araucanía, Chile (Table 1).

The cultivars used differing in fruiting season; Bluegold and Elliot are early- and late-season cultivars, respectively, whereas Legacy and Brigitta are mid-season cultivars. Plants of the orchards ranged from three to five-year-old and were established at a spacing of $3 \times 0.9\text{m}$, in north–south oriented rows on an Andisol. In each orchard, the average fertilization was 70 kg N ha^{-1} , $40 \text{ kg P}_2\text{O}_5 \text{ ha}^{-1}$ and $75 \text{ kg K}_2\text{O ha}^{-1}$, the irrigation was applied as needed, and pests and diseases were controlled according to the locally recommended practices. Winter pruning was annually done in each orchard, removing old wood, damaged canes, unproductive branches, and short shoots that develop from the base of the plant.

Fruits were weekly picked from fruit coloring phase (stage in which berries are changing from green to pink to blue in the cluster) up to beginning of commercial harvest for each season. After picking, samples were placed in a cooler and then rapidly transported to the laboratory for evaluating equatorial diameter and fresh weight by using digital caliper (Mitutoyo, Japan, accuracy $\pm 0.01\text{mm}$) and technical balance (Gibertini Elettronica S.r.l., Italy), respectively.

The data set was randomly divided to obtain data for both calibrating ($n= 200$) and validating ($n= 216$) model (external validation).

2.2. Allometric model calibration

Preliminary regression analysis showed that the relationships between fruit weight and diameter followed a power fit typical of allometric models, for all situations in which fruit was obtained (Eq. 1):

$$FW = aFD^b \quad (\text{Eq. 1})$$

where FW is berry weight, FD is berry diameter, a and b are allometric coefficients.

In order to discriminate the use of cultivar-specific models (including the season effect) or a common model for four cultivars, an Analysis of Covariance was performed on data linearized by natural logarithm transformation (Eq. 2)

$$\ln FW = \ln a + b \times \ln FD \quad (\text{Eq. 2})$$

Thus, when interaction between logarithm transformed diameter (continuous variable) and cultivar/season (categorical variable) is not significant, the slopes among groups are similar, so a common model for four cultivars can be constructed (Warton and Weber, 2002; Chen et al., 2010).

2.3. Model goodness-of-fit and validation

Model results were compared with independent data through graphical procedures. The Relative Root Mean Squared Error (RRMSE, Eq. 3), a common criterion to quantify the mean difference between estimated data and observed data (Kobayashi and Us Salam, 2000) was calculated:

$$RRMSE = \left(\sqrt{\frac{\sum n_i(x_i - y_i)^2}{N}} \right) / \bar{y} \quad (\text{Eq. 3})$$

where n_i , x_i , and y_i are the number of observed data, the estimated data, and the observed data at the measurement date i , respectively, N is the total number of observed data and $\bar{y}$ the mean of observed values. The smaller the value of RRMSE, the closer to the measurement the estimation is (Grechi et al., 2008).

Model residuals were also calculated and plotted against data estimated by the model to check their normal distribution (i.e. homoscedastic behaviour).

2.4. Data analysis

In order to characterize the data set for calibrating model, results obtained from cultivars/season were subjected to Multiple Comparison of Mean (MCM) by using Scheffé's test ($\alpha = 0.05$). This test was selected because the sample size among groups was different (Ruxton and Beauchamp, 2008). The MSM, analysis of covariance and model fitting were performed with R software version 2.15.0 (R Development Core Team, 2012).

3. Results

3.1. Berry diameter and weight

The mean fruit diameter and weight from different cultivars/seasons ranged from 13.35 to 17.75 mm and from 1.19 to 2.58 g, respectively (Table 2). As observed, the highest values were measured for cv. Legacy in the season 2015-2016, while the lowest values for cv. Brigitta. When comparing cultivars, the mean diameter and weight of the fruit was significantly higher in Legacy and Bluegold than in Elliot and Brigitta (Table 2).

3.2. Evaluation of regression slopes and allometric model

Diameter and weight of the fruits were regressed on logarithm transformed data in order to evaluate whether the interactive effect of berry diameter (continuous variable) and cultivar/season (categorical variable) is significant for estimating berry weight (Table 3). Regression slopes of logarithm transformed data were no statistically different among cultivars/season, thereafter whole data set for calibration procedure was used for constructing a common model for four cultivars. Then, fruit weight data were plotted against fruit diameter data, giving a power functional relationship as shown in the Fig. 1. The relation between these variables was highly significant with p -values of the coefficients lower than 2×10^{-16} .

3.3. Model goodness-of-fit and validation

For model validation, estimated versus actual berry weights were plotted (Fig. 2). As observed, the values were closely distributed alongside the 1:1 line, although there was some tendency to overestimate the weight, which can be noticed on regression slope, being slightly higher than one (Fig. 2A). The external validation of the model showed that the model goodness-of-fit (RRMSE) decreased from cv. Bluegold (4.7%) to cv. Elliot (6.9%), with cv. Legacy and Brigitta showing

intermediate values (Fig. 2A). Residual were randomly distributed along estimated data, which indicates absence of heteroscedasticity (Fig. 2B).

4. Discussion

Empirical models are required for the non-destructive estimation of biomass of a plant organ of interest from easily obtainable measurements. These models, which are based on allometric theory, use the statistic relations between a variable of interest and one or several variables used as predictor (Niklas, 1994).

Many abiotic and biotic factors may potentially affect allometric relationships, including genetics, ontogeny, size, age, structure, site, climate, and their interactions (Misle et al., 2013; Niklas, 1994), however our results indicated that the relationship between fresh weight and diameter was sufficiently stable irrespectively to the genotype and growth seasons (Table 3). From this, a common allometric model was calibrated, giving a power functional relationship as showed in the Fig. 1. On the other hand, according to the principle of parsimony and the objective of proposing easy-to-use models, this finding was convenient. In the model, as a fruit grows, its dimensions change in ways that maintain their functional balance, but not in the ways that maintain constant ratios between dimensions (Castellan-Estrada et al., 2002). Similar allometric relationships have been reported in studies of fruit physiology. In most of these, fruit diameter was included as predictor of fresh and dry weight (e.g. Gautier et al., 2001; Gilbert et al., 2005; Díaz-Pérez et al., 2007; Grechi et al., 2008). Our allometric model was validated with independent data, giving higher goodness-of-fit for evaluated cultivars (Fig. 2A). This showed that a common model was adequate for providing a reliable means of estimating blueberry fruit weight of different cultivars. On the other hand, residuals follow a normal distribution, with data randomly distributed, which indicates absence of heteroscedasticity (Fig. 2B). When the dispersion pattern of residuals follows a

heteroscedastic behavior, the estimated standard errors of the regression coefficients are often underestimated, giving an incorrect perception of accuracy (Chatterjee and Hadi, 2006).

In conclusion, blueberry fruit weight was satisfactorily estimated on the basis of diameter measurements via allometric approach. Despite that model showed to be reliable for estimating berry weight of four blueberry cultivars, it must be validated when applied to other genetic materials. Further studies must be done in blueberry fruits to establish allometric equations for relating other fruit quality traits to easily obtainable non-destructive measurements

5. Acknowledgments

The first author acknowledges to the Convenio Marco 2016, Ministerio de Educación, Chile – Universidad de La Frontera, Chile.

6. References

- Antunes W.C., Pompelli M.F., Carretero D.M., DaMatta F.M. 2008. Allometric models for non-destructive leaf area estimation in coffee (*Coffea arabica* and *C.canephora*). *Annals of Applied Biology*, 153: 33–40.
- Castelan-Estrada M., Vivin P. Gaudillere J.P. 2002. Allometric relationships to estimate seasonal above ground vegetative and reproductive biomass of *Vitis vinifera* L. *Annals of Botany*, 89: 401-408.
- Chatterjee S., Hadi A.S. 2006. *Regression analysis by example*. Hoboken, NJ, USA, Wiley.
- Chen H., Felker S., Sun S. 2010. Allometry of within-fruit reproductive allocation in subtropical dicot woody species. *American Journal of Botany*, 97: 611–619.
- Coombe B.G. 1976. The development of fleshy fruits. *Annual Review of Plant Physiology*, 27: 207–228.

- 206 Díaz-Pérez J.C., Muy-Rangel M.D., Mascorro A.G. 2007. Fruit size and stage of ripeness affect
 207 postharvest water loss in bell pepper fruit (*Capsicum annuum* L.). Journal of Science and Food
 208 Agriculture, 87:68–73.
- 209 Kobayashi K., Us Salam M. 2000. Comparing simulated and measured values using mean squared
 210 deviation and its components. Agronomy Journal, 92:345–352.
- 211 Gautier H., Guichard S., Tchamitchian M. 2001. Modulation of competition between fruits and
 212 leaves by flower pruning and water fogging, and consequences on tomato leaf and fruit growth.
 213 Annals of Botany, 88: 645–652.
- 214 Gilbert C., Lescourret F., Génard M., Vercambre G., Pérez Pastor A. 2005. Modelling the effect of
 215 fruit growth on surface conductance to water vapour diffusion. Annals of Botany 95: 673–683.
- 216 Godoy C., Monterubbianesi G., Tognetti J. 2008. Analysis of highbush blueberry (*Vaccinium*
 217 *corymbosum* L.) fruit growth with exponential mixed models. Scientia Horticulturae, 115: 368–
 218 376.
- 219 Grechi I., Hilgerts N., Génard M., Lescourret F. 2008. Assessing the peach fruit refractometric
 220 index at harvest with a simple model based on fruit growth. Journal of the American Society of
 221 Horticultural Science 133:178–187.
- 222 Grossman Y.L., DeJong T.M. 1995. Maximum fruit growth potential and seasonal patterns of
 223 resource dynamic during peach growth. Annals of Botany, 75, 553–560
- 224 Minchin P.E.H., Snelgar W.P., Blattmann P., Hall A.J. 2010. Competition between fruit and
 225 vegetative growth in Hayward kiwifruit. New Zealand Journal Crop and Horticulture, 38: 101-
 226 112.
- 227 Misle E., Kahlaoui B., Hachicha M., Alvarado P. 2013. Leaf area estimation in muskmelon by
 228 allometry. Photosynthetica, 51: 613–620.
- 229 Niklas K.J. 1994. Plant Allometry: The scaling of form and process; University Chicago Press:
 230 Chicago, IL, USA.

231 R Development Core Team, 2012. R: A Language and Environment for Statistical Computing. R
 232 Foundation for Statistical Computing, Vienna, Austria.

233 Roxburgh SH, Paul KI, Clifford D, England, JR, Raison RJ (2015) Guidelines for constructing
 234 allometric models for the prediction of woody biomass: How many individuals to harvest?
 235 Ecosphere 6: 38 doi: 10.1890/ES14-00251.1.

236 Ruxton G.D., Beauchamp G. 2008. Time for some a priori thinking about post hoc testing.
 237 Behavioral Ecology, 19: 690–693.

238 Spreer W., Ongprasert S., Hegele M., Wünsche J.N., Muller J. 2009. Yield and fruit development in
 239 mango (*Mangifera indica* L. cv. Chok Anan) under different irrigation regimes. Agricultural
 240 Water Management, 96: 574-584.

241 Tuccio L., Remorini D., Pinelli P., Fierini E., Tonutti P., Scalabrelli G., Agati G. 2011. Rapid and
 242 non-destructive method to assess in the vineyard grape berry anthocyanins under different
 243 seasonal and water conditions. Australian Journal of Grape and Wine Research, 17: 181–189.

244 Warton D. I., Weber N.C. 2002. Common slope tests for bivariate errors-in-variables models.
 245 Biometrical Journal, 44: 161 – 174.

246

247

248

249

250

251

252

253

254

255

256

257 **Table 1.** Cultivars, season and orchard locations

Cultivar	Season	Orchard location
Bluegold	2015-2016	38°84' S 72°69' W
Brigitta	2010-2011	38°29' S 72° 23'
Elliot	2008-2009	39°19' S 72°25' W
Elliot	2015-2016	38°84' S 72°69' W
Legacy	2010-2011	38°29' S 72° 23'
Legacy	2015-2016	38°84' S 72°69' W

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

Table 2. Mean fruit diameter and weight for four field-grown blueberry cultivars on data set used for calibrating model. Standard deviation (in brackets) and letters indicating statistic differences by Scheffe's test ($\alpha = 0.05$) are shown.

Cultivar	Season	Berry diameter (mm)	Berry weight (g)	N
Bluegold	2015-2016	16.00 (± 2.25) b	1.95 (± 0.74) b	50
Brigitta	2010-2011	13.35 (± 3.24) d	1.19 (± 0.85) c	50
Elliot	2008-2009	15.52 (± 1.30) bc	1.65 (± 0.36) bc	21
Elliot	2015-2016	15.83 (± 1.31) cd	1.14 (± 0.31) c	27
Legacy	2010-2011	14.97 (± 1.81) bcd	1.55 (± 0.57) bc	16
Legacy	2015-2016	17.75 (± 2.09) a	2.58 (± 0.79) a	33
<i>Comparison among cultivars</i>				
Bluegold		16.00 (± 2.25) A	1.95 (± 0.74) A	53
Brigitta		13.35 (± 3.24) B	1.19 (± 0.85) B	50
Elliot		14.57 (± 1.54) B	1.36 (± 0.41) B	48
Legacy		16.84 (± 2.39) A	2.24 (± 0.86) A	49

Table 3. Linear regression between log-transformed weight and log-transformed diameter data, and ANCOVA for log-transformed diameter and cultivar/season variables. Determination coefficient (R^2) and significance values are shown. Abbreviations: $\ln$ = natural logarithm, FW = fresh weight, FD = fruit diameter.

Cultivar / season	Linear regressions on	
	logarithmic transformed data	R^2
Legacy / 2015-16	$\ln(FW) = 2.68 \ln(FD) - 6.79$	0.98
Legacy / 2008-09	$\ln(FW) = 2.77 \ln(FD) - 7.10$	0.97
Bluegold / 2015-16	$\ln(FW) = 2.74 \ln(FD) - 6.97$	0.98
Elliot / 2015-16	$\ln(FW) = 2.77 \ln(FD) - 7.17$	0.97
Elliot / 2008-09	$\ln(FW) = 2.97 \ln(FD) - 7.68$	0.98
Brigitta / 2010-11	$\ln(FW) = 2.87 \ln(FD) - 7.43$	0.99
ANCOVA		
<i>Variable</i>	<i>p-value</i>	
$\ln(FD)$	2.20×10^{-16}	
Cultivar / season	2.67×10^{-15}	
$\ln(FD) \times \text{cultivar / season}$	0.07

Fig. 1. Allometric relationship between berry diameter (*FD*) and berry weight (*FW*). Fitted model, *p*-value and *n* are shown.

Fig. 2. Estimated versus observed berry weight (A), and residuals versus estimated fruit weight (B) for four field-grown blueberry cultivars. In A, Relative Root Mean Squared Error (RRMSE), line 1:1 (dotted line) and regression between plotted data are showed.