Sensory-motor relationships in speech production in post-lingually deaf cochlear-implanted adults and normal-hearing seniors: Evidence from phonetic convergence and speech imitation

Lucie Scarbel, Denis Beaudet, Jean-Luc Schwartz, Marc Sato

► To cite this version:

Lucie Scarbel, Denis Beaudet, Jean-Luc Schwartz, Marc Sato. Sensory-motor relationships in speech production in post-lingually deaf cochlear-implanted adults and normal-hearing seniors: Evidence from phonetic convergence and speech imitation. *Neuropsychologia*, 2017, *Neuropsychologia*, 101, pp.39 - 46. 10.1016/j.neuropsychologia.2017.05.005 . hal-01546755

HAL Id: hal-01546755

<https://hal.science/hal-01546755>

Submitted on 26 Jun 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Sensory-motor relationships in speech production in post-lingually deaf cochlear-implanted adults and normal-hearing seniors: evidence from phonetic convergence and speech imitation

Lucie Scarbel¹, Denis Beautemps¹, Jean-Luc Schwartz¹, & Marc Sato²

¹ GIPSA-LAB, Département Parole & Cognition, CNRS & Grenoble Université, Grenoble, France

² Laboratoire Parole & Langage, CNRS & Aix-Marseille Université, Aix-en-Provence, France

Authors' note:

Correspondence can be addressed to Lucie Scarbel, GIPSA-LAB, UMR CNRS 5216, Grenoble Université, Domaine Universitaire, BP 46, 38402 Saint Martin d'Hères, France. Tel: (+33) (0)476 824 128. Fax: (+33) (0)476 574 710. Email: lucie.scarbel@gipsa-lab.grenoble-inp.fr.

The authors declare no competing financial interests.

Abstract:

Speech communication can be viewed as an interactive process involving a functional coupling between sensory and motor systems. One striking example comes from phonetic convergence, when speakers automatically tend to mimic their interlocutor's speech during communicative interaction. The goal of this study was to investigate sensory-motor linkage in speech production in postlingually deaf cochlear implanted participants and normal hearing elderly adults through phonetic convergence and imitation. To this aim, two vowel production tasks, with or without instruction to imitate an acoustic vowel, were proposed to three groups of young adults with normal hearing, elderly adults with normal hearing and post-lingually deaf cochlear-implanted patients. Measure of the deviation of each participant's f_0 from their own mean f_0 was measured to evaluate the ability to converge to each acoustic target.

Results showed that cochlear-implanted participants have the ability to converge to an acoustic target, both intentionally and unintentionally, albeit with a lower degree than young and elderly participants with normal hearing. By providing evidence for phonetic convergence and speech imitation, these results suggest that, as in young adults, perceptuo-motor relationships are efficient in elderly adults with normal hearing and that cochlear-implanted adults recovered significant perceptuo-motor abilities following cochlear implantation.

Key-words: Phonetic convergence, imitation, cochlear implant, elderly, sensory-motor interactions, speech production

1. Introduction

1.1. Sensory-motor interactions in speech perception and speech production

In the speech communication domain, an old and fundamental debate concerns the nature of processes and representations involved in speech perception and production. Concerning speech perception, auditory theories assume that speech perceptual processing and categorization are based on acoustic features and auditory representations (Stevens and Blumstein 1978, 1979; Lindblom et al. 1988, 1990). Conversely, the motor theory of speech perception (Liberman et al., 1985) or the direct realist theory (Fowler, 2005) respectively claim that speech perception involves the recovery of the speaker's motor intentions or articulatory gestures. More recently, various perceptuo-motor theories introduced syntheses of arguments by tenants of both auditory and motor theories, and proposed that implicit motor knowledge and motor representations are used in relationship with auditory representations and processes to elaborate phonetic decisions (Skipper et al., 2007; Schwartz et al., 2012). These theories capitalize on the increasing amount of evidence for the role of motor representations and processes in speech perception (see a recent review in Skipper et al., 2017).

Concerning speech production, various theories have also been introduced to characterize the speaker's goals. Motor or articulatory theories, like Task dynamics (Saltzman, 1986) and the associated Articulatory Phonology (Browman & Goldstein, 1992), consider that speech targets are defined in the articulatory space. On the contrary, auditory theories like Stevens' Quantal Theory (Stevens 1972, 1988) and Perkell's speech production control model (Perkell et al. 1995, 2000), suggest that targets are specified in auditory terms. Finally, sensory-motor models claim that speech production control is multimodal and combines auditory and somatosensory information (see Perrier, 2005; and the DIVA model, Guenther et al. 1998, Guenther & Vladusich, 2012). A number of data about the effect of auditory (e.g. Lametti et al., 2014; Shiller & Rochin, 2014) or somatosensory (Tremblay et al., 2003) perturbations applied to speech production in adults or children are in agreement with this sensory-motor framework.

From a number of these theories, sensory-motor relationships seem to play an important role in both speech perception and speech production in normal conditions. The present study deals with subjects for which some degradation of the sensory-motor link could be expected, because of sensory deficits. The first and primary situation is the case of post-lingually deaf subjects equipped with cochlear implants. Indeed, post-lingually deaf subjects are expected to have acquired efficient sensory-motor relationships before deafness. Their ability to achieve intelligible speech production is interpreted as evidence that they have maintained a stable and rather accurate internal model of these sensory-motor relationships all along their deafness life (Perkell et al., 2000). Cochlear implantation then results in providing them with a new kind of auditory input, quite different from the one they had acquired before deafness considering the very special nature of cochlear-implant coding of acoustic information. The question is to know how the internal relationships between these “new” auditory inputs, the “old” ones acquired at the first stages of development, and the motor representations are established and organized, and how efficient they are.

A number of data on speech production after cochlear implantation display an increase in acoustic dispersion of vowels after a few months of implantation (e.g. Langereis et al., 1998; Lane et al., 2005; Ménard et al., 2007), suggesting that the internal model has indeed benefited from the new auditory input provided by cochlear implantation. Interestingly however, some studies explored the response to articulatory perturbation using a bite-block task (Lane et al., 2005), a robotic device displacing the jaw during speech (Nasir & Ostry, 2008) or a lip-tube task (Turgeon, Prémont, Trudeau-Fisette, & Ménard, 2015), with or without auditory feedback. Results showed that post-lingual cochlear-implanted participants (CI) are able to adapt their articulatory trajectory when it is perturbed in order to reach their auditory goals and make their production intelligible, even when the implant is turned off. Such compensatory strategies show that perceptuo-motor abilities acquired during speech acquisition in CI subjects are still at work after deafness, though the addition of auditory information provided by the cochlear-implant does result in enhanced precision and efficiency of the sensory-motor internal model. Finally, a PET-scanning study on visual speech perception in post-lingually deaf CI patients (Rouger et al., 2012) showed that after a short adaptation period with the implant, there was a decrease of the

initially abnormal activity in the superior temporal sulcus, a cross-modal brain area, accompanied by a progressive reactivation of frontal premotor speech areas. This suggests that sensorimotor neuroplasticity after cochlear implantation provides a progressive reactivation of the audio-visuo-motor linkage in CI subjects.

The second situation considered in the present study is the case of elderly subjects. They are of interest for us for two reasons. The first one is the interest of assessing the consequences of the potential decline of cognitive and language abilities in relation with decline in sensory and motor accuracy on the efficiency of sensory-motor relationships. The second reason is that a number of the cochlear-implanted post-lingually deaf participants are rather old, and we considered important to include senior subjects with no severe auditory deficit as a control population. As a matter of fact, very few studies investigated perceptuo-motor relationships in elderly population. One of them shows that elderly people adapt their production in case of degraded auditory feedback (Liu et al., 2010, 2011). Sensorimotor neuroplasticity was also observed in elderly people, linked with age-dependent intelligibility effects mainly found in auditory and motor cortical areas (Tremblay, Dick, & Small, 2013; Bilodeau-Mercure, Ouellet, & Tremblay, 2015). Taken together, these behavioral and neuro-imaging studies suggest that sensory-motor relationships are well preserved in aging.

1.2. Phonetic convergence and imitation, a paradigm for studying sensory-motor relationships in speech

Imitation is a quite widespread phenomenon in speech communication and can be viewed as a key mechanism in the acquisition of human language. In interactive situations, adult speakers tend to continuously adapt their productions to those of their interlocutor, in order to facilitate communicative exchanges. These adaptive changes in speech production can be voluntary, that is when the speaker consciously imitates his interlocutor, but also unintentional. Indeed, unintentional imitation, or phonetic convergence, that is the tendency to automatically imitate a number of acoustic-phonetic characteristics in the productions of an interacting speaker, has been displayed in various studies (see a recent collection of papers on this topic in Nguyen et al., 2013, and for recent reviews see Babel, 2009; Aubanel, 2011; Lelong, 2012). In these studies, convergence effects have been observed both in

paralinguistic features, such as gestures, and in speech acoustic parameters, such as intensity, fundamental frequency f_0 or formants. These phonetic convergence effects may be related to the social component of communication, assuming that they would contribute to setting a common ground between speakers (Giles, Coupland, & Coupland, 1991) and that they could be associated to the human desire of affiliation to a social group.

Interestingly however, while most reports on phonetic convergence are based on conversational exchanges in natural conditions, a few studies showed that phonetic convergence can also be observed using laboratory settings involving non-interactive situations of communication (Goldinger & Azuma, 2004; Gentilucci & Cattaneo, 2005; Delvaux & Soquet, 2007; Garnier, Lamalle, & Sato, 2013; Sato et al., 2013). For example, Delvaux & Soquet (2007) obtained phonetic convergence effects during the production of auditorily presented words without interaction, and they also reported offline adaptation to the auditory targets in post-tests following stimulus exposure. These studies hence suggest that phonetic convergence is not only a matter of social attunement, but could also involve a more basic stage of continuous automatic adaptation of the speech production system to the external speech sounds environment (for a recent review, see Sato et al., 2013).

Such an automatic adaptation mechanism requires the existence of a sensory-motor coupling process in which variations in the external environment provide sensory targets that drive motor control procedures to adapt and produce stimuli closer to these external targets. More in detail, the speaker would program motor commands to achieve an articulatory and ultimately an auditory (or more generally sensory) goal according to the linguistic message to be conveyed to the listener. Then, in the course of speech production, the speaker would compare sensory feedback to the initial sensory objective. Variations in the external environment would shift the sensory targets and accordingly result in modifications of motor commands to converge towards speech sounds in the environment. Hence, the convergence/imitation paradigm is a natural tool to study sensory-motor coupling in speech communication.

Phonetic convergence and imitation in laboratory settings can involve various kinds of acoustic features, though fundamental frequency effects seem to be larger and easier to obtain than variations in

e.g. formant values (Sato et al., 2013). The present study therefore focus on convergence and imitation on f_0 variations.

1.3.f0 perception and motor control in cochlear-implanted adults and normal-hearing seniors

Past studies on pitch control in speech production by CI subjects reported higher f_0 values for CI adults than for adults with normal hearing and, more importantly, larger variations in f_0 for CI than for adults with normal hearing (e.g. Lane & Webster, 1991). Langereis (1998) and Kishon-Rabin et al. (1999) also reported that while pitch production is generally altered in CI patients soon after implantation, almost normal f_0 values were observed one year post-implantation for two participants in Kishon-Rabin et al. (1999) study. Notice that technological evolutions led to important progress in the coding of pitch in cochlear implants since these studies.

Concerning speech perception, several factors appear to influence performance in CI patients. In Blamey et al. (2012), 2251 CI patients participated to a battery of auditory tests, where they had to recognize phonemes, words and sentences. The experimenters reported that both duration of implant experience, age at onset of severe to profound hearing loss, age at cochlear implantation and duration of deafness influence speech perception to a certain extent – though inter-subject variability was quite large in this kind of study. Regarding f_0 , for cochlear-implanted patients, pitch is difficult to estimate because of different reasons. First, as the implant is composed of 12 to 22 electrodes, the merging of frequencies in a given electrode likely impacts frequency discrimination by patients. Furthermore, because of their proximity, electrodes can stimulate nerve fibers, which do not correspond to the intended frequency, which leads to an imperfect tonotopy restitution.

Regarding elderly people, it is well known that auditory capacities decline with aging. More specifically, even with close-to-normal auditory thresholds, elderly people with normal hearing present difficulties in adverse listening conditions, e.g. in a noisy environment (Gelfand, Piper, & Silman, 1985; Ohde & Abou-Khalil, 2001; Fullgrabe, 2013). The role of cognitive functions in this decline of speech perception is still under debate (Cienkovski & Camey, 2002; Cienkovski & Vasil-Dilaj, 2010; Moore & Fullgrabe, 2013; see a review in Fullgrabe & Rosen, 2016). On the motor side, elderly people also show deficits in orofacial movements, which might impact speech production.

Indeed, a majority of studies demonstrate a degradation of speech production in elderly people, with more variable and less stable speech utterances than observed in younger adults. An alteration of voice control is also observed in elderly people, notably with higher pitch variability and less stable pitch production at the level of individual subjects (Morgan & Rastatter, 1986; Russell, Penny, & Pemberton, 1995; Lortie, Thibeault, Guitton, & Tremblay, 2015).

1.4.Goal of the study

In the present study, voluntary and unintentional imitative changes were tested during speech production in both post-lingually deaf or hearing impaired adults equipped with a cochlear implant and in seniors with normal hearing. Our goal was to explore in these populations the underlying perceptuo-motor mechanisms at work in phonetic convergence and speech imitation. To this aim, we capitalized on two recent studies by Sato et al. (2013) and Garnier et al. (2013) displaying both unintentional and voluntary imitative changes on fundamental frequency of auditorily presented vowel targets during speech production in a non-interactive situation of communication. In these studies, participants were asked to produce different vowels according to acoustic targets with their pitch varying around the averaged pitch of their own voice, with or without instruction to imitate the target. Results showed that participants strongly converged to the vowel target not only in the imitative task, but also, at a lower degree, in the production task even if no instruction to imitate the vowel target was given.

To our knowledge, convergence and imitation abilities have never been studied either in CI subjects or in elderly people with normal hearing. We here report online imitative changes on the fundamental frequency in relation to acoustic vowel targets in a non-interactive situation of communication during both unintentional and voluntary imitative production tasks in the two studied populations. Our goal was to determine to which extent CI and elderly participants display convergence and imitation abilities, and to compare these abilities with those observed in young adults with normal hearing (NHY), with the larger aim to evaluate sensory-motor linkage during speech production in CI and elderly population. Three populations have been contrasted: post-lingually deaf or hearing-impaired cochlear-implanted adults (generally not very young), senior adults with no specific auditory deficit apart from aging, and normal-hearing young adults as a control. The same

162 paradigm as the one developed by Garnier et al. (2013) and Sato et al. (2013) was exploited in this
163 study.

164

165

2. Methods

2.1. Participants

Three groups of participants performed the experiment. The first group consisted of fifteen young participants with normal hearing (NHY) (10 females and 5 males, mean age: 30 years old, range: 20-40) who reported no history of speaking, hearing or motor disorders. The second group consisted of ten elderly participants with normal hearing (NHE) (4 females and 6 males, mean age: 69 years old, range: 63-78). The third group consisted of ten post-lingually deaf cochlear-implanted (CI) participants (7 males and 3 females, mean age: 58.9 years old, range: 27-76). As indicated in Table 1, the CI group was heterogeneous in several aspects: age at onset of deafness varied from 7 to 65 years, duration of deafness varied from 1 month to 58 years and duration of cochlear implant experience varied from 1 month to 9 years. In addition, seven of the ten participants wore classical hearing aid in the non-implanted ear, and one participant was bilaterally implanted. The experiment was performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki.

Insert Table 1 about here

2.2. Stimuli

A vowel database was created from /e/, /oe/, /o/ French vowels produced by one male and one female speaker. For each vowel, one clearly articulated occurrence was selected and digitized at 44.1 kHz. From these stimuli, f_0 was artificially shifted by steps of ± 5 Hz (from 80Hz to 180Hz for the male vowels, and from 150 to 350Hz for the female vowels) using the PSOLA module integrated in the Praat software (Boersma and Weenink, 2013). These stimuli allowed us to present to each participant nine distinct stimuli per vowel ranging from -20% and +20% from the mean participant's f_0 by steps of 5% (-20%, -15%, -10%, -5%, 0%, +5%, +10%, +15%, +20%).

2.3. Experimental procedure

The experiment was carried out in a sound-proof room. Participants sat in front of a computer monitor at a distance of approximately 50 cm. The acoustic stimuli were presented at a comfortable sound level

through a loudspeaker, with the same sound level set for all participants. The Presentation software (Neurobehavioral Systems, Albany, CA) was used to control the stimulus presentation during all experiments. All participants' productions were recorded for off-line analyses.

The experiment consisted of three vowel production tasks. First, participants had to individually produce /e/, /æ/ and /o/ vowels, according to a visual orthographic target. This allowed the experimenter to measure the participant's f_0 . In the subsequent task (which will be referred by "convergence task"), participants were asked to produce the three vowels according to an acoustic target. Importantly, no instruction to "repeat" or to "imitate" the acoustic targets was given to the participants. Finally, the third task (which will be referred by "imitative task") was the same as the second task except that participants were explicitly asked to imitate the acoustic targets. The only indication given to participants was to imitate the voice characteristics of the perceived speaker. Importantly, we hence kept a fixed order of the "convergence" and "imitative" tasks so as to avoid cases in which prior explicit imitation instructions could modify further convergence effects and add conscious imitation strategies to automatic convergence processes.

Acoustic targets in the "convergence" and "imitative" tasks for each participant consisted in the 27 stimuli selected from the vowel database (9 per vowel category), repeated three times each so as to obtain 81 trials altogether. Each participant was presented with one model talker of the same sex, with the 9 quantified f_0 frequencies varying from -20% to +20% by steps of 5% around his/her own pitch, as measured in the first task.

At the end of the experiment, participants were asked to perform a frequency discrimination test to estimate their pitch just noticeable difference (JND), based on an experimental paradigm proposed by Vinay & Moore (2010). Two groups of four sounds were presented to participants in a random order, composed either of identical sounds (AAAA) or of two sounds A and B differing in pitch (ABAB), and participants had to determine in which group sounds were different among each other. We used synthesized stimuli, obtained by addition of harmonics of a given f_0 value, with 0 phase and amplitude based on the spectrum of a vowel /e/ produced by either a male or a female, depending of the participant's gender. For sound A, the f_0 value was set at the participant's mean production value. For sound B, the selected f_0 value was systematically higher than f_0 for sound A, beginning with a 5%

deviation further varying along the test. Moreover, small random intensity variations were applied to the two groups of sounds from one trial to the other, in order to force the participant to focus on pitch rather than timbre or intensity perception (Vinay & Moore, 2010). Variations of the f_0 values for sound B were driven by an adaptive two-alternative forced choice procedure: after two consecutive correct answers, the frequency difference between sounds A and B decreased, whereas one mistake drove an increase of this difference. The test result corresponded to the mean of the difference between f_0 values for sound A and sound B for the last eight trials.

2.4.Data analysis

All acoustic analyses of participants' productions were performed using the Praat software (Boersma & Weenink, 2013). In the second and third tasks which both involved an acoustic target (/e/, /œ/ or /o/) that could possibly be misunderstood by the participant, we annotated the participant productions in order to estimate the percentage of errors in production in relation to the target.

Analysis began by a semi-automatic segmentation procedure aiming at segmenting each individual vowel produced by a given participant in a given task. Using intensity and duration criteria, the algorithm automatically identified pauses between each vowel and segmented individual productions accordingly. Segmentation was hand-corrected when appropriate, checking waveform and spectrogram. Only correctly produced vowels (that is, checked by the experimenter as corresponding to the phonemic target) were further analyzed. For each correctly produced vowel, f_0 was determined within a time frame around ± 25 ms of the maximum intensity of the sound file.

2.5.Statistical analysis

For each subject and task, there were altogether 9 values of f_0 production per target f_0 value between -20% and 20% (with 3 vowel types and 3 utterances per type). The mean of the 9 produced f_0 values was computed, and linear regression between mean produced f_0 values and f_0 targets was computed for each participant separately in the convergence and imitation tasks,. The slope and correlation coefficients were then determined for each task and each participant. Individual t-tests were done on these correlation coefficients to determine if they were significant.

For each group, one-tailed t-tests were then performed on individual slope and correlation coefficients compared to zero in order to determine significant imitative changes in each task. To test

for possible differences in imitative changes from one task to the other, additional t-tests were performed on slope and correlation coefficients between the two tasks. However, it is important to remind that the order between the two tasks being not counterbalanced, any difference between them can be interpreted both as a consequence of difference in tasks and order effect.

In addition, in order to test whether imitative changes in the convergence and imitation tasks may correlate among speakers, a Pearson's correlation analysis was performed between slope coefficients. Finally, Pearson's correlation analyses were performed between slope coefficients and JND values within each group of participants and for each task, as well as between slope coefficients and age, duration of deafness, age at implantation, age at the beginning of deafness and duration of implant experience for CI participants in each task.

To compare the results of the three groups, a repeated-measures ANOVA was finally performed on slope coefficients with the group (NHY vs. NHE vs. CI) as between-subject variable and the task (convergence vs. imitation) as within-subject variable. Additionally, a one-factor ANOVA was performed on JND values, with group as categorical factor. The sphericity assumption was tested using a Mauchly test and, when necessary, Greenhouse-Geiser corrections were applied (Greenhouse and Geiser, 1959).

3. Results

3.1. Young participants with normal hearing (see Figure 1)

For NHY participants, imitative changes were observed in both tasks, though stronger during voluntary imitation. Slope coefficients differed significantly from zero in both the convergence ($t(14)=5.98$; $p<0.001$) and imitation ($t(14)=35.78$; $p<0.001$) tasks. In addition, slope coefficients were higher in the imitation compared to the convergence tasks (on average: 0.87 vs. 0.45; $t(14)=6.02$; $p<.001$). No significant correlation was observed between the slope coefficients in the convergence and imitation tasks ($r^2=0.12$).

Similarly, correlation coefficients differed significantly from zero in both the convergence ($t(14)=8.3$; $p<0.001$) and imitation ($t(14)=93.34$; $p<0.001$) tasks, and were higher in the imitation compared to the convergence task (on average: 0.94 vs. 0.64; $t(14)=4.3$; $p<0.001$). Individual analyses showed that fourteen of the sixteen participants converged to the target in the convergence task (r^2 from 0.05 to 0.99), and in the imitative task all of the participants converged to the target (r^2 from 0.91 to 1).

JND values for NHY participants were at 1.06 Hz on average (range: 0.14 to 3.88 Hz), with no significant correlation between the slope coefficients in the convergence or the imitation tasks and JND values (convergence: $r^2=0.00$ $p=.82$, imitation: $r^2=0.00$ $p=.9$).

Insert Figure 1 about here

3.2. Elderly participants with normal hearing (see Figure 2)

For NHE participants, imitative changes were observed in both tasks, though stronger in voluntary imitation. Slope coefficients differed significantly from zero in both the convergence ($t(9)=3.52$; $p<0.01$) and imitation ($t(9)=8.5$; $p<0.001$) tasks. In addition, slope coefficients were higher in the imitation compared to the convergence task (on average: 0.75 vs. 0.33; $t(9)=4.97$; $p<.001$). There was no significant correlation between convergence and imitation tasks ($r^2=0.59$).

Similarly, correlation coefficients differed significantly from zero in both the convergence ($t(9)=5.96$; $p<0.001$) and imitation ($t(9)=11.78$; $p<0.001$) tasks, and were higher in the imitation compared to the

convergence tasks (on average: 0.84 vs. 0.54; $t(9)=4.12$; $p<0.001$). Individual analyses showed that eight of the ten participants converged to the target in the convergence task (r^2 from 0.35 to 0.99), and in the imitative task all of the participants except one converged to the target (r^2 from 0.28 to 0.99).

JND values for NHE participants were at 0.95 Hz on average (range: 0.43 to 1.8 Hz) with no correlation with either the convergence or the imitation slopes (convergence: $r^2=0.03$ $p=.63$ imitation: $r^2=0.00$ $p=.79$).

Insert Figure 2 about here

3.3. Cochlear-implanted participants (see Figure 3)

As for NHY and NHE participants, imitative changes were also observed in both tasks for CI participants. Slope coefficients differed significantly from zero in both the convergence ($t(9)=3.24$; $p<0.02$) and imitation ($t(9)=4.84$; $p<0.001$) tasks. In addition, slope coefficients were higher in the imitation compared to the convergence tasks (on average: 0.39 vs. 0.14; $t(9)=3.53$; $p<0.001$). As for NH participants, there was no significant correlation between convergence and imitation tasks ($r^2=0.28$).

Similarly, correlation coefficients differed significantly from zero in both the convergence ($t(9)=4.08$; $p<0.01$) and imitation ($t(9)=5.53$; $p<0.001$) tasks, and were higher in the imitation compared to the convergence tasks (on average: 0.49 vs. 0.24; $t(9)=2.95$; $p<0.02$).

JND values for CI participants were at 14.06 Hz on average (range: 1.71 to 45.35 Hz). There was also no correlation between the convergence or imitation tasks and the others factors, that are JND values, age, deafness duration, age at implantation, age at the beginning of deafness and duration of implant experiment (all $r^2<0.3$).

While young and elderly participants with normal hearing made no errors in both tasks, cochlear-implanted participants made a number of errors in both tasks, with similar error percentage in the convergence (15%) and imitation tasks (13%). In order to verify if these errors influence convergence effects, correlations between production and target were measured in both tasks on error trials only, for participants presenting at least 15% of errors (four participants in each task). The correlation values are displayed in Table 2. For those participants, correlations between production and target for error

324 trials appear similar to those obtained for correct trials in both convergence and imitation tasks. As a
325 matter of fact, the correlation values for correct trials and for error trials (when they are sufficient to do
326 computations) are highly correlated among the 8 cases where both can be computed ($r^2=0,71$,
327 $t(6)=3,81$, $p<0,005$).

328 To relate convergence effects with etiology variability and JND values, we report determination
329 coefficients in both tasks for each participant (see Table 2). In the convergence task, we obtained
330 significant correlation between target and production for five participants, with determination
331 coefficients from 0.05 and 0.75. In the imitative task, correlation is significant for seven participants,
332 with determination coefficients from 0.09 and 0.94. As shown in Table 2, the participants with no
333 significant convergence in one or the other tasks do not display obvious similarity in etiology in e.g.
334 age at deafness onset or deafness duration. Importantly, clear convergence in both tasks is obtained for
335 some subjects after a quite reduced duration of implantation: see the cases of subjects CI1, CI2, CI5 or
336 CI9 with significant convergence or imitation on f_0 after 1 to 5 months of implantation.

337 Concerning JND values, the participants with no significant convergence in one or the other tasks
338 do not present the worst JND values, and the participant with the highest JND value obtains significant
339 convergence effect in both tasks. More strikingly, the corresponding JND value for this subject, at 45
340 Hz, is actually larger than the largest variation in F_0 applied to the stimuli in both convergence tasks.
341 As a matter of fact, F_0 for this subject equals 125 Hz, and the maximal variation of F_0 at 20% results
342 in a 25 Hz modification. Altogether, these facts converge to suggest that the task used for JND
343 estimation, well adapted for normal-hearing subjects (Vinay & Moore, 2010), is probably too complex
344 for cochlear-implanted subjects.

345 Insert Table 2 about here

347 Insert Figure 3 about here

349 3.4. Comparison between groups (see Figure 4)

The main effect of group was significant ($F(2,32)=31.3$; $p<.001$) with lower mean slope coefficient for CI participants than for NHY and NHE participants (CI: 0.26 ; NHY: 0.65 ; NHE: 0.55). No difference was obtained between NHY and NHE groups. The main effect of task was also significant ($F(1,32)=67.29$; $p<.001$), with lower slope coefficients for the convergence task than for the imitation task (convergence: 0.33 ; imitation: 0.69). No significant interaction was found between the group and the task.

For the JND test, the effect of group was significant ($F(2,32)=11.03$; $p<.001$), with larger values for CI participants than for NHY and NHE participants (CI: 14.06; NHY: 1.06; NHE: 0.95) without significant difference between NHY and NHE participants. This shows that CI participants stay largely impaired in f_0 discrimination.

Insert Figure 4 about here

4. Discussion

The goal of this study was to investigate sensory-motor linkage in speech production on post-lingually deaf cochlear-implanted patients, in comparison with young and elderly adults with normal hearing, through the abilities of f_0 convergence and imitation. To this aim, we used a paradigm of intentional and non-intentional imitation of vowels with modified fundamental frequency.

4.1. Young adults

We firstly replicated the previous findings by Sato et al. (2013) and Garnier et al. (2013) for young adults with normal hearing. Imitative changes towards the acoustic target were indeed observed in both tasks, with stronger convergence in the task with direct instruction than in the convergence task. It is likely that this is mainly due to the effect of the explicit instruction, though we cannot discard the possibility that order intervened here, with a trend to increase convergence in the second task just becomes it comes after the first one. As in the two previous studies, no correlation was found between the two tasks, that is individual f_0 changes in the convergence task were not related to those observed in the imitative task among subjects. This lack of correlation could be due to ceiling effects provided by the very high degree of convergence in the imitative task and the very low variability of these imitative changes for almost all participants. These results confirm that convergence can occur even in a non-interactive situation of communication, in line with previous studies (Goldinger & Azuma, 2004; Gentilucci & Cattaneo, 2005; Delvaux & Soquet, 2007; Garnier et al. 2013; Sato et al. 2013). Such non-interactive phonetic convergence effects likely rely on low-level sensory-motor mechanisms described in the Introduction section, according to which participants would first analyze the target stimulus to elaborate their motor and sensory goals, resulting in an adaptation of their production to the acoustic target.

4.2. Elderly population

The results for elderly participants with normal hearing were similar to those of young adults. Indeed, NHE imitated the auditory target in the imitative task and converged towards the acoustic target in a probably subconscious way in the convergence task. In addition, as for young participants with normal

hearing, no correlation was observed between the two tasks. The observed convergence effects in both voluntary and unintentional imitation in this elderly population, as high as in young subjects, suggests that sensory-motor relationships during speech production are efficient in these participants, which is in line with previous studies suggesting that sensory-motor relationships are still active and efficient in seniors (Liu et al., 2010, 2011, Tremblay et al. 2013).

4.3. Post-lingually deaf cochlear-implanted participants

In spite of their strongly impaired ability to discriminate frequencies, as displayed by JND values 10 to 15 times larger than those of subjects with normal hearing, it is striking that CI participants were also found to be able to imitate and to converge towards the acoustic targets. This suggests that they are able to estimate the pitch of a vowel target to a certain extent and to monitor their own vocal source to attempt to get closer to this target. Even more strikingly, they do it even in the convergence task where no explicit imitation instruction was provided – hence the importance to know that all participants performed this task before the “imitative task”, reducing the risk that they could have considered that imitation mattered in the task.

Although these results demonstrate that cochlear-implanted patients are able to perceive and imitate the fundamental frequency of the vowel targets, no correlation was however found between their imitative abilities and their auditory discrimination scores in the JND test. This suggests that their imitation abilities are not related to their auditory capacity of perceiving frequencies in a simple way. Nevertheless, cochlear-implanted participants showed less convergence and imitation than adults with normal hearing whatever their age, and higher pitch variability between participants. At this stage, it is unclear if pitch variability in production is due to inaccurate estimation of the target pitch or to degraded voice control, but interestingly, the lack of correlation between convergence and JND suggests that apart from sensory ability, the sensory-motor relationships could differ from one subject to the other. Considering the importance and efficiency of somatosensory control for deaf subjects (Nasir & Ostry, 2008), together with the range of inter-individual variations in the relative importance of auditory and somatosensory control (Lametti et al., 2012; Feng et al., 2011), it is likely that variations in somatosensory dependence could be at hand in these data.

Contrary to Blamey et al. (2012) who showed that deafness duration and age at deafness onset are two factors influencing speech perception performance, we did not find any correlation between these factors and CI abilities to imitate or to converge towards an acoustic target. However, it is important to note that in the Blamey et al. study (2012), residual variability not explained by CI- and deafness-related factors was considerable (representing 90% of variation in their study). It seems therefore likely that the limited sample of our CI participants impeded any clear correlation with CI- and deafness-related characteristics to appear in the present study.

Finally, contrary to NH participants, CI participant's responses were not always correct (with up to 13-15% errors in each task), which suggests some auditory difficulties in the decoding of isolated vowels. Actually, it is known that CI listeners need formant transition to accurately categorize a vowel, as shown by Hanna (2011). The isolated vowels used in our study are hence probably difficult to categorize efficiently by the CI participants. Moreover, convergence effects tested in error trials appear to be similar to those of correct trials, which suggests that timbre and pitch estimations seem to be relatively independent for CI participants. Nevertheless, the major result of this study is that a number of CI participants have already recovered a good ability to associate auditory with motor parameters even a short time after implantation (with significant convergence after only one, two or three months of implantation for some subjects). Indeed, results displayed both automatic and conscious imitation abilities in post-lingually deaf cochlear-implanted patients, displaying their ability to adapt their production to the environment, which is uniquely due to a functional sensory-motor linkage. This should be crucial for the retuning of their speech production system (see Perkell, Lane, Svirsky, & Webster, 1992), enabling them to improve their internal model (Perkell et al., 2000). Interestingly, recovering perceptuo-motor abilities could also be of importance for their speech perception abilities, considering the proposals about the role of the motor system in speech perception (e.g. Skipper, vanWassenhove, Nusbaum, & Small, 2007; Schwartz, Basirat, Ménard, & Sato, 2012).

5. Conclusion

The present study reports convergence and imitation abilities in adults with normal hearing, both young and elderly, and in cochlear-implanted subjects. Indeed, in the three groups, we obtain

444 significant and generally strong imitation of the acoustic target during a task with explicit instruction,
445 and also convergence effects in a task without direct instruction. These results first confirm that
446 sensory-motor relationships during speech production are still efficient in elderly subjects, with
447 basically no difference with younger ones. Moreover, crucially, we also show for the first time that
448 cochlear-implanted participants have the ability to converge to an acoustic target, intentionally and
449 unintentionally. Therefore, they have recovered significant sensory-motor abilities, which could be
450 crucial for improving both their speech production and speech perception abilities.

451

452

453 **6. Acknowledgments**

454 This work was supported by the French National Research Agency (ANR) through funding from
455 the PlasMody project (Plasticity and Multimodality in Oral Communication for the Deaf).

456

References

- Aubanel V. (2011). Variation phonologique régionale en interaction conversationnelle. *Doctoral Dissertation, Aix-Marseille University.*
- Babel M. (2009). Phonetic and social selectivity in speech accommodation. *Doctoral Dissertation, University of California, Berkeley.*
- Bilodeau-Mercure, M., Ouellet, C., & Tremblay, P. (2015). Vieillissement oro-facial et production de la parole. *Affiche présentée au Congrès québécois de recherche en adaptation-réadaptation. Boucherville, Qc, Canada.*
- Blamey, P., Artieres, F., Baskent, D., Bergeron, F., Beynon, A., Burke, E., Dillier, N., Dowell, R., Fraysse, B., Gallégo, S., Govaerts, P.J., Green, K., Huber, A.M., Kleine-Punte, A., Maat, B., Marx, M., Mawman, D., Mosnier, I., Fitzgerald O'Connor, A., O'Leary, S., Rousset, A., Schauwers, K., Skarzynski, H., Skarzynski, P.H., Sterkers, O., Terranti, A., Truy, E., Van de Heyning, P., Venail, F., Vincent, C., & Lazard, D.S. (2012). Factors affecting auditory performance of postlinguistically deaf adults using cochlear implant: an update with 2251 patients. *Audiology & Neurology*, 18, 36-47.
- Blumstein, S. E. & Stevens, K. N. (1979). Acoustic invariance in speech production: Evidence from measurements of the spectral characteristics of stop consonants. *Journal Of the Acoustical Society of America*, 66, 1001-1017
- Boersma, P. (2001). Praat, a system for doing phonetics by computer, *Glott International*, 5(9/10), 341-345.
- Browman, C. P., & Goldstein, L. (1992). Articulatory phonology: an overview. *Phonetica*, 49(3-4), 155-180.
- Cienkowski, K. M. & Carney, A. E. (2002). Auditory-visual speech perception and aging. *Ear and Hearing*, 23(5), 439-449.
- Cienkowski, K. M., & Vasil-Dilaj, K. (2010). The Role of Central Auditory and Cognitive Processing Ability in Speech Perception for Elderly Listeners. *Perspectives on Hearing and Hearing Disorders Research and Diagnostics*, 14(2), 30.

- 484 Delvaux, V. & Soquet, A. (2007). The influence of ambient speech on adult speech productions
485 through unintentional imitation. *Phonetica*, 64, 145–173.
- 486 Feng, Y., Gracco, V. L., & Max, L. (2011). Integration of auditory and somatosensory error signals in
487 the neural control of speech movements. *Journal of neurophysiology*, 106(2), 667-679.
- 488 Fowler, C. A. (1986). An event approach to the study of speech perception for a direct-realist
489 perspective. *Journal of Phonetics*, 14.
- 490 Füllgrabe, C. (2013). Age-Dependent Changes in Temporal-Fine-Structure Processing in the Absence
491 of Peripheral Hearing Loss. *American Journal of Audiology*, 22(2), 313.
- 492 Füllgrabe C, Rosen S (2016) On the (un)importance of working memory in speech-in-noise processing
493 for listeners with normal hearing thresholds. *Frontiers in Psychology*
- 494 Garnier, M., Lamalle, L., & Sato, M. (2013). Neural correlates of phonetic convergence and speech
495 imitation. *Frontiers in psychology*, 4, 600.
- 496 Gentilucci, M. & Cattaneo, L. (2005). Automatic audiovisual integration in speech perception.
497 *Experimental Brain Research*, 167:66-75.
- 498 Goldinger, S.D. & Azuma, T. (2004). Episodic memory reflected in printed word naming.
499 *Psychonomic Bulletin and Review*, 11(4): 716-722.
- 500 Gelfand, S. A., Piper, N., & Silman, S. (1985). Consonant recognition in quiet as a function of aging
501 among normal hearing subjects. *The Journal of the Acoustical Society of America*, 78(4),
502 1198-1206.
- 503 Giles H., Coupland N., & Coupland J. (1991). Accommodation theory: Communication, context, and
504 consequence. In Giles H, Coupland N, Coupland J (Editors) *Contexts of Accommodation:*
505 *Developments in Applied Sociolinguistic*, Cambridge University Press pp. 1-68.
- 506 Guenther, F. H., Hampson, M., & Johnson, D. (1998). A theoretical investigation of reference frames
507 for the planning of speech movements. *Psychological Review*, 105(4), 611-633.
- 508 Guenther, F.H. and Vladusich, T. (2012). A neural theory of speech acquisition and
509 production. *Journal of Neurolinguistics*. 25, pp. 408-422.
- 510 Hanna, N. (2011). Vowel identification by cochlear implant users: comparison on vowel edges and
511 vowel centers. Honors College Thesis, University of South Florida.

- 512 Kishon-Rabin, L., Taitelbaum, R., Tobin, Y., & Hildesheimer, M. (1999). The effect of partially
513 restored hearing on speech production of postlingually deafened adults with multichannel cochlear
514 implants. *The Journal of the Acoustical Society of America*, 106(5), 2843-2857.
- 515 Lametti, D.R., Sazzad, Nasir, R.M., & Ostry, D.J. (2012). Sensory preference in speech production
516 revealed by simultaneous alteration of auditory and somatosensory feedback. *Journal of*
517 *Neuroscience*, 32(27), 9351-9358.
- 518 Lametti, D.R., Rochet-Capellan, A., Neufeld, E., Shiller, D.M., & Ostry, D.J. (2014) Plasticity in the
519 human speech motor system drives changes in speech perception. *Journal of Neuroscience*,
520 30;34(31), 10339-46.
- 521 Lane, H. & Webster, J.W. (1991). Speech deterioration in postlingually deafened adults. *The Journal*
522 *of the Acoustical Society of America*, 89, 859-866.
- 523 Lane, H., Denny, M., Guenther, F. H., Matthies, M. L., Menard, L., Perkell, J. S., Stockmann, E.,
524 Tiede, M., Vic, J. & Zandipour, M. (2005). Effects of bite blocks and hearing status on vowel
525 production. *The Journal of the Acoustical Society of America*, 118(3 ; 1), 1636-1646.
- 526 Langereis, M.C., Bosman, A.J., Van Olphen, A.F., & Smoorenburd, G.F. (1998). Effect of cochlear
527 implantation on voice fundamental frequency in post-lingually deafened adults. *Audiology*, 37,
528 219-230.
- 529 Lelong, A. (2012). *Convergence Phonétique en Interaction*. Doctoral Dissertation, Grenoble
530 University.
- 531 Liberman, A. M., & Mattingly, I. G. (1985). The Motor Theory of Speech Perception Revised.
532 *Cognition*, 21(1), 1–36.
- 533 Lindblom, B., & Maddieson, I. (1988). Phonetic Universals in Consonant Systems. In *Language,*
534 *Speech and Mind* (p. 62-78). London: Routledge.
- 535 Lindblom, B. (1990). Explaining Phonetic Variation: A Sketch of the H&H Theory. In W. J.
536 Hardcastle & A. Marchal (éd.), *Speech Production and Speech Modelling* (p. 403-439). Springer
537 Netherlands.
- 538 Linville, S. E., & Fisher, H. B. (1985). Acoustic characteristics of women's voices with advancing
539 age. *Journal of Gerontology*, 40(3), 324-330.

- 540 Linville, S. E., & Korabic, E. W. (1987). Fundamental frequency stability characteristics of elderly
541 women's voices. *The Journal of the Acoustical Society of America*, 81(4), 1196-1199.
- 542 Liu, H., Wang, E. Q., Chen, Z., Liu, P., Larson, C. R., & Huang, D. (2010). Effect of tonal native
543 language on voice fundamental frequency responses to pitch feedback perturbations during
544 sustained vocalizations. *The Journal of the Acoustical Society of America*, 128(6), 3739-3746.
- 545 Liu, H., Meshman, M., Behroozmand, R., & Larson, C. R. (2011). Differential effects of perturbation
546 direction and magnitude on the neural processing of voice pitch feedback. *Clinical*
547 *neurophysiology : official journal of the International Federation of Clinical Neurophysiology*,
548 122(5), 951-957.
- 549 Liu, P., Chen, Z., Jones, J. A., Huang, D., & Liu, H. (2011). Auditory Feedback Control of Vocal Pitch
550 during Sustained Vocalization: A Cross-Sectional Study of Adult Aging. *PLoS ONE*, 6(7), e22791.
- 551 Lortie, C. L., Thibeault, M., Guitton, M. J., & Tremblay, P. (2015). Effects of age on the amplitude,
552 frequency and perceived quality of voice. *Age (Dordrecht, Netherlands)*, 37(6), 117.
- 553 Nasir, S. M., & Ostry, D. J. (2008). Speech motor learning in profoundly deaf adults. *Nature*
554 *Neuroscience* 11, 1217 - 1222 (2008).
- 555 Nguyen, N., Sato, M., Postma, M., Pardo, J., Babel, M. & Postma, E. (2013). Speech imitation: the
556 cognitive underpinnings of adaptive vocal behavior. *Frontiers in Psychology*.
- 557 Max, L., Daniels, J., Curet, K., & Cronin, K. (2008) Modulation of auditory and somatosensory
558 processing during the planning of speech movements *Proceedings of the 8th International Seminar*
559 *on Speech Production*, 41-44, Strasbourg, France.
- 560 Ménard, L., Polak, M., Denny, M., Burton, E., Lane, H., Matthies, M. L., ... Vick, J. (2007).
561 Interactions of speaking condition and auditory feedback on vowel production in postlingually deaf
562 adults with cochlear implants. *The Journal of the Acoustical Society of America*, 121(6),
563 3790-3801.
- 564 Moore, D. R., & Fullgräbe, C. (2013). Cognitive contributions to hearing in older people. *Journal of*
565 *Hearing Science*, 2(4), 58-60.
- 566 Morgan, E. E., & Rastatter, M. (1986). Variability of voice fundamental frequency in elderly female
567 speakers. *Perceptual and Motor Skills*, 63(1), 215-218.

- 568 Ohde, R. N., & Abou-Khalil, R. (2001). Age differences for stop-consonant and vowel perception in
569 adults. *The Journal of the Acoustical Society of America*, 110(4), 2156-2166.
- 570 Perkell, J.S., Lane, H., Svirsky, M., & Webster, J. (1992). Speech of cochlear implant patients: A
571 longitudinal study of vowel production. *The Journal of the Acoustical Society of America*, 91,
572 2961-2979.
- 573 Perkell, J. S., Matthies, M. L., Svirsky, M. A., & Jordan, M. I. (1995). Goal-based speech motor
574 control: A theoretical framework and some preliminary data. *Journal of Phonetics*, 23(1), 23-35.
- 575 Perkell, J.S., Guenther, F.H., Lane, H., Matthies, L.M., Perrier, P., Vick, J., Wilhelms-Tricarico, R., &
576 Sandipour, M. (2000). A theory of speech motor control and supporting data from speakers with
577 normal hearing and with profound hearing loss. *Journal of phonetics*, 28, 233-272.
- 578 Perrier, P. (2005). Control and representations in speech production. *ZAS Papers in Linguistics*, 40,
579 109-132.
- 580 Rouger, J., Lagleyre, S., Démonet, J.-F., Fraysse, B., Deguine, O., & Barone, P. (2012). Evolution of
581 crossmodal reorganization of the voice area in cochlear-implanted deaf patients. *Human Brain*
582 *Mapping*, 33(8), 1929-1940.
- 583 Russell, A., Penny, L., & Pemberton, C. (1995). Speaking fundamental frequency changes over time
584 in women: a longitudinal study. *Journal of Speech and Hearing Research*, 38(1), 101-109.
- 585 Saltzman, E. (1986). Task Dynamic Coordination of the Speech Articulators: A preliminary model.
586 *Experimental Brain Research Series*, 15, 130 - 144.
- 587 Sato, M., Schwartz, J.-L. & Perrier, P. (2014). Phonemic auditory and somatosensory goals in speech
588 production. *Language, Cognition and Neuroscience*, 29(1): 41-43.
- 589 Sato, M., Grabski, K., Garnier, M., Granjon, L., Schwartz, J.L., & Nguyen, N. (2013). Converging to a
590 common speech code: imitative and perceptuo-motor recalibration processes in speech production.
591 *Frontiers in Psychology*, 4, 422.
- 592 Savariaux, C., Perrier, P. & Orliaguet, J. P. (1995) Compensation strategies for the perturbation of the
593 rounded vowel [u] using a lip tube: A study of the control space in speech production. *Journal of*
594 *the Acoustical Society of America*, 98 (5), 2428-2442.

- Schwartz, J.L., Basirat, A., Ménard, L. & Sato, M. (2012). The Perception-for-Action-Control Theory (PACT): A perceptuo-motor theory of speech perception. *Journal of Neurolinguistics*, 25, 336-354.
- Shiller, D.M., & Rochon, A.M. (2014) Auditory-perceptual learning improves speech motor adaptation in children. *Journal of Experimental Psychology. Human Perception and Performance*, 40(4):1308-15.
- Skipper, J. I., vanWassenhove, V., Nusbaum, H. C., & Small, S. L. (2007). Hearing lips and seeing voices: How cortical areas supporting speech production mediate audiovisual speech perception. *Cerebral Cortex*, 17, 2387-2399.
- Skipper, J. I., Devlin, J. T., & Lametti, D. R. (2016). The hearing ear is always found close to the speaking tongue: Review of the role of the motor system in speech perception. *Brain Lang*, 164, 77-105.
- Stevens, K. N. (1972). The quantal nature of speech: Evidence from articulatory-acoustic data. In *Human communication: A unified view* (p. 51-66). EE David et PB Denes editors.
- Stevens, K. N., & Blumstein, S. E. (1978). Invariant cues for place of articulation in stop consonants. *The Journal of the Acoustical Society of America*, 64(5), 1358-1368.
- Stevens, K. N. (1989). On the quantal nature of speech. *Journal of Phonetics*, 17, 3-46.
- Tremblay, S., Shiller, D.M., & Ostry, D.J. (2003) Somatosensory basis of speech production. *Nature*, 19;423(6942):866-9.
- Tremblay, P., Dick, A. S., & Small, S. L. (2013). Functional and structural aging of the speech sensorimotor neural system: functional magnetic resonance imaging evidence. *Neurobiology of Aging*, 34(8), 1935-1951.
- Turgeon, C., Prémont, A., Trudeau-Fisette, P., & Ménard, L. (2015). Exploring production-perception relationships in normal hearing and cochlear implant adults: a lip-tube perturbation study. *Clinical Linguistics and Phonetics*, 29(5), 378-400.
- Vinay, S.N. & Moore, B.C. (2010). Effects of the use of personal music players on amplitude modulation detection and frequency discrimination. *The Journal of the Acoustical Society of America*, 128(6), 3634-41.

Figures

Figure 1: Phonetic convergence and voluntary imitative changes on fundamental frequency observed in young participants with normal hearing (X axis: deviation percentage with respect to mean participant' f_0 , Y axis: mean production for all participants, error bars corresponding to standard deviation

Figure 2: Phonetic convergence and voluntary imitative changes observed on fundamental frequency in elderly participants with normal hearing (X axis: deviation percentage with respect to mean participant' f_0 , Y axis: mean production for all participants, error bars corresponding to standard deviation)

Figure 3: Phonetic convergence and voluntary imitative changes observed on fundamental frequency in cochlear-implanted participants (X axis: deviation percentage with respect to mean participant' f_0 , Y axis: mean production for all participants, error bars corresponding to standard deviation)

644

645 Figure 4: Phonetic convergence and voluntary imitative slope changes on fundamental frequency in
 646 young (NHY) and elderly (NHE) participants with normal hearing and cochlear-implanted participants
 647 (CI) (error bars corresponding to standard deviation)

648

649

650

651

652

Tables

Table 1: Characteristics of participants with cochlear implants

	Gender	Age (y.)	Age at onset of deafness (y.)	Hearing aid	Duration of deafness	Duration of CI experiment
CI1	M	65	7	No	58 years	1 month
CI2	M	56	20	Yes	35 years	3 months
CI3	F	66	32	Yes	25 years	9 years
CI4	M	60	59	Yes	1 month	1 years 4 months
CI5	F	43	20	Yes	13 years	2 months
CI7	M	27	25	Yes	2 months	2 years 6 months
CI8	F	67	65	Yes	2 years	7 months
CI9	M	72	40	Yes	30 years	5 months
CI10	M	76	48	No	27 years	3 years 4 months
CI11	M	57	48	Yes	8 years	10 months

Table 2: Determination coefficients in both tasks for each participant of the cochlear-implanted group (CI), normal-hearing elderly adults group (NHE) and normal-hearing young adults (NHY). For the cochlear-implanted group, determination coefficients measured on errors trials only are indicated in parentheses for participants with at least 15% errors (enabling to obtain a sufficient number of errors for computing correlations).

CI			NHE		NHY	
convergence	imitation	JND	convergence	imitation	convergence	imitation
0,25 (0,33)	0,51(0,32)	15,22	0,72	0,95	0,96	0,95
0,51	0,70	45,35	0,36	0,28	0,99	1,00
0,78 (0,51)	0,77 (0,73)	1,71	0,98	0,99	0,69	0,99
0,04 (0,17)	0,39	9,76	0,49	0,93	0,99	0,97
0,27	0,94	1,79	0,99	0,96	0,87	0,99
0,12	0,77 (0,41)	4,55	0,88	0,97	0,71	0,91
0,49	0,62	18,52	0,50	0,98	0,05	0,97
0,75	0,92	10,55	0,92	0,98	0,96	0,99
0,11 (0,11)	0,22 (0,03)	16,21	0,99	0,99	0,99	0,99
0,60	0,09	28,54	0,35	0,91	0,10	0,98
					0,95	0,94
					0,98	0,99
					0,84	0,95
					0,63	0,97
					0,97	0,99