

A decomposition theorem for projective manifolds with nef anticanonical bundle

Junyan Cao, Andreas Höring

► To cite this version:

Junyan Cao, Andreas Höring. A decomposition theorem for projective manifolds with nef anticanonical bundle. 2017. hal-01547826

HAL Id: hal-01547826

<https://hal.archives-ouvertes.fr/hal-01547826>

Preprint submitted on 27 Jun 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A DECOMPOSITION THEOREM FOR PROJECTIVE MANIFOLDS WITH NEF ANTICANONICAL BUNDLE

JUNYAN CAO AND ANDREAS HÖRING

ABSTRACT. Let X be a simply connected projective manifold with nef anticanonical bundle. We prove that X is a product of a rationally connected manifold and a manifold with trivial canonical bundle. As an application we describe the MRC fibration of any projective manifold with nef anticanonical bundle.

1. INTRODUCTION

1.A. Main results. Manifolds with nef anticanonical class naturally appear as an interpolation of Fano manifolds and compact Kähler manifolds with trivial canonical class, but they come with many new features: in general the anticanonical bundle is not semiample (blow-up \mathbb{P}^2 in nine general points), nor hermitian semipositive [DPS94, Ex.1.7]. Initiated by the fundamental papers of Demailly, Peternell and Schneider [DPS93, DPS94, DPS96], a central goal of the theory is to understand the natural maps attached to X , e.g. the Albanese map or the MRC-fibration. The expected outcome of this study is contained in the following conjecture which should be understood as an analogue of the Beauville-Bogomolov decomposition theorem [Bea83]:

1.1. Conjecture. *Let X be a compact Kähler manifold with nef anticanonical class. Then the universal cover \tilde{X} of X decomposes as a product*

$$\tilde{X} \simeq \mathbb{C}^q \times \prod Y_j \times \prod S_k \times Z,$$

where Y_j are irreducible Calabi-Yau manifolds, S_k are irreducible hyperkähler manifolds, and Z is a rationally connected manifold.

This conjecture has been proven under the stronger assumption that T_X is nef [CP91, DPS94], $-K_X$ is hermitian semipositive [DPS96, CDP15] or the general fibre of the Albanese map is weak Fano [CH17].

In this paper we focus on the case where X is a projective manifold. Very recently the first-named author proved that for any projective manifold X with nef anticanonical bundle the Albanese map $X \rightarrow \text{Alb}(X)$ is a locally trivial fibration [Cao16]. We know that the fundamental group is almost abelian [Pau97], so the next step is to study X when it is simply connected. We show that the structure of X is as simple as possible:

Date: June 27, 2017.

2010 Mathematics Subject Classification. 14D06, 14J40, 14E30, 32J25, 32J27.

Key words and phrases. anticanonical bundle, positivity of direct images, MRC fibration.

1.2. Theorem. *Let X be a projective manifold such that $-K_X$ is nef and $\pi_1(X) = 1$. Then $X \simeq Y \times F$ such that $K_Y \sim 0$ and F is a rationally connected manifold.*

By [Cao16, Cor.1.4] this immediately implies:

1.3. Corollary. *Conjecture 1.1 holds if X is projective.*

Moreover we obtain a precise description of the MRC-fibration:

1.4. Theorem. *Let X be a projective manifold such that $-K_X$ is nef. Then there exists a locally trivial fibration*

$$X \rightarrow Y$$

such that the fibre F is rationally connected and $K_Y \equiv 0$.

Using Mori theory and explicit computations this statement was shown for threefolds in [BP04, Sect.3]. For Kähler manifolds even this low-dimensional case is still open.

1.B. Strategy and organisation of the paper. By a result of Qi Zhang [Zha05] the base of the MRC fibration has Kodaira dimension zero, so the situation of Theorem 1.2 looks similar to the case of the Albanese fibration studied in [Cao16]. However, even by using techniques from the MMP, it seems a priori not clear that the MRC-fibration can be represented by a *holomorphic* map $X \rightarrow Y$. Therefore we have to proceed in a less direct way: the MRC-fibration is an almost holomorphic map, so it determines a unique component of the Chow space parametrising its general fibres. We denote by Y a resolution of this component, and by $\varphi : \Gamma \rightarrow Y$ a resolution of the universal family over this component.

The natural map $\pi : \Gamma \rightarrow X$ is birational with exceptional locus E , and our first goal is to describe the structure of the fibre space $\varphi : \Gamma \rightarrow Y$. The tool for this description is the positivity of certain direct image sheaves: for some sufficiently ample line bundle A on X , we set $L := \pi^*A$. The main technical result of this paper (Proposition 3.10) is that for some $m \gg 0$ and all p sufficiently divisible, the direct image sheaves

$$\mathcal{V}_p := \varphi_*(\mathcal{O}_\Gamma(pL + pmE - \frac{p}{r}\varphi^*\det\varphi_*(\mathcal{O}_\Gamma(L + mE))))$$

are trivial vector bundles over a certain open subset $Y_0 \subset Y$ which is simply connected and has only constant holomorphic functions (cf. Setup 3.1 for the definition of Y_0), where r is the rank of $\varphi_*(\mathcal{O}_\Gamma(L + mE))$. Following an argument going back to [DPS94] this implies that we have a birational map

$$\varphi^{-1}(Y_0) \dashrightarrow Y_0 \times F,$$

where F is a general fibre of the MRC fibration.

The second step is to see how the product structure of some birational model yields a product structure on X . This typically requires some control of the birational map $X \dashrightarrow Y \times F$, in our case we simply observe that the product structure on $Y_0 \times F$ induces a splitting of the tangent bundle of $\varphi^{-1}(Y_0) \setminus E$. Since (by the construction of Y_0) the complement of $\varphi^{-1}(Y_0) \setminus E \subset X$ has codimension at least two, we obtain a splitting of the tangent bundle T_X defining two algebraically integrable foliations. Now standard arguments for manifolds with split tangent bundle yield the theorem.

The technical core of this paper is to study the direct image sheaves on the non-compact quasi-projective set $Y_0 \subset Y$, making only assumptions of a birational

nature on the (relative) anticanonical divisor. This setup has not been studied in earlier papers, we therefore establish some basic results in Section 2 (and the appendix) before applying them to our situation in Subsection 3.B.

Acknowledgements. It is a pleasure to thank S. Boucksom, J.-P. Demailly and M. Păun for very helpful discussions. In particular, Lemma 2.9 comes from a nice idea of J.-P. Demailly. A part of the article was done during the first-named author's visit to the Institut for Mathematical Sciences of National University of Singapore for the program "Complex Geometry, Dynamical Systems and Foliation Theory". He is very grateful to the organizers for the invitation and the support. This work was partially supported by the Agence Nationale de la Recherche grant project Foliage¹ and the Agence Nationale de la Recherche grant "Convergence de Gromov-Hausdorff en géométrie kählérienne" (ANR-GRACK).

Notation and basic results

For general definitions we refer to [Har77, KK83, Dem12]. We use the terminology of [Deb01] for birational geometry and [Laz04a] for (algebraic) notions of positivity. Manifolds and varieties will always be supposed to be irreducible. A fibration is a proper surjective map with connected fibers $\varphi : X \rightarrow Y$ between normal complex spaces.

Let us recall some basic facts about reflexive sheaves:

1.5. Proposition.

- 1) [Har80, Cor.1.4] *Let M be a manifold, and let \mathcal{F} be a reflexive sheaf on M . Then there exists a subset Z of codimension at least three such that $\mathcal{F} \otimes \mathcal{O}_{M \setminus Z}$ is locally free.*
- 2) [Har80, Cor.1.7] *Let $\varphi : M \rightarrow N$ be a proper, equidimensional, dominant morphism between normal varieties. If \mathcal{F} is a reflexive sheaf on M , then $\varphi_*(\mathcal{F})$ is a reflexive sheaf on N .*
- 3) [Har80, Prop.1.6] *Let M be a normal variety, and let \mathcal{F}_1 and \mathcal{F}_2 be reflexive sheaves on M . Suppose that there exists a closed subset $Z \subset M$ of codimension at least two such that $\mathcal{F}_1 \otimes \mathcal{O}_{M \setminus Z} \simeq \mathcal{F}_2 \otimes \mathcal{O}_{M \setminus Z}$. Then we have $\mathcal{F}_1 \simeq \mathcal{F}_2$.*

2. POSITIVITY OF VECTOR BUNDLES

2.A. Preliminaries. We first recall the definition of possibly singular Hermitian metrics on vector bundles and some basic properties. We refer to [BP08, PT14, Rau15, Pău16, HPS16] for more details. Let $E \rightarrow X$ be a holomorphic vector bundle of rank r on a complex manifold X . We denote by

$$H_r := \{A = (a_{i,\bar{j}})\}$$

the set of $r \times r$, semi-positive definite Hermitian matrices. Let \overline{H}_r be the space of semi-positive, possibly unbounded Hermitian forms on \mathbb{C}^r . A possibly singular Hermitian metric h on E is locally given by a measurable map with values in \overline{H}_r such that

$$(1) \quad 0 < \det h < +\infty \quad \text{almost everywhere.}$$

In the above definition, a matrix valued function $h = (h_{i,\bar{j}})$ is measurable provided that all entries $h_{i,\bar{j}}$ are measurable.

¹ANR-16-CE40-0008

2.1. Definition. Let X be a (not necessarily compact) complex manifold, and let (E, h) be a holomorphic vector bundle on X with a possibly singular hermitian metric h . Let θ be a smooth closed $(1, 1)$ -form on X .

We say that $i\Theta_h(E) \succcurlyeq \theta \otimes \text{Id}_{\text{End}(E)}$, if for any open set $U \subset X$ and $s \in H^0(U, E^*)$,

$$dd^c \ln |s|_{h^*} - \theta \geq 0 \quad \text{on } U \text{ in the sense of currents.}$$

We say that (E, h) is positively curved on X if $\Theta_h(E) \succcurlyeq 0$.

We introduce here a weaker notion of positivity which will be important for us.

2.2. Definition. Let X be a Kähler manifold with a Kähler metric ω_X . Let E be a holomorphic vector bundle on X . We say that E is weakly positively curved on X , if for every $\epsilon > 0$, there exists a possibly singular hermitian metric h_ϵ on E such that $i\Theta_{h_\epsilon}(E) \succcurlyeq -\epsilon\omega_X \otimes \text{Id}_{\text{End}(E)}$.

2.3. Remark.

(i): If $i\Theta_h(E) \succcurlyeq \theta \otimes \text{Id}_{\text{End}(E)}$ for some smooth form θ , by the definition of quasi-psh functions, for every subset $U \Subset X$, we have $h \geq c \cdot \text{Id}$ on U for some constant $c > 0$ depending on U .

(ii): If X is projective, by using the same argument as in [PT14, Sect 2][Pău16, Thm 2.21], we can prove that the notion of "weakly positively curved vector bundle" implies the weak positivity in the sense of Viehweg. However, the inverse seems unclear.

Now we recall two basic properties of possibly singular hermitian metrics proved in [PT14, Pău16] which are extremely useful.

2.4. Proposition. Let E be a holomorphic vector bundle on a complex manifold X and let θ be a smooth $(1, 1)$ -form on X . Let $X_0 \subset X$ be a Zariski open set and let $(E|_{X_0}, h_0)$ be a possibly singular hermitian metric h_0 defined only on $E|_{X_0}$ such that $i\Theta_{h_0}(V) \succcurlyeq \theta \otimes \text{Id}_{\text{End}(E)}$ on X_0 .

(i): If $h_0 \geq c \cdot \text{Id}$ on X_0 for some constant $c > 0$, then h_0 can be extended as a possibly singular hermitian metric h on the total space E over X such that

$$i\Theta_h(E) \succcurlyeq \theta \otimes \text{Id}_{\text{End}(E)} \quad \text{on } X.$$

(ii): If $\text{codim}_X(X \setminus X_0) \geq 2$, then h_0 can be extended as a possibly singular hermitian metric h on the total space E over X such that

$$i\Theta_h(E) \succcurlyeq \theta \otimes \text{Id}_{\text{End}(E)} \quad \text{on } X.$$

Proof. Let $s \in H^0(U, E^*)$ for some open set $U \subset X$. By assumption, $dd^c \ln |s|_{h_0^*} \geq \theta$ on $U \cap X_0$. The condition $h_0 \geq c \cdot \text{Id}$ implies that $\ln |s|_{h_0^*}$ is upper bounded on any compact set in U . By Hartogs, $\ln |s|_{h_0^*}$ can be extended as a quasi-psh function on U with $dd^c \ln |s|_{h_0^*} \geq \theta$ on U . This proves (i). As for (ii), we can directly apply Hartogs' theorem. \square

As a consequence of Proposition 2.4, we have the following important *curvature increasing* property.

2.5. Proposition. [PT14][Pău16, Lemma 2.19] *Let $\pi : E_1 \rightarrow E_2$ be a generically surjective morphism between two holomorphic vector bundles on a complex manifold X . Let h be a possibly singular metric on E_1 such that $i\Theta_h(E_1) \succeq \theta \otimes \text{Id}_{\text{End}(E_1)}$ for some smooth $(1, 1)$ -form θ on X . Then h induces a quotient metric h_{quo} on E_2 such that $i\Theta_{h_{\text{quo}}}(E_2) \succeq \theta \otimes \text{Id}_{\text{End}(E_2)}$ on X .*

In particular, if E_1 is (resp. weakly) positively curved, E_2 is also (resp. weakly) positively curved.

Proof. Let $X_0 \subset X$ be the Zariski open subset such that π is surjective over X_0 . Then h induces a quotient metric h_2 on $E_2|_{X_0}$ such that $i\Theta_{h_2}(E_2) \succeq \theta \otimes \text{Id}_{\text{End}(E_2)}$ on X_0 . Let $x \in X \setminus X_0$ be a generic point. Thanks to Remark 2.3, there exists a small neighborhood U of x such that $h \geq c \cdot \text{Id}$ on U for some constant $c > 0$. Therefore $h_2 \geq c \cdot \text{Id}$ on $U \cap X_0$. By applying Proposition 2.4, we can know that h_2 can be extended as a metric h_{quo} on the total space E_2 such that $i\Theta_{h_{\text{quo}}}(E_2) \succeq \theta \otimes \text{Id}_{\text{End}(E)}$ on X . \square

Thanks to the fundamental works like [Ber09, BP08, PT14] among many others, we know that the notion of possibly singular hermitian metrics fits well with the positivity of direct images. We will use the following version in the article.

2.6. Theorem. [PT14, Section 5] *Let $f : X \rightarrow Y$ be a fibration between two projective manifolds and let L be a line bundle on X with a possibly singular metric h_L such that $\Theta_{h_L}(L) \geq 0$. Let $m \in \mathbb{N}$ such that the multiplier ideal sheaf $\mathcal{I}(h_L^{\frac{1}{m}}|_{X_y})$ is trivial over a generic fiber X_y , namely $\int_{X_y} |e_L|_{h_L}^{\frac{2}{m}} < +\infty$, where e_L is a basis of L .*

Let Y_1 be the locally free locus of $f_(\mathcal{O}_X(mK_{X/Y} + L))$. Then the vector bundle $f_*(\mathcal{O}_X(mK_{X/Y} + L)) \otimes \mathcal{O}_{Y_1}$ admits a possibly singular hermitian metric h such that $(f_*(\mathcal{O}_X(mK_{X/Y} + L)) \otimes \mathcal{O}_{Y_1}, h)$ is positively curved on Y_1 .*

2.7. Remark. We recall briefly the idea of the proof. In fact, by using [Ber09, BP08], over some Zariski open set Y_0 of Y , we can construct a continuous metric h on $f_*(\mathcal{O}_X(mK_{X/Y} + L)) \otimes \mathcal{O}_{Y_0}$ such that it is positively curved. Thanks to the Ohsawa-Takegoshi extension theorem and the semistable reduction, we know that $h \geq c \cdot \text{Id}$ for some constant $c > 0$. The theorem is thus proved by using Proposition 2.4,

As a consequence of Theorem 2.6 and an easy trick [CP17, Lemma 5.25], we have the following variant.

2.8. Proposition. *Let $f : X \rightarrow Y$ be a fibration between two projective manifolds and let L be a line bundle on X with a possibly singular metric h_L such that $\Theta_{h_L}(L) \geq f^*\theta$ for some smooth d -closed $(1, 1)$ -form θ (not necessarily positive) on Y . Let $m \in \mathbb{N}$ such that $\mathcal{I}(h_L^{\frac{1}{m}}|_{X_y}) = \mathcal{O}_{X_y}$ for a generic fiber X_y .*

Let Y_1 be the locally free locus of $f_(\mathcal{O}_X(mK_{X/Y} + L))$. Then the vector bundle $f_*(\mathcal{O}_X(mK_{X/Y} + L)) \otimes \mathcal{O}_{Y_1}$ admits a possibly singular hermitian metric h such that*

$$i\Theta_h(f_*(\mathcal{O}_X(mK_{X/Y} + L))) \succcurlyeq \theta \otimes \text{Id}_{\text{End}(f_*(\mathcal{O}_X(mK_{X/Y} + L)))} \quad \text{on } Y_1.$$

2.B. Numerical flatness. Let X be a compact Kähler manifold and let E be a holomorphic vector bundle on X . We say that E is numerically flat if E and its dual E^* are nef (in the sense of [DPS94]), one sees easily that this is equivalent to

$$E \text{ is nef and } c_1(E) = 0 \in H^{1,1}(X, \mathbb{Q}).$$

The aim of this subsection is to give a sufficient condition for a vector bundle E with $c_1(E) = 0$ to be numerically flat.

2.9. Lemma. *Let C be a smooth curve and let E be a numerically flat vector bundle of rank n on C . Let $\pi : \mathbb{P}(E) \rightarrow C$ be the natural projection. Fix an ample line bundle A over C . Let $T_m \geq 0$ be a positive current with analytic singularities in the class of $c_1(\mathcal{O}_{\mathbb{P}(E)}(m) + \pi^*A)$ such that T_m is smooth in the neighbourhood of a general π -fibre. Set*

$$a_m := \max_{x \in \mathbb{P}(E)} \nu(T_m, x),$$

where $\nu(T_m, x)$ is the Lelong number of T_m over x . Then

$$(2) \quad \lim_{m \rightarrow 0} \frac{a_m}{m} = 0.$$

Proof. Thanks to Proposition A.2, for the current T_m , we can find a $k \in \{1, 2, \dots, n\}$ depending on m , a closed (k, k) -positive current Θ_k in the same class of $(c_1(\mathcal{O}_{\mathbb{P}(E)}(m) + \pi^*A))^k$ and a subspace Z_m of codimension k contained in some π -fibers such that

$$(3) \quad \Theta_k \geq \left(\frac{a_m}{n}\right)^k \cdot [Z_m]$$

in the sense of currents. Set $\alpha := c_1(\mathcal{O}_{\mathbb{P}(E)}(1))$, then α is a nef cohomology class. Thus (3) implies

$$(m\alpha + \pi^*c_1(A))^k \cdot \alpha^{n-k} = \Theta_k \cdot \alpha^{n-k} \geq \left(\frac{a_m}{n}\right)^k \int_{[Z_m]} \alpha^{n-k}.$$

Note that Z_m is contained in some fiber of π , we have $\int_{[Z_m]} \alpha^{n-k} \geq 1$. Therefore

$$(4) \quad (m\alpha + \pi^*c_1(A))^k \cdot \alpha^{n-k} \geq \left(\frac{a_m}{n}\right)^k$$

Since $c_1(E) = 0$, we have $\alpha^n = 0 \in H^{n,n}(\mathbb{P}(E))$. Therefore for $m \gg 0$ the left side of (4) is less or equal than $C_1 m^{k-1} + C_2$ for some uniform constants C_1 and C_2 . Then (4) implies that $\left(\frac{a_m}{n}\right)^k \leq C_1 m^{k-1} + C_2$. As $k \leq n$, we obtain finally

$$a_m \leq C_3 m^{\frac{n-1}{n}} + C_4$$

for some uniform constants C_3 and C_4 independent of m . This proves (2). \square

The following lemma is essentially shown in [PT14, Prop 2.3.5][Pău16, Thm 2.21].

2.10. Lemma. *Let X be a n -dimensional projective manifold and let E be a holomorphic vector bundle on X . Let $\pi : \mathbb{P}(E) \rightarrow X$ be the natural projection and let ω be a Kähler metric on $\mathbb{P}(E)$. Then the following three conditions are equivalent:*

(i) *There exists a very ample line bundle A on X such that the restriction*

$$H^0(\mathbb{P}(E), \mathcal{O}_{\mathbb{P}(E)}(m) + \pi^*A) \rightarrow H^0(\mathbb{P}(E)_x, \mathcal{O}_{\mathbb{P}(E)}(m) + \pi^*A)$$

is surjective over a generic point $x \in X$ for $m \gg 1$.

(ii) $\pi_*(\mathbb{B}_-(\mathcal{O}_{\mathbb{P}(E)}(1))) \subsetneq X$.²

(iii) For every $\epsilon > 0$, there exists a possibly singular metric h_ϵ on $\mathcal{O}_{\mathbb{P}(E)}(1)$ such that $i\Theta_{h_\epsilon}(\mathcal{O}_{\mathbb{P}(E)}(1)) \geq -\epsilon\omega$ and the multiplier ideal $\mathcal{I}(h_\epsilon^{\otimes m}|_{\mathbb{P}(E)_x})$ of the restriction of $h_\epsilon^{\otimes m}$ to the generic fiber $\mathbb{P}(E)_x$ is trivial for every $m \in \mathbb{N}$.

Proof. (i) \implies (ii): It comes from the definition.

(ii) \implies (iii): If $\pi_*(\mathbb{B}_-(\mathcal{O}_{\mathbb{P}(E)}(1))) \subsetneq X$, for every $\epsilon > 0$, we have

$$\pi_*(\mathbb{B}(\mathcal{O}_{\mathbb{P}(E)}(1) + \epsilon A)) \subsetneq X,$$

where A is some ample line bundle on $\mathbb{P}(E)$. Then for some m large enough, $\pi_*(\text{Bs}(\mathcal{O}_{\mathbb{P}(E)}(m) + \epsilon mA)) \subsetneq X$. Therefore, a basis of $H^0(\mathbb{P}(E), \mathcal{O}_{\mathbb{P}(E)}(m) + \epsilon mA)$ induces a metric h_ϵ on $\mathcal{O}_{\mathbb{P}(E)}(1)$ with analytic singularities such that

$$(5) \quad i\Theta_{h_\epsilon}(\mathcal{O}_{\mathbb{P}(E)}(1)) \geq -\epsilon\omega_A$$

and h_ϵ is smooth on the generic fiber of π . (iii) is proved.

(iii) \implies (i): Since $\mathcal{O}_{\mathbb{P}(E)}(1)$ is relatively ample, there exists some $m_0 \in \mathbb{N}$ and some ample line bundle A_1 on X such that $-K_{\mathbb{P}(E)} + \mathcal{O}_{\mathbb{P}(E)}(m_0) + \pi^*A_1$ is ample. Let A_2 be a sufficiently ample line bundle on X . For every $m \in \mathbb{N}$ and ϵ small enough, since $\mathcal{I}(h_\epsilon^{\otimes m}|_{\mathbb{P}(E)_x})$ is trivial for a generic $x \in X$, Ohsawa-Takegoshi extension theorem implies that

$$\begin{aligned} & H^0(\mathbb{P}(E), K_{\mathbb{P}(E)} + (-K_{\mathbb{P}(E)} + \mathcal{O}_{\mathbb{P}(E)}(m_0) + \pi^*A_1) + \mathcal{O}_{\mathbb{P}(E)}(m) + \pi^*A_2) \\ & \rightarrow H^0(\mathbb{P}(E)_x, K_{\mathbb{P}(E)} + (-K_{\mathbb{P}(E)} + \mathcal{O}_{\mathbb{P}(E)}(m_0) + \pi^*A_1) + \mathcal{O}_{\mathbb{P}(E)}(m) + \pi^*A_2) \end{aligned}$$

is surjective for a generic $x \in X$. This proves (i). \square

We can now prove the main result in this subsection:

2.11. Proposition. *Let X be a n -dimensional projective manifold and let E be a holomorphic vector bundle on X with $c_1(E) = 0$ and $\pi_*(\mathbb{B}_-(\mathcal{O}_{\mathbb{P}(E)}(1))) \subsetneq X$. Then E is numerically flat.*

In particular, let E be a vector bundle on X . If E is weakly positively curved on X and $c_1(E) = 0$, then E is numerically flat.

Proof. Let B be an arbitrary curve in $\mathbb{P}(E)$ such that $B \rightarrow \pi(B)$ is a finite morphism. We are done if we prove that

$$c_1(\mathcal{O}_{\mathbb{P}(E)}(1)) \cdot B \geq 0.$$

Step 1: The current T_m . Let A be a sufficiently ample line bundle on X and $\{s_1, \dots, s_{k_m}\}$ be a basis of $H^0(\mathbb{P}(E), \mathcal{O}_{\mathbb{P}(E)}(m) + \pi^*A)$ and set $T_m := dd^c \ln \sum_{i=1}^{k_m} |s_i|^2$. Then $T_m \geq 0$ is a current in the class of $c_1(\mathcal{O}_{\mathbb{P}(E)}(m) + \pi^*A)$. Set $a_m := \nu(T_m, B)$ for the Lelong number of T_m over a generic point of B . We will prove that

$$(6) \quad \lim_{m \rightarrow +\infty} \frac{a_m}{m} = 0,$$

Note first that, thanks to Lemma 2.10, T_m is smooth on the generic fiber of π . Let $C \subset X$ be a curve defined by a complete intersection of general elements of $|A|$ such

²Recall that, for a line bundle L , $\mathbb{B}(L)$ is the stable base locus of L , namely $\mathbb{B}(L) = \bigcap_{m \geq 1} \text{Bs}(mL)$. $\mathbb{B}_-(L) := \mathbb{B}(L + \epsilon A)$ for any ample A and sufficiently small $\epsilon > 0$.

that $\pi_*(B) \cap C$ is a finite, non-empty set. Since C is general, the restricted bundle $E|_C$ is weakly positively curved and $c_1(E|_C) = 0$. As $\dim C = 1$, the bundle $E|_C$ is numerically flat.

We consider the restriction of T_m on $\pi^{-1}(C)$. Thanks to Lemma 2.10, $T_m|_{\pi^{-1}(C)}$ is smooth on the fiber $\mathbb{P}(E)_x$, for a generic point $x \in C$. We can thus apply Lemma 2.9 to $(E|_C, T_m|_{\pi^{-1}(C)})$ and obtain

$$(7) \quad \lim_{m \rightarrow +\infty} \frac{\nu(T_m|_{\pi^{-1}(C)}, x)}{m} = 0,$$

for any point $x \in B \cap \pi^{-1}(C)$. Together with the fact that $\nu(T_m|_{\pi^{-1}(C)}, x) \geq \nu(T_m, B)$ for every $x \in B$ we obtain (6).

Step 2: Final conclusion. Fix a Kähler metric ω on $\mathbb{P}(E)$. Thanks to [Dem92, thm 1.1], we can find a current \tilde{T}_m in the same class of T_m such that

$$\tilde{T}_m \geq -Ca_m \cdot \omega \quad \text{and} \quad \nu(\tilde{T}_m, x) = \max(\nu(T_m, x) - a_m, 0)$$

for every $x \in \mathbb{P}(E)$, where C is a uniform constant. In particular, $\nu(\tilde{T}_m, B) = 0$. Therefore

$$c_1(\mathcal{O}_{\mathbb{P}(E)}(m) + \pi^*A) \cdot B = \tilde{T}_m \cdot B \geq -Ca_m \int_B \omega.$$

Combining this with (6), by letting $m \rightarrow +\infty$, we get $c_1(\mathcal{O}_{\mathbb{P}(E)}(1)) \cdot B \geq 0$.

For the second part of the proposition, if E is weakly positively curved, for every $\epsilon > 0$, we can find a possible singular metric $h_{\epsilon,1}$ such that

$$i\Theta_{h_{\epsilon,1}}(E) \succcurlyeq -\epsilon\omega_X \otimes \text{Id}_{\text{End}(E)}.$$

Then $h_{\epsilon,1}$ induces a possibly singular metric h_ϵ on $\mathcal{O}_{\mathbb{P}(E)}(1)$ such that $i\Theta_{h_\epsilon}(\mathcal{O}_{\mathbb{P}(E)}(1)) \geq -\epsilon\pi^*\omega_X$. Moreover, the condition (1) implies that h_ϵ is bounded over of generic fiber of π . In particular, $\mathcal{I}(h_\epsilon^{\otimes m}|_{\mathbb{P}(E)_x})$ is trivial for every $m \in \mathbb{N}$, where $\mathbb{P}(E)_x$ is a generic fiber. Thanks to Lemma 2.10, we have $\pi_*(\mathbb{B}_-(\mathcal{O}_{\mathbb{P}(E)}(1))) \subsetneq X$. Therefore E is numerically flat. \square

As a corollary, we obtain

2.12. Corollary. *Let S be a smooth projective surface, and let $Z \subset S$ be a finite set. Let V be a vector bundle on $S \setminus Z$ such that $c_1(V) = 0$ and V is weakly positively curved on $S \setminus Z$. Denote by $j : S \setminus Z \rightarrow S$ the inclusion and set $\tilde{V} := (j_*V)^{**}$.*

Then \tilde{V} is a numerically flat vector bundle.

Proof. The sheaf \tilde{V} is reflexive, hence locally free on the surface S . Moreover $c_1(\tilde{V}) = 0$, since Z has codimension two. By Proposition 2.4, \tilde{V} is also weakly positively curved on S . By Proposition 2.11 it is thus numerically flat. \square

3. SIMPLY CONNECTED PROJECTIVE MANIFOLDS WITH NEF ANTICANONICAL
BUNDLE

This whole section is devoted to the proof of Theorem 1.2. After fixing the notation for the whole section, we establish some basic geometric properties of the ‘universal family’ $\Gamma \rightarrow Y$ in Subsection 3.A. The results and proofs are already contained in [Zha05], but we give the details for the convenience of the reader. In Subsection 3.B contains the proof of the key technical result on direct image sheaves. Finally, in Subsection 3.C we use these results to describe first the birational structure of $\Gamma \rightarrow Y$ and translate it later into a biregular statement for X .

3.1. Setup. *Let X be a uniruled projective manifold such that $-K_X$ nef and $\pi_1(X) = 1$. There exists a unique irreducible component of the Chow space $\mathcal{C}(X)$ such that a very general point corresponds to a very general fibre of the MRC fibration. We denote by Y a resolution of singularities of this irreducible component, and by $f : X' \rightarrow Y$ the normalisation of the pull-back of the universal family over it. Let $p : X' \rightarrow X$ be the natural map, then p is birational and, since the MRC fibration is almost holomorphic, the p -exceptional locus does not dominate Y .*

By [Zha05, Cor.1] the Kodaira dimension $\kappa(Y)$ is equal to zero, and we denote by N_Y an effective \mathbb{Q} -Cartier divisor on Y such that $N_Y \sim_{\mathbb{Q}} K_Y$.

Let A be a very ample line bundle on X . Up to replacing A by some multiple, we can suppose that

$$(8) \quad \text{Sym}^p H^0(X_y, A) \rightarrow H^0(X_y, pA)$$

is surjective for every $p \in \mathbb{N}$ where $y \in Y$ is a general point.

*Since X' is obtained by base change from a universal family of cycles, the pull-back p^*A is globally generated and ample on every fibre of f .*

*Choose now a resolution of singularities $\mu : \Gamma \rightarrow X'$ and denote by $\varphi : \Gamma \rightarrow Y$ and $\pi : \Gamma \rightarrow X$ the induced maps. Set $L := \pi^*A$ and let r be the rank of $\varphi_*(\mathcal{O}_{\Gamma}(L))$.*

$$\begin{array}{ccccc}
 & & \pi & & \\
 & & \curvearrowright & & \\
 \Gamma & \xrightarrow{\mu} & X' & \xrightarrow{p} & X \\
 & \searrow \varphi & \downarrow f & \swarrow & \\
 & & Y & &
 \end{array}$$

Since X is \mathbb{Q} -factorial, we know that the π -exceptional locus has pure codimension one, and we denote it by E .

*Let $Y_0 \subset Y$ be the largest Zariski open subset such that $\Gamma_0 := \varphi^{-1}(Y_0) \rightarrow Y_0$ is equidimensional and for every prime divisor $B \subset Y_0$, the pull-back φ^*B is not contained in the π -exceptional locus E . Denote by $P \subset Y$ the largest reduced divisor that is contained in $Y \setminus Y_0$.*

3.A. Birational geometry of the universal family. The following basic statement is a birational variant of the situation in [LTZZ10], the proof follows their strategy.

3.2. Lemma. *Let X be a projective manifold such that $-K_X$ is nef. Let $\psi : Z \rightarrow X$ be a generically finite morphism from a projective manifold Z , and let $\varphi' : Z \rightarrow Y'$ be a fibration onto a projective manifold Y' . Suppose that we have*

$$K_{Z/Y'} \sim_{\mathbb{Q}} \psi^* K_X + E_1 - E_2$$

where E_1 and E_2 are effective \mathbb{Q} -divisors such that $\text{Supp}(E_1 + E_2)$ does not surject onto Y' . Then the following holds:

If $\text{Supp}(E_1)$ is ψ -exceptional, then $\text{Supp}(E_2)$ is ψ -exceptional.

Proof. Fix a very ample divisor H on X . We argue by contradiction and suppose that E_2 is not ψ -exceptional.

We claim that $E_1 - E_2$ is pseudoeffective. Assuming this for the time being let us see how to conclude: and let $C \subset Z$ be a curve that is a complete intersection of pull-backs of general elements of $|H|$. Since $\text{Supp}(E_1)$ is ψ -exceptional, we have $E_1 \cdot C = 0$. Since the support of $-E_2$ is not ψ -exceptional and C is cut out by pull-backs of ample divisors, we have $(E_1 - E_2) \cdot C = -E_2 \cdot C < 0$, a contradiction.

Proof of the claim: It is sufficient to show that

$$K_{Z/Y'} + \psi^*(-K_X + \delta H) \sim_{\mathbb{Q}} E_1 - E_2 + \psi^*(\delta H)$$

is pseudoeffective for all $\delta > 0$. Since $\text{Supp}(E_1 + E_2)$ does not surject onto Y' , we know that K_Z and $\psi^*(K_X)$ coincide along the general φ' -fibre. Thus, since ψ is generically finite, the \mathbb{Q} -divisor $K_{Z/Y'} + \psi^*(-K_X + \delta H)$ is big on the general φ' -fibre. In particular for a sufficiently divisible $m \gg 0$ the higher direct images

$$\varphi'_*(\mathcal{O}_Z(m(K_{Z/Y'} + \psi^*(-K_X + \delta H))))$$

are not zero sheaves. Moreover since H is ample and $\delta > 0$ the divisor $\psi^*(-K_X + \delta H)$ is semiample. Thus we have

$$\psi^*(-K_X + \delta H) \sim_{\mathbb{Q}} B_\delta$$

where B_δ is an effective \mathbb{Q} -divisor such that the pair (Z, B_δ) is klt. By [Cam04, Thm.4.13] the direct images $\varphi'_*(\mathcal{O}_Z(m(K_{Z/Y'} + B_\delta)))$ are weakly positive. Since the relative evaluation map

$$(\varphi')^* \varphi'_*(\mathcal{O}_Z(m(K_{Z/Y'} + B_\delta))) \rightarrow \mathcal{O}_Z(m(K_{Z/Y'} + B_\delta))$$

is generically surjective, this shows that $m(K_{Z/Y'} + B_\delta)$ is weakly positive, hence pseudoeffective. \square

3.3. Proposition. *In the situation of Setup 3.1, the following holds:*

- 1) We have $\text{Supp}(N_Y) \subset P$.
- 2) Every irreducible component of $\pi(\varphi^{-1}(Y \setminus Y_0))$ has codimension at least two. In particular every φ -exceptional divisor is π -exceptional.
- 3) Let $B \subset Y_0$ be a prime divisor, and let $\varphi^*B = \sum_{j=1}^l m_j B_j$ be the decomposition in prime divisors. If $m_j > 1$ then B_j is π -exceptional.

Proof. Denote by $(E_j)_{j=1, \dots, l}$ the irreducible components of the π -exceptional locus E . The map π is birational and X is smooth, so we have

$$K_\Gamma \sim \pi^*K_X + \sum_{j=1}^l a_j E_j$$

where $a_j > 0$ for all j . Since the MRC fibration is almost holomorphic, we know that $E = \text{Supp}(\sum_{j=1}^l a_j E_j)$ does not surject onto Y .

Proof of 1) We have

$$(9) \quad K_{\Gamma/Y} \sim_{\mathbb{Q}} \pi^* K_X + \sum_{j=1}^l a_j E_j - \varphi^* N_Y,$$

so by Lemma 3.2 the divisor $\varphi^* N_Y$ is π -exceptional. By the definition of Y_0 this implies that $\text{Supp}(N_Y) \subset P$.

Proof of 2) Assume that there exists a prime divisor $D \subset \Gamma$ that is not π -exceptional and such that $\varphi(D) \subset Y \setminus Y_0$. By the definition of Y_0 this implies that $\varphi(D)$ has codimension at least two. Denote by $\tau : Y' \rightarrow Y$ the composition of the blowup of $\varphi(D)$ with a resolution of singularities. Since Y is smooth, we have

$$K_{Y'} \sim \tau^* K_Y + E_{Y'} \sim_{\mathbb{Q}} \tau^* N_Y + E_{Y'}$$

with $E_{Y'}$ an effective divisor whose support is equal to the τ -exceptional locus. In particular $\tau(\text{Supp } E_{Y'})$ contains $\varphi(D)$.

Let now $\tau' : Z \rightarrow \Gamma$ be a resolution of indeterminacies of $\Gamma \dashrightarrow Y'$ such that Z is smooth, and denote by $\varphi' : Z \rightarrow Y'$ the induced fibration. Let $D' \subset Z$ be the strict transform of D . By construction we have $\varphi'(D') \subset \text{Supp}(\tau^* N_Y + E_{Y'})$. Since $\tau^* N_Y + E_{Y'}$ is a canonical divisor, we can argue as in the proof of 1) to see that the support of $(\varphi')^*(\tau^* N_Y + E_{Y'})$ is $\pi \circ \tau'$ -exceptional. Thus D' is $\pi \circ \tau'$ -exceptional. Since D' is not τ' -exceptional, the divisor $D = \tau'(D')$ is π -exceptional, a contradiction.

Proof of 3) By [Laz04a, Prop.4.1.12] we can choose a generically finite covering $\tau' : Y' \rightarrow Y$ by some projective manifold Y' that ramifies with multiplicity m_j over B and over every other prime divisor contained in the branch locus the general φ -fibre is smooth. The fibre product $\Gamma \times_Y Y'$ is not normal along a divisor mapping onto B_j , and we denote by Z its normalisation. By [Rei94, Prop.2.3] we have

$$(10) \quad K_{Z/Y'} = (\tau')^* K_{\Gamma/Y} - A + C_1 - C_2$$

where $\tau' : Z \rightarrow \Gamma$ is the natural map, the divisor A is effective s.t. $B_j \subset \tau'(A)$ and C_1 and C_2 are effective divisors taking into account that the base change formula for relative canonical sheaves does not hold over the non-flat locus. In particular $\tau'(\text{Supp}(C_1 - C_2))$ is contained in φ -exceptional divisors, hence $\text{Supp}(C_1 - C_2)$ is $\pi \circ \tau'$ -exceptional by 2). These properties do not change if we replace Z by a resolution of singularities, so we suppose without loss of generality that Z is smooth. Combining (10) and (9) we obtain

$$K_{Z/Y'} = (\tau')^* \pi^* K_X + \pi^* \left(\sum a_j E_j \right) + C_1 - (\varphi^* N_Y + A + C_2).$$

By what precedes the support of $\pi^* \left(\sum a_j E_j \right) + C_1$ is $\pi \circ \tau'$ -exceptional, so the support of $\varphi^* N_Y + A + C_2$ is $\pi \circ \tau'$ -exceptional by Lemma 3.2. Since $\tau'(A)$ contains B_j , this implies that B_j is π -exceptional. \square

3.4. Remark. The following elementary observation will be quite useful: any holomorphic function $s : Y_0 \rightarrow \mathbb{C}$ is constant. Indeed the pull-back $\varphi^* s$ is holomorphic on $(\Gamma \setminus E) \cap \varphi^{-1}(Y_0)$ which we identify to a Zariski open subset of X . By Proposition 3.3, 2) the complement of this set has codimension at least two, so $\varphi^* s$ extends to X . Thus it is constant.

Since the rank of a linear map is determined by the nonvanishing of its minors, this implies the following: let

$$\mathcal{O}_{Y_0}^{\oplus r_1} \rightarrow \mathcal{O}_{Y_0}^{\oplus r_2}$$

be a morphism between trivial vector bundles on Y_0 . Then the rank of this map is constant.

3.B. Triviality of direct images. Following the terminology in Setup 3.1, the aim of this subsection is to prove that, for m large enough,

$$\mathcal{V}_p := \varphi_*(\mathcal{O}_\Gamma(pL + pmE - \frac{p}{r}\varphi^*\det\varphi_*(\mathcal{O}_\Gamma(L + mE))))$$

is a trivial vector bundle over Y_0 for every $p \in \mathbb{N}$ sufficiently divisible.

We first sketch the idea of the proof and describe the outline of the subsection. Our argument is very close to [Cao16]. In fact, thanks to Proposition 3.3, we know that $-K_{\Gamma/Y}$ can be written as the sum of a nef line bundle and a divisor supported in the exceptional locus of π . Then most arguments in [Cao16] still work for our case cf. Lemma 3.5 and Lemma 3.6.

The main difference is that, because of the exceptional locus $E \cap \varphi^{-1}(Y_0)$, we can à priori only get the flatness of \mathcal{V}_p over $Y_0 \setminus \varphi_*(E)$. To overcome this difficulty, we need a new observation: Lemma 3.7, namely, for m large enough, there is an isomorphism over Y_0 :

$$(11) \quad (\det\varphi_*(\mathcal{O}_\Gamma(L + mE))) \otimes \mathcal{O}_{Y_0} \simeq (\det\varphi_*(\mathcal{O}_\Gamma(L + (m+1)E))) \otimes \mathcal{O}_{Y_0}.$$

Together with Lemma 3.5 and Lemma 3.6, we prove in Proposition 3.9 that \mathcal{V}_p is weakly positively curved over Y_0 and $\pi_*\varphi^*c_1(\mathcal{V}_p) = 0$. Finally, thanks to Corollary 2.12 and the property of Y_0 (cf. Proposition 3.3), we prove in Proposition 3.10 that \mathcal{V}_p is a trivial vector bundle on Y_0 .

Before proving the main proposition 3.10 in this subsection, we need a series of technical lemmas. The first one comes from Viehweg's diagonal trick [Vie95, Thm 6.24] and Q. Zhang's argument [Zha05]. We refer to [Cao16, Prop 3.12] [CP17, Thm 3.13] for similar argument.

3.5. Lemma. *In the situation of Setup 3.1, for every $m \in \mathbb{N}$,*

$$(12) \quad A - \frac{1}{r}\pi_*\varphi^*c_1(\varphi_*(\mathcal{O}_\Gamma(L + mE))) \quad \text{is pseudo-effective on } X.$$

Proof. Let $E' \subset \Gamma$ be the union of non reduced and non flat locus of φ . Let $\Gamma^{(r)}$ be a desingularisation of the r -times φ -fibrewise product $\Gamma \times_Y \cdots \times_Y \Gamma$, and let $\text{pr}_i : \Gamma^{(r)} \rightarrow \Gamma$ be the i -th directional projection. Let $\varphi_r : \Gamma^{(r)} \rightarrow Y$ be the natural fibration. Set $L^{(r)} := \sum_{i=1}^r \text{pr}_i^*L$ and $E^{(r)} := \sum_{i=1}^r \text{pr}_i^*E$. We have a natural morphism

$$(13) \quad \det\varphi_*(\mathcal{O}_\Gamma(L + mE)) \rightarrow (\varphi_r)_*(\mathcal{O}_{\Gamma^{(r)}}(L^{(r)} + mE^{(r)} + \Delta))$$

for some divisor Δ supported in $\sum_{i=1}^r \text{pr}_i^*(E')$. Set

$$L' := L^{(r)} + mE^{(r)} + \Delta - \varphi_r^*c_1(\varphi_*(\mathcal{O}_\Gamma(L + mE))).$$

As the restriction of (13) on a generic point $y \in Y$ is non zero, (13) induces a nontrivial section in $H^0(\Gamma^{(r)}, L')$. Therefore L' is an effective line bundle on $\Gamma^{(r)}$.

We now compare $K_{\Gamma^{(r)}/Y}$ with $\sum_{i=1}^r \text{pr}_i^*(\pi^* K_X)$. Let $\varphi^{(r)} : \Gamma^{(r)} \rightarrow Y$ be the natural morphism. Then

$$K_{\Gamma^{(r)}/Y} - \sum_{i=1}^r \text{pr}_i^*(\pi^* K_X) = -(\varphi^{(r)})^* K_Y + (K_{\Gamma^{(r)}} - \sum_{i=1}^r \text{pr}_i^* K_\Gamma) + \sum_{i=1}^r \text{pr}_i^*(K_\Gamma - \pi^* K_X)$$

By Proposition 3.3, $(\varphi^{(r)})^* K_Y$ is equivalent to a divisor supported in $\sum_{i=1}^r \text{pr}_i^* E$. By construction, $K_{\Gamma^{(r)}} - \sum_{i=1}^r \text{pr}_i^* K_\Gamma$ is supported in $\sum_{i=1}^r \text{pr}_i^* E'$, and $\sum_{i=1}^r \text{pr}_i^*(K_\Gamma - \pi^* K_X)$ is supported in $\sum_{i=1}^r \text{pr}_i^* E$. As a consequence, we can find a divisor (not necessary effective) Δ' supported in $\sum_{i=1}^r \text{pr}_i^*(E' + E)$ such that

$$-K_{\Gamma^{(r)}/Y} + \Delta' = -\sum_{i=1}^r \text{pr}_i^*(\pi^* K_X).$$

In particular, $-K_{\Gamma^{(r)}/Y} + \Delta'$ is nef.

By using [Cao16, Prop 2.10], there exists an ample line bundle A_Y on Y such that for every $p \in \mathbb{N}$ and for q large enough (with respect to p),

$$\begin{aligned} & H^0(\Gamma^{(r)}, pqK_{\Gamma^{(r)}/Y} + pq(-K_{\Gamma^{(r)}/Y} + \Delta') + pL' + \varphi_r^* A_Y) \\ & \rightarrow H^0(\Gamma_y^{(r)}, pqK_{\Gamma^{(r)}/Y} + pq(-K_{\Gamma^{(r)}/Y} + \Delta') + pL' + \varphi_r^* A_Y) \end{aligned}$$

is surjective for a generic $y \in Y$.

By restricting the above morphism on diagonal³, we can find some effective divisor $F_{p,q}$ (depending on p and q) supported in $E \cup E'$ such that

$$\begin{aligned} & H^0(\Gamma, F_{p,q} + prL - p\varphi^* c_1(\varphi_*(\mathcal{O}_\Gamma(L + mE))) + \varphi^* A_Y) \\ & \rightarrow H^0(\Gamma_y, F_{p,q} + prL - p\varphi^* c_1(\varphi_*(\mathcal{O}_\Gamma(L + mE))) + \varphi^* A_Y) \end{aligned}$$

is surjective. In particular, $F_{p,q} + prL - p\varphi^* c_1(\varphi_*(\mathcal{O}_\Gamma(L + mE))) + \varphi^* A_Y$ is effective. Thanks to Proposition 3.3, $\pi_*(F_{p,q}) \subset \pi_*(E \cup E')$ has codimension at least two. Therefore

$$\pi_* \left(c_1 \left(rL - \varphi^* c_1(\varphi_*(\mathcal{O}_\Gamma(L + mE))) + \frac{1}{p} \varphi^* A_Y \right) \right)$$

is pseudoeffective on X . The lemma is proved by letting $p \rightarrow +\infty$. \square

The second lemma is very close in spirit to Lemma 3.2, namely, the proof uses the positivity of direct images as well as Q. Zhang's technique [Zha05] in the study of manifolds with nef anticanonical bundles.

3.6. Lemma. *Let E_1 be an effective \mathbb{Q} -divisor satisfying $\varphi_*(E_1) \subsetneq Y$ and let L_Y be a \mathbb{Q} -line bundle on Y . Let $a > 0$ be some constant such that the line bundle*

$$L_1 := aL + E_1 + \varphi^* L_Y$$

³It is only well defined on $\Gamma \setminus E'$, but by adding some divisor supported in E' , we can extend it on Γ .

is pseudo-effective. Then for $c \in \mathbb{N}$ large enough, $\varphi_*(L_1 + cE)$ is weakly positive curved (cf. Definition 2.2) over Y_1 , where Y_1 is the locally free locus of $\varphi_*(L_1 + cE)$. In particular, $\det \varphi_*(L_1 + cE)$ is pseudoeffective on Y .

Proof. By construction, we can find some effective divisor E_2 supported in E such that $-K_\Gamma + E_2 = -\pi^*K_X$. Let A_Y be an ample line bundle on Y such that $A_Y + L_Y$ is still ample.

For every $\epsilon \in \mathbb{Q}^+$,

$$-K_\Gamma + E_2 + \epsilon aL + \epsilon \varphi^*(L_Y + A_Y) = \pi^*(-K_X + \epsilon aA) + \epsilon \varphi^*(L_Y + A_Y)$$

is semi-ample. Note that $\epsilon E_1 + \varphi^*K_Y \sim_{\mathbb{Q}} D$ for some \mathbb{Q} -effective divisor D satisfying $\varphi_*(D) \subsetneq Y$. Then the \mathbb{Q} -line bundle

$$-K_{\Gamma/Y} + E_2 + \epsilon L_1 = (-K_\Gamma + E_2 + \epsilon aL + \epsilon \varphi^*(L_Y + A_Y)) + (\epsilon E_1 + \varphi^*K_Y) - \epsilon \varphi^*A_Y.$$

can be equipped with a metric $h_{1,\epsilon}$ such that

$$i\Theta_{h_{1,\epsilon}}(-K_{\Gamma/Y} + E_2 + \epsilon L_1) \geq -\epsilon \varphi^*\omega_Y$$

and $h_{1,\epsilon}|_{\Gamma_y}$ is smooth for a generic fiber Γ_y , where ω_Y is a Kähler metric representing A_Y .

Since L_1 is pseudo-effective, there exists a possibly singular metric h on L_1 such that $i\Theta_h(L_1) \geq 0$. Then we can find a m (depends only on h) such that $\mathcal{I}(h^{\frac{1}{m}}|_{\Gamma_y}) = \mathcal{O}_{\Gamma_y}$ for a generic fiber Γ_y . Therefore, for every $0 < \epsilon < \frac{1}{m}$, the metric $h_\epsilon := mh_{1,\epsilon} + (1 - m\epsilon)h$ defines a metric on

$$-mK_{\Gamma/Y} + mE_2 + L_1 = m(-K_{\Gamma/Y} + E_2 + \epsilon L_1) + (1 - m\epsilon)L_1$$

with $i\Theta_{h_\epsilon}(-mK_{\Gamma/Y} + mE_2 + L_1) \geq -m\epsilon \varphi^*\omega_Y$ on Γ and $\mathcal{I}(h_\epsilon^{\frac{1}{m}}|_{\Gamma_y}) = \mathcal{O}_{\Gamma_y}$.

By applying Proposition 2.8 to the line bundle $(-mK_{\Gamma/Y} + mE_2 + L_1, h_\epsilon)$, we can find a metric \tilde{h}_ϵ on

$$\varphi_*(\mathcal{O}_\Gamma(mE_2 + L_1)) = \varphi_*(\mathcal{O}_\Gamma(mK_{\Gamma/Y} + (-mK_{\Gamma/Y} + mE_2 + L_1)))$$

such that

$$(14) \quad i\Theta_{\tilde{h}_\epsilon}(\varphi_*(\mathcal{O}_\Gamma(mE_2 + L_1))) \succeq -m\epsilon \omega_Y \otimes \text{Id}_{\text{End}(\varphi_*(\mathcal{O}_\Gamma(mE_2 + L_1)))} \quad \text{on } Y_{1,m},$$

where $Y_{1,m}$ is the locally free locus of $\varphi_*(\mathcal{O}_\Gamma(mE_2 + L_1))$.

As m depends only on h , we can find a constant c independent of ϵ such that $cE \geq mE_2$. Then (14) implies the existence of possible singular metrics $\tilde{h}_{c,\epsilon}$ such that $i\Theta_{\tilde{h}_{c,\epsilon}}(\varphi_*(\mathcal{O}_\Gamma(cE + L_1))) \succeq -m\epsilon \omega_Y \otimes \text{Id}_{\text{End}(\varphi_*(\mathcal{O}_\Gamma(cE + L_1)))}$ on Y_1 . The first part of the lemma is thus proved.

The second part of lemma comes from the similar argument in [Pău16, Cor 2.26, first part]. \square

Before proving the main proposition in this subsection, we need two more lemmas. The first lemma is to compare $\varphi_*\mathcal{O}_\Gamma(L + mE)$ with $\varphi_*\mathcal{O}_\Gamma(L + (m+1)E)$ over Y_0 :

3.7. Lemma. *In the situation of Setup 3.1, there exists a $m_0 \in \mathbb{N}$ such that for all $m \geq m_0$ the natural inclusions*

$$(15) \quad \det \varphi_*(\mathcal{O}_\Gamma(L + mE)) \rightarrow \det \varphi_*(\mathcal{O}_\Gamma(L + (m+1)E))$$

induce an isomorphism

$$(16) \quad (\det \varphi_*(\mathcal{O}_\Gamma(L + mE))) \otimes \mathcal{O}_{Y_0} \simeq (\det \varphi_*(\mathcal{O}_\Gamma(L + (m+1)E))) \otimes \mathcal{O}_{Y_0}.$$

Proof. Note that, the effective divisor

$$c_1(\varphi_*(\mathcal{O}_\Gamma(L + (m+1)E))) - c_1(\varphi_*(\mathcal{O}_\Gamma(L + mE)))$$

is supported in $\varphi_*(E)$, which is a fixed strict subvariety of Y independent of m .

If we assume by contradiction that (16) is not an isomorphism on Y_0 for infinitely many m , we can find an effective divisor \tilde{D} in Y such that \tilde{D} is not contained in $Y \setminus Y_0$ and a sequence $a_m \rightarrow +\infty$ such that

$$c_1(\varphi_*(\mathcal{O}_\Gamma(L + mE))) - c_1(\varphi_*(\mathcal{O}_\Gamma(L))) \geq a_m \cdot [\tilde{D}].$$

Combining this with Lemma 3.5, we know that

$$A - \frac{1}{r} \pi_* \varphi^* c_1(\varphi_*(\mathcal{O}_\Gamma(L))) - \frac{a_m}{r} \pi_* \varphi^* [\tilde{D}]$$

is a pseudoeffective class on X . By the definition of Y_0 and \tilde{D} , the class $\pi_* \varphi^* [\tilde{D}]$ is a non trivial effective class on X . We get thus a contradiction as $a_m \rightarrow +\infty$. \square

The following lemma tells us that, for every $c > 0$, $L + cE - \frac{1}{r} \varphi^* c_1(\varphi_*(\mathcal{O}_\Gamma(L + mE)))$ has no positivity in the horizontal direction. More precisely, we have

3.8. Lemma. *In the situation of Setup 3.1, let $m_0 \in \mathbb{N}$ as in Lemma 3.7, and let $m \geq m_0$ be an integer.*

Let θ be a closed $(1, 1)$ -current on Y such that $\theta \cdot \mathcal{C} > 0$ for some movable compact curve \mathcal{C} contained in Y_0 . Then the class

$$L + cE - \frac{1}{r} \varphi^* c_1(\varphi_*(\mathcal{O}_\Gamma(L + mE))) - \varphi^* \theta$$

is not pseudoeffective for any constant $c > 0$.

Remark. Note that we *do not* assume that θ is a positive current.

Proof. We assume by contradiction that there exists some $c \geq m$ such that

$$T := L + cE - \frac{1}{r} \varphi^* c_1(\varphi_*(\mathcal{O}_\Gamma(L + mE))) - \varphi^* \theta$$

is pseudoeffective on Γ . We can thus apply Lemma 3.6 to T and find a $c_1 > 0$ such that

$$c_1(\varphi_*(\mathcal{O}_\Gamma(T + c_1E))) \quad \text{is pseudoeffective on } Y.$$

Combining this with the definition of T , we have

$$(17) \quad \det \varphi_*(\mathcal{O}_\Gamma(L + cE + c_1E)) \geq \det \varphi_*(\mathcal{O}_\Gamma(L + mE)) + r\theta \quad \text{on } Y.$$

Thanks to Lemma 3.7, the effective class

$$c_1(\varphi_*(\mathcal{O}_\Gamma(L + cE + c_1E))) - c_1(\varphi_*(\mathcal{O}_\Gamma(L + mE)))$$

is supported in $Y \setminus Y_0$. As \mathcal{C} is contained in Y_0 , we have

$$c_1(\varphi_*(\mathcal{O}_\Gamma(L + cE + c_1E))) \cdot \mathcal{C} = c_1(\varphi_*(\mathcal{O}_\Gamma(L + mE))) \cdot \mathcal{C}.$$

We get thus a contradiction to (17) since $\theta \cdot \mathcal{C} > 0$ and \mathcal{C} is movable by assumption. \square

By combining Lemma 3.5, 3.6 and 3.8, we can prove the following key proposition.

3.9. Proposition. *In the situation of Setup 3.1, let $m_0 \in \mathbb{N}$ as in Lemma 3.7, and let $m \geq m_0$ be an integer. Fix a Kähler metric ω_X on X and let β be a smooth form representing $\frac{1}{r} \det \varphi_*(\mathcal{O}_\Gamma(L + mE))$ on Y .*

(i): *For every $p \in \mathbb{N}$, let $Y_{1,p}$ be the locally free locus of $\varphi_*(\mathcal{O}_\Gamma(pL + pmE))$. Then for every $\epsilon > 0$, we can find a metric $h_{\epsilon,p}$ on $\varphi_*(\mathcal{O}_\Gamma(pL + pmE))$ over $Y_0 \cap Y_{1,p}$ such that*

$$i\Theta_{h_{\epsilon,p}}(\varphi_*(\mathcal{O}_\Gamma(pL + pmE))) \succcurlyeq (-\epsilon\omega_X + p\beta) \otimes \text{Id}_{\text{End}(\varphi_*(\mathcal{O}_\Gamma(pL + pmE)))} \text{ on } Y_0 \cap Y_{1,p}.$$

(ii): *For every $p \in \mathbb{N}$ divisible by r , we have*

$$\frac{1}{r_p} \pi_* \varphi^*(c_1(\varphi_* \mathcal{O}_\Gamma(pL + pmE))) = \frac{p}{r} \pi_* \varphi^*(c_1(\varphi_* \mathcal{O}_\Gamma(L + mE)))$$

in $H^{1,1}(X, \mathbb{R})$, where r_p is the rank of $\varphi_(\mathcal{O}_\Gamma(pL + pmE))$.*

Proof of (i): Thanks to Lemma 3.5, we can find a constant $c > 0$ such that $L + cE - \varphi^*\beta$ is pseudoeffective on Γ . It means that we can find a possible singular metric h_L on $L + cE$ such that $i\Theta_{h_L}(L + cE) \geq \varphi^*\beta$. As a consequence, by applying Lemma 3.6 to $L + cE - \varphi^*\beta$, there exists a constant $c' \geq m$ such that, for every $\epsilon > 0$, we can find a possible singular hermitian metric h_ϵ on $\varphi_*(L + c'E)$ such that

$$(18) \quad i\Theta_{h_\epsilon \varphi_*(L + c'E)} \succcurlyeq (-\epsilon\omega_X + \beta) \otimes \text{Id}_{\text{End}(\varphi_*(L + c'E))} \quad \text{on } Y_{1,1,c'},$$

where $Y_{1,1,c'}$ is the locally free locus of $\varphi_*(L + c'E)$. Thanks to (16), we have

$$\varphi_*(\mathcal{O}_\Gamma(L + c'E)) \otimes \mathcal{O}_{Y_0} = \varphi_*(\mathcal{O}_\Gamma(L + mE)) \otimes \mathcal{O}_{Y_0}.$$

Together with (18), for every $\epsilon > 0$, we can find a metric h_ϵ on $\varphi_*(\mathcal{O}_\Gamma(L + mE))$ such that

$$(19) \quad i\Theta_{h_\epsilon}(\varphi_*(\mathcal{O}_\Gamma(L + mE))) \succcurlyeq (-\epsilon\omega_X + \beta) \otimes \text{Id}_{\text{End}(\varphi_*(\mathcal{O}_\Gamma(L + mE)))} \text{ on } Y_{1,1} \cap Y_0.$$

We consider the natural morphism

$$(20) \quad \text{Sym}^p \varphi_* \mathcal{O}_\Gamma(L + mE) \otimes \mathcal{O}_{Y_0} \rightarrow \varphi_* \mathcal{O}_\Gamma(pL + pmE) \otimes \mathcal{O}_{Y_0}.$$

By (8), it is surjective on a generic point. By applying Proposition 2.5 to (20), (19) induces thus a possible singular metric $h_{\epsilon,p}$ on $\varphi_*(\mathcal{O}_\Gamma(pL + pmE))$ such that

$$(21) \quad i\Theta_{h_{\epsilon,p}}(\varphi_*(\mathcal{O}_\Gamma(pL + pmE))) \succcurlyeq (-\epsilon\omega_X + p\beta) \otimes \text{Id}_{\text{End}(\varphi_* \mathcal{O}_\Gamma(pL + pmE))} \text{ on } Y_{1,p} \cap Y_0.$$

The part (i) is thus proved. \square

For every $p \in \mathbb{N}$ divisible by r , set

$$\mathcal{E}_p := (\text{Sym}^p \varphi_* \mathcal{O}_\Gamma(L + mE)) \otimes (\det \varphi_* \mathcal{O}_\Gamma(L + mE))^{\otimes \frac{-p}{r}}$$

and

$$(22) \quad \mathcal{V}_p := \varphi_* \mathcal{O}_\Gamma(pL + pmE) \otimes (\det \varphi_* \mathcal{O}_\Gamma(L + mE))^{\otimes \frac{-p}{r}}.$$

Before proving the second statement, we observe the following: let $\tau : \tilde{C} \rightarrow Y_0 \cap Y_{1,1}$ be a morphism from a smooth compact curve to Y_0 that is very general (i.e. the image is not contained in the locus where the metrics constructed above are singular). Then, for every $p \in \mathbb{N}$ divisible by r , $\tau^* \mathcal{E}_p$ is numerically flat. Indeed its first Chern class is zero by construction and $\tau^* \mathcal{E}_p$ is nef by (i).

Proof of (ii): Let α be a smooth form representing $\frac{1}{r_p}c_1(\varphi_*(\mathcal{O}_\Gamma(pL + pmE)))$. Let $C \subset X$ be a very general complete intersection curve cut out by sections of a very ample line bundle A_X on X . Since $C \cap Z = \emptyset$, we identify with its strict transform in Γ . Since $\varphi_*\mathcal{O}_\Gamma(L+mE)$ is reflexive on Y_0 , we know that $Y_0 \setminus Y_{1,1}$ has codimension at least two in Y_0 . Thus it follows by Proposition 3.3 that $\pi(\varphi^{-1}(Y \setminus Y_{1,1}))$ has codimension at least two. Thus we can assume without loss of generality that the movable curve $\varphi(C)$ is contained in $Y_{1,1} \cap Y_0$. Let $\tau : \tilde{C} \rightarrow \varphi(C)$ be the normalisation.

Since $N_1(X)$ is generated by cohomology classes of the form $A_X^{\dim X - 1}$ with A_X a very ample line bundle on X , it is sufficient, by duality and the projection formula, to prove that

$$(23) \quad \int_{\tilde{C}} \tau^*(\alpha - p\beta) = 0.$$

The map (20) induces a generically surjective morphism

$$\tau^*\mathcal{E}_p \rightarrow \tau^*\mathcal{V}_p \quad \text{on } \tilde{C}.$$

We observed above that $\tau^*\mathcal{E}_p$ is numerically flat on \tilde{C} , hence also $\tau^*(\mathcal{V}_p)$ [Laz04b, Ex.6.4.17]. Yet by definition $c_1(\mathcal{V}_p)$ is represented by $r_p(\alpha - p\beta)$. Thus we get

$$(24) \quad \int_{\tilde{C}} \tau^*(\alpha - p\beta) \geq 0.$$

In the other direction, as pL is also the pull-back of some very ample bundle on X , we can replace L by pL in Lemma 3.5. Therefore $pL + c_3E - \varphi^*\alpha$ is pseudoeffective on Γ for some c_3 large enough. Hence

$$L + \frac{c_3}{p}E - \frac{\varphi^*\alpha}{p} = L + \frac{c_3}{p}E - \varphi^*\beta - \varphi^*\left(\frac{\alpha - p\beta}{p}\right)$$

is pseudoeffective. By applying Lemma 3.8, we get

$$\int_{\tilde{C}} \tau^*(\alpha - p\beta) \leq 0.$$

Together with (24) this proves (23). \square

Now we can prove the main result in this subsection.

3.10. Proposition. *In the situation of Setup 3.1, let $m_0 \in \mathbb{N}$ as in Lemma 3.7, and let $m \geq m_0$ be an integer. Fix a sufficiently divisible $p \in \mathbb{N}$ and let \mathcal{V}_p be the direct image defined in (22). Then \mathcal{V}_p is a trivial vector bundle on Y_0 .*

Proof. Denote by $Y_{1,p} \subset Y$ the locally free locus of \mathcal{V}_p . Since \mathcal{V}_p is reflexive on Y_0 we know that the complement of $Y_0 \cap Y_{1,p}$ in Y_0 has codimension at least two. By Proposition 1.5, 3) it is sufficient to show that $\mathcal{V}_p \otimes \mathcal{O}_{Y_0 \cap Y_{1,p}}$ is a trivial vector bundle.

The image of the π -exceptional locus $Z := \pi(E)$ has codimension at least two. Denote by $j : X \setminus Z \rightarrow X$ the inclusion.

Let A_X be a very ample line bundle on X , and let

$$S = H_1 \cap \cdots \cap H_{n-2}$$

be a surface cut out by general elements $H_1, \dots, H_{n-2} \in |A_X|$.

Then $Z_S := Z \cap S$ is a finite set and $S \setminus Z_S$ is simply connected, since by the Lefschetz theorem $\pi_1(S) \simeq \pi_1(X)$ is trivial. Let $\pi^{-1}(S \setminus Z_S)$ be the strict transform of $S \setminus Z_S$. By using Proposition 3.3, we see that $\pi^{-1}(S \setminus Z_S) \setminus \varphi^{-1}(Y_0)$ is also finite.

The vector bundle $\varphi^* \mathcal{V}_p|_{\pi^{-1}(S \setminus Z_S)}$ is weakly positively curved by Proposition 3.9(i) and $c_1(\varphi^* \mathcal{V}_p|_{\pi^{-1}(S \setminus Z_S)}) = 0$ Proposition 3.9(ii). Therefore, by Proposition 2.12 its extension to S is numerically flat. Since $\pi_1(S) = 1$ this implies that this extension, hence $\varphi^* \mathcal{V}_p|_{\pi^{-1}(S \setminus Z_S)}$ is a trivial vector bundle.

Since Z has codimension at least two, we have isomorphisms

$$(25) \quad H^0(\Gamma \setminus E, \varphi^* \mathcal{V}_p) \simeq H^0(X \setminus Z, \varphi^* \mathcal{V}_p) \simeq H^0(X, (j_*(\varphi^* \mathcal{V}_p))^{**}).$$

Consider now a curve

$$C = S \cap H_{n-1}$$

where $H_{n-1} \in |A_X|$ is general. Up to replacing A_X by some positive multiple, we can suppose that we have a surjective morphism

$$H^0(\Gamma \setminus E, \varphi^* \mathcal{V}_p) \rightarrow H^0(C, \varphi^* \mathcal{V}_p).$$

As $Z \cap S = \emptyset$ we know that $(\varphi^* \mathcal{V}_p)|_C$ is a trivial vector bundle. Thus the surjectivity above implies the existence of section

$$s_1, s_2, \dots, s_{r_p} \in H^0(\Gamma \setminus E, \varphi^* \mathcal{V}_p)$$

such that

$$s_1 \wedge \dots \wedge s_{r_p} \in H^0(\Gamma \setminus E, \det \varphi^* \mathcal{V}_p) \text{ is a non trivial section.}$$

Thanks to (25), it induces a section of $H^0(X, (j_*(\det \varphi^* \mathcal{V}_p))^{**})$. Yet X is compact and $c_1(j_*(\det \varphi^* \mathcal{V}_p)) = 0$ by using Proposition 3.9. So the nonvanishing section is a non-zero constant. In particular $s_1 \wedge \dots \wedge s_{r_p}$ is a non-zero constant.

We claim that $s_i|_{\varphi^{-1}(Y_0) \setminus E} = \varphi^* \tau_i$ for some $\tau_i \in H^0(Y_0, \mathcal{V}_p)$. If the claim is proved, since

$$s_1 \wedge \dots \wedge s_{r_p} = \varphi^*(\tau_1 \wedge \dots \wedge \tau_{r_p})$$

is a non-zero constant, these sections define the trivialisation of \mathcal{V}_p . The proposition is proved.

Proof of the claim. Since $\Gamma \setminus E \rightarrow Y$ is not proper, this is not totally obvious. However, since φ has connected fibers, the sections $s_i \in H^0(\Gamma \setminus E, \varphi^* \mathcal{V}_p)$ induce sections $\tilde{\tau}_i \in H^0(Y_0 \setminus \varphi(E), \mathcal{V}_p)$.

We need to prove that $\tilde{\tau}_i$ extends to a section $\tau_i \in H^0(Y_0, \mathcal{V}_p)$: since $\mathcal{V}_p \otimes \mathcal{O}_{Y_0}$ is reflexive, it is enough to show that $\tilde{\tau}_i$ extends in the general point of every divisor $B \subset Y$. By definition of Y_0 the morphism $\Gamma \setminus E \rightarrow Y$ dominates every divisor $B \subset Y_0$. Moreover, by Proposition 3.3,3) the general fibre over B has at least one reduced component. Thus there exists a local section of $\Gamma \setminus E \rightarrow Y$ in a general point of B . Using this section we see that $\tilde{\tau}_i$ is bounded in a general point of B , hence it extends by Riemann's extension theorem. \square

3.C. Proof of Theorem 1.2. In the situation of Setup 3.1 we use the notation of Proposition 3.9. Moreover we choose $m \geq m_0$ as in Proposition 3.10. For a $p_0 \in \mathbb{N}$ sufficiently divisible we know by Proposition 3.10 that the vector bundle $\mathcal{V}_{p_0} \otimes \mathcal{O}_{Y_0}$ is trivial. Thus we have

$$\varphi_*(\mathcal{O}_\Gamma(p_0(L + mE))) \otimes \mathcal{O}_{Y_0} \simeq ((\det \varphi_*(\mathcal{O}_\Gamma(L + mE)))^{\otimes \frac{p_0}{r}})^{\oplus r_{p_0}} \otimes \mathcal{O}_{Y_0},$$

In particular $\det \varphi_*(\mathcal{O}_\Gamma(p_0(L + mE))) \otimes \mathcal{O}_{Y_0}$ is divisible by r_{p_0} in the Picard group. In order to simplify the notation we replace L by p_0L and mE by p_0mE , hence, up to this change of notation, the line bundle $\det \varphi_*(\mathcal{O}_\Gamma(L + mE)) \otimes \mathcal{O}_{Y_0}$ is divisible by r and Proposition 3.10 holds for all $p \in \mathbb{N}$.

By (8) the natural map

$$(26) \quad \text{Sym}^p(\varphi_*(\mathcal{O}_\Gamma(L + mE))) \otimes \mathcal{O}_{Y_0} \rightarrow \varphi_*(\mathcal{O}_\Gamma(p(L + mE))) \otimes \mathcal{O}_{Y_0}$$

is generically surjective. Using the notation of Proposition 3.9 the twist of this map by $(\det \varphi_*(\mathcal{O}_\Gamma(L + mE)))^{\otimes \frac{p}{r}}$ is the map

$$\text{Sym}^p \mathcal{V}_1 \otimes \mathcal{O}_{Y_0} \rightarrow \mathcal{V}_p \otimes \mathcal{O}_{Y_0}.$$

By Proposition 3.10 these are trivial vector bundles, so the rank of the map is constant by Remark 3.4. Thus (26) is surjective.

While the line bundle L is globally generated and φ -big, the line bundle $L + mE$ is in general only φ -globally generated on a general fibre. Hence the natural map

$$\varphi^* \varphi_*(\mathcal{O}_\Gamma(L + mE)) \rightarrow L + mE$$

is not necessarily surjective. However, up to taking a resolution of the indeterminacies $\Gamma' \rightarrow \Gamma$ by some smooth projective manifold Γ' , and replacing L and E by their pull-backs, we can suppose without loss of generality that we have

$$\varphi^* \varphi_*(\mathcal{O}_\Gamma(L + mE)) \twoheadrightarrow M \hookrightarrow L + mE$$

where M is a line bundle. Pushing down to Y gives morphisms

$$\varphi_*(\mathcal{O}_\Gamma(L + mE)) \rightarrow \varphi_*(\mathcal{O}_\Gamma(M)) \hookrightarrow \varphi_*(\mathcal{O}_\Gamma(L + mE))$$

and the composition is simply the identity. Thus the injection $\varphi_*(\mathcal{O}_\Gamma(M)) \hookrightarrow \varphi_*(\mathcal{O}_\Gamma(L + mE))$ is an isomorphism. In particular we obtain

$$(27) \quad L + mE = M + N$$

where M is φ -globally generated and N is an effective divisor that does not surject onto Y . For a general φ -fibre F we have

$$L|_F = (L + mE)|_F = (M + N)|_F = M|_F,$$

hence $M|_F$ is very ample.

The surjective map $\varphi^* \varphi_*(\mathcal{O}_\Gamma(M)) \twoheadrightarrow M$ defines a morphism $\Gamma \rightarrow \mathbb{P}(\varphi_*(\mathcal{O}_\Gamma(M)))$, and we denote by $\mathcal{U}_M \subset \mathbb{P}(\varphi_*(\mathcal{O}_\Gamma(M)))$ its image. We summarise the situation and notation in a commutative diagram:

$$\begin{array}{ccccc} \Gamma & \xrightarrow{\psi_M} & \mathcal{U}_M & \hookrightarrow & \mathbb{P}(\varphi_*(\mathcal{O}_\Gamma(M))) \\ & \searrow \varphi & \downarrow \varphi_M & & \swarrow \text{pr} \\ & & Y & & \end{array}$$

Step 1. We prove that $\varphi_M^{-1}(Y_0) \simeq Y_0 \times F$ with F a general φ -fibre. For every $p \in \mathbb{N}$ the inclusion $pM \hookrightarrow p(L + mE)$ induces a commutative diagram

$$\begin{array}{ccc} \mathrm{Sym}^p(\varphi_*(\mathcal{O}_\Gamma(M))) \otimes \mathcal{O}_{Y_0} & \longrightarrow & \varphi_*(\mathcal{O}_\Gamma(pM)) \otimes \mathcal{O}_{Y_0} \\ \downarrow \simeq & & \downarrow \\ \mathrm{Sym}^p(\varphi_*(\mathcal{O}_\Gamma(L + mE))) \otimes \mathcal{O}_{Y_0} & \twoheadrightarrow & \varphi_*(\mathcal{O}_\Gamma(p(L + mE))) \otimes \mathcal{O}_{Y_0} \end{array}$$

The isomorphism in the left column and the surjectivity of the bottom line were established above, so the right column is also an isomorphism. If we set

$$G := (\det \varphi_*(\mathcal{O}_\Gamma(L + mE)) \otimes \mathcal{O}_{Y_0})^{\otimes \frac{1}{r}} \simeq (\det \varphi_*(\mathcal{O}_\Gamma(M)) \otimes \mathcal{O}_{Y_0})^{\otimes \frac{1}{r}},$$

we obtain by Proposition 3.10 that

$$(28) \quad \varphi_*(\mathcal{O}_\Gamma(pM)) \otimes \mathcal{O}_{Y_0} \simeq (G \otimes \mathcal{O}_{Y_0})^{\oplus r_p}$$

for every $p \in \mathbb{N}$.

Denote by $\mathcal{O}_{\mathbb{P}(\varphi_*(\mathcal{O}_\Gamma(M)))}(1)$ the tautological bundle on $\mathbb{P}(\varphi_*(\mathcal{O}_\Gamma(M)))$, and by $\mathcal{I}_{\mathcal{U}_M}$ the ideal sheaf of $\mathcal{U}_M \subset \mathbb{P}(\varphi_*(\mathcal{O}_\Gamma(M)))$. For $p \in \mathbb{N}$ sufficiently high, the direct image $\mathrm{pr}_*(\mathcal{I}_{\mathcal{U}_M}(p))$ is pr -globally generated, and the higher direct images $R^j \mathrm{pr}_*(\mathcal{I}_{\mathcal{U}_M}(p))$ vanish. Thus we have an exact sequence

$$(29) \quad 0 \rightarrow \mathrm{pr}_*(\mathcal{I}_{\mathcal{U}_M}(p)) \rightarrow \mathrm{pr}_*(\mathcal{O}_{\mathbb{P}(\varphi_*(\mathcal{O}_\Gamma(M)))}(p)) \simeq \mathrm{Sym}^p(\varphi_*(\mathcal{O}_\Gamma(M))) \rightarrow (\varphi_M)_*(\mathcal{O}_{\mathcal{U}_M}(p)) \rightarrow 0,$$

where $\mathcal{O}_{\mathcal{U}_M}(p)$ denotes the restriction of the tautological bundle to \mathcal{U}_M . By construction we have $M = \psi_M^* \mathcal{O}_{\mathbb{P}(\varphi_*(\mathcal{O}_\Gamma(M)))}(1)$, so by the projection formula there is a natural inclusion

$$(30) \quad (\varphi_M)_*(\mathcal{O}_{\mathcal{U}_M}(p)) \rightarrow \varphi_*(\mathcal{O}_\Gamma(pM)).$$

This morphism is an isomorphism in the generic point, since ψ_M is an isomorphism on the general φ -fibre. Twisting (29) and (30) with G^{-p} we obtain morphisms

$$\mathrm{Sym}^p(\varphi_*(\mathcal{O}_\Gamma(M)) \otimes G^{-p}) \rightarrow (\varphi_M)_*(\mathcal{O}_{\mathcal{U}_M}(p)) \otimes G^{-p} \rightarrow \varphi_*(\mathcal{O}_\Gamma(pM)) \otimes G^{-p}.$$

By (28) the restriction of the first and the last sheaf to Y_0 are trivial vector bundles, so by Remark 3.4 the rank of the map is constant. In particular the inclusion $(\varphi_M)_*(\mathcal{O}_{\mathcal{U}_M}(p)) \rightarrow \varphi_*(\mathcal{O}_\Gamma(pM))$ is an isomorphism on Y_0 . Hence

$$(\varphi_M)_*(\mathcal{O}_{\mathcal{U}_M}(p)) \otimes G^{-p} \otimes \mathcal{O}_{Y_0}$$

is a trivial vector bundle, and by the exact sequence (29)

$$\mathrm{pr}_*(\mathcal{I}_{\mathcal{U}_M}(p)) \otimes G^{-p} \otimes \mathcal{O}_{Y_0}$$

is also trivial. Finally, this proves that the equations defining \mathcal{U}_M in the product $\mathbb{P}(\varphi_*(\mathcal{O}_\Gamma(M)) \otimes \mathcal{O}_{Y_0}) \simeq Y_0 \times \mathbb{P}^{r-1}$ are constant, hence \mathcal{U}_M is a product.

Step 2. We prove that the tangent bundle of $\varphi^{-1}(Y_0) \cap (\Gamma \setminus E)$ splits. Recall that by (27) we have $L + mE = M + N$ with N an effective divisor corresponding to φ -fixed components of the $L + mE$. The line bundle L is φ -globally generated, so any relative fixed component of $L + mE$ is contained in mE . Thus $mE - N$ is an effective divisor.

Now observe the following: given any fibre $\varphi^{-1}(y)$, let F_0 be an irreducible component of $\varphi^{-1}(y)$ that is not contained in E . Then F_0 maps birationally onto its image in X , so $L|_{F_0} = (\pi^*A)|_{F_0}$ is nef and big. Since

$$M|_{F_0} = L|_{F_0} + (mE - N)|_{F_0}$$

we obtain that $M|_{F_0}$ is also big. Thus F_0 is not contracted by ψ_M . Consider now the birational morphism

$$\psi_M^0 : \varphi^{-1}(Y_0) \rightarrow \varphi_M^{-1}(Y_0) \simeq Y_0 \times F.$$

Since $Y_0 \times F$ is smooth, the ψ_M^0 -exceptional locus has pure codimension one, and we denote by D one of its irreducible components. Since $\varphi^{-1}(Y_0) \rightarrow Y_0$ is flat, the image $\varphi(D)$ is a divisor, and we denote by F_0 an irreducible component of a general fibre of $D \rightarrow \varphi(D)$. If $F_0 \not\subset E$, then we have just shown that F_0 is not contracted by ψ_M . Thus D would not be contracted by ψ_M^0 . This shows that the exceptional locus of ψ_M^0 is contained in E . In particular

$$(31) \quad T_{\varphi^{-1}(Y_0) \cap (\Gamma \setminus E)} \simeq V_1^0 \oplus V_2^0,$$

where V_1^0 (resp. V_2^0) is the pull-back of $T_{Y_0 \times F / Y_0}$ (resp. $T_{Y_0 \times F / F}$) under the embedding

$$\varphi^{-1}(Y_0) \cap \Gamma \setminus E \hookrightarrow \varphi_M^{-1}(Y_0) \simeq Y_0 \times F.$$

Note that V_1^0 and V_2^0 are both algebraically integrable foliations and the general leaves of V_1^0 are φ -fibres.

Step 3. Splitting of T_X and conclusion. Since $\Gamma \setminus E \rightarrow X$ is an embedding, we can identify $\varphi^{-1}(Y_0) \cap (\Gamma \setminus E)$ to a Zariski open subset X_0 of X , and we denote by $j : X_0 \hookrightarrow X$ the inclusion. Then

$$V_1 := (j_*(V_1^0))^{**}, \quad V_2 := (j_*(V_2^0))^{**}$$

are reflexive sheaves on X . By (31) we have

$$(V_1 \oplus V_2) \otimes \mathcal{O}_{X_0} = T_{X_0}.$$

By Proposition 3.3,2) the complement of $X \setminus X_0$ has codimension at least two. Thus the equality above extends by Proposition 1.5,3) to an isomorphism of reflexive sheaves

$$V_1 \oplus V_2 \simeq T_X.$$

Yet this implies that the upper semicontinuous functions $\dim(V_{j,x} \otimes_{\mathcal{O}_{X,x}} \mathbb{C}(x))$ are constant, since their sum is equal to $\text{rk} T_X$. Thus V_1 and V_2 are locally free [Har77, II, Ex.5.8.c)] and, identifying V_1 and V_2 to their images in T_X , they define regular foliations on X . We have seen above that a general V_1 -leaf is a general fibre of the MRC-fibration, hence rationally connected. By [Hör07, Cor.2.11] this implies that there exists a smooth morphism $X \rightarrow Y$ such that $T_{X/Y} = V_1$. Since $T_X = V_1 \oplus V_2$, the relative tangent sequence of this submersion is split by V_2 . Since V_2 is integrable, the classical Ehresmann theorem (e.g. [CLN85, V., §2, Prop.1 and Thm.3]) implies that the universal cover of X is a product. Yet X is simply connected, so X itself is a product. \square

Proof of Theorem 1.4. If X is not uniruled, then $K_X \equiv 0$ by [BDPP13]. Thus we can suppose that X is uniruled, and denote by $\mathcal{F} \subset T_X$ the unique integrable saturated subsheaf such that for a very general point $x \in X$, the \mathcal{F} -leaf through x is a fibre of the MRC-fibration. Note that a saturated subsheaf is reflexive.

Step 1. We prove that $\mathcal{F} \subset T_X$ is a subbundle. Note first that if $f : X' \rightarrow X$ is a finite étale cover, then, since f^* is exact for flat morphisms, the pull-back $f^*\mathcal{F} \subset f^*T_X = T_{X'}$ is the unique integrable saturated subsheaf corresponding to the MRC-fibration of X' . Moreover if $f^*\mathcal{F} \subset T_{X'}$ is a subbundle, then $\mathcal{F} \subset T_X$ is

a subbundle. Thus the claim is invariant under finite étale covers, in particular it holds for manifolds with finite fundamental group by Theorem 1.2.

By [Cao16, Cor.1.4] we know that, maybe after finite étale cover, we have a locally trivial Albanese map $\alpha : X \rightarrow \text{Alb}(X)$ such that the fibre G has finite fundamental group. Since $\text{Alb}(X)$ is not uniruled, we have a factorisation

$$\mathcal{F} \subset T_{X/\text{Alb}(X)} \subset T_X.$$

Fix any point $y \in \text{Alb}(X)$ and choose an analytic neighborhood $U \subset \text{Alb}(X)$ such that $\alpha^{-1}(U) \simeq U \times G$. Since G has finite fundamental group, we already know that the MRC-fibration can be represented by $U \times G \rightarrow U \times Z$, where $G \rightarrow Z$ is a smooth fibration. By uniqueness of $\mathcal{F}|_{\alpha^{-1}(U)}$ this shows that $\mathcal{F}|_{\alpha^{-1}(U)} \subset T_X|_{\alpha^{-1}(U)}$ is a subbundle.

Step 2. Structure of MRC-fibration. By Step 1 we know that $\mathcal{F} \subset T_X$ is a regular foliation. Since it has a rationally connected leaf, we know by [Hör07, Cor.2.11] that there exists a smooth morphism $\varphi : X \rightarrow Y$ such that $T_{X/Y} = \mathcal{F}$. Since $-K_X$ is nef and φ is smooth, we know that $-K_Y$ is nef [Deb01, Cor.3.15]. Since Y is not uniruled [GHS03], this proves that $K_Y \equiv 0$.

We claim that φ is locally trivial. By the Fischer-Grauert theorem [FG65] it is enough to show that all the fibres are biholomorphic. Note again that this statement is invariant under étale cover: rationally connected manifolds are simply connected, so a φ -fibre lifts to a φ' -fibre if $X' \rightarrow X$ is étale. Thus by [Cao16, Cor.1.4] we reduce to the case where we have a locally trivial Albanese map $\alpha : X \rightarrow \text{Alb}(X)$ such that the fibre G has finite fundamental group. By invariance under étale covers and Theorem 1.2 we know that the MRC-fibration of G is locally trivial, so φ -fibres being contained in the same α -fibre are biholomorphic. Yet α is locally trivial, and automorphisms preserve the MRC-fibration, so all the φ -fibres are biholomorphic. \square

APPENDIX A. SOME ANALYTIC RESULTS

To begin with, we first prove an approximation lemma which is a direct consequence of [Dem92, thm 1.1] and a standard gluing lemma [Dem92, section 3].

A.1. Lemma. *Let C be a 1-dimensional projective manifold and let E be a vector bundle of rank n on C . Let $\pi : \mathbb{P}(E) \rightarrow C$ be the natural projection. Let $T = \alpha + dd^c\psi \geq 0$ be a positive current of analytic singularity, where α is smooth form and ψ is a quasi-psh function. We suppose moreover that ψ is smooth over the generic fiber of π .*

Then for every $c > 0$, there exists a sequence of closed almost positive $(1,1)$ -currents $T_{c,k} = \alpha + dd^c\psi_{c,k}$ such that $\psi_{c,k}$ is smooth on $\mathbb{P}(E) \setminus E_c(T)$, decreases to ψ as k tends to $+\infty$, and

$$T_{c,k} \geq -\epsilon_k \omega,$$

where $\lim_{k \rightarrow +\infty} \epsilon_k = 0$ and $\nu(T_{c,k}, x) = (\nu(T, x) - c)_+$ at every point $x \in X$. Here $E_c(T) := \{x \mid \nu(T, x) \geq c\}$ and $(\nu(T, x) - c)_+ := \max\{\nu(T, x) - c, 0\}$.

Proof. As ψ is smooth over the generic fiber of π , we can take a Stein cover $\{U_i\}$ of C such that T is smooth over $\pi^{-1}(U_i \cap U_j)$ for every $i \neq j$. Over $\pi^{-1}(U_i)$, since $T_{\mathbb{P}(E)/C}$ is relatively ample, we can find a smooth metric on $\mathcal{O}_{\mathbb{P}(T_{\pi^{-1}(U_i)})}(1)$ such

that the curvature form is positive. By applying [Dem92, thm 1.1], we can find a sequence of closed almost positive $(1, 1)$ -currents $T_{c,k,i} = \alpha + dd^c \psi_{c,k,i}$ on $\pi^{-1}(U_i)$, such that $\psi_{c,k,i}$ is smooth on $\pi^{-1}(U_i) \setminus E_c(T)$, decreases to $\psi|_{\pi^{-1}(U_i)}$ as k tends to $+\infty$, and

$$T_{c,k,i} \geq -\epsilon_k \omega \quad \text{on } \pi^{-1}(U_i),$$

where $\epsilon_k \rightarrow 0$ and $\nu(T_{c,k,i}, x) = (\nu(T, x) - c)_+$ at every point $x \in \pi^{-1}(U_i)$.

Since ψ is smooth over $\pi^{-1}(U_i \cap U_j)$ for every $i \neq j$, for any set $V \Subset U_i \cap U_j$, we know that $\lim_{k \rightarrow +\infty} \|\psi_{c,k,i} - \psi_{c,k,j}\|_{\mathcal{C}^0(\pi^{-1}(V))} = 0$. By applying the standard gluing process [Dem92, lemma 3.5], we can glue $\{T_{c,k,i}\}_i$ together by losing a bit of positivity. The lemma is thus proved. \square

Thanks to Lemma A.1, we can get an upper estimate of the Lelong numbers of T by using the same arguments in [Dem92, thm 1.7].

A.2. Proposition. *In the setting of Lemma A.1, set $a := \max_{x \in \mathbb{P}(E)} \nu(T, x)$. Then there exists a $k \in \{1, \dots, n\}$, a closed (k, k) -positive current Θ_k in the same class of $(c_1(T))^k \in H^{k,k}(\mathbb{P}(E))$ and a k -codimensional subvariety Z contained in some fiber of π such that*

$$\Theta_k \geq \left(\frac{a}{n}\right)^k \cdot [Z]$$

in the sense of current.

Proof. Thanks to Lemma A.1, the proposition can be proved by the same argument as in [Dem92, thm 1.7]. For the convenience of readers, we give the proof here.

We follow the notations introduced [Dem92, thm 1.7]: let $0 = b_1 \leq b_2 \leq \dots \leq b_n \leq b_{n+1}$ be the sequence of "jumping values" b_p , namely $\text{codim} E_c(T) = p$ when $c \in (b_p, b_{p+1}]$. Here, $\text{codim} E_c(T) = p$ means that all components of $E_c(T)$ is of codimension $\geq p$ and at least some component of $E_c(T)$ is of codimension p . By the definition of a , there exists a b_k such that $b_{k+1} - b_k \geq \frac{a}{n}$. Let Z be a k -codimensional component of $E_{b_{k+1}}(T)$ such that the generic Lelong of T along Z is b_{k+1} . Since ψ is smooth on the generic fiber of π , Z is contained in some fiber of π .

We now prove by induction the following claim: for every $1 \leq p \leq k$, we can find (p, p) -positive current Θ_p in the same class of $(c_1(T))^k$ such that

$$(32) \quad \nu(\Theta_p, x) \geq (b_{k+1} - b_1) \cdots (b_{k+1} - b_p) \quad \text{for every } x \in Z.$$

For $p = 1$, we can take $\Theta_1 = T$ and (32) holds. Now, suppose that Θ_{p-1} has been constructed. For $c > b_p$, the current $T_{c,k}$ produced by Lemma A.1 satisfies $T_{c,k} \geq -\epsilon_k \omega$ and the singular locus of $T_{c,k}$ is contained in $E_c(T)$ which is of codimension at least p . Then [Dem92, Lemma 7.4] shows that $\Theta_{p-1} \wedge (T_{c,k} + \epsilon_k \omega)$ is a well defined positive current. We define

$$\Theta_p = \lim_{c \rightarrow b_p+0} \lim_{k \rightarrow +\infty} \Theta_{p-1} \wedge (T_{c,k} + \epsilon_k \omega)$$

possibly after extracting some weakly convergent subsequence. Moreover, we have

$$\nu(\Theta_p, x) \geq \nu(\Theta_{p-1}, x) \times \overline{\lim}_{c \rightarrow b_p+0} \overline{\lim}_{k \rightarrow +\infty} \nu(T_{c,k}, x) \geq \nu(\Theta_{p-1}, x) \times (\nu(T, x) - b_p)_+.$$

In particular, for every $x \in Z$, we have $\nu(\Theta_p, x) \geq (b_{k+1} - b_1) \cdots (b_{k+1} - b_p)$. (32) is proved.

We now prove the proposition. Thanks to (32), we can find a closed (k, k) -positive current Θ_k in the same class of $(c_1(T))^k$ such that

$$\nu(\Theta_k, x) \geq (b_{k+1} - b_1) \cdots (b_{k+1} - b_k) \geq (b_{k+1} - b_k)^k \geq \left(\frac{a}{n}\right)^k \quad \text{for every } x \in Z.$$

As Z is of codimension k , Siu's decomposition theorem implies

$$\Theta_k \geq \left(\frac{a}{n}\right)^k \cdot [Z]$$

in the sense of current. The proposition is proved. \square

REFERENCES

- [BDPP13] Sébastien Boucksom, Jean-Pierre Demailly, Mihai Păun, and Thomas Peternell. The pseudo-effective cone of a compact Kähler manifold and varieties of negative Kodaira dimension. *Journal of Algebraic Geometry*, 22:201–248, 2013.
- [Bea83] Arnaud Beauville. Variétés Kähleriennes dont la première classe de Chern est nulle. *J. Differential Geom.*, 18(4):755–782 (1984), 1983.
- [Ber09] Bo Berndtsson. Curvature of vector bundles associated to holomorphic fibrations. *Annals of Mathematics*, 169(2):531–560, 2009.
- [BP04] Thomas Bauer and Thomas Peternell. Nef reduction and anticanonical bundles. *Asian J. Math.*, 8(2):315–352, 2004.
- [BP08] Bo Berndtsson and Mihai Păun. Bergman kernels and the pseudoeffectivity of relative canonical bundles. *Duke Math. J.*, 145(2):341–378, 2008.
- [Cam04] Frédéric Campana. Orbifolds, special varieties and classification theory. *Ann. Inst. Fourier (Grenoble)*, 54(3):499–630, 2004.
- [Cao16] Junyan Cao. Albanese maps of projective manifolds with nef anticanonical bundles. *arXiv preprint*, 1612.05921, 2016.
- [CDP15] F. Campana, J.-P. Demailly, and Th. Peternell. Rationally connected manifolds and semipositivity of the Ricci curvature. In *Recent advances in algebraic geometry*, volume 417 of *London Math. Soc. Lecture Note Ser.*, pages 71–91. Cambridge Univ. Press, Cambridge, 2015.
- [CH17] Junyan Cao and Andreas Höring. Manifolds with nef anticanonical bundle. *J. Reine Angew. Math.*, 724:203–244, 2017.
- [CLN85] César Camacho and Alcides Lins Neto. *Geometric theory of foliations*. Birkhäuser Boston Inc., Boston, MA, 1985. Translated from the Portuguese by Sue E. Goodman.
- [CP91] Frédéric Campana and Thomas Peternell. Projective manifolds whose tangent bundles are numerically effective. *Math. Ann.*, 289(1):169–187, 1991.
- [CP17] Junyan Cao and Mihai Păun. Kodaira dimension of algebraic fiber spaces over abelian varieties. *Invent. Math.*, 207(1):345–387, 2017.
- [Deb01] Olivier Debarre. *Higher-dimensional algebraic geometry*. Universitext. Springer-Verlag, New York, 2001.
- [Dem92] Jean-Pierre Demailly. Regularization of closed positive currents and intersection theory. *J. Algebraic Geom.*, 1(3):361–409, 1992.
- [Dem12] Jean-Pierre Demailly. *Analytic methods in algebraic geometry*, volume 1 of *Surveys of Modern Mathematics*. International Press, Somerville, MA; Higher Education Press, Beijing, 2012.
- [DPS93] Jean-Pierre Demailly, Thomas Peternell, and Michael Schneider. Kähler manifolds with numerically effective Ricci class. *Compositio Math.*, 89(2):217–240, 1993.
- [DPS94] Jean-Pierre Demailly, Thomas Peternell, and Michael Schneider. Compact complex manifolds with numerically effective tangent bundles. *J. Algebraic Geom.*, 3(2):295–345, 1994.
- [DPS96] Jean-Pierre Demailly, Thomas Peternell, and Michael Schneider. Compact Kähler manifolds with Hermitian semipositive anticanonical bundle. *Compositio Math.*, 101(2):217–224, 1996.
- [FG65] Wolfgang Fischer and Hans Grauert. Lokal-triviale Familien kompakter komplexer Mannigfaltigkeiten. *Nachr. Akad. Wiss. Göttingen Math.-Phys. Kl. II*, 1965:89–94, 1965.

- [GHS03] Tom Graber, Joe Harris, and Jason Starr. Families of rationally connected varieties. *J. Amer. Math. Soc.*, 16(1):57–67 (electronic), 2003.
- [Har77] Robin Hartshorne. *Algebraic geometry*. Springer-Verlag, New York, 1977. Graduate Texts in Mathematics, No. 52.
- [Har80] Robin Hartshorne. Stable reflexive sheaves. *Math. Ann.*, 254(2):121–176, 1980.
- [Hör07] Andreas Höring. Uniruled varieties with split tangent bundle. *Math. Z.*, 256(3):465–479, 2007.
- [HPS16] C. Hacon, M. Popa, and C. Schnell. Algebraic fiber spaces over abelian varieties: around a recent theorem by Cao and Paun. *ArXiv e-prints*, November 2016.
- [KK83] Ludger Kaup and Burchard Kaup. *Holomorphic functions of several variables*, volume 3 of *de Gruyter Studies in Mathematics*. Walter de Gruyter & Co., Berlin, 1983. An introduction to the fundamental theory, With the assistance of Gottfried Barthel, Translated from the German by Michael Bridgland.
- [Laz04a] Robert Lazarsfeld. *Positivity in algebraic geometry. I*, volume 48 of *Ergebnisse der Mathematik und ihrer Grenzgebiete*. Springer-Verlag, Berlin, 2004. Classical setting: line bundles and linear series.
- [Laz04b] Robert Lazarsfeld. *Positivity in algebraic geometry. II*, volume 49 of *Ergebnisse der Mathematik und ihrer Grenzgebiete*. Springer-Verlag, Berlin, 2004. Positivity for vector bundles, and multiplier ideals.
- [LTZZ10] Steven Lu, Yuping Tu, Qi Zhang, and Quan Zheng. On semistability of Albanese maps. *Manuscripta Math.*, 131(3-4):531–535, 2010.
- [Pau97] Mihai Paun. Sur le groupe fondamental des variétés kählériennes compactes à classe de Ricci numériquement effective. *C. R. Acad. Sci. Paris Sér. I Math.*, 324(11):1249–1254, 1997.
- [Pău16] Mihai Păun. Singular hermitian metrics and positivity of direct images of pluricanonical bundles. *arXiv preprint*, 1606.00174, 2016.
- [PT14] Mihai Păun and Shigeharu Takayama. Positivity of twisted relative pluricanonical bundles and their direct images. *arXiv preprint*, 1409.5504, 2014.
- [Rau15] Hossein Raufi. Singular hermitian metrics on holomorphic vector bundles. *Ark. Mat.*, 53(2):359–382, 2015.
- [Rei94] Miles Reid. Nonnormal del Pezzo surfaces. *Publ. Res. Inst. Math. Sci.*, 30(5):695–727, 1994.
- [Vie95] Eckart Viehweg. *Quasi-projective moduli for polarized manifolds*, volume 30 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1995.
- [Zha05] Qi Zhang. On projective varieties with nef anticanonical divisors. *Math. Ann.*, 332(3):697–703, 2005.

ANDREAS HÖRING, UNIVERSITÉ CÔTE D’AZUR, CNRS, LJAD, FRANCE

E-mail address: Andreas.Hoering@unice.fr

JUNYAN CAO, UNIVERSITÉ PARIS 6, INSTITUT DE MATHÉMATIQUES DE JUSSIEU, 4, PLACE JUSSIEU, PARIS 75252, FRANCE

E-mail address: junyan.cao@imj-prg.fr