Identification and assessment of ecotoxicological hazards attributable to pollutants in urban wet weather discharges

Antoine Gosset, Claude Durrieu, Frédéric Orias, Rémy Bayard, Yves Perrodin

► To cite this version:

Antoine Gosset, Claude Durrieu, Frédéric Orias, Rémy Bayard, Yves Perrodin. Identification and assessment of ecotoxicological hazards attributable to pollutants in urban wet weather discharges. Environ. Sci.: Processes Impacts, 2017, 19, pp.1150-1168. 10.1039/C7EM00159B . hal-01567078

HAL Id: hal-01567078

<https://hal.science/hal-01567078>

Submitted on 21 Jul 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Identification and assessment of ecotoxicological hazards attributable to pollutants in urban wet weather discharges

Antoine Gosset^{1, 2*}, Claude Durrieu¹, Frédéric Orias¹, Rémy Bayard², Yves Perrodin¹

¹Université de Lyon, ENTPE, CNRS, UMR 5023 LEHNA, 3 Rue Maurice Audin, 69518 Vaulx-en-Velin, France

²Université de Lyon, INSA-Lyon, Laboratoire DEEP, Bât. S. Carnot, 9 rue de la Physique, F-69621 Villeurbanne, France

**corresponding author.* Université de Lyon, ENTPE, CNRS, UMR 5023 LEHNA, 3 Rue Maurice Audin, 69518 Vaulx-en-Velin, France. Tel.: +33 4 72 04 72 90; Fax: +33 4 72 04 77 43.

E-mail address: antoine.gosset@entpe.fr

Abstract

The phenomenon of urbanization leads to considerable pressure on urban/suburban aquatic ecosystems. Urban Wet Weather Discharges (UWWDs) during rainfall events are a major source of pollutants leached onto and into urban surfaces and sewers, which in turn affect aquatic ecosystems. The aim of this study is to assess the ecotoxicity of the different compounds identified in UWWDs and identify the hazard represented by each of them. To this end, hazard quotients (HQ) were calculated for each compound detected in UWWDs, based on their Predicted No Effect Concentration (PNEC) values and their maximum concentration (MEC_{max}) found in the dissolved part of UWWDs. For the 207 compounds identified in UWWDs, sufficient data existed for 165 of them to calculate their PNEC. The ecotoxicity of these compounds is greatly variable. Pesticides represent a high proportion of the wide variety of hazardous compounds whose HQ were calculated (a total of 94 HQ), and they are among the most hazardous pollutants ($HQ > 1000$) transported by stormwater. On the contrary, the hazard of Combined Sewer Overflows (CSO) is mainly linked to heavy metals and pharmaceutical compounds. Consequently, the monitoring of these pollutants is a priority in the future. Lastly, it is to bear in mind that the hazard level of certain pollutants could be underestimated, due to their adsorption on particles, leading their low presence in the dissolved phase of UWWDs. So, an in-depth study of these pollutants will be required to clarify their effects on aquatic organisms.

Introduction

Urban Wet Weather Discharges (UWWDs) are complex and variable polluted matrices, notably due to the different types of catchments leached by the rain, and the variable flows for different events.¹ Many micropollutants have been detected in these effluents as a function of population density and anthropogenic activities. Among them, we can cite here: (i) European priority substances such as Polycyclic Aromatic Hydrocarbons (PAHs),²⁻⁵ diethylexylphthalate,^{4,6} 4-nonylphenol,^{4,6} bisphenol A and alkylphenolethoxylate⁷, (ii) other micropollutants like heavy metals (copper, lead, zinc, etc.), polychlorobiphenyls (PCBs), pesticides, phthalates^{3,5,8,9,10}, and (iii) pharmaceutical residues (PC) in the particular case of the Combined Sewer Overflows (CSO)^{11,12}. These micropollutants have also already been detected in receiving river water having received these discharges.^{13,14} Many of them are potentially hazardous for aquatic organisms. That is why UWWDs have toxic effects on aquatic species,^{15,16,17} and can thus modify biological communities.^{15,18,19,20,21} Treatment to diminish UWWD pollutants through abatement are nonexistent or very limited depending on municipal sewer systems, and raw UWWD are often release into (sub)urban aquatic ecosystems. It is therefore necessary to optimize their management. This optimization can concern: (i) a better monitoring of toxic pollutants and their effects, including continuous monitoring with biosensors;^{22,23} (ii) a specific treatment of pollutants;²⁴ and (iii) the suppression or the substitution at source of the toxic products at the origin of these pollutants (in particular, certain detergents, disinfectants and pharmaceuticals) by less toxic but equally efficient products. For that, it is necessary to identify and prioritize the hazardous pollutants present in UWWDs.

To establish the list of the compounds most concerned by the ecotoxicological hazard represented by a specific urban effluent, many authors have proposed to formulate and apply a hazard quotient based on the ratio “Concentration of the selected pollutant in filtered polluted matrices”/“Predicted No Effect Concentration (PNEC) of the molecules selected”.^{25,26} This quotient is used to distinguish the hazardousness of the different compounds and families of compounds found in UWWDs. Orias and Perrodin²⁵ proposed comparing the maximum Measured Effluent Concentrations (MECmax) measured in hospital effluents to their PNEC value, which is the method used in the current study. This method was also applied by many other authors for contaminated matrices such as rivers polluted by raw or treated municipal wastewaters polluted by various pharmaceuticals,²⁶ pesticides,²⁷ or

lakes by antibiotics.²⁸ The choice of using the maximum value of MEC (MEC_{max}) is to place us in the worst (extreme conditions) case scenario^{25,27} and in the context of the precautionary principle. This method was applied here for the first time to the UWWDs, translating its originality.

Consequently, the main objectives of our study were to: (i) conduct a review of the ecotoxicity data available for the different compounds commonly detected in UWWDs; (ii) assess the concentrations of these pollutants in UWWDs performing a literature review of UWWDs pollutant compositions; and (iii) calculate the potential hazard of these pollutants for aquatic ecosystems. Several studies have already proposed to identify and prioritize hazardous pollutants in UWWDs.²⁹⁻³² Nevertheless, they generally considered only the stormwater (no studies were reported for the CSO). Moreover, these pollutant hazard assessments were based only on the intrinsic properties of the pollutants (bioaccumulation, biodegradability, physicochemistry, ecotoxicity) and did not take into account the concentrations found *in situ*.

In this article, we present the PNEC values inventoried for all the pollutants identified in UWWDs, and the MEC_{max} found in these discharges. Lastly, we calculated the HQ for each pollutant found in UWWDs. We also chose to focus on combined and separate stormwater systems (named here “CSO” and “stormwater” systems) due to the very different sources and types of pollutants they contain.

2. Methodology

All the pollutants used and analyzed (on dissolved and particles phases of UWWDs) in this study were classified into families of chemical compounds, as detailed in Table 1. Scientific research papers for UWWD pollutants identification were obtained using databases such as the Web of Knowledge, Science Direct and Google Scholar, using key search terms (e.g. “stormwater”, “CSO”, “UWWD”, “pollutants”, “chemicals”). An identified paper’s bibliography and citations were also checked to find other suitable studies. Finally, papers used in this study were restricted to peer-reviewed research articles published between 1995 and 2016. Other scientific productions (e.g. conference papers, non-peer reviewed procedures) were not retained for this study.

2.1 Obtaining and determining the PNEC values of micropollutants in UWWDs

To define the hazard of the different pollutants identified in UWWD physico-chemical studies, the first step of our method was to determine their ecotoxicity by seeking or calculating their PNEC. PNEC values were found/calculated for all pollutants sought in physico-chemical studies (on particle and dissolved phases). In this study, the PNEC values were obtained for all the pollutants using the following approach: (i) When they were available, PNEC values were obtained from free European databases^{33,34} (INERIS and AGRITOX) containing ecotoxicological data from international scientific literature, (ii) When values were not available in these databases, PNEC values were calculated using experimental ecotoxicological data found in the EPA ECOTOX database³⁵, following the method proposed by Orias et Perrodin,³⁶ and in agreement with the guidelines proposed by the EU (European Union) in the Technical Guidance Document (TGD)³⁷ (see Table 2). This method is the following: for each compound, bibliographical research using the EPA ECOTOX database allowed identifying all the bioassays performed, the trophic level concerned (decomposers, primary producers, primary consumers and secondary consumers) and the chronicity (or not) of the bioassay (e.g. reproduction). Using all the ecotoxicity data, an Extrapolation Factor (EF), calculated following the criteria presented in Table S1, was applied to the lowest toxicity values to establish the PNEC of the compound concerned. No differentiation was done between the ecotoxicity data used (e.g. EC₅₀, EC₁₀) for the EF and PNEC calculation. However, several minor modifications in comparison to TGD were made: ecotoxicity data on fungi were considered as data on decomposers while data on periphyton were considered to originate from chronic bioassays on decomposers. In the case where only acute ecotoxicity data were available, but not for all trophic levels, PNEC values were calculated with the ECOSAR model, by applying an EF to the lowest ecotoxicological value. In all cases, no PNEC values were calculated if at least one acute ecotoxicity value was available for each trophic level. The chronic algae and macrophyte bioassays did not permit lowering the EF when they were the only ecotoxicological data available. Finally, the EF was not reduced when rotifer (*Brachionus calyciflorus*, noted “Bc”) bioassays were the only tests performed on the trophic level of secondary consumers.³⁷ Finally, when no PNEC and ecotoxicity data were available in databases and literature, PNEC values were estimated with modelled ecotoxicological data obtained using the ECOSAR method.³⁸

It is important to point out that the authors of several studies did not indicate the speciation or the isomer of the compounds measured. This is the case of chromium, for which we chose to keep the two speciation forms (chromium III and chromium VI), as it has been reported that the annual concentrations of these two chromium species in rainwater samples are similar.³⁹ In the case of dichloroethane, we chose to keep the two isomers (1-1 and 1-2-dichloroethane), while the hazard quotient was calculated twice for this pollutant, using two PNECs.

2.2 Obtaining micropollutant MEC_{max}

The second step for UWWD pollutant hazard assessment was the MEC_{max} determination for each of them. As described in Orias and Perrodin²⁵, two different approaches can be used to obtain the concentration of micropollutants in UWWDs: (i) a theoretical method based on the calculation of PEC (Predicted Effluent Concentration) with the design of a scenario, and (ii) an experimental approach based on analyses of the pollutants in the dissolved phase of the UWWDs. The second method (MEC_{max} determination) is more realistic, because it is based on real measured concentrations in UWWDs, contrary to the calculation of PEC values presenting a high level of uncertainty due to the parameters used (e.g. product consumption). Therefore, this option was selected although not all the compounds were analyzed in the discharges.

In this study, we will discuss the different choices made to select concentration data (see section 3.5). Firstly, we considered studies providing compound concentration values only in the soluble phase (<1.2 µm). Indeed, ecotoxicity assays performed to calculate PNEC values for the different pollutants have relied only on pure liquid solutions. Moreover, it is impossible to evaluate the bioavailability of the contaminants in the particulate phase for aquatic organisms. Additionally, taking into account the concentrations on particles could lead to overestimating the hazard (see section 3.5). Regarding the formulation of the Hazard Quotient (HQ), we also chose to separate concentration values for separate and combined sewer systems, thus we calculated the hazard for both cases. The hazards of these two systems differ, as the source and type of pollutants (i.e. the presence of pharmaceuticals in discharges from combined sewer overflows) are also different.

For each compound sought in the dissolved part of the stormwater/CSO samples, the MEC of the events observed was reported for each study. Finally, the highest effluent concentration ever measured in UWWDs (MEC_{max}) was determined by seeking the highest MEC reported previously, with the aim to be situated in the worst-case scenario, as explained in the introduction. In further research, it would be interesting to make the same work with the median and the lowest values, but that will be limited by the fact that for some compounds, only one concentration value was available in scientific literature.

2.3 Calculation of the hazard quotients

Finally, for each compound detected in discharges (section 2.2) with an available PNEC value (section 2.1), we calculated an hazard quotient using the equation (1):

$$(1) \text{ HQ} = \text{MEC}_{\text{max}} / \text{PNEC}$$

With:

MEC_{max} : the highest effluent concentration ever measured in UWWDs in $\mu\text{g.L}^{-1}$;

PNEC: Predicted No Effect Concentration in $\mu\text{g L}^{-1}$.

It was considered that compounds with an HQ up to 1 are potentially hazardous for the environment, and that compounds with the highest HQ are the most hazardous for aquatic ecosystems. On the contrary, compounds with an HQ lower than 1 were considered slightly hazardous.

In this study, 207 compounds were identified in UWWDs. We found or calculated/modelled a PNEC value representing the ecotoxicity of 165 of them. By seeking the dissolved proportion analyzed in UWWDs, we were finally able to determine a MEC_{max} for only 61 compounds in “stormwater” and 49 in “CSO”. Consequently, we calculated 94 hazard quotients using equation (1) to prioritize their hazardousness for freshwater ecosystems.

3. Results and discussion

3.1 Pollutant classification and available data.

Pollutants analyzed in UWWDs were classified as shown in Table 1. All the PNECs, concentrations and hazard data were analyzed as per the pollutant ranking presented in Table 1.

Table 1: Families of compounds considered in this study and colour codes associated in the various figures.

Abbreviation	Family of pollutants	Figure colour code
ALK	Alkylphenols	Blue
FR	Flame Retardants	Orange
M	Metals	Red
O	Others	Black
ORG	Organotins	Grey
P	Pesticides	Green
PAH	Polycyclic Aromatic Hydrocarbons and derivatives	Brown
PBDE	Polybrominated Diphenyl Ethers	White
PC	Pharmaceutical Compounds	Cyan
PCB	Polychlorobiphenyls	Orange
PFAA	Perfluoroalkyl Acids	Yellow
PHT	Phthalates	Dark Grey
VOC	Volatile Organic Compounds	Purple

3.2 Ecotoxicity of the pollutants sought in UWWDs

Of the 207 compounds sought in the UWWD samples (total, dissolved and/or particle phases), 161 were obtained by calculation using data from the European database and the international literature. In addition, we formulated ourselves 52 new PNEC values based on experimental data and 4 were modelled using ECOSAR model, as detailed in Table 2. Finally, we listed a total of 165 PNEC values. Figure 1 presents the distribution of available PNEC values for each pollutant family. This figure shows that 5 families present a large number of molecules for which PNEC data are not available: PAH, PC, ALK, O and PCB. This lack of data poses a problem for estimating UWWD compound hazards.

Figure 1: Distribution of the number of molecules and their PNEC available for the different families of pollutants identified in UWWDs.

The 165 PNEC values obtained have been separated into three categories according to the ranking of Orias and Perrodin (2014): a PNEC less than $0.1 \mu\text{g L}^{-1}$ (the most hazardous, the most ecotoxic) (Figure 2), a PNEC between 1 and $0.1 \mu\text{g L}^{-1}$ (intermediate ecotoxicity) (Figure 3) and a PNEC $> 1 \mu\text{g L}^{-1}$ (less ecotoxic) (Figure 4). Finally, due to the results obtained, 59 of the compounds were ranked as very ecotoxic, 30 as ecotoxic and 76 as slightly ecotoxic.

Among the 59 very ecotoxic compounds, pesticides feature predominantly in the most hazardous category of pollutants sought in UWWDs. Indeed, 33 of the compounds with a PNEC less than $0.1 \mu\text{g L}^{-1}$ are pesticides. Moreover, regarding the PNEC values of a very large number of pesticides sought in UWWDs (54 compounds), they represent a potentially considerable source of hazard for this kind of matrix: 43 compounds have a PNEC value under $1.0 \mu\text{g L}^{-1}$ and 33 have a PNEC value under $0.1 \mu\text{g L}^{-1}$. Two other families represent a significant share of the most ecotoxic pollutants: PCs (7 compounds) and PAHs (also 7). The share of the other families is lower. The most ecotoxic compound is Endrin, a pesticide with a PNEC of $1.22 \cdot 10^{-8} \mu\text{g.L}^{-1}$. However, when examining slightly ecotoxic compounds ($\text{PNEC} > 1 \mu\text{g.L}^{-1}$), VOCs are largely represented, with 24 of the compounds listed in this

category. The gap between the share of the other families is narrower: FR (3 compounds), PHT (6), PFAA (3), PAH (5), P (11), O (5), M (6), PC (10), ALK (1). The least toxic compound sought in UWWDs is Iohexol, a PC with a PNEC value of 45705 $\mu\text{g.L}^{-1}$.

The large number of very ecotoxic pollutants sought in UWWDs can be explained by the fact that many substances found in UWWDs were developed to eliminate certain families of organisms, thus they have a considerable impact on many aquatic species belonging to the same family (the most relevant example is pesticides). Nevertheless, a large number of pollutants of this family present low toxicity regarding PNEC values. This observation may be explained by three possibilities: the pollutant specifically affects the organism targeted without having any effect on the others (the case of similar pesticides), it is only slightly toxic for aquatic organisms, or there are not enough data to lower the PNEC value.

We also note that no or insufficient ecotoxicity data are available for 13 of the compounds (3 PC, 6 PAH, 2 ALK, 1 ORG, 1 PFAA) detected in UWWDs. These compounds include a high number of PCs and PAHs. More ecotoxicity data are necessary for these families of compounds to avoid an underestimation of the hazard represented by UWWDs. Moreover, as it can be seen in Table 2, the PNECs of certain pollutants, such as PCBs, propylparaben and cotinine, were calculated using a small number of ecotoxicological data, and consequently their toxicity could be underestimated. However, we should bear in mind that these PNEC values were created with high safety factors (1000).

How to cite this paper: Gosset, A., Durrieu, C., Orias, F., Bayard, R., Perrodin, Y. (2017) Identification and assessment of ecotoxicological hazards attributable to pollutants in urban wet weather discharges. *Environmental Science: Processes & Impacts* DOI: 10.1039/C7EM00159B

$0,1 \mu\text{g.L}^{-1} < \text{UWWD pollutant PNECs} \leq 1 \mu\text{g.L}^{-1}$

Figure 3: PNEC values between $0.1 \mu\text{g L}^{-1}$ and $1 \mu\text{g L}^{-1}$ for the different compounds sought in UWWDs. Letters in brackets correspond to the classification in Table 1. PNEC values are ranked in ascending order into each separated pollutant family.

229 Figure 4: PNEC > 1 $\mu\text{g L}^{-1}$ for the different compounds sought in UWWDs. Letters in brackets correspond to the classification in Table 1. PNEC values are
230 ranked in ascending order into each separated pollutant family.

231

232 Table 2: Summary of ecotoxicity data and PNEC formulated by calculation or modelling for 60 pollutants sought in UWWDs. (Legend: D: Decomposers; P:
233 Primary producers; 1C: Primary consumers; 2C: Secondary consumers; C: ecotoxicity data available for organism community bioassays; Bc: *Brachionus*
234 *calyciflorus*; EF: Extrapolation factor).

N°CAS Compound		Family	Experimental ecotoxicity										Data used for PNEC calculation				EF	PNEC (ug/L)	Ref.
			Test species	Trophic level				Chronic test				Specie	Parameter	Endpoint	Value (ug/L)				
				D	P	1C	2C	D	P	1C	2C								
104-40-5	4-(para)-nonylphenol	ALK	77+C	0	5	19	36+17 (Bc)	0	5	6	11+17 (Bc)	<i>Pimephales promelas</i>	Reproduction	LOEC	3.31E-02	10	3.31E-03	35	
80-46-6	4-tert-pentylphenol	ALK	4	0	0	1	3	0	0	1	2	<i>Danio rerio</i>	Development	LOEC	3.20E+01	50	6.40E-01	35	
78-51-3	Tris(2-butoxyethyl)phosphate	FR	3	0	0	0	3	0	0	0	0	<i>Danio rerio</i>	Mortality	NOEC	4.00E+03			35	
7439-89-6	Iron	M	6	0	1	2	3	0	1	0	0	<i>Cyprinus carpio</i>	Mortality	LC50	5.60E+02	1000	5.60E-01	35	
86-74-8	Carbazole	O	4	0	1	2	1	0	1	1	0	<i>Pimephales promelas</i>	Mortality	LC50	9.30E+02	50	1.86E+01	35	
789-02-6	2,4'-DDT	P	4	0	0	0	4	0	0	0	2	<i>Oryzias latipes</i>	Reproduction (fertility)	LOEC	2.30E-01	100	2.30E-03	35	
50-29-3	4,4'-DDT	P	40	0	11	21	8	0	11	5	0	<i>Hyallela azteca</i>	Mortality	LC50	7.00E-02	50	1.40E-03	35	
309-00-2	Aldrin	P	55	0	0	18	37	0	0	0	0	<i>Paratelphusa jacquemontii</i>	Mortality	LC50	9.70E-02			35	
218-01-9	Chrysene	P	3	0	0	2	1	0	0	0	0	<i>Aedes aegypti</i>	Mortality	LC50	1.70E+03			35	
1319-77-3	Cresol	P	6	0	1	2	3	0	1	0	0	<i>Gammarus fasciatus</i>	Immobilization	EC50	7.00E+03	1000	7.00E+00	35	
52315-07-8	Cypermethrin	P	68	0	6	28	33+1 (Bc)	0	6	7	2+1 (Bc)	<i>Palaemonetes argentinus</i>	Mortality	LOEC	1.00E-04	10	1.00E-05	35	
6190-65-4	Desethylatrazine	P	11	0	8	3	0	0	7	1	0	<i>Stagnicola elodes</i>	Mortality	NOEC	3.30E-01	50	6.60E-03	35	
1007-28-9	Desethylsimazine	P	10	0	7	2	1	0	7	0	0	<i>Pseudokirchneriella subcapitata</i>	Growth	NOEC	6.00E-01	1000	6.00E-04	35	
333-41-5	Diazinon	P	131	0	9	53	69	0	9	8	18	<i>Daphnia magna</i>	Reproduction	EC50	2.00E-04	10	2.00E-05	35	
60-57-1	Dieldrin	P	98+C	0	7	43	45+1 (Bc)	0	7	2	1+1 (Bc)	<i>Brachionus calyciflorus</i>	Population growth	LOEC	1.00E-03	10	1.00E-04	35	
72-20-8	Endrin	P	89	0	1	22	66	0	1	0	1	<i>Asellus aquaticus</i>	Mortality	LC50	6.10E-07	50	1.22E-08	35	

How to cite this paper: Gosset, A., Durrieu, C., Orias, F., Bayard, R., Perrodin, Y. (2017) Identification and assessment of ecotoxicological hazards attributable to pollutants in urban wet weather discharges. Environmental Science: Processes & Impacts DOI: 10.1039/C7EM00159B

2682-20-4	Methylisothiazolinone	P	3	0	0	1	2	0	0	0	0	<i>Oncorhynchus mykiss</i>	Mortality	LC50	7.00E+01			35
51218-45-2	Metolachlor	P	63	0	37	12	14	0	35	6	5	<i>Stuckenia pectinata</i>	Photosynthesis disturbance	LOEC	5.00E+00	10	5.00E-01	35
134-62-3	N,N-diethyl-3-méthylbenzamide	P	6	0	0	3	3	0	0	1	0	<i>Pimephales promelas</i>	Growth	NOEC	6.00E-01	100	6.00E-03	35
134-62-3	N,N-diethyl-meta-toluamide	P	6	0	0	3	3	0	0	0	0	<i>Pimephales promelas</i>	Growth (condition index)	NOEC	6.00E-01	1000	6.00E-04	35
26530-20-1	N-Octylisothiazolinone	P	6	0	0	1	5	0	0	0	1	<i>Pimephales promelas</i>	Growth	NOEC	8.50E+00	100	8.50E-02	35
52645-53-1	Permethrine	P	4	0	1	2	1	0	0	0	0	Daphnid	Mortality	LC50	3.00E-02	1000	3.00E-05	38
1610-18-0	Prometon	P	15	0	5	1	9	0	5	1	1	<i>Chlorella fusca var. vacuolata</i>	Population growth (growth rate)	EC01	5.00E+00	10	5.00E-01	35
886-50-0	Terbutryn	P	21	0	9	2	10	0	6	0	0	<i>Craticula accomoda</i>	Growth	EC10	1.90E-02	1000	1.90E-05	35
101-20-2	Triclocarban	P	11	0	1	5	5	0	1	2	1	<i>Daphnia magna</i>	Reproduction	NOEC	2.50E-01	10	2.50E-02	35
319-84-6	α -HCH	P	21	0	1	8	12	0	1	2	4	<i>Lymnaea stagnalis</i>	Reproduction	EC50	6.50E+01	10	6.50E+00	35
33213-65-9	β -endosulfan	P	15	1	0	6	8	0	0	0	0	<i>Oncorhynchus mykiss</i>	Mortality	LC50	3.30E+00	1000	3.30E-03	35
319-85-7	β -HCH	P	4	0	0	0	4	0	0	0	0	<i>Oryzias latipes</i>	Behavioural changes (general)	NOEC	3.20E+01			35
90-12-0	1-Methylnaphthalene	PAH	6	0	3	1	2	0	1	0	0	<i>Daphnia magna</i>	Immobilization	EC50	1.42E+03	1000	1.42E+00	35
91-57-6	2-Methylnaphthalene	PAH	5	0	2	1	2	0	0	0	0	<i>Oncorhynchus mykiss</i>	Mortality	EC50	1.46E+03	1000	1.46E+00	35
84-65-1	Anthraquinone	PAH	6	0	2	2	2	0	2	0	0	<i>Lemna gibba</i>	Photosynthesis	IC50	1.00E-02	1000	1.00E-05	35
132-65-0	Dibenzothiophene	PAH	3	0	0	1	2	0	0	0	0	<i>Daphnia magna</i>	Mortality	LC50	4.20E+02			35
119-61-9	Benzophenone	PC	5	0	0	4	1	0	0	0	1	<i>Daphnia magna</i>	Immobilization	EC50	2.80E+02	100	2.80E+00	35
486-56-6	Cotinine	PC	1	0	1	0	0	0	1	0	0	<i>Lemna gibba</i>	Biomass	NOEC	1.00E+03	1000	1.00E+00	35
42399-41-7	Diltiazem	PC	3	0	1	1	1	0	0	0	0	<i>Paspalum vaginatum</i>	Calcium uptake	LOEC	4.16E+03	1000	4.16E+00	35
58-73-1	Diphenhydramine	PC	2	0	0	1	1	0	0	1	0	<i>Ceriodaphnia dubia</i>	Reproduction	LC50	3.94E+03	100	3.94E+01	35
119-65-3	Isoquinoline	PC	5	0	1	2	2	0	1	0	0	<i>Poecilia reticulata</i>	Mortality	LC50	1.40E+04	1000	1.40E+01	35
94-13-3	Propylparaben	PC	2	0	0	2	0	0	0	1	0	<i>Daphnia magna</i>	Immobilization	NOEC	1.00E+03	100	1.00E+01	35

How to cite this paper: Gosset, A., Durrieu, C., Orias, F., Bayard, R., Perrodin, Y. (2017) Identification and assessment of ecotoxicological hazards attributable to pollutants in urban wet weather discharges. Environmental Science: Processes & Impacts DOI: 10.1039/C7EM00159B

37680-73-2	PCB101	PCB	4	0	0	3	1	0	0	0	1	<i>Pimephales promelas</i>	Growth	LOEC	2.50E+00	100	2.50E-02	35
35065-28-2	PCB138	PCB	2	0	0	0	2	0	0	0	1	<i>Pimephales promelas</i>	Growth	LOEC	2.50E+00	100	2.50E-02	35
35065-27-1	PCB153	PCB	4	0	0	1	3	0	0	1	1	<i>Pimephales promelas</i>	Growth	LOEC	2.50E+00	50	5.00E-02	35
35065-29-3	PCB180	PCB	1	0	0	0	1	0	0	0	1	<i>Pimephales promelas</i>	Growth	LOEC	2.50E+00	100	2.50E-02	35
35693-99-3	PCB52	PCB	2	0	0	1	1	0	0	0	1	<i>Pimephales promelas</i>	Reproduction	NOEC	2.50E+01	100	2.50E-01	35
335-67-1	Pentadécafluorooctanoic acid	PFAA	12	0	2	3	6 + 1 (Be)	0	2	2	2	<i>Oryzias latipes</i>	Mortality	LOEC	1.00E+02	10	1.00E+01	35
335-76-2	Perfluorodecanoic acid	PFAA	3	0	1	1	1	0	1	0	0	<i>Danio rerio</i>	Mortality	LC50	8.40E+03	1000	8.40E+00	35
375-85-9	Perfluoroheptanoic acid	PFAA	1	0	1	0	0	0	1	0	0	<i>Chlorella vulgaris</i>	Growth	EC50	1.90E+06	1000	1.90E+03	35
375-95-1	Perfluorononanoic acid	PFAA	4	0	2	1	1	0	2	0	0	<i>Danio rerio</i>	Growth	LOEC	1.00E+01	1000	1.00E-02	35
1763-23-1	Perfluorooctanesulfonic acid	PFAA	4	0	0	2	2	0	0	1	0	<i>Oryzias latipes</i>	Mortality	LOEC	1.00E+01	100	1.00E-01	35
2058-94-8	Perfluoroundecanoic acid	PFAA	2	0	0	1	1	0	0	0	0	<i>Daphnia magna</i>	Immobilization	NOEC	5.64E+04			35
117-81-7	Diethylexyl phthalate	PHT	29	0	3	5	21	0	3	2	3	<i>Danio rerio</i>	Reproduction	NOEC	2.00E+00	10	2.00E-01	35
26761-40-0	Diisodecyl phthalate	PHT	7	0	1	1	5	0	1	1	0	<i>Daphnia magna</i>	Mortality	LOEC	6.00E+01	50	1.20E+00	35
27554-26-3	Diisononyl phthalate	PHT	7	0	1	2	4	0	1	1	0	<i>Daphnia magna</i>	Reproduction	NOEC	6.20E+01	50	1.24E+00	35
131-11-3	Dimethyl phthalate	PHT	9	0	1	2	6	0	1	0	0	<i>Cyprinodon variegatus</i>	Mortality	NOEC	3.20E+03	1000	3.20E+00	35
98-86-2	Acetophenone	VOC	4	0	1	2	1	0	0	0	0	<i>Tetrahymena pyriformis</i>	Population Growth	EC50	4.28E+04	1000	4.28E+01	35
119-61-9	Benzophenone	VOC	5	0	0	4	1	0	0	1	0	<i>Daphnia magna</i>	Immobilization	EC50	2.80E+02	100	2.80E+00	35
75-25-2	Bromoform	VOC	5	0	1	2	2	0	1	0	0	<i>Cyprinodon variegatus</i>	Mortality	NOEC	2.90E+03	1000	2.90E+00	35
138-86-3	D-Limonene	VOC	11	0	1	7	3	0	1	0	0	<i>Lymnaea acuminata</i>	Mortality	LC50	1.13E+03	1000	1.13E+00	35
107-21-1	Ethylene glycol	VOC	20	1	1	7	11	0	1	1	1	<i>Ceriodaphnia dubia</i>	Reproduction	NOEC	4.20E+03	10	4.20E+02	35
120-72-9	Indole	VOC	3	0	1	2	0	0	1	1	0	<i>Oryzias latipes</i>	Mortality	LC50	1.00E+03	100	1.00E+01	35
124-76-5	Isoborneol	VOC	1	0	0	0	1	0	0	0	0	Fish	Mortality	LC50	5.90E+04			38

3.3 Concentration of pollutants in separate and combined systems

Maximum pollutant concentrations found in UWWDs were assessed using available literature and are presented in Tables 3 and 4. MEC_{max} were very variable, with some pollutants (e.g. hexachlorobenzene) found at several ng.L⁻¹, while others (e.g. copper) were found at a few mg.L⁻¹. However, the compound concentrations generally ranged between ng.L⁻¹ and several hundred µg.L⁻¹.

On one hand, as observed in Table 3, 61 compounds were detected in stormwater samples, and 50 were sought but not detected. We found the highest concentrations for zinc (2.12 mg.L⁻¹). High concentrations of pollutants such as other heavy metals (copper, nickel) and pesticides (such as glyphosate and AMPA) and PAH (e.g. pyrene) were also detected. The main families of pollutants detected at high concentrations were PAH, VOC, ALK, PHT, P and M.

On the other hand, out of the 81 chemicals of interest sought in the dissolved part of the CSO samples, 49 were reported (see Table 4). The highest concentration was obtained for zinc (236 µg.L⁻¹). The main families found at high concentrations were M, P and PC. The occurrence of high concentrations of heavy metals in separate systems and CSO is not surprising due to the source of wastewater and rainfall runoff. More precisely, tyre wear, vehicle brake deposition, conveyance system infrastructure (e.g. downspouts and culverts), and atmospheric deposition can lead to the presence of heavy metals.⁴ Moreover, the remobilization of wastewater and in-sewer deposits (due to pollutant water/particle partition) can represent a large share of the pollutant load in CSO, such as Cu and PAH.⁵⁸ Launay et al.⁶⁴ observed the same phenomenon for pharmaceutical compounds and industrial chemicals. Contrary to stormwater, we note the presence of PC compounds sought and detected in the dissolved phase (18 compounds), ~~probably~~ due to the contribution of urban wastewater to the pollution of CSO.

If we compare the Maximum concentrations of UWWD pollutants with other urban effluents (e.g. urban wastewater and hospital wastewater), such as phthalates, heavy metals, pharmaceutical compounds and pesticides, MEC_{max} were close to those detected in these other matrices.^{60,61,62,63} Some of them, such as Bisphenol A, concentrations in UWWDs (max. 130 µg L⁻¹) can be even higher than in wastewater (max. 2.14 µg L⁻¹). This fact was also observed with heavy metals such as copper and zinc.

Consequently, since they undergo hardly any or no treatment, and that most of few current treatment

How to cite this paper: Gosset, A., Durrieu, C., Orias, F., Bayard, R., Perrodin, Y. (2017) Identification and assessment of ecotoxicological hazards attributable to pollutants in urban wet weather discharges. Environmental Science: Processes & Impacts DOI: 10.1039/C7EM00159B

methods employed do not reduce concentration of many identified pollutants, considerable amounts of these substances may enter surface waters.

Nevertheless, it is important to point out that the data collected and discussed above have some limitations. Firstly, few studies have analyzed the dissolved phase of discharges and thus many compounds identified and sought in UWWDs are not taken into account here because they are found in only the particle or total phase (due to their low solubility and high affinity for the particle phase). Consequently, the number of compounds detected here is limited (see Table 3 and Table 4). Secondly, concentrations found in UWWDs need to be considered with caution because, as stated in the study of Huber et al.,⁵² concentrations measured in stormwater samples can be influenced by different parameters such as sample collection, preparation and analysis, which can also affect the concentrations observed in discharges.

286 Table 3: Maximum Measured Effluent Concentrations (MECmax) of micropollutants in the dissolved part sought in stormwater samples (ND: Not detected, *:
287 Unspecified).

Pollutant family	Pollutant	MECmax (µg/L)	Country	Ref.	Pollutant family	Pollutant	MECmax (µg/L)	Country	Ref.
ALK	4-n-octylphenol (ALK)	ND	France	4	PAH	Dibenzothiophene (PAH)	0.0417	USA (Washington D.C.)	2
ALK	4-nonylphenol di-ethoxylate (ALK)	ND	France	4	PAH	Acenaphthylene (PAH)	0.070	Australia; France; Sweden; USA (Alabama); USA (Washington D.C.)	2 ; 4 ; 5 ; 47 ; 48
ALK	4-tert-pentylphenol (ALK)	0.7	Sweden	5	PAH	Perylene (PAH)	0.12	USA (Washington D.C.)	2
ALK	Para-tert-octylphenol (ALK)	0.99	France, Sweden	4 ; 5	PAH	Dibenzo[a,h]anthracene (PAH)	0.13	France; Sweden; USA (Alabama); USA (Washington D.C.)	2 ; 4 ; 5 ; 47
ALK	4-Octylphenol monoethoxylate (ALK)	4.39	Sweden	5	PAH	Benzo[a]anthracene (PAH)	0.34	Australia; France; Sweden; USA (Alabama); USA (South Carolina); USA (Washington D.C.)	2 ; 4 ; 5 ; 47 ; 48 ; 49
ALK	4-(para)-nonylphenol (ALK)	7.7	France, Sweden	4 ; 5	PAH	Benzo[k]fluoranthene (PAH)	0.40	France; Sweden; USA (Alabama); USA (Washington D.C.)	2 ; 4 ; 5 ; 47
ALK	4-tert-butylphenol (ALK)	7.82	France, Sweden	4 ; 5	PAH	Benzo[e]pyrene (PAH)	0.42	USA (South Carolina); USA (Washington D.C.)	2 ; 48
ALK	4-nonylphenol mono-ethoxylate (ALK)	8.36	Sweden	5	PAH	Indeno[1,2,3-cd]pyrene (PAH)	505	Australia; France; Sweden; USA (Alabama); USA (South Carolina); USA (Washington D.C.)	2 ; 4 ; 5 ; 47 ; 48 ; 49
M	Platinum (M)	ND	France	4	PAH	Benzo[a]pyrene (PAH)	0.54	Australia; France; Sweden; USA (Alabama); USA (South Carolina); USA (Washington D.C.)	2 ; 4 ; 5 ; 47 ; 48 ; 49
M	Mercury (M)	0.01	Canada; France	4 ; 41	PAH	Benzo[g,h,i]perylene (PAH)	0.55	Australia; Sweden; USA (Alabama); USA (South Carolina); USA (Washington D.C.)	2 ; 5 ; 47 ; 48 ; 49
M	Lead (M)	15.9	Australia; China; France; USA (Alabama); USA (California); *	4 ; 5 ; 47 ; 50 ; 51 ; 52	PAH	Anthracene (PAH)	0.56	Australia; France; Sweden; USA (Alabama); USA (South Carolina); USA (Washington D.C.)	2 ; 4 ; 5 ; 47 ; 48 ; 49
M	Cadmium (M)	31.1	Australia; France; USA (Alabama); USA (California); *	4 ; 47 ; 50 ; 52 ; 55	PAH	Chrysene (PAH)	0.62	Australia; France; Sweden; USA (Alabama); USA (South Carolina); USA (Washington D.C.)	2 ; 4 ; 5 ; 47 ; 48 ; 49
M	Chromium (M)	32	Australia; France; USA (Alabama); USA (California); *	4 ; 47 ; 50 ; 52 ; 55	PAH	Acenaphthene (PAH)	0.72	Australia; France; Sweden; USA (Alabama); USA (Washington D.C.)	4 ; 5 ; 47 ; 49
M	Nickel (M)	49	France; USA (Alabama); USA (California); *	4 ; 5 ; 47 ; 52	PAH	Benzo[b]fluoranthene (PAH)	0.98	Australia; France; USA (Alabama); USA (South Carolina); USA (Washington D.C.)	2 ; 4 ; 47 ; 48 ; 49

How to cite this paper: Gosset, A., Durrieu, C., Orias, F., Bayard, R., Perrodin, Y. (2017) Identification and assessment of ecotoxicological hazards attributable to pollutants in urban wet weather discharges. Environmental Science: Processes & Impacts DOI: 10.1039/C7EM00159B

M	Copper (M)	1520	Australia; China; France; USA (Alabama); USA (California); *	4 ; 5 ; 47 ; 50 ; 51 ; 52	PAH	Phenanthrene (PAH)	1.32	Australia; France; Sweden; USA (Alabama); USA (South Carolina); USA (Washington D.C.)	2 ; 4 ; 5 ; 47 ; 48 ; 49
M	Zinc (M)	2118	Australia; China; France; USA (Alabama); USA (California); *	4 ; 5 ; 47 ; 50 ; 51 ; 52	PAH	Fluorene (PAH)	1.37	Australia; France; Sweden; USA (Alabama); USA (South Carolina); USA (Washington D.C.)	2 ; 4 ; 5 ; 47 ; 48 ; 49
O	Bisphenol A (O)	130	Sweden	5	PAH	Fluoranthene (PAH)	14	Australia; France; Sweden; USA (Alabama); USA (South Carolina); USA (Washington D.C.)	2 ; 4 ; 5 ; 47 ; 48 ; 49
ORG	Dibutyltin (ORG)	ND	France	4	PAH	Naphthalene (PAH)	17	Australia; France; Sweden; USA (Alabama); USA (Washington D.C.)	2 ; 4 ; 5 ; 47 ; 49
ORG	Tributyltin (ORG)	ND	France	4	PAH	Pyrene (PAH)	20	Australia; France; Sweden; USA (Alabama); USA (South Carolina); USA (Washington D.C.)	2 ; 4 ; 5 ; 47 ; 48 ; 49
ORG	Monobutyltin (ORG)	0.025	France	4	PBDE	Decabromodiphenyl ether (PBDE)	ND	France	4
P	2,4'-DDT (P)	ND	France	4	PBDE	Octabromodiphenyl ether (PBDE)	ND	France	4
P	4,4'-DDT (P)	ND	France	4	PBDE	Pentabromodiphenyl ether (PBDE)	ND	France	4
P	Alachlor (P)	ND	France	4	PCB	PCB101 (PCB)	ND	France	4
P	Aldrin (P)	ND	France; USA (Alabama)	4 ; 47	PCB	PCB118 (PCB)	ND	France	4
P	Atrazine (P)	ND	France	4	PCB	PCB138 (PCB)	ND	France	4
P	Chlordane (P)	ND	France; USA (Alabama)	4 ; 47	PCB	PCB153 (PCB)	ND	France	4
P	Chlorfenvinphos (P)	ND	France	4	PCB	PCB180 (PCB)	ND	France	4
P	Chlorpyrifos (P)	ND	France	4	PCB	PCB194 (PCB)	ND	France	4
P	Desethylsimazine (P)	ND	France	4	PCB	PCB28 (PCB)	ND	France	4
P	Dieldrin (P)	ND	France	4	PCB	PCB52 (PCB)	ND	France	4
P	Endrin (P)	ND	France; USA (Alabama)	4 ; 47	PFAA	Perfluoroheptanoic acid (PFAA)	0.011	Japan; Singapour; Switzerland; USA (Minnesota)	42 ; 43 ; 44 ; 45
P	Esfenvalerate (P)	ND	France; USA (Alabama)	4 ; 47	PFAA	Perfluoroundecanoic acid (PFAA)	0.045	Japan; Singapour; Switzerland; USA (Minnesota)	42 ; 43 ; 44 ; 45 ; 46
P	Lindane (P)	ND	France	4	PFAA	Perfluorodecanoic acid (PFAA)	0.077	Japan; Singapour; Switzerland; USA (Minnesota)	42 ; 43 ; 44 ; 45 ; 46
P	Simazine (P)	ND	France	4	PFAA	Perfluorononanoic acid (PFAA)	0.093	Japan; Singapour; Switzerland; USA (Minnesota)	42 ; 43 ; 44 ; 45 ; 46

P	Trifluralin (P)	ND	France	4	PFAA	Perfluorooctanesulfonic acid (PFAA)	0.16	Japan; Singapour; Switzerland; USA (Minnesota)	42 ; 43 ; 44 ; 45 ; 46
P	α -endosulfan (P)	ND	France	4	PFAA	Pentadecafluorooctanoic acid (PFAA)	0.17	Japan; Singapour; Switzerland; USA (Minnesota)	42 ; 43 ; 44 ; 45 ; 46
P	β -endosulfan (P)	ND	France	4	PHT	Dimethyl phthalate (PHT)	ND	USA (Alabama)	47
P	Lambda-cyhalothrin (P)	0.007	USA (California)	40	PHT	Di-n-octyl phthalate (PHT)	ND	France	4
P	Cyfluthrin (P)	0.023	USA (California)	40	PHT	Di-n-butyl phthalate (PHT)	1	Sweden; USA (Alabama)	5 ; 47
P	Cypermethrin (P)	0.026	USA (California)	40	PHT	Diisodecyl phthalate (PHT)	1.2	Sweden	5
P	Bifenthrin (P)	0.034	USA (California)	40	PHT	Diisononyl phthalate (PHT)	2	Sweden	5
P	Metaldehyde (P)	0.05	France	4	PHT	Diethyl phthalate (PHT)	15.9	Sweden; USA (Alabama)	5 ; 47
P	Permethrin (P)	0.13	USA (California)	40	PHT	Benzyl butyl phthalate (PHT)	16	Sweden; USA (Alabama)	5 ; 47
P	Deltamethrin (P)	0.25	USA (California)	40	PHT	Diethylhexyl phthalate (PHT)	45	France; Sweden; USA (Alabama)	4 ; 5 ; 47
P	Aminotriazole (P)	0.5	France	4 ; 47 ; 48	VOC	1,2,3-Trichlorobenzene (VOC)	ND	France	4
P	Isoproturon (P)	0.6	France	4	VOC	1,2,4-Trichlorobenzene (VOC)	ND	France	4
P	Diuron (P)	0.64	France	4	VOC	1,3,5-Trichlorobenzene (VOC)	ND	France	4
P	AMPA (P)	19	France; Switzerland	4 ; 53 ; 54	VOC	1-2 dichloroéthane (VOC)	ND	France	4
P	Glyphosate (P)	150	France; Switzerland	4 ; 53 ; 54	VOC	Benzène (VOC)	ND	France	4
					VOC	Chloroform (VOC)	ND	France	4
					VOC	Ethylbenzene (VOC)	ND	France	4
					VOC	Hexachlorobenzene (VOC)	ND	France	4
					VOC	Hexachlorobutadiene (VOC)	ND	France	4
					VOC	Isopropylbenzene (VOC)	ND	France	4
					VOC	Pentachlorobenzene (VOC)	ND	France	4
					VOC	Tetrachloroethylene (VOC)	ND	France	4
					VOC	Toluene (VOC)	ND	France	4
					VOC	Trichloroethylene (VOC)	ND	France	4
					VOC	Xylene (VOC)	ND	France	4
					VOC	Pentachlorophenol (VOC)	0.2	France	4
					VOC	Dichloromethane (VOC)	13	France	4
					VOC	1,4-Dichlorobenzene (VOC)	47	USA (Alabama)	48

290 Table 4: Maximum Measured Effluent Concentrations (MECmax) of the micropollutants sought in the dissolved part of CSO samples (ND: Not detected, *:
291 Unspecified).

Pollutant family	Pollutant	MECmax (µg/L)	Country	Ref.	Pollutant family	Pollutant	MECmax (µg/L)	Country	Ref.
ALK	4-n-octylphenol (ALK)	ND	France	57	PC	Estrone (PC)	ND	South Korea	12
ALK	Para-tert-octylphenol (ALK)	ND	France	57	PC	Diltiazem (PC)	ND	South Korea	12
ALK	4-tert-butylphenol (ALK)	0.080	France	57	PC	Trimethoprim (PC)	ND	South Korea	12
ALK	Nonylphenol (ALK)	0.390	France	57	PC	Meprobamate (PC)	ND	South Korea	12
FR	Tris(2-chloro-1-methylethyl) phosphate (FR)	1.140	South Korea	12	PC	Diclofenac (PC)	ND	South Korea	12
M	Nickel (M)	1.1	France	15 ; 57 ; 58	PC	Gemfibrozil (PC)	0.006	South Korea	12
M	Chromium (M)	1.8	France	15 ; 57 ; 58	PC	Diphenhydramine (PC)	0.023	South Korea	12
M	Cadmium (M)	9.9	France, United Kingdom	15 ; 57 ; 58 ; 59	PC	Primidone (PC)	0.040	South Korea	12
M	Copper (M)	65.4	France, United Kingdom	15 ; 57 ; 58 ; 59	PC	Triclosan (PC)	0.055	South Korea	12
M	Lead (M)	143.4	France, United Kingdom	15 ; 57 ; 58 ; 59	PC	Carbamazepine (PC)	0.060	South Korea	12
M	Zinc (M)	236.1	France, United Kingdom	15 ; 57 ; 58 ; 59	PC	Propylparaben (PC)	0.069	South Korea	12
O	1H-benzotriazole (O)	0.078	South Korea	12	PC	Sulfamethoxazole (PC)	0.070	South Korea	12
ORG	Dibutyltin (ORG)	ND	France	57	PC	Trimethoprim (PC)	0.073	South Korea	12
ORG	Tributyltin (ORG)	ND	France	57	PC	Benzophenone (PC)	0.136	South Korea	12
ORG	Monobutyltin (ORG)	0.013	France	57	PC	Atenolol (PC)	0.363	South Korea	12
P	Aldrin (P)	ND	France	57	PC	Acesulfame (PC)	0.727	South Korea	12
P	Dieldrin (P)	ND	France	57	PC	Naproxen (PC)	0.750	South Korea	12
P	Desethylatrazine (P)	0.017	France	57	PC	Ibuprofen (PC)	0.773	South Korea	12
P	Atrazine (P)	0.030	France	57	PC	Sucralose (PC)	0.818	South Korea	12
P	N,N-diéthyl-3-méthylbenzamide (P)	0.035	South Korea	12	PC	Iopamidol (PC)	1.550	South Korea	12
P	Isoproturon (P)	0.040	France	57	PC	Ioexol (PC)	1.770	South Korea	12
P	Triclocarban (P)	0.061	South Korea	12	PC	Caffeine (PC)	2,54	South Korea	12

How to cite this paper: Gosset, A., Durrieu, C., Orias, F., Bayard, R., Perrodin, Y. (2017) Identification and assessment of ecotoxicological hazards attributable to pollutants in urban wet weather discharges. Environmental Science: Processes & Impacts DOI: 10.1039/C7EM00159B

P	Aminotriazole (P)	0.460	France	57	PC	Iopromide (PC)	2,66	South Korea	12
P	Diuron (P)	0.470	France	57	PCB	PCB101 (PCB)	ND	France	57
P	AMPA (P)	1.600	France; Switzerland	53 ; 57	PCB	PCB118 (PCB)	ND	France	57
P	Glyphosate (P)	3.4	France; Switzerland	53 ; 57	PCB	PCB138 (PCB)	ND	France	57
PAH	Acenaphthene (PAH)	ND	France	15 ; 57	PCB	PCB153 (PCB)	ND	France	57
PAH	Benzo[a]anthracene (PAH)	ND	France	15 ; 57	PCB	PCB180 (PCB)	ND	France	57
PAH	Benzo[b]fluoranthene (PAH)	ND	France	15 ; 57	PCB	PCB194 (PCB)	ND	France	57
PAH	Benzo[g,h,i]perylene (PAH)	ND	France	15 ; 57	PCB	PCB28 (PCB)	ND	France	57
PAH	Benzo[k]fluoranthene (PAH)	ND	France	15 ; 57	PCB	PCB52 (PCB)	ND	France	57
PAH	Chrysene (PAH)	ND	France	15 ; 57	PHT	Diethylhexyl phthalate (PHT)	ND	France	57
PAH	Dibenzo[a,h]anthracene (PAH)	ND	France	15 ; 57	VOC	Ethylbenzene (VOC)	ND	France	57
PAH	Benzo[a]pyrene (PAH)	0.005	France	15 ; 57	VOC	Tetrachloroethylene (VOC)	ND	France	57
PAH	Benzo[e]pyrene (PAH)	0.006	France	15 ; 57	VOC	Toluene (VOC)	ND	France	57
PAH	Indeno[1,2,3-cd]pyrene (PAH)	0.008	France	15 ; 57	VOC	Trichloroethylene (VOC)	ND	France	57
PAH	Anthracene (PAH)	0.009	France	15 ; 57	VOC	Xylene (VOC)	ND	France	57
PAH	Acenaphthylene (PAH)	0.020	France	15 ; 57	VOC	Hexachlorobenzene (VOC)	0.002	USA (New York)	56
PAH	Pyrene (PAH)	0.021	France	15 ; 57					
PAH	Fluoranthene (PAH)	0.025	France	15 ; 57					
PAH	Fluorene (PAH)	0.027	France	15 ; 57					
PAH	Phenanthrene (PAH)	0.085	France	15 ; 57					
PAH	Naphthalene (PAH)	0.180	France	15 ; 57					

292

293

3.4 Hazard of pollutants in separate and combined systems

Using ecotoxicological (PNEC) and chemical (MEC_{max}) results previously enunciated we were able to calculate a very wide range of HQs in discharges (see Figures 5 and 6). The high dispersion of HQ values calculated could be explained by the heterogeneity of PNEC values observed in Figures 2 to 4 and the heterogeneity of pollutant concentrations due to the high variability of anthropic activities, and thus that of the catchment leached by rain, and the variability of rainfall events (length, height, flow rate, etc.). Due to the relatively different nature and composition of stormwater and CSO, pollutant hazards associated to these two matrices are discussed separately in the following sections.

3.4.1 Hazard quotients in separate systems

Concerning stormwater samples, only 51 hazard quotients were calculated due to the low number of compounds found in the dissolved part (Figure 5). We found 14 compounds with an HQ lower than 1. The lowest hazard quotient was calculated for Metaldehyde (P) with an HQ of $6.7 \cdot 10^{-5}$. On the contrary, the majority of the micropollutants detected in stormwater samples had an HQ as high as 1 and are therefore potentially hazardous for aquatic ecosystems. Among these 37 compounds, 10 (27.1%) were pesticides, and 9 PAHs (24.32%). It is also noteworthy that 4 of the 6 most hazardous compounds ($HQ > 1000$) were pesticides. Among these very hazardous compounds, we also observed the presence of one ALK (4-p-nonylphenol) and one PAH (fluoranthene). The most hazardous compound found was permethrin, with a hazard quotient of 4166.6. The important hazard from pesticides (pyrethroids) linked to high HQ could be explained by the high ecotoxicity of these pesticides rather than by the concentrations found in discharges (around 100 ng.L^{-1}). As pointed out before, most of these pesticides (such as bifenthrin, cypermethrin) have a very low PNEC value, lower than 0.1 ng.L^{-1} .

3.4.2 Hazard quotients in combined systems

Regarding CSO, only 43 HQs were calculated, also due to the low number of molecules measured in dissolved phase in CSO (Figure 6). The HQs obtained were lower than for separate systems.

Indeed, the highest HQ was found for lead, with a value of 68.29. Contrary to separate systems, the highest hazards were due to heavy metals, pesticides and pharmaceutical compounds. Only five families were found in this ranking ($HQ > 1$): PC (3 compounds), M (4), PAH (1), P (5) and ALK (1), and were potentially hazardous for aquatic ecosystems. Results highlighted for heavy metals can be explained by the fact that certain heavy metals in these effluents were in high concentrations (e.g. $2.1 \mu\text{g.L}^{-1}$ for lead and $1.6 \mu\text{g.L}^{-1}$ for copper), and not because of their low ecotoxicity, as shown by their relatively high PNEC values. The presence of hazardous PC in CSO can be easily explained by the discharge of large amounts of these often very ecotoxic molecules coming from urban wastewater in stormwater, without any treatment. Comparing hazard quotients from CSO samples to those of separate systems, we can observe that calculated HQs are often lower (e.g. 65.4 in CSO against 950 in stormwater for copper). We can simply explain these results by a dilution of stormwater pollutants by wastewater in CSO.

3.4.3 The case of pharmaceutical compounds in CSO studies

With the hazardous pharmaceutical compounds that have been highlighted above, 15 HQs have been calculated only and presented in Figure 6. As urban wastewater is mixed with stormwater in combined sewers, all the pharmaceutical compounds classically consumed in cities could be found in this type of discharge. However, few of them (28) have been taken into account to date in scientific literature. Consequently, our study and all the other ecotoxicological studies dealing with CSO contain a bias attributable to the fact that the potentially considerable hazard represented by these types of molecules is not taken into account. Indeed, as shown by Orias et al.³⁶, many of them can have very low PNEC values (e.g. 17β -estradiol: 8 pg.L^{-1} , clotrimazole: 0.5 pg.L^{-1}), and/or are widely consumed (e.g. ibuprofen and salicylic acid).

3.4.4 The case of molecules (e.g., pesticides) forbidden by European legislation

Some of the pesticides involved in the hazard of UWWDs (found in both stormwater and CSO, such as diuron) were forbidden in 2008 by European legislation. However, several European studies^{65,66,67} have detected the presence of these pesticides (with concentrations reaching $0.64 \mu\text{g L}^{-1}$ in discharges). This observation can be notably explained by the fact that farmers have stored pesticides

345 purchased in the past and that they continue to use. Moreover, some of them linger on in the environment
346 for long periods (due to their high half-life times) and their release can continue due to rainfall events.
347 Consequently, it is true to say that the hazard represented by these pesticides continues to exist even if
348 they are no longer on the market.

349 Figure 5: Hazard quotients calculated for micropollutants detected in stormwater samples (the red line delimits the compounds with an HQ up to 1000,
 350 considered very hazardous). Hazard quotients are ranked in ascending order into each separated pollutant family.

How to cite this paper: Gosset, A., Durrieu, C., Orias, F., Bayard, R., Perrodin, Y. (2017) Identification and assessment of ecotoxicological hazards attributable to pollutants in urban wet weather discharges. Environmental Science: Processes & Impacts DOI: 10.1039/C7EM00159B

351 Figure 6: Hazard quotients calculated for micropollutants detected in CSO samples. Hazard quotients are ranked in ascending order into each separated
 352 pollutant family.

353

How to cite this paper: Gosset, A., Durrieu, C., Orias, F., Bayard, R., Perrodin, Y. (2017) Identification and assessment of ecotoxicological hazards attributable to pollutants in urban wet weather discharges. Environmental Science: Processes & Impacts DOI: 10.1039/C7EM00159B

3.5 Discussion of the method and data used

The method presented in this study proved to be relevant to detect UWWD pollutant hazards. It could be applied for many other matrices, and included in many risk assessment scenarios or chemical prioritization programs. Nevertheless, our method and results are subject to limitations that need to be developed. The first one is the non-consideration of certain parameters, such as the bioconcentration, bioaccumulation³⁰ and biomagnification of pollutants in the aquatic food web, which could amplify considerably the hazard of UWWDs pollutants. Indeed, these phenomena have already been demonstrated for many families of pollutants, such as the most hazardous found in UWWDs: heavy metals⁶⁸, pesticides^{69,70} and pharmaceutical compounds.^{71,72,73,74} Other parameters such as pH, temperature, water chemistry (pH, hardness) can also modify pollutant toxicity. The “mixture” effect, linked to antagonisms or synergisms effects between sought, but also non-potentially thousand molecules not yet identified in UWWDs, are not considered into this study and can modify their hazardousness. Secondly, the use of NOECs for PNEC values construction can be discussed. The authors are aware that ECx values are more accurate than NOEC. ECx values are calculated from the modeling of the dose-response curve, which is itself based on the results of the whole range of concentration, whereas NOEC corresponds only to the identification of the last concentration of the range without effect. However, in the context of this work, it has been chosen to use to the largest set of values available in the literature and in international databases for the two reasons: (1) In the specific context of the PNEC calculation (i.e., a no-effect value for all aquatic organisms), considering the scarcity of existing data in the literature, it seems more useful to consider values for one or several additional species than using accurate values for a limited number of species. In other words, inter-species variability is more important to take into account than intra-species variability, (2) This is the most common approach adopted by international Agencies in charge of the establishment of PNECs for substances, such as INERIS in France, or the European Chemical Bureau in Europe, through the TGD guidelines. Thirdly, our method considers only the maximum concentrations of the different pollutants in UWWDs. Consequently, these values represent only the maximum hazard for each compound. This work could also have been done with median values if sufficient data had been available. This problem

How to cite this paper: Gosset, A., Durrieu, C., Orias, F., Bayard, R., Perrodin, Y. (2017) Identification and assessment of ecotoxicological hazards attributable to pollutants in urban wet weather discharges. Environmental Science: Processes & Impacts DOI: 10.1039/C7EM00159B

of limited data collection is at the origin of another limit: most of the chemical studies focusing on dissolved compound concentrations found in stormwater and CSO have dealt with relatively small areas of the world and many of them were published in Europe. Thus data are lacking for some continents, a factor which limits our studies due to differences in, for example, PC consumption, pesticide use (e.g. glyphosate is widely used in South America), sewer systems, etc. Consequently, we can confidently assert that the HQs calculated here are underestimated and would probably increase if more data were available. Fourthly, dissolved part of UWWDs were only considered in this study because the PNEC values were established for dissolved substances and it was impossible to extend them to pollutants whose bioavailability is unknown but present on particles in UWWDs. Indeed, pollutant bioavailability is linked to particle desorption which depends on multiple parameters like pH, molar mass, solubility/lipophily, and the number of carboxyl groups (for organic pollutants).⁷⁵ It is therefore possible that our method underestimated the hazard of these UWWD pollutants for aquatic environments. For example, as in the study of Zgheib et al.⁴ if we had considered the concentration of fluoranthene in the dissolved phase ($0.018 \mu\text{g L}^{-1}$) and the total effluent ($0.832 \mu\text{g L}^{-1}$), assuming that all the pollutants are available, the hazard index of the pollutants would have increased from 1.5 to 69.33, making it one of the most hazardous compounds for aquatic organisms. This point is illustrated in Figure 7. Similar findings were recorded by Baun et al.³⁰ who prioritized 56 organic pollutants in the water phase of stormwater, against 121 in the solid phase. Fifthly, many chemical studies have considered only mineral and organic priority pollutants from the EU Water Framework Directive and other international legislations. However, many emerging compounds currently used present proven ecotoxic effects but have not yet been analyzed. For example, pesticides used massively in Europe like metam-sodium and epoxiconazole present relatively low PNEC values (respectively 0.79 and $0.18 \mu\text{g L}^{-1}$). We can therefore expect high hazard quotients linked to these kinds of molecule. Finally, the presence of high concentrations of bacteria in CSO has already been detected, for example by Passerat et al. (with concentrations up to 10^7 cell/L for *E.coli*).⁷⁶ It is now known that bacteria can transform pollutants and create by-products that are more or less toxic than their parent compounds detected in CSO. These bacteria can therefore influence the composition of CSO and their toxicity with time.

Despite these limits, our method enabled us to highlight new outlooks in terms of prioritization and management of UWWDD pollutants. Comparing our results to previous works done with the same purpose, we can observe that considering only intrinsic characteristics (e.g. particle solubility, sorption potential, ecotoxicity) of the pollutants selected, Baun et al.³⁰ classified 56 organic compounds as priority pollutants dissolved in water using their RICH (Ranking and Identification of Chemical Hazards) method against 37 in our case (of which 31 are organic). The difference between the two studies is particularly notable for pesticides (10 against 33 pesticides for the method of Baun et al.³⁰). This difference can be explained by non-detected or low environmental concentrations of many pesticides dissolved in water of our studied discharges. On the contrary, we revealed here 9 hazardous PAH and 3 hazardous PHT instead of respectively 5 and 2 in Baun et al.³⁰ An explanation of this discrepancy could be the multiplication of decision filters (sorption/volatilization, technical problems, persistence, bioaccumulation, toxicity) applied by Baun et al. when selecting their compounds, and the lack of data for some pollutants that can lead to the rejection or selection of the priority list, as suggested by these authors. In the case of CSO linked hazard, no studies have investigated a similar prioritization method. Consequently, no comparison can be done.

Looking for the scientific literature, our study enabled us to find new hazardous and priority pollutants. The same observation could be made for the European (Urban Wastewater Directive (UWD 91/271/EEC)) and American (USA EPA) priority pollutant lists. In fact, as stated in Table S2, some pollutant identified here as hazardous such as diuron or fluoranthene are already under the regulation of international instances. On the contrary, some pollutants are absent from these lists. For example, some heavy metals (e.g. copper and zinc in UWD list), pesticides as glyphosate or pyrethrinoids in stormwater, phthalates (e.g. diisononyl phthalate) and pharmaceuticals such as ibuprofen or trimethoprim in CSO absent in UWD and EPA priority lists, are revealed to be at the origin of an important hazard for aquatic receiving ecosystems, and thus should be so the object of reinforced controls and could be proposed to be added to these lists.

433 Figure 7: Example of hazard quotients (HQ) calculated with dissolved and total concentrations of pollutants in stormwater samples from Zgheib et al.⁴

434

How to cite this paper: Gosset, A., Durrieu, C., Orias, F., Bayard, R., Perrodin, Y. (2017) Identification and assessment of ecotoxicological hazards attributable to pollutants in urban wet weather discharges. Environmental Science: Processes & Impacts DOI: 10.1039/C7EM00159B

Conclusion and perspectives

The aim of our study was to develop and apply a methodology to determine and calculate hazard quotients associated with the different pollutants sought and found in UWWDs. A wide range of pollutants was sought in the latter, with highly variable ecotoxicity and concentrations. Although many families of pollutants (PC, PAH, VOC, ALK, M, P, O, PFAA and PHT) play a role in the UWWD hazards identified, we emphasize that they are comprised of a large number of pesticides and heavy metals (e.g. pyrethroids, copper and cadmium). The hazards of the pollutants identified in this study have different origins: (i) the presence of high concentrations of certain highly ecotoxic compounds (e.g. Pyrene in stormwater), (ii) the presence of high concentrations of slightly ecotoxic compounds (e.g. heavy metals in CSO), (iii) the presence of weak concentrations of certain highly ecotoxic compounds (e.g. pesticides like Bifenthrin and Permethrin in stormwater samples), (iv) the presence of intermediate concentrations of compounds with intermediate ecotoxicity (e.g. alkylphenols in UWWDs). We showed that the hazard linked to CSO was lower than that linked to stormwater from separate systems. However, these results should be considered with caution because only a few pharmaceutical compounds (just 28 compounds) were sought in CSO, compared to the huge range of drugs consumed by human beings. Moreover, no ecotoxicity data were available for 13 of the compounds (3 PC, 6 PAH, 2 ALK, 1 ORG, 1 PFAA) detected in UWWDs, and therefore no PNEC or HQ values could be calculated. Consequently, it will be necessary in further studies to obtain more ecotoxicity data on these compounds. Moreover, current hazard and risk assessment methods must be improved to take into account pollutant bioconcentration/accumulation, which can represent a serious risk for higher trophic levels.

To conclude, this work enabled us to identify the potential hazard of UWWDs and the pollutants involved. We now know which pollutants require an in-depth monitoring. We have also identified different points which require in-depth studies. In particular, it is now necessary to better assess the hazard of certain pollutants bound to the particulate phase. Lastly, this work should help managers to identify the largest and most hazardous pollution sources in order to treat them in priority.

Acknowledgements

This work was funded by the French Minister of Ecology and Sustainable Development, the French Minister of Higher Education and Research and the University of Lyon (Faculté des Sciences et Technologies). The authors thank Dr. Laurent Lassabatère for helpful discussions about the article. They also thank the anonymous referees for their constructive reviews.

References

- 1 M. Kafi, J. Gasperi, R. Moilleron, M. C. Gromaire and G. Chebbo, *Water Research*, 2008, **42**, 539–549.
- 2 H.-M. Hwang and G. D. Foster, *Environmental Pollution*, 2006, **140**, 416–426.
- 3 S. Terzakis, M. S. Fountoulakis, I. Georgaki, D. Albantakis, I. Sabathianakis, A. D. Karathanasis, N. Kalogerakis and T. Manios, *Chemosphere*, 2008, **72**, 141–149.
- 4 S. Zgheib, R. Moilleron, M. Saad and G. Chebbo, *Water Research*, 2011, **45**, 913–925.
- 5 Y. Kalmykova, K. Björklund, A.-M. Strömvall and L. Blom, *Water Research*, 2013, **47**, 1317–1328.
- 6 K. Björklund, A. P. Cousins, A.-M. Strömvall and P.-A. Malmqvist, *Science of The Total Environment*, 2009, **407**, 4665–4672.
- 7 M. Cladière, J. Gasperi, C. Lorgeoux, C. Bonhomme, V. Rocher and B. Tassin, *Environ Sci Pollut Res*, 2013, **20**, 2973–2983.
- 8 J. Gasperi, S. Garnaud, V. Rocher and R. Moilleron, *Science of The Total Environment*, 2008, **407**, 263–272.
- 9 E. Gooré Bi, F. Monette, J. Gasperi and Y. Perrodin, *Environmental Science and Pollution Research*, 2015, **22**, 4460–4474.
- 10 C. Becouze-Lareure, A. Dembélé, M. Coquery, C. Cren-Olivé, B. Barillon and J.-L. Bertrand-Krajewski, *Urban Water Journal*, 2016, **13**, 600–617.
- 11 S. Deffontis, A. Breton, C. Vialle, M. Montréjaud-Vignoles, C. Vignoles and C. Sablayrolles, *Science of The Total Environment*, 2013, **452**, 394–403.
- 12 J. Ryu, J. Oh, S. A. Snyder and Y. Yoon, *Environ Monit Assess*, 2014, **186**, 3239–3251.
- 13 X. Peng, Y. Yu, C. Tang, J. Tan, Q. Huang and Z. Wang, *Science of The Total Environment*, 2008, **397**, 158–166.
- 14 J.-Y. Pailler, A. Krein, L. Pfister, L. Hoffmann and C. Guignard, *Science of The Total*

- 492 *Environment*, 2009, **407**, 4736–4743.
- 493 15 R. Angerville, Y. Perrodin, C. Bazin and E. Emmanuel, *International Journal of*
- 494 *Environmental Research and Public Health*, 2013, **10**, 2670–2687.
- 495 16 L. Wu, Y. Jiang, L. Zhang, L. Chen and H. Zhang, *Environmental Science and Pollution*
- 496 *Research*, 2014, **21**, 2663–2676.
- 497 17 R.-H. Chen, F.-P. Li, H.-P. Zhang, Y. Jiang, L.-C. Mao, L.-L. Wu and L. Chen, *Science of*
- 498 *The Total Environment*, 2016, **553**, 519–523.
- 499 18 M. Lafont, L. Grapentine, Q. Rochfort, J. Marsalek, G. Tixier and P. Breil, *Water Science*
- 500 *and Technology*, 2007, **56**, 13–20.
- 501 19 C. Parent-Raoult and J.-C. Boisson, *Revue des sciences de l'eau / Journal of Water*
- 502 *Science*, 2007, **20**, 229–239.
- 503 20 K. A. Johnson, A. D. Steinman, W. D. Keiper and C. R. Ruetz, *Journal of the North*
- 504 *American Benthological Society*, 2011, **30**, 710–727.
- 505 21 C. M. Kinsella and T. P. Crowe, *Marine Environmental Research*, 2015, **105**, 20–29.
- 506 22 Y. Ferro, M. Perullini, M. Jobbagy, S. Bilmes and C. Durrieu, *Sensors*, 2012, **12**, 16879–
- 507 16891.
- 508 23 A. Gosset, Y. Ferro and C. Durrieu, *Water Research*, 2016, **89**, 330–354.
- 509 24 A. Wigh, A. Devaux, V. Brosselin, A. Gonzalez-Ospina, B. Domenjoud, S. Aït-Aïssa, N.
- 510 Creusot, A. Gosset, C. Bazin and S. Bony, *Environmental Science and Pollution Research*,
- 511 2016, **23**, 3008–3017.
- 512 25 F. Orias and Y. Perrodin, *Chemosphere*, 2014, **115**, 31–39.
- 513 26 M. Gros, M. Petrović, A. Ginebreda and D. Barceló, *Environment International*, 2010, **36**,
- 514 15–26.
- 515 27 A. Tsaboula, E.-N. Papadakis, Z. Vryzas, A. Kotopoulou, K. Kintzikoglou and E.
- 516 Papadopoulou-Mourkidou, *Environment International*, 2016, **91**, 78–93.
- 517 28 W. Li, Y. Shi, L. Gao, J. Liu and Y. Cai, *Chemosphere*, 2012, **89**, 1307–1315.
- 518 29 E. Eriksson, A. Baun, P. S. Mikkelsen and A. Ledin, *Water Science and Technology*, 2005,
- 519 **51**, 47–55.
- 520 30 A. Baun, E. Eriksson, A. Ledin and P. S. Mikkelsen, *Science of The Total Environment*,
- 521 2006, **370**, 29–38.
- 522 31 E. Eriksson, A. Baun, L. Scholes, A. Ledin, S. Ahlman, M. Revitt, C. Noutsopoulos and P.
- 523 S. Mikkelsen, *Science of The Total Environment*, 2007, **383**, 41–51.
- 524 32 E. Eriksson, A. Baun, P. S. Mikkelsen and A. Ledin, *Desalination*, 2007, **215**, 187–197.

How to cite this paper: Gosset, A., Durrieu, C., Orias, F., Bayard, R., Perrodin, Y. (2017) Identification and assessment of ecotoxicological hazards attributable to pollutants in urban wet weather discharges. *Environmental Science: Processes & Impacts* DOI: 10.1039/C7EM00159B

- 33 AGRITOX database on plant protection substances. <http://www.agritox.anses.fr/> (December 2016)
- 34 INERIS database, <http://www.ineris.fr/rapports-d%C3%A9tude/toxicologie-et-environnement/fiches-de-donn%C3%A9es-toxicologiques-et-environnementales>, (December 2016)
- 35 EPA ECOTOX database, <http://www.epa.gov/ecotox/>, (December 2016)
- 36 F. Orias and Y. Perrodin, *Science of The Total Environment*, 2013, **454–455**, 250–276.
- 37 TGD, Technical Guidance Document in Support of Commission Directive 93/67/EEC on Risk Assessment for New Notified Substances and Commission Regulation (EC), 2003.
- 38 H. Sanderson, D. J. Johnson, C. J. Wilson, R. A. Brain and K. R. Solomon, *Toxicology Letters*, 2003, **144**, 383–395.
- 39 R. J. Kieber, J. D. Willey and S. D. Zvalaren, *Environ. Sci. Technol.*, 2002, **36**, 5321–5327.
- 40 D. P. Weston, R. W. Holmes and M. J. Lydy, *Environmental Pollution*, 2009, **157**, 287–294.
- 41 C. S. Eckley and B. Branfireun, *Science of The Total Environment*, 2008, **403**, 164–177.
- 42 S.-K. Kim and K. Kannan, *Environ. Sci. Technol.*, 2007, **41**, 8328–8334.
- 43 Y. Zushi and S. Masunaga, *Environmental Toxicology and Chemistry*, 2009, **28**, 691–700.
- 44 V. T. Nguyen, M. Reinhard and G. Y.-H. Karina, *Chemosphere*, 2011, **82**, 1277–1285.
- 45 F. Xiao, M. F. Simcik and J. S. Gulliver, *Water Research*, 2012, **46**, 6601–6608.
- 46 M. Murakami, H. Shinohara and H. Takada, *Chemosphere*, 2009, **74**, 487–493.
- 47 R. Pitt, R. Field, M. Lalor, M. Brown, *Water Environment Research*, **67**, 260–275.
- 48 B. Ngabe, T.F. Bidleman and G.I. Scott, *Science of The Total Environment*, 2000, **255**, 1–9.
- 49 L. Herngren, A. Goonetilleke, G. A. Ayoko and M. M. M. Mostert, *Environmental Pollution*, 2010, **158**, 2848–2856.
- 50 M. N. Chong, J. Sidhu, R. Aryal, J. Tang, W. Gernjak, B. Escher and S. Toze, *Environmental Monitoring and Assessment*, 2013, **185**, 6645–6652.
- 51 Q. Zhang, X. Wang, P. Hou, W. Wan, Y. Ren, Z. Ouyang and L. Yang, *Environmental Monitoring and Assessment*, 2013, **185**, 9763–9775.
- 52 M. Huber, A. Welker and B. Helmreich, *Science of The Total Environment*, 2016, **541**, 895–919.
- 53 I. Hanke, I. Wittmer, S. Bischofberger, C. Stamm and H. Singer, *Chemosphere*, 2010, **81**, 422–429.
- 54 G. Imfeld, M. Lefrancq, E. Maillard and S. Payraudeau, *Chemosphere*, 2013, **90**, 1333–1339.
- 55 M. Kayhanian, C. Suverkropp, A. Ruby and K. Tsay, *Journal of Environmental Management*, 2007, **85**, 279–295.
- 56 E. Pratt, K. Irvine, S. Marshall, B. G. Loganathan, S. Kumar and H. C. Sikka, *Journal of Water Management Modeling*, 1995.

How to cite this paper: Gosset, A., Durrieu, C., Orias, F., Bayard, R., Perrodin, Y. (2017) Identification and assessment of ecotoxicological hazards attributable to pollutants in urban wet weather discharges. *Environmental Science: Processes & Impacts* DOI: 10.1039/C7EM00159B

- 559 57 J. Gasperi, S. Zgheib, M. Cladière, V. Rocher, R. Moilleron and G. Chebbo, *Water Research*,
560 2012, **46**, 6693–6703.
- 561 58 J. Gasperi, M. C. Gromaire, M. Kafi, R. Moilleron and G. Chebbo, *Water Research*, 2010,
562 **44**, 5875–5886.
- 563 59 R. Mulliss, D. M. Revitt and R. B. E. Shutes, *Water Science and Technology*, 1997, **36**, 195–
564 199.
- 565 60 M. Karvelas, A. Katsoyiannis and C. Samara, *Chemosphere*, 2003, **53**, 1201–1210.
- 566 61 N. Bolong, A. F. Ismail, M. R. Salim and T. Matsuura, *Desalination*, 2009, **239**, 229–246.
- 567 62 T. Deblonde, C. Cossu-Leguille and P. Hartemann, *International Journal of Hygiene and*
568 *Environmental Health*, 2011, **214**, 442–448.63 Luo et al., 2014
- 569 64 M. A. Launay, U. Dittmer and H. Steinmetz, *Water Research*, 2016, **104**, 82–92.
- 570 65 Stachel, J.-U. Holthuis, W. Schulz, W. Seitz, W. H. Weber, K.-T. Tegge and I. Dobner, in
571 *Xenobiotics in the Urban Water Cycle*, eds. D. Fatta-Kassinos, K. Bester and K.
572 Kümmerer, Springer Netherlands, 2010, pp. 445–461.
- 573 66 Zgheib, R. Moilleron and G. Chebbo, *Water Research*, 2012, **46**, 6683–6692.
- 574 67 M. P. Ensminger, R. Budd, K. C. Kelley and K. S. Goh, *Environ Monit Assess*, 2013, **185**,
575 3697–3710.
- 576 68 Y. Tao, Z. Yuan, H. Xiaona and M. Wei, *Ecotoxicology and Environmental Safety*, 2012,
577 **81**, 55–64.
- 578 69 S. Ruangsomboon and L. Wongrat, *Aquatic Toxicology*, 2006, **78**, 15–20.
- 579 70 T. Katagi, in *Review of Environmental Contamination and Toxicology Volume 204*, ed. D.
580 M. Whitacre, Springer New York, New York, NY, 2010, vol. 204, pp. 1–132.
- 581 71 Coogan et al., 2007
- 582 72 F. Orias, L. Simon, G. Mialdea, A. Clair, V. Brosselin and Y. Perrodin, *Ecotoxicology and*
583 *Environmental Safety*, 2015, **120**, 457–462.
- 584 73 F. Orias, L. Simon and Y. Perrodin, *Chemosphere*, 2015, **122**, 251–256.
- 585 74 A. Zenker, M. R. Cicero, F. Prestinaci, P. Bottoni and M. Carere, *Journal of*
586 *Environmental Management*, 2014, **133**, 378–387.
- 587 75 D. Lachassagne, M. Casellas, A. Gonzalez-Ospina and C. Dagot, *TSM*, 2016, 31–43.
- 588 76 J. Passerat, N. K. Ouattara, J.-M. Mouchel, V. Rocher and P. Servais, *Water Research*,
589 2011, **45**, 893–903.
- 590
- 591

Table S1: Extrapolation Factor assessment criteria derivate from Orias and Perrodin³⁶ and TDG³⁷.

Available data	EF (Extrapolation Factor)
At least one short-term L(E)C50 from each of three trophic levels of the base-set (e.g. algae, daphnids and fishes)	1000
One long term chronic data (e.g. invertebrate or fish)	100
Two long term chronic data from species representing two trophic levels (e.g.algae and/or daphnids and/or fishes)	50
Two long term chronic data from species representing three trophic levels (e.g. algae, daphnids and fishes)	10
Statistics method (e.g. species sensitivity distribution (SSD)), field data or model ecosystem	5-1 (Reviewed case by case)

615

616

Pollutant	Identified as hazardous		Identification as a priority substance	
	In stormwater	In combined sewer overflows	In European Water Framework Directive (WFD, 2000/60/EC) pollutant priority list	In Environmental Protection Agency pollutant priority list
4-(para)-nonylphenol (ALK)	Yes	No	Yes	No
4-tert-pentylphenol (ALK)	Yes	No	Yes	No
4-tert-butylphenol (ALK)	Yes	No	Yes	No
Nonylphenol (ALK)	No	Yes	Yes	No
Para-tert-octylphenol (ALK)	Yes	No	Yes	No
Cadmium (M)	Yes	Yes	Yes	Yes
Chromium III (M)	Yes	No	No	Yes
Copper (M)	Yes	Yes	No	Yes
Lead (M)	Yes	Yes	Yes	Yes
Nickel (M)	Yes	No	Yes	Yes
Zinc (M)	Yes	Yes	No	Yes
Bisphenol A (O)	Yes	No	No	No
Aminotriazole (P)	Yes	Yes	No	No
Bifenthrin (P)	Yes	No	No	No
Cyfluthrin (P)	Yes	No	No	No
Cypermethrin (P)	Yes	No	Yes	No
Deltamethrin (P)	Yes	No	No	No
Desethylatrazine (P)	Yes	Yes	No	No
Diuron (P)	Yes	Yes	Yes	No
Glyphosate (P)	Yes	No	No	No
Isoproturon (P)	Yes	No	Yes	No
Lambda-cyhalothrin (P)	Yes	No	No	No
N,N-diethyl-3-méthylbenzamide (P)	No	Yes	No	No
Permethrin (P)	Yes	No	No	No
Triclocarban (P)	No	Yes	No	No
Anthracene (PAH)	Yes	No	Yes	Yes
Benzo[a]anthracene (PAH)	Yes	No	Yes	Yes
Benzo[k]fluoranthene (PAH)	Yes	No	Yes	Yes
Benzo[g,h,i]perylene (PAH)	Yes	No	Yes	Yes
Benzo[a]pyrene (PAH)	Yes	No	Yes	Yes
Fluoranthene (PAH)	Yes	Yes	Yes	Yes
Fluorene (PAH)	Yes	No	Yes	Yes
Naphthalene (PAH)	Yes	No	Yes	Yes
Pyrene (PAH)	Yes	No	Yes	Yes
Ibuprofen (PC)	No	Yes	No	No
Triclosan (PC)	No	Yes	No	No
Trimethoprim (PC)	No	Yes	No	No
Perfluorooctanesulfonic acid (PFAA)	Yes	No	Yes	No
Perfluorononanoic acid (PFAA)	Yes	No	Yes	No
Benzyl butyl phthalate (PHT)	Yes	No	No	Yes
Diethylhexyl phthalate (PHT)	Yes	No	Yes	Yes
Diisononyl phthalate (PHT)	Yes	No	No	No
1,4-Dichlorobenzene (VOC)	Yes	No	No	Yes

How to cite this paper: Gosset, A., Durrieu, C., Orias, F., Bayard, R., Perrodin, Y. (2017) Identification and assessment of ecotoxicological hazards attributable to pollutants in urban wet weather discharges. Environmental Science: Processes & Impacts DOI: 10.1039/C7EM00159B

Pentachlorophenol (VOC)	Yes	No	Yes	Yes
-------------------------	-----	----	-----	-----

617 Table S2: Comparison of hazardous compounds list identified in UWWDs in this study with the
618 classification of these compounds as priority pollutants or not by European Water Framework
619 Directive and In Environmental Protection Agency.