Modifications of the soluble proteome of a mediterranean strain of the invasive neurotoxic dinoflagellate *Alexandrium catenella* under metal stress conditions

Natacha Jean, Estelle Dumont, Faouzi Herzi, Thierry Balliau, Mohamed Laabir, Estelle Masseret, Stéphane Mounier

► To cite this version:

Natacha Jean, Estelle Dumont, Faouzi Herzi, Thierry Balliau, Mohamed Laabir, et al.. Modifications of the soluble proteome of a mediterranean strain of the invasive neurotoxic dinoflagellate *Alexandrium catenella* under metal stress conditions. *Aquatic Toxicology*, 2017, 188, pp.80-91. 10.1016/j.aquatox.2017.04.011 . hal-01569507v2

HAL Id: hal-01569507

<https://hal.science/hal-01569507v2>

Submitted on 23 Sep 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Modifications of the soluble proteome of a mediterranean strain of the invasive neurotoxic dinoflagellate *Alexandrium catenella* under metallic stress conditions

Natacha JEAN^{1 (*)}, Estelle DUMONT¹, Faouzi HERZI¹, Thierry BALLIAU², Mohamed LAABIR³, Estelle MASSERET³, Stéphane MOUNIER¹

¹ Université de Toulon, PROTEE, EA 3819, 83957 La Garde, France

² Plate-forme d'Analyse Protéomique Paris Sud-Ouest (PAPPSO), INRA, Unité Mixte de Recherche en Génétique Végétale (UMR 320), Ferme du Moulon, 91190 Gif-sur-Yvette, France

³ MARBEC UMR 9190 IRD-Ifremer-CNRS-Université de Montpellier, Place Eugène Bataillon, Case 093, 34095 Montpellier Cedex 5, France

E-mail addresses: jean@univ-tln.fr (N. JEAN); estelle.dumont1@gmail.com (E. DUMONT); herzi_faouzi@yahoo.fr (F. HERZI); balliau@moulon.inra.fr (T. BALLIAU); Mohamed.Laabir@umontpellier.fr (M. LAABIR); estelle.masseret@umontpellier.fr (E. MASSERET); mounier@univ-tln.fr (S. MOUNIER)

* Corresponding author. Tel.: +33 4 94 14 25 29; E-mail address: jean@univ-tln.fr (N. JEAN)

Abstract

The soluble proteome of the mediterranean strain ACT03 of the invasive neurotoxic dinoflagellate *Alexandrium catenella* **exposed to** lead or zinc at 6, 12 or 18 M (total concentrations), or grown **under** control conditions, has been characterized by two-dimensional gel electrophoresis (2-DE). **Concentrations of zinc increasingly reduced** ($P < 0.05$) the total number of protein spots constituting the soluble proteome (-41%, -52% and -60%, at 6, 12 or 18 M, respectively) of *A. catenella* ACT03. **Besides, most of the proteins** constituting the soluble proteome **were down-regulated in response to lead or zinc stresses**. **These proteins** were involved mainly in photosynthesis (20-37% for lead; 36-50% for zinc) (ribulose-1,5-bisphosphate carboxylase/oxygenase: RUBISCO; ferredoxin-NADP⁺ reductase: FNR; peridinin-chlorophyll *a*-protein: PCP) and in the oxidative stress response (29-34% for lead; 17-36% for zinc) (superoxide dismutase: SOD; proteasome α/β subunits). **These** harmful effects could be partly compensated by the up-regulation of specific proteins **as** ATP-synthase β subunit (+16.3 fold **after exposure to** lead at 12 M). **Indeed**, increase in abundance of ATP-synthase could enrich the ATP pool and provide more energy available for the cells to survive under **metallic** stress conditions, **and** make more efficient the ATP-synthase transport of metal cations out **of** the cells. Finally, this study **shows** that **exposure to lead or zinc have** a harmful **effect** on the soluble proteome of *A. catenella* ACT03, **but** also suggests **a proteomic adaptative response to metallic stresses**, which could contribute to sustain this dinoflagellate development in some trace metal-contaminated ecosystems.

Keywords: *Alexandrium catenella*; harmful algal bloom; proteomics; stress proteins; trace metals; two-dimensional electrophoresis

64 Abbreviations

1	65	BiP	Binding immunoglobulin protein
2	66	CALR	Calreticulin
3	67	CaM	Calmodulin
4	68	CHAPS	3-[3-(cholamidopropyl)-dimethylammonio]1-propanesulfonate
5	69	2-DE	Two-dimensional gel electrophoresis
6	70	DTT	Dithiothreitol
7	71	DHAP	Dihydroxyacetone phosphate
8	72	FNR	Ferredoxin-NADP ⁺ reductase
9	73	G3P	Glyceraldehyde 3-phosphate
10	74	HAB	Harmful algal bloom
11	75	HSP	Heat shock protein
12	76	IEF	Isoelectric focusing
13	77	IPG	Immobilized pH gradient
14	78	LC-MS/MS	Liquid chromatography coupled to tandem mass spectrometry
15	79	MS	Mass spectrometry
16	80	MS/MS	Tandem mass spectrometry
17	81	MW	Molecular weight
18	82	NCBI	National centre for biotechnology information
19	83	NADPH	Nicotinamide adenine dinucleotide phosphate
20	84	PCP	Peridinin-chlorophyll <i>a</i> -protein
21	85	pI	Isoelectric point
22	86	PSP	Paralytic shellfish poisoning
23	87	RBP	Rubisco binding protein
24	88	ROS	Reactive oxygen species
25	89	RPI	Ribose-5-phosphate isomerase
26	90	R5P	Ribose-5-phosphate
27	91	RUBISCO	Ribulose-1,5-bisphosphate carboxylase/oxygenase
28	92	RuBP	Ribulose-1,5-bisphosphate
29	93	Ru5P	Ribose-5-phosphate isomerase
30	94	SDS	Sodium dodecylsulfate
31	95	SDS-PAGE	Sodium dodecylsulfate-polyacrylamide gel electrophoresis
32	96	SOD	Superoxide dismutase

1. Introduction

Harmful Algal Blooms (HABs) are proliferations of noxious phytoplankton including toxic species, able to (i) threaten human health after the consumption of seafood contaminated by toxins (ii) affect negatively many components of the marine life (shellfish, fishes, seabirds and mammals) (iii) impact socioeconomic activity due to the closure of aquaculture areas during HAB events (Anderson et al., 2012).

Since the last decades, frequency and distribution of HABs have increased in coastal mediterranean ecosystems subject to growing anthropisation (Ferrante et al., 2013). ***Alexandrium* genus, belonging the dinoflagellates**, is involved in severe, diverse and widely distributed HABs (Anderson et al., 2012). Indeed, the toxic *Alexandrium* species (*A. catenella*, *A. fundyense*, *A. minutum*, *A. tamarense*, *A. taylori*) can produce saxitoxins **and congeners, inducing** the Paralytic Shellfish Poisoning (PSP) syndrome.

Occurrences of the *Alexandrium* genus have been reported in several mediterranean areas receiving trace metal contaminations, as Catalan and Balearic coasts (Bravo et al., 2006; Penna et al., 2005), Tyrrhenian coast (Italy), Thau lagoon (France) (Laabir et al., 2013; Péna and Picot, 1991), **Bizerte lagoon (Tunisia) (Fertouna et al., 2015)** and Toulon Bay (western Mediterranean, France) (Abadie, personal communication; Jean et al., 2005, 2006). This latter is a severely trace metal contaminated ecosystem, both in the water column and the sediments (Jean et al., 2012; Tessier et al., 2011). Thus, occurrence of *Alexandrium* in such **area**, suggests that it could be potentially tolerant to trace metal contaminations, **although** this has not been demonstrated so far. Consequently, studies should be carried out to characterize the potential tolerance of *Alexandrium* to metallic stresses, which could contribute to explain its development in metal polluted anthropized environments.

In response to changes in the environment, an organism can modify its proteome by increasing or decreasing the expression of some proteins, making proteomics a dynamic field of investigation. These modifications, activating or slowing down some metabolic pathways, allow the organism to adapt to environmental changes. Therefore, modifications of the proteome are at the basis of the adaptive organism response to environmental stresses, as suggested in studies reporting higher protein expression (for actin and calreticulin) in zooplankton communities sampled in the most metal polluted stations of Toulon Bay (Jean et al., 2012).

Many studies have characterized the proteomic responses of aquatic organisms exposed to metal stresses, whether these are marine pluricellulars, as *Chlamys farreri*

(scallop) (Gao et al., 2007), *Eriocheir sinensis* (crab) (Silvestre et al., 2006), *Montastraea franksi* (coral) (Venn et al., 2009), *Mytilus edulis* (mussel) (Sanders and Martin, 1991), *Perna veridis* (bivalve) (Leung et al., 2011), *Saccostrea glomerata* and *Crassostrea gigas* (oyster) (Choi et al., 2008; Thompson et al., 2011, 2012), *Ectocarpus siliculosus*, *Fucus serratus*, *Lessonia nigrescens*, *Sargassum fusiforme* and *Scytosiphon gracilis* (brown algae) (Contreras et al., 2010; Ireland et al., 2004, **Lovazzano et al., 2013; Ritter et al., 2010; Zou et al., 2015**), or **aquatic unicellulars, as** *Anabaena* sp. (Pandey et al., 2012), *Klebsiella pneumoniae*, *Pseudomonas aeruginosa* and *Pseudomonas fluorescens* (Daware et al., 2012; Kiliç et al., 2010; Poirier et al., 2008) or *Tetrahymena pyriformis* (Zhang et al., 2012). **However, only one study has been devoted to the characterization of the proteomic response of a dinoflagellate, *Karenia brevis*, notably submitted to lead contamination, and in which some proteins (heat shock 60 kDa protein and superoxide dismutases) have been induced by the metal stress (Miller-Morey and Van Dolah, 2004).**

Under an ecotoxicoproteomic approach, the purpose of the present study is to characterize the proteomic response to metallic stresses, of the invasive neurotoxic dinoflagellate *Alexandrium catenella*. **A monoclonal mediterranean strain of *A. catenella* exposed to metallic contaminations, either by lead (toxic for the cells whatever its concentration), or by zinc (first required by the cells as a trace element, then harmful at higher concentrations), has been used. Then, the soluble proteomes of this strain, grown respectively under control and metallic stress conditions, have been characterized by two-dimensional gel electrophoresis (2-DE), and compared. This comparison** has generated a list of proteins of interest whose expression **was** significantly modified in response to the metallic stresses. The proteins of interest can be considered as potential biomarkers of **metallic** stresses in the *A. catenella* cells. Besides, they provide informations about the **metabolic pathways which were modified in *A. catenella* under metallic stress conditions**. Some of these proteins could be candidates **as potential** detoxification and protection processes in *A. catenella*, thus contributing to its adaptive response and tolerance capacity towards metallic stresses. Consequently, the overall results obtained could **allow** to understand the biochemical mechanisms **making** *A. catenella* **able** to develop in anthropized coastal marine ecosystems contaminated by trace metals.

2. Materials and Methods

2.1. Cultures of *Alexandrium catenella* ACT03

A. catenella strain **ACT03** was obtained after isolation of a single vegetative cell from seawater sample **collected** during a toxic bloom event in October 2003, in Thau lagoon (Laabir et al., 2013). It was maintained **in** enriched f/2 culture medium (Guillard and Ryther, 1962), at 20°C in sterile 250 mL flasks (75 cm² Greiner, Dominique Dutscher SAS), with a 12h: 12h photoperiod and under a light intensity of 135 mol photons m⁻² s⁻¹ (Herzi et al., 2013, 2014). Natural seawater filtered through a GF/F glass fibre filter (Ø = 47 mm, Whatman) and then sterilized **was** used as a basis for the culture medium.

To study **modifications of the soluble proteome** of *A. catenella* ACT03 under **metallic** stress, the culture medium was **supplemented with** sterile stock solutions of trace metals prepared with ZnSO₄, 7H₂O, or with Pb(CH₃COO)₂, 3H₂O salts dissolved **in ultrapure water**. For each **metal**, **three total concentrations** (6, 12 and 18 M) were **tested**. **A metal-free culture medium was considered as the control**. Based on the MINEQL-program and on the known composition of the f/2 culture medium, the corresponding free metal concentrations (Zn²⁺ or Pb²⁺) bioavailable for toxicity towards cells, were calculated (Table I) (Herzi et al., 2014).

Table I. Total (M²⁺_T) and free (M²⁺_F) trace metal concentrations (expressed in M), used for lead and zinc contaminations of the *Alexandrium catenella* ACT03 cultures

[M ²⁺] _T (M)	[Pb ²⁺] _F (M)	[Zn ²⁺] _F (M)
6 × 10 ⁻⁶	1.1 × 10 ⁻⁸	2.8 × 10 ⁻⁷
12 × 10 ⁻⁶	2.6 × 10 ⁻⁸	6.2 × 10 ⁻⁷
18 × 10 ⁻⁶	4.4 × 10 ⁻⁸	1.0 × 10 ⁻⁶

2.2. Preparation of the protein extracts

Control and metal-contaminated cultures were **studied** at the end of their exponential phase-early stationary phase of growth. Cell density of these cultures was estimated by counting (in triplicates) all the cells in 50 L **sub-sample** of a 1 mL Lugol fixed **culture**, with an inverted microscope (magnification: 100 ×). Taking the obtained density into account, a volume of culture containing 8 × 10⁶ cells **was** used for proteomic **analysis**.

The sample was centrifuged for 15 min at 1 500 g, **at** 15°C. The obtained cell pellet was washed twice in natural seawater (prepared as described in 2.1.), for 10 min at 15 000 g, to be immediately **used**. As recommended in Wang et al. (2008), the washed pellet **was then** re-suspended in an extraction solution, containing 2 mL 40 mM Tris **at** pH 8.7, 2.4 L

benzonase nuclease (Sigma-Aldrich) and 10 L of protease inhibitor cocktail (Sigma-Aldrich). Then, protein extraction **was** achieved by sonication of the obtained suspension in an ice-water bath, using a microtip Vibra Cell 734 24 (Bioblock Scientific) during 3 min at 50 W and 25 kHz, with 10 s on/ 10 s off cycles. Then, the solution was centrifuged for 30 min at 15 000 g, to obtain a supernatant which **contains** the soluble protein extract. This **fraction was further** concentrated by ultrafiltration at 1 500 g, **at** 15°C, through Vivaspın concentrator tubes with molecular weight cutoff membranes of 15 kDa (15R Hydrosart 10 kDa, Thermo Fisher Scientific), until reaching a volume of 150 L. To obtain the protein extract ready **to be used** for proteomics, the concentrate had to be finally mixed with 400 L of a sample solution consisting of urea (7 M), thiourea (2 M), CHAPS (1% w/v), Triton X-100 (3% v/v), DTT (1% w/v), carrier ampholytes (0.2% v/v) and **bromophenol blue** (0.002% v/v).

2.3. Protein determination

Protein determination of the extracts was achieved according to Lowry et al. (1951), using the Reagent Compatible Detergent Compatible Protein Assay (RC DC Protein Assay, Bio-Rad), and bovine serum albumin (BSA) as standard.

2.4. Two-dimensional electrophoresis

A pre-prepared immobilized pH gradient (IPG) strip (17 cm length, linear gradient, pH 3-10, Bio-Rad) was used, on which the extract sample (350 L containing 200 g proteins) was transferred (Linares et al., 2016). The rehydration of the IPG strip in presence of the sample, and **then isoelectric focusing** (IEF), were performed in the horizontal electrophoresis system PROTEAN IEF Cell (Bio-Rad), at 20°C, in the following manner: 18 h at 50 V (active rehydration), 2 h at 100 V, 2 h at 250 V, 2 h at 500 V, 2 h at 1 000 V, 2 h at 4 000 V and 5 h at 10 000 V, so as to reach a total minimal value of 60 000 Vh for each loaded IPG strip. After IEF, the IPG strip was successively equilibrated for 10 min at room temperature in equilibration buffers 1 (6 M urea, 2% w/v SDS, 0.375 M Tris pH 8.8, 20% v/v glycerol, 2% w/v DTT) and 2 (6 M urea, 2% w/v SDS, 0.375 M Tris pH 8.8, 20% v/v glycerol, 2.5% w/v iodoacetamide).

Sodium dodecylsulfate polyacrylamide gel electrophoresis (SDS-PAGE) was performed according to Laemmli (1970). The equilibrated IPG strip was placed in dyed (BPB) melted agarose, across a handmade polyacrylamide gel (18 cm × 18 cm × 1 mm; stacking gel: 5%, resolving gel: 11%). A volume of 20 L of molecular weight marker solution (10-250 kDa range, Precision Plus Protein Standards Dual Color, Bio-Rad) was

loaded at the left top of the gel. The gels were run at 4°C in a Protean II XL (Bio-Rad), with a constant current of 20 mA per gel for 1 h, and then, with a constant current of 30 mA per gel until the dye reached the bottom of the gel. After SDS-PAGE, gels were washed three times for 5 min in ultrapure water, then stained with the Imperial Protein Stain (Thermo Fisher Scientific) under orbital shaking for 1 h 30 min, to be destained in ultrapure water until visualization of separated protein spots. Each gel presented (Figures 1 & 2) is representative of three different biological gel replicates. The protein spots on the gels were analysed using the PD Quest 2-D Analysis Software 8.0.1 version (Bio-Rad). Abundance of a given protein spot has been obtained after normalization, as the *ratio* (in %) of its individual abundance on the abundance of all the marker bands.

2.5. Tracking of protein of interest in the soluble proteome of *A. catenella* ACT03

Among the proteins making the soluble proteome of *A. catenella* ACT03, the proteins of interest are those whose expression was significantly affected by the metallic stresses, through up- or down-regulation, or through appearance or disappearance. The up- or down-regulated proteins were respectively tracked thanks to the *ratio* of the abundance of each protein spot on the 2D-gel after metallic stress, on the abundance on the 2D-gel under control conditions (for up-regulation), and the *ratio* of the abundance of each protein spot on the 2D-gel under control conditions, on the abundance on the 2D-gel after metallic stress (for down-regulation). **First**, to be considered as a protein of interest, a protein had a *ratio* greater than or equal to 2 (Bae et al., 2003). Then, a statistical Student test was applied between the abundances of this protein on the 2D-gel replicates obtained under metallic stress conditions, and the abundances of this same protein on the 2D-gel replicates under control condition. This statistical test allowed to determine if the fold difference in up-regulation or down-regulation was significant (very significant: $P < 0.01$; significant: $0.01 < P < 0.05$; rather significant: $0.05 < P < 0.10$), in which case, the concerned protein was significantly up-regulated or down-regulated, and considered as a protein of interest. The other proteins of interest were those appearing or disappearing on at least two 2D-gels among the three obtained replicates.

2.6. Protein identification by liquid chromatography tandem mass spectrometry

Proteins of interest were picked on gel in order to be identified by mass spectrometry. In-gel digestion of the picked 2-D spots was performed with the Progest system (Genomic Solution) according to a standard trypsin protocol (Page et al., 2010). HPLC was performed on a NanoLC-Ultra system (Eksigent). A 4 L sample of the peptide solution was loaded at 7.5 L min⁻¹ on a precolumn cartridge (stationary phase: C18 Biosphere, 5 m; column: 100 m inner diameter, 2 cm; Nanoseparations) and desalted with 0.1% HCOOH. After 3 min, the precolumn cartridge was connected to the separating PepMap C18 column (stationary phase: C18 Biosphere, 3 m; column: 75 m inner diameter, 150 mm; Nanoseparations). Buffers A and B respectively were prepared with 0.1% HCOOH in water, and with 0.1% HCOOH in acetonitrile. The peptide separation was achieved with a linear gradient from 5 to 30% B for 28 min at 0.3 L min⁻¹. Including the regeneration step at 95% B and the equilibration step at 95% A, one run took 45 min.

Eluted peptides were on-line analyzed with an LTQ XL ion trap (Thermo Electron) using a nanoelectrospray interface. Ionization (1.5 kV ionization potential) was performed with liquid junction and a non-coated capillary probe (10 m inner diameter; New Objective). Peptide ions were analysed using Xcalibur 2.07 with the following data-dependent acquisition steps: (1) full MS scan (mass to-charge ratio (m/z) 300-1 400, centroid mode) and (2) MS/MS ($q_z = 0.25$, activation time = 30 ms, and collision energy = 35%; centroid mode). Step 2 was repeated for the three major ions detected in step 1. Dynamic exclusion was set to 30 s.

A database search was performed with XTandem (version 2010.12.01.1) (<http://www.thegpm.org/TANDEM/>). Enzymatic cleavage was declared as a trypsin digestion with one possible miscleavage. Cys carboxyamidomethylation and Met oxidation were set to static and possible modifications, respectively. Precursor mass and fragment mass tolerance were 2.0 and 0.8, respectively. A refinement search was added with similar parameters except that semi-trypsinic peptide, and possible N-ter proteins acetylation were searched. The UniprotKB database (<http://www.uniprot.org/>) restricted to Metazoa excepted Mammalia (1755330 entries, version 201201), and a contaminant database (trypsin, keratins.) were used. Only peptides with an E value smaller than 0.1 were reported.

Identified proteins were filtered and grouped using XTandem Pipeline (<http://pappso.inra.fr/bioinfo/xtandempipeline/>) according to: (1) A minimum of two different peptides required with an E value smaller than 0.05, (2) a protein E value (calculated as the product of unique peptide E values) smaller than 10^{-4} . In the case of identification with only two or three MS/MS spectra, similarity between the experimental and the theoretical MS/MS

spectra was visually checked. Identification by database searching was completed by a *de novo* approach. Peptides sequences were determined by automatic *de novo* interpretation from MS/MS spectra, using PepNovo software (version 2010225). This analysis was performed on MS/MS spectra with a quality score smaller than 0.1. Trypsin digestion, Cys carboxyamidomethylation and Met oxidation were set to enzymatic cleavage, static and possible modifications, respectively. Only sequences with a score greater than 70 were selected.

Homology searches were performed by Fasts software (version 36.06) using the MD20-MS matrix. Sequences corresponding to keratins or trypsin were firstly removed by interrogating a homemade contaminant database. Secondly, the search computing process was carried out on the same database. Homologies with a minimum of two independent peptides and an *E* value smaller than 0.001, were only selected. In **all** cases, the automatic *de novo* interpretation of MS/MS spectra was visually confirmed. Since the genome of *A. catenella* has not been fully **sequenced**, protein identification has been performed by homology with the known peptidic sequences from the National Centre for Biotechnology Information (NCBI) protein database (<http://www.ncbi.nlm.nih.gov/>), preferentially coming from, in this order: (i) other *Alexandrium* species (*A. fundyense*, *A. minutum*, *A. ostenfeldii*) (ii) other dinoflagellates (*Amphidinium carterae*, *Gonyaulax polyedra*, *Heterocapsa triquetra*, *Karlodinium micrum*, *Oxyrrhis marina*) (iii) other microalgae (*Cryptothecodinium cohnii*, *Symbiodinium* sp.) (iv) Protozoans (*Neospora caninum*, *Paramecium tetraurelia*, *Plasmodinium berghei*, *Tetrahymena thermophila*, *Toxoplasma gondii*).

3. Results

The results of mass spectrometry protein identification are shown in Table II (see also SuppData 1).

Table II. LC-MS/MS identification of the proteins up- or down-regulated in the soluble proteome of *Alexandrium catenella* ACT03 under metallic stresses (lead or zinc at 6, 12 or 18 M)

Spot	Peptide number	MW (kDa) / pI		Protein name	Species Accession number	Putative function
		Exp	Theo			
28	13	19.8/5.6	33.2/5.9	Ribose-5-phosphate isomerase	<i>Heterocapsa triquetra</i> Q5ENN9	Pentose phosphate pathway (02.07)
9	25	38.8/5.9	59.5/5.4	Ribulose-1,5-bisphosphate carboxylase	<i>Gonyaulax polyedra</i> Q42813	Photosynthesis (02.30)
29	16	21.6/5.7	79.1/5.8	Ribulose-1,5-bisphosphate carboxylase	<i>Heterocapsa triquetra</i> Q5ENN5	Photosynthesis (02.30)
4	15	25.0/5.9	20.6/8.8	Peridinin-chlorophyll <i>a</i> -binding protein	<i>Heterocapsa pygmaea</i> Q9FEY3	Light absorption (02.45.03)
34	17	25.7/6.9	37.8/6.6	Peridinin-chlorophyll <i>a</i> -binding protein	<i>Symbiodinium</i> sp. P51874	Light absorption (02.45.03)
36	3	24.8/6.4	38.0/9.1	Peridinin-chlorophyll <i>a</i> -binding protein	<i>Symbiodinium</i> sp. AFH88375	Light absorption (02.45.03)
41	18	24.6/6.7	38.9/9.0	Peridinin-chlorophyll <i>a</i> -binding protein	<i>Gonyaulax polyedra</i> O00941	Light absorption (02.45.03)
43	17	25.7/6.7	38.2/8.7	Peridinin-chlorophyll <i>a</i> -binding protein	<i>Amphidinium carterae</i> P80484	Light absorption (02.45.03)
44	20	25.8/6.5	37.8/6.6	Peridinin-chlorophyll <i>a</i> -binding protein	<i>Symbiodinium</i> sp. P51874	Light absorption (02.45.03)
46	7	25.8/6.2	38.0/9.1	Peridinin-chlorophyll <i>a</i> -binding protein	<i>Symbiodinium</i> sp. AFH88375	Light absorption (02.45.03)
16	3	49.4/5.1	60.0/6.4	ATP beta synthase subunit	<i>Neospora caninum</i> F0VGD5	Energy generation (02.45.15)
6	11	24.3/5.3	74.3/5.1	BiP	<i>Cryptocodinium cohnii</i> Q8S4R0	Chaperone (14.01)
47	5	25.0/6.2	74.3/5.1	BiP	<i>Cryptocodinium cohnii</i> Q8S4R0	Chaperone (14.01)
10	2	18.9/5.4	70.6/5.1	HSP70	<i>Cryptocodinium cohnii</i> Q8S4Q8	Chaperone (14.01)
40	7	22.9/5.3	70.6/5.1	HSP70	<i>Cryptocodinium cohnii</i> Q8S4Q8	Chaperone (14.01)
5	2	24.5/5.8	29.0/5.4	Proteasome alpha subunit	<i>Perkinsus marinus</i> XP_002775560	Proteasomal degradation (14.13.01.01)
30	4	20.4/6.0	22.6/6.1	Proteasome beta subunit	<i>Daphnia magna</i> JAN48661	Proteasomal degradation (14.13.01.01)
33	3	15.6/5.8	27.9/5.8	Proteasome alpha subunit	<i>Perkinsus marinus</i> XP_002764909	Proteasomal degradation (14.13.01.01)
35	4	24.8/6.5	27.9/5.8	Proteasome alpha subunit	<i>Perkinsus marinus</i> XP_002764909	Proteasomal degradation (14.13.01.01)
37	7	23.8/6.6	27.9/5.8	Proteasome alpha subunit	<i>Perkinsus marinus</i> XP_002764909	Proteasomal degradation (14.13.01.01)
45	8	25.8/6.4	27.9/5.8	Proteasome alpha subunit	<i>Alexandrium fundyense</i> A4UHA5	Proteasomal degradation (14.13.01.01)
19	10	18.1/6.2	27.2/5.0	Calreticulin	<i>Heterocapsa triquetra</i> Q5ENL5	Calcium binding (16.17.01)
26	3	14.1/4.5	16.8/4.1	Calmodulin	<i>Prorocentrum minimum</i> AAL61535	Calcium binding (16.17.01)
7	17	31.7/6.4	45.0/7.0	Ferredoxin-NADP ⁺ - reductase	<i>Cryptocodinium cohnii</i> Q5ENS9	Electron transport (20.01.15)
8	19	31.7/6.3	45.0/7.0	Ferredoxin-NADP ⁺ - reductase	<i>Heterocapsa triquetra</i> Q5ENS9	Electron transport (20.01.15)
21	23	32.8/5.9	45.0/7.0	Ferredoxin-NADP ⁺ - reductase	<i>Heterocapsa triquetra</i> Q5ENS9	Electron transport (20.01.15)
39	2	34.1/5.7	45.0/7.0	Ferredoxin-NADP ⁺ - reductase	<i>Heterocapsa triquetra</i> Q5ENS9	Electron transport (20.01.15)
25	11	18.2/5.3	17.9/5.9	Superoxide dismutase [Cu-Zn]	<i>Tetrahymena thermophila</i> Q22H00	Detoxification (32.07)
1	22	29.2/4.8		Chromosome undetermined scaffold_2	<i>Paramecium tetraurelia</i> AOCJD5	Unclassified protein (99)
12	16	31.7/4.3	50.3/5.1	Hypothetical protein	<i>Plasmodium berghei</i> Q4YAV6	Unclassified protein (99)
31	3	19.4/4.8		Unknown	<i>Alexandrium minutum</i> GW796469	Unclassified protein (99)
32	2	14.1/5.4		Unknown	<i>Oxyrrhis marina</i> EG736358	Unclassified protein (99)

3.1. Modifications of the soluble proteome under lead exposure

Whatever its concentration, lead did not induce a significant ($P > 0.05$) decrease in the total number of protein spots constituting the soluble proteome of *A. catenella* ACT03 (6 M: 240 ± 74 spots; 12 M: 202 ± 98 spots; 18 M: 239 ± 92 spots), compared to the control (339 ± 89 spots) (Figures 1A & 1B).

Spot n°	Protein name	6 M	12 M	18 M
1	Unknown protein	+2.0**	+8.2***	
2	Unidentified protein	-4.1**		
4	Peridinin -chlorophyll <i>a</i> -protein	4.0***		-3.0**
5	Proteasome alpha subunit	D	D	D
6	Binding immunoglobulin protein		D	
7	Ferredoxin -NADP ⁺ reductase	-7.3*		
9	RUBISCO	D	D	D
10	HSP 70	-2.9*	D	
11	Unidentified protein			-3.9*
12	Unknown protein		+18.5***	
13	Unidentified protein		+13.5**	
14	Unidentified protein		+17.7***	+2.6*
15	Unidentified protein		+16.4**	+1.7*
16	ATP synthase beta subunit		+16.3***	
17	Unidentified protein		+19.2***	
18	Unidentified protein		+16.7***	
19	Calreticulin		D	D
21	Ferredoxin -NADP ⁺ reductase	-7.3***	D	
25	Superoxide dismutase	-6.5*	D	D
28	Ribose -5-phosphate isomerase		D	
29	RUBISCO	-2.8**	D	D
30	Proteasome beta subunit		D	
31	Unknown protein		D	D
32	Unknown protein		D	
33	Proteasome beta subunit	-2.3***	D	
34	Peridinin -chlorophyll <i>a</i> -protein	-2.2*	-2.3*	
36	Peridinin -chlorophyll <i>a</i> -protein	-3.2**		-1.5*
37	Proteasome alpha subunit		D	
39	Ferredoxin-NADP ⁺ reductase	-6.0**	D	
40	HSP 70	-2.4**	D	

Figure 1. Representative 2-DE maps comparing the soluble proteomes of *Alexandrium catenella* ACT03 grown under **exposure** to lead 6, 12 or 18 M ($[Pb^{2+}]_F = 11, 26$ and 44 nM respectively) and **control** conditions. (A): Protein profiles of *Alexandrium catenella* ACT03 grown under **exposure** to lead 6, 12 or 18 M (B): Protein profile of *Alexandrium catenella* ACT03 grown under **control** condition. Proteins of interest are identified by their spot numbers (C): Names of the proteins of interest are shown with the fold differences in their expression, according to the level of **lead exposure**. In black, up-regulated proteins; in grey, down-regulated proteins; D, disappeared proteins. *, ** and ***: proteins whose expression was different under lead **exposure**, with $0.05 < P < 0.10$; $0.01 < P < 0.05$ and $P < 0.01$, respectively

3.1.a. Exposure to lead 6 M

Exposure to lead 6 M led to 46 proteins of interest (*i.e.* whose expression was significantly modified in comparison with **control**: $P < 0.10$) (**Figures 1B & 1C**). Among these differentially expressed proteins, 12 proteins could be identified, which were significantly **down-regulated**, whereas only one protein, not recognized in the protein database, was significantly up-regulated. **The down-regulated proteins** are: (i) involved in photosynthesis, **for** RUBISCO (spots 9 and 29), PCP (spots 4, 34 and 36) and FNR (spots 7, 21 and 39) (ii) involved in the response to oxidative stress, as proteasome α subunit (spots 5 and 33) and SOD (spot 25) (iii) the chaperone HSP 70 (spots 10 and 40). The fold differences in expression of these **proteins ranged** from -7.3 to +2.0 fold, with the highest harmful **effect** on FNR (-7.3 fold) (spots 7 and 21) (**Figure 1C**).

3.1.b. Exposure to lead 12 M

Exposure to lead 12 M led to 82 proteins of interest ($P < 0.10$). Eight proteins were significantly up-regulated and 17 proteins were significantly **down-regulated** (**Figures 1B & 1C**). Only one protein **up-regulated could be** identified: it was the ATP-synthase β subunit (spot 16), involved in energy metabolism. Fold differences in expression of the up-regulated proteins reached the highest values observed in this study, *i.e.* exceeding +16.0 fold (for spots 12 and 15-18) (**Figure 1C**). These results showed the **strong effect** of lead 12 M on this protein up-regulation. This **effect** was opposite regarding the **down-regulated proteins**: (i) PCP (spot 34), RUBISCO (spots 9 and 29) and FNR (spots 21 and 39) involved in photosynthesis (ii) proteasome α/β subunits (spots 5, 30, 33 and 37) and SOD (spot 25) implied in the response to oxidative stress (iii) CALR (spot 19) chaperone **also** involved in Ca^{2+} binding (iv) RPI (spot 28) taking part in carbohydrate metabolism (v) HSP 70 (spots 10 and 40) and BiP (spot 6), two chaperones.

3.1.c. *Exposure to lead 18 M*

A number of 59 proteins of interest ($P < 0.10$) have been detected in response to exposure to lead 18 M (Figures 1B & 1C). These proteins of interest were mainly down-regulated. Among them, we have identified: (i) RUBISCO (spots 9 and 29), enzyme at the root of photosynthesis (ii) proteasome α subunit (spot 5) and SOD (spot 25), involved in the response to oxidative stress (iii) CALR (spot 19), chaperone also involved in Ca^{2+} binding (iv) PCP (spots 4 and 36), implied in photosynthetic activity. Finally, two proteins have been significantly up-regulated (spots 14 and 15), with a moderate fold difference around +2.0 fold, but their identification has not been successful (Figure 1C).

3.1.d. *Intercomparison of the proteomic modifications under lead stresses*

No significant ($P > 0.05$) difference was observed between the total numbers of proteins constituting the soluble proteomes of *A. catenella* respectively exposed to lead 6, 12 or 18 M (Figure 1A). However, in comparison with the other concentrations, lead at 12 M induced the greatest number of modifications in protein expression (82 proteins of interest), with the highest numbers of up-regulated and down-regulated proteins (Figures 1B & 1C). Besides, lead at 12 M induced the maximal fold differences in up-regulation, reaching between +13.5 (spot 13) and +19.2 (spot 17) fold (Figures 1B & 1C).

Whatever the lead concentration, the proteins RUBISCO (spot 9) and proteasome α subunit (spot 5) disappeared (Figures 1B & 1C, Figure 1S in SuppData 2). On the other hand, SOD (spot 25) ended up disappearing after exposure to lead at 12 M and 18 M. Such a harmful effect was also observed for HSP 70 (spots 10 and 40), FNR (spots 21 and 39), RUBISCO (spot 29) and proteasome α subunit (spot 33). In contrast, only ATP-synthase β subunit (spot 16) was up-regulated after exposure to lead at 12 M.

3.2. Modifications of the soluble proteome under zinc exposure

In comparison with control (339 ± 89 spots), **systematical significant** ($P < 0.05$) decreases in the total numbers of protein spots **were observed in the soluble proteomes** of *A. catenella* ACT03 exposed to: zinc 6 M: 200 ± 11 spots (-41%); zinc 12 M: 162 ± 90 spots (-52%); zinc 18 M: 136 ± 37 spots (-60%) (Figures 2A & 2B).

Spot n°	Protein name	6 M	12 M	18 M
1	Unknown protein		+2.0**	+6.0***
4	Peridinin-chlorophyll <i>a</i> -protein	-3.2**	-3.4**	D
5	Proteasome alpha subunit	-3.9***	D	D
8	Ferredoxin -NADP ⁺ reductase	D	D	
9	RUBISCO	D	D	D
10	HSP 70		D	D
12	Unknown protein		D	
13	Unidentified protein			+6.8***
14	Unidentified protein	+2.1*		+2.3*
16	ATP synthase beta subunit			D
17	Unidentified protein		D	D
18	Unidentified protein	-2.7*	D	D
19	Calreticulin		D	
21	Ferredoxin -NADP ⁺ reductase	D	-4.4**	D
22	Unidentified protein		D	
24	Unidentified protein			D
25	Superoxide dismutase	-28.0**	D	D
26	Calmodulin		-21.8*	D
28	Ribose -5-phosphate isomerase			D
29	RUBISCO		-13.2**	D
30	Proteasome beta subunit	D		D
31	Unknown protein		-3.1**	D
34	Peridinin-chlorophyll <i>a</i> -protein	-2.7*		D
35	Proteasome alpha subunit	-6.5**	-2.7*	D
36	Peridinin-chlorophyll <i>a</i> -protein	-3.3**	-2.1*	-2.0*
37	Proteasome alpha subunit			D
39	Ferredoxin -NADP ⁺ reductase		-8.0**	D
40	Heat shock protein 70			D
41	Peridinin-chlorophyll <i>a</i> -protein	-2.4**	-2.2*	-4.4**
43	Peridinin-chlorophyll <i>a</i> -protein		-2.0*	-2.0*
44	Proteasome alpha subunit			D
45	Proteasome alpha subunit			D
46	Peridinin-chlorophyll <i>a</i> -protein			D
47	Binding immunoglobulin protein			D

Figure 2. Representative 2-DE maps comparing the soluble proteomes of *Alexandrium catenella* ACT03 grown under **exposure to** zinc 6, 12 **or** 18 M ($[Zn^{2+}]_F = 280, 620$ and $1\,000$ nM respectively) **and control** conditions. (A): Protein profiles of *Alexandrium catenella* ACT03 grown under **exposure to** zinc 6, 12 **or** 18 M (B): Protein profile of *Alexandrium catenella* ACT03 grown **in** control conditions. Proteins of interest are identified by their spot numbers (C): Names of the proteins of interest are shown with the fold differences in their expression, according **to** the level of **zinc exposure**. In black, up-regulated proteins; in grey, down-regulated proteins; D, disappeared proteins. *, ** and ***: proteins whose expression was different under zinc **exposure**, with $0.05 < P < 0.10$; $0.01 < P < 0.05$ and $P < 0.01$, respectively

3.2.a. *Exposure to zinc 6 M*

With 6 M, 54 proteins of interest **were detected** ($P < 0.10$). Among these differentially expressed proteins, only one protein (spot 14) was significantly up-regulated (its identification failed), whereas 12 proteins were significantly **down-regulated** (Figures 2B & 2C). The proteins **down-regulated** were **involved**: (i) in photosynthesis for RUBISCO (spot 9), PCP (spots 4, 34, 36 and 41) and FNR (spot 8) (ii) **in the** response to oxidative stress for proteasome α/β subunits (spots 5, 30 and 35) and SOD (spot 25). Regarding the fold differences in expression of these proteins in comparison with control, the main result was the very strong **down-regulation** of SOD (spot 25) (-28.0 fold) (Figure 2C).

3.2.b. *Exposure to zinc 12 M*

With 12 M, 68 proteins of interest **were detected** ($P < 0.10$), among which 20 proteins were significantly down-regulated, whereas one protein was significantly up-regulated (not recognized in the protein database) (Figures 2B & 2C). The proteins down-regulated were **involved**: (i) in photosynthesis for RUBISCO (spots 9 and 29), PCP (spots 4, 36, 41 and 43) and FNR (spots 8, 21 and 39) (ii) **in the** response to oxidative stress for proteasome α subunit (spots 5 and 35) and SOD (spot 25) (iii) in cell signaling, activation of protein kinases and Ca^{2+} binding for CaM (spot 26) (iv) **with** the chaperones HSP 70 (spot 10), and CALR (spot 19) **which is** also involved in Ca^{2+} binding. **Effect** of zinc 12 M was particularly harmful on the protein CaM, as indicated by the maximal fold difference in its **down-regulation** (-21.8 fold) with regards to control (Figure 2C).

3.2.c. *Exposure to zinc (18 M)*

With 18 M, 111 proteins of interest were detected ($P < 0.10$), among which 25 proteins were identified (Figures 2B & 2C). As observed for lead at the same concentration, zinc induced the disappearance of many proteins. These proteins are involved (i) in photosynthesis for RUBISCO (spots 9 and 29), FNR (spots 21 and 39) and PCP (spots 4, 34, 36, 41, 43 and 46) (ii) in the response to oxidative stress for proteasome α/β subunits (spots 5, 30, 35, 37, 44 and 45, 46) and SOD (spot 25) (iii) in energy metabolism for ATP-synthase β subunit (spot 16) (iv) take part in cell signaling, activation of protein kinases and Ca^{2+} binding for CaM (spot 26) (v) is the chaperone HSP 70 (spots 10 and 40) (vi) in carbohydrate metabolism for RPI (spot 28). Three proteins were significantly up-regulated (spots 1, 13 and 14), but their identification failed, or led to unknown peptide sequences. The fold differences in expression of the up- and down-regulated proteins in comparison with control, ranged between -4.4 (PCP: spot 41) and +6.8 (spot 13) fold (Figure 2C).

3.2.d. *Intercomparison of the proteomic modifications under zinc stresses*

The total number of proteins constituting the soluble proteome of *A. catenella* ACT03 exposed to 18 M of zinc was significantly lower (of 32%) than with 6 M ($P = 0.04$), confirming the harmful effect of zinc (Figure 2A). In comparison with the other concentrations, zinc 18 M induced the most drastic modifications in the protein expression, with the highest numbers of up- and down-regulated proteins (Figures 2B & 2C). Besides, zinc at 18 M led to the strongest up-regulation of some proteins, reaching +6.0 (spot 1: unknown protein) and +6.8 (spot 13: unidentified protein) fold (Figure 2C). Zinc at 12 M and 18 M resulted in the up-regulation of a same unknown protein (spot 1) (at 18 M: +6.0 fold; at 12 M: +2.0 fold), whereas an unidentified protein (spot 14) was up-regulated in common for zinc at 6 M and 18 M (+2.0 fold) (Figure 2C). Whatever the zinc concentration, RUBISCO (spot 9) disappeared and PCP (spots 36 and 41) was down-regulated (Figures 2B & 2C, Figure 2S in SuppData 2). Finally, zinc concentration had an increasing effect on the down-regulation of proteasome α subunit (spots 5 and 35), FNR (spots 21 and 39), SOD (spot 25), CaM (spot 26) and PCP (spots 4, 34, 41 and 43).

3.3. Comparison of the proteomic modifications under lead or zinc stresses

Exposure to lead or zinc led to the up-regulation of three proteins (spots 1, 13 and 14) (Figures 1C & 2C). However, this up-regulation, observed at a lower concentration for lead than for zinc, was stronger for lead (**higher values of the fold differences**). Only lead resulted in the up-regulation of five supplementary proteins (spots 12, 15, 16, 17 and 18), among which ATP-synthase β subunit has been identified. Whatever the trace **metal and its concentration**, the disappearance of RUBISCO (spot 9) **was observed**. Similarly, zinc or lead at 12 M and 18 M, induced disappearance of SOD (spot 25). **However**, higher concentrations **of zinc were** required to induce the disappearance of RUBISCO (spot 29). **Only lead** had a harmful effect on BiP (spot 6) and on proteasome β subunit (spot 33), **and** HSP 70 (spot 40) was **perturbed in expression** from 6 M for lead, against 18 M for zinc.

3.4. Biological functions assigned to the proteins of interest

Biological functions have been assigned to the identified proteins of interest, **in order** to characterize the metabolic pathways implied in the proteomic response of *A. catenella* ACT03 exposed to metallic stresses (Figure 3). These biological functions **are**: calcium binding (CALR, CaM), carbohydrate metabolism (RPI); cell signaling and activation of protein kinases (CaM), chaperone activity (BiP, CALR, HSP 70), energy metabolism (ATP-synthase β subunit; FNR), oxidative stress response (BiP, CALR, HSP 70, proteasome α/β subunits, SOD), photorespiration (RUBISCO) and photosynthesis (ATP-synthase β subunit, FNR, PCP, RPI, RUBISCO). **The most affected proteins were** involved in the photosynthesis (from 20 to 37% of the proteins of interest identified, for lead **at** 6, 12 or 18 M and between 36 and 50% of the proteins of interest identified for zinc **at** 6, 12 or 18 M) and in the oxidative stress response (from 29 to 34% for lead **at** 6, 12 or 18 M, and between 17 and 36% for zinc **at** 6, 12 or 18 M). **The** metabolic pathways potentially **concerned** by modifications in protein expression were (i) energy metabolism (8-9% for lead **at** 6 and 12 M 6-9% for zinc **at** 6 and 18 M) (ii) photorespiration (5-13% for lead **at** 6, 12 or 18 M ; 4-6% for zinc **at** 6, 12 or 18 M) (iii) chaperone activity (4-8% for lead **at** 6 and 12 M ; 6% for zinc at 12 and 18 M) (iv) calcium binding (2-4% for lead **at** 12 and 18 M ; 2-5% for zinc **at** 12 and 18 M) **and** (v) carbohydrate metabolism (3% for lead **at** 12 M and 2% for zinc 18 M), whereas cell signaling and activation of protein kinases (CaM) could be only **perturbed** in response to zinc stresses (2-3% for zinc **at** 12 and 18 M).

Figure 3. Biological functions assigned to the proteins of interest identified in the soluble proteome of *Alexandrium catenella* ACT03 exposed to lead or zinc at 6, 12 or 18 M

4. Discussion

4.1. Proteomic modifications and oxidative stress response

Through **down-regulation** of key proteins such as SOD and proteasome α/β subunits, the oxidative stress response of *A. catenella* ACT03 **could be a metabolic pathway particularly perturbed** in response to **the** metallic stresses.

The induction of SOD is known to minimize oxidative cell damages among organisms living in polluted environments (Okamoto and Colepicolo, 1998). Indeed, SOD participates to the first antioxidant response to the ROS generated by metal stress (Bareen et al., 2012). In the dinoflagellate *Karenia brevis*, a fast induction of [Mn] SOD has been observed in response to contamination by lead at 18 M (Miller-Morey and Van Dolah, 2004). Similarly, in the marine bivalve *Perna viridis*, a significant expression of [Cu-Zn] SOD has been recorded, following a contamination by cadmium (Leung et al., 2011). In the dinoflagellate *Gonyaulax polyedra* exposed to mercury, cadmium, lead and copper, activity of SOD increased during the first day of exposure to stress, with induction of [Fe] SOD and [Mn] SOD, which was not observed for [Cu-Zn] SOD (Okamoto and Colepicolo, 1998). In the **chlorophyte** *Chlamydomonas reinhardtii* exposed to cadmium, increase **in** [Mn] SOD **co-occurred with a** decrease **in** [Fe] SOD (Gillet et al., 2006). A strong stimulation in **the** expression of [Mn] SOD was recorded at the same time as a slight decrease in expression of [Cu-Zn] SOD, in response to cadmium stress in *Saccharomyces cerevisiae* (Vido et al., 2001). However, **lead or zinc at 12 and 18 M**, systematically **induced the disappearance of SOD from** the soluble proteome of *A. catenella* ACT03, **whereas** zinc **at 6 M led to a** fold difference in down-regulation **reaching** the very low value of -28.0 (Figures 1C & 2C). Trace metals are known to generate oxidative stress producing ROS **which** damages polypeptide sequences (Stohs and Bagchi, 1995). Then, **these** damaged proteins are targeted for the proteasomal degradation (Delattre et al., 2005), which could **explain** the decreased SOD abundances measured in the soluble proteome of *A. catenella* ACT03, **induced by** metallic stresses.

The proteasome contributes to cell proteolytic activity, and thus, plays a decisive role in the oxidative stress response, by eliminating oxidized proteins, which have been beforehand targeted by ubiquitination (Flick and Kaiser, 2012). Hence, some studies reported a significant increase in abundance of ubiquitinated proteins, in cells exposed to metal stress conditions (Marzano et al., 2012). Often, increases in proteasomal degradation observed under oxidative stress conditions, correspond to prior increases in abundance of the

proteasome subunits (Rodrigues et al., 2012). For instance, an over-expression of the proteasome α subunits, **with a difference of +2.1 fold**, was observed in the crab *Eriocheir sinensis* exposed to chronic stress by cadmium (Silvestre et al., 2006). Similarly, proteasome α subunits were over-expressed in the macroalgae *Scytosiphon gracilis* and in the fish *Oryzias melanostigma*, under copper and mercury contaminated conditions, respectively (Contreras et al., 2010; Wang et al., 2011). Nevertheless, other studies also mentioned that oxidative stress targets the proteasome itself, during reactions of glutathionylation or nitrosylation, which leads to a dysfunctioning of the proteasomal activity (Dasuri et al., 2013). In the present study, zinc or lead often induced **the decrease** in abundance of proteasome subunits (Figures 1C & 2C), which could indicate such a proteasomal degradation **in** oxidative stress conditions.

4.2. Proteomic modifications and photosynthesis

Important decreases in abundances of key **photosynthetic** proteins (RUBISCO, PCP and FNR) have been recorded, making photosynthesis **a metabolic pathway potentially and severely perturbed in response to the metallic stresses tested**. These results are in agreement with many studies reporting **harmful effects of trace metals on photosynthesis, as** chlorophyll degradation, destruction of chloroplast ultrastructure and down-regulation of many proteins involved in the Calvin cycle (Ahsan et al., 2009; Führes et al. 2008; Gillet et al., 2006; Kieffer et al., 2008).

Concerning RUBISCO, which catalyses carboxylation of the substrate ribulose-1,5-bisphosphate (RuBP) during the second step of photosynthesis (Reuman and Weber, 2006), trace metals generally decrease its abundance in the chloroplasts, as observed in the microalgae *Pseudokirchneriella subcapitata* exposed to chromium and palladium (Vannini et al., 2009), or in the **chlorophyte** *Chlamydomonas reinhardtii* stressed by cadmium (Gillet et al., 2006). This could result from high sensitivity of RUBISCO methionine amino acids to the metal-induced oxidative stress, which leads to methionine sulfoxide residues responsible for the targeting of proteins intended to the proteolytic degradation (Brot and Weissbach, 2000; Desimone et al., 1996; Ge et al., 2009; Stadtman et al., 2005). The oxidative degradation of the RUBISCO Binding Protein (RBP), which contributes to the RUBISCO assembly (Roy et al., 1988), as suggested in a study on *P. subcapitata* exposed to stress by chromium (Vannini et al., 2009), **could** also explain this result. Other harmful effects of metals on photosynthesis have been reported, as a significant decrease in RUBISCO activity (around 50%) in the aquatic plant *Salvinia natans* submitted to metal stresses by cobalt, zinc or copper (Dhir et al.,

2011). This could be due, either to the harmful substitution of cation Mg^{2+} (organizing centre of the active sites of RUBISCO) by the metallic contaminant cations (Van Assche and Clijsters, 1990), or to the irreversible bonds likely to form between metal contaminant cations and thiol groups in RUBISCO (Pankovic et al., 2000). The degradation, in metal stress conditions, of the RUBISCO activase involved in the RUBISCO activation (Salvucci et al., 1987), for example observed in *P. subcapitata* after a stress by chromium (Vannini et al., 2009), also could explain decrease in RUBISCO activity. Similarly, in the present study, RUBISCO in *A. catenella* ACT03 either disappeared whatever the metal tested and its concentration, or was less abundant in the soluble proteome as the metallic concentrations increased, with a stronger harmful effect of lead **compared to** zinc (Figures 1C & 2C). Among the diverse consequences of this decrease in RUBISCO, a lower CO_2 assimilation by the cells (Dhir et al., 2011) and a loss in efficiency of the RuBP recycling which is required by the Calvin cycle (Kieffer et al., 2008), could be **expected** in the cells of *A. catenella* ACT03.

Light energy input for photosynthesis in dinoflagellates is linked to the PCP complex (Hofmann et al., 1996). **This PCP** contains a carotenoid pigment, called peridinin, **which** is probably used as a quencher for the cells, deactivating by energy transfer, the excited state of a chemical species, and thus, acting as an antioxidant (Sigaud-Kutner et al., 2002). For instance, in the dinoflagellate *Lingulodinium polyedrum*, **production** of peridinin increased **up to** 60% in response to **an** oxidative stress (Pinto et al., 2000). However, opposite results have been obtained **with** the dinoflagellate *Gonyaulax polyedra* for which peridinin abundance decreased after cell **exposure** to mercury, cadmium or copper (Okamoto et al., 2001). Even though this **process** reflects decrease in photosynthetic activity, the reduced peridinin **concentration** could also contribute to a regulation of the electron flux between the photosystems I and II, avoiding O_2^- superoxide anion formation, and consequently reducing oxidative damages (Okamoto et al., 2001). Here, PCP abundances decreased in response to most of the metallic **levels** (Figures 1C & 2C), **suggesting** such a form of antioxidant response towards metal stress.

Similarly, the FNR enzyme, involved in the photophosphorylation step during photosynthesis, was down-regulated in the soluble proteome of *A. catenella* ACT03 exposed to zinc or lead. **Gillet et al. (2006)** have mentioned a toxic effect of cadmium **at** 150 μM on **the** expression of FNR by *C. reinhardtii* (fold difference: -1.7), **showing the** extreme FNR sensitivity to trace metals. **As shown in higher plants**, the FNR degradation by solubilisation can take place under oxidative conditions (Palatnik et al., 1997), which **could also** occur in

the present study. Globally, these results confirm the harmful **effects** of metallic stresses on **the** photosynthesis, suggesting *in fine* a potential loss of energy available for the *A. catenella* ACT03 cells. This could be compensated by the up-regulation of some proteins, as **the** ATP-synthase β subunit, as suggested below.

4.3. Proteomic modifications and energy metabolism

Exposure to lead at 12 M **induced the** up-regulation (+16.3 fold) of ATP-synthase β subunit (Figure 1C). Such a metal-induced up-regulation of ATP-synthase was also observed in the autotrophs: *Typha angustifolia* exposed to millimolar concentrations of chromium (+1.5 fold) (Bah et al., 2010), *Anabaena* sp. and *Pteris vittata* submitted to arsenic contaminations (Bona et al., 2010; Pandey et al., 2012), *C. reinhardtii* and lichens in response to mercury and cadmium stresses (Gillet et al., 2006; Nicolardi et al., 2012). Associated with the up-regulation of ATP-synthase, the potential increase in the ATP pool could contribute to an adaptive response of these organisms, by bringing more energy to survive under metal stress conditions (Kosova et al., 2011) and sustaining their perturbed photosynthetic activity (Bona et al., 2010). This energy could be used for detoxication processes, for example making more efficient the ATP-synthase active transport of contaminant metal cations out of the cells (Gillet et al., 2006). Moreover, ATP-synthase is known to assume peptidase function and chaperone activity, degrading abnormal and damaged proteins, or repairing them, so as to ensure the survival of the cells (Suzuki et al., 1997).

4.4. Proteomic modifications and photorespiration

Despite its essential contribution to photosynthesis, RUBISCO also catalyses oxygenation of the RuBP substrate during the **first** step of the photorespiratory pathway in autotrophs (Reumann and Weber, 2006). As a result, **down-regulation** of RUBISCO recorded **in the present study** could also suggest a weakening of photorespiration in the cells of *A. catenella* ACT03. Since photorespiration produces CO₂ and NH₃, and consumes O₂ and ATP (Zhang et al., 2012), it is *a priori* considered as an energy counter-productive pathway for autotrophs, **contrarily** to photosynthesis. However, some studies highlight that consumption of O₂ and ATP associated with photorespiration could contribute to the protection of photosynthetic organisms living in oxidative conditions (Wingler et al., 2000). As a result, in regards to photosynthesis, the photorespiratory pathway would be used by stressed autotrophs as wells of energy avoiding the hyper-reduction of the photosynthetic chain of electron transport (Wingler et al., 2000). In addition, the photorespiratory pathway

leads to metabolites as glycine amino acid which is used in the biosynthesis of glutathion involved in the cellular response to oxidative stress (Noctor et al., 1999).

4.5. Proteomic modifications and chaperone activity

In *A. catenella* ACT03, lead at 6 and 12 M, and zinc at 12 and 18 M, perturbed expression of diverse chaperones belonging to the HSPs 70 family, for example BiP, also called “heat shock 70 kDa protein 5 (HSPA5)” (Figures 1C & 2C). These results are in accordance with those reported for *Phanerochaete chrysosporium* exposed to cadmium or copper at 50 M, in which abundances in HSP 20 were reduced from 70 to 80% (Ozcan et al., 2007). Similarly, in the Sydney Rock oysters, HSP 70 was the most affected protein after contamination by cadmium (50 g L⁻¹), lead (5 g L⁻¹) or zinc (5 g L⁻¹) (Thompson et al., 2012), whereas in the soya beans, HSP 70 abundances were four times lower under cadmium stress (Hossain et al., 2012). Besides, it is known that regulation in HSP expression is metal and concentration depending, as observed for HSP 90 in human lung cells, which was down-regulated by cadmium, nickel and chromium, and up-regulated by arsenic (Andrew et al., 2003), and for HSP 22 in *Argopectens irradians* which decreased in response to stress by cadmium at 200 g L⁻¹, while it was over-expressed in this bivalve exposed to cadmium at 100 g L⁻¹ (Leung et al., 2011).

Heat shock proteins (HSPs) represent up to 2% of the cellular proteins (Gao et al., 2007). These chaperones are known to (i) help protein maturation and folding, optimizing their metabolic functionalities (Beere, 2005) (ii) avoid protein aggregation and ensuring protein membrane transport (Hossain et al., 2012). On the other hand, HSPs take part to the response to many environmental stresses, which explains increases in their abundances observed after exposure to extreme temperatures, UV, trace metals, and as a result, their use as general biomarkers under stress conditions (Timperio et al., 2008; Venn et al., 2009). During stresses, HSPs are required to (i) target the damaged polypeptides so as to allow their subsequent proteasomal degradation (ii) maintain the less damaged polypeptides in their folded conformation so as to avoid their proteasomal degradation, and then, to repair and reactivate them (Miller-Morey and Van Dolah, 2004). Consequently, recovery of the cellular metabolic processes after stress is generally associated with an increase in HSP abundances (Thompson et al., 2012). For example, studies have mentioned that exposure of the dinoflagellate *Karenia brevis* to lead at 18 M, induced the up-regulation of HSP 60 (Miller-Morey and Van Dolah, 2004), whereas HSP 70 was up-regulated in marine macroalgae and aquatic plants exposed to cadmium (Ireland et al., 2004).

In the present study, **the chaperone calreticulin was down-regulated** by the metallic stresses tested (Figures 1C & 2C). CALR is known as an endoplasmic reticulum (ER) molecular chaperone, and an ubiquitous multifunctional calcium-binding protein (Ryu et al., 2012). Calreticulin is needed in Ca^{2+} buffering and in quality control processes during protein synthesis and folding, in particular during refolding of misfolded proteins and ER stress responses (Leung et al., 2011). In addition to bind Ca^{2+} , CALR **would also** be able to bind Zn^{2+} (Michalak et al., 1992), and requires zinc for its interaction with the ERp57 thiol oxidoreductase (Leach et al., 2002). Studies showed evidence that cadmium stress induced a calreticulin up-regulation in hepatopancreas of *Perna viridis* (Leung et al., 2011), and in leafs of *Phytolacca americana* (Zhao et al., 2011). Thus, it has been suggested that elevated levels of CALR could serve to protect the cells against cadmium cytotoxicity (Zhao et al., 2011). The metal stress inducing ER stress, the resulting misfolded proteins would need the up-regulation of ER chaperone, like calreticulin, to be refolded (Leung et al., 2011).

4.6. Proteomic modifications and carbohydrate metabolism

Carbohydrate metabolism **could also** be **perturbed in *A. catenella* ACT03 exposed to metal stresses**, due to **down-regulation** of the ribose-5-phosphate isomerase (RPI), which catalyses the interconversion of Ru5P (or R5P) into R5P (or Ru5P) during (i) the Calvin cycle (photosynthesis), in the penultimate step leading to final regeneration of RuBP (ii) the pentose phosphate pathway, in its non oxidative second phase (Zhang et al., 2003). This modification in RPI expression in the soluble proteome of *A. catenella* ACT03 was observed in response to contaminations by lead at 12 M and by zinc at 18 M. This result agrees with the decrease in abundance also reported for RPI in: poplar leaves exposed at cadmium 20 M, *C. reinhardtii* stressed by cadmium at 150 M (fold difference: -1.9), tomato roots in response to contamination by aluminium at 20 M (Gillet et al., 2006; Kieffer et al., 2008; Zhou et al., 2009), confirming the **harmful** effects of trace metals.

4.7. Proteomic modifications and cellular signalling/activation of protein kinases

In the present study, calmodulin was **down-regulated after exposure to** zinc at 12 M (-21.8 fold), to finally disappear at zinc 18 M. Thanks to the formation of a complex with Ca^{2+} , CaM is activated to take part in cellular signaling and regulate activity of proteins called “CaM kinases”, with which it interacts (Alves et al., 2010; Stefan et al., 2008). Indeed, kinases are known to regulate activity of transcription factors, which trigger or repress the ADN transcription rate into proteins (Mittler, 2002). More generally, kinases stimulate

metabolic pathways *via* activation of many enzymes (ex: phosphorylase kinase involved in glycogenolysis) (Maksymiec et al., 2007). It is known that stress conditions induce a series of reactions increasing the Ca^{2+} levels, which induce CaM kinase activation. Yet, as observed here, a CaM decrease (-3.8) was observed in the liver of sea-bream exposed to chronic stress (Alves et al., 2010). Some studies have reported a possible competition between Ca^{2+} and metal cations on the CaM binding site, as observed during germination of radish plants contaminated by Cd^{2+} (Rivetta et al., 1997). The trace metal binding to CaM could lead to perturbations of the cellular signaling (Clemens, 2006), through inhibition, by the Ca^{2+} -calmodulin complex, of the CaM kinase phosphodiesterase. Taking these informations into account, the **down-regulated** CaM **could mean** that **the** synthesis of some proteins would be inhibited **after exposure to** zinc *via* the repression of transcription factors, which **could also** explain the decreases in abundance (**rather than down-regulation**) of many proteins of the soluble proteome of *A. catenella* ACT03 **exposed to metallic stress** conditions.

Conclusion

This study showed the harmful effects of lead or zinc on the soluble proteome of *A. catenella* ACT03, and thus potentially, on key metabolic pathways **in** this dinoflagellate, through down-regulation of many proteins to be considered as potential biomarkers of metal stress. However, an adaptive proteomic response could also take place in the cells, **at the same time**, thanks to the up-regulation of other proteins as ATP-synthase implied in production of energy, which could be used to sustain detoxification processes and cell survival in presence of the metals. This suggests that, despite the clear harmful **effect** of trace metals, differences in the expression of some proteins **could also** help to maintain the development of *A. catenella* ACT03 in metal-contaminated ecosystems.

Acknowledgments

This research **was supported** by the GdR « PHYCOTOX: Des micro-algues aux risques pour l'Homme et l'écosystème », the Conseil Général du Var (CG 83), Toulon Provence Méditerranée (TPM) and the research programme ARCUS CERES (PACA-MAE). **It also benefited from the financial help of LAGUNOTOX project funded by Fondation TOTAL.**

References

- Ahsan, N., Renaut, J., Komatsu, S., 2009. Recent developments in the application of proteomics to the analysis of plant responses to heavy metals. *Proteomics* 9, 2602–2621.
- Alves, R.N., Cordeiro, O., Silva, T.S., Richard, N., De Vareilles, M., Marino, G., Di Marco, P., Rodrigues, P.M., Conceicao, L.E.C., 2010. Metabolic molecular indicators of chronic stress in gilthead seabream (*Sparus aurata*) using comparative proteomics. *Aquaculture* 299, 57–66.
- Anderson, D.M., Alpermann, T.J., Cembella, A.D., Collos, Y., Masseret, E., Montresor, M., 2012. The globally distributed genus *Alexandrium*: Multifaceted roles in marine ecosystems and impacts on human health. *Harmful Algae* 14, 10–35.
- Andrew, A.S., Warren, A.J., Barchowsky, A., Temple, K.A., Klei, L., Soucy, N.V., O'Hara, K.A., Hamilton, J.W., 2003. Genomic and proteomic profiling of responses to toxic metals in human lung cells. *Environ. Health Perspect.* 111, 825–35.
- Bae, M.S., Cho, E.J., Choi, E.Y., Park, O.K., 2003. Analysis of the *Arabidopsis* nuclear proteome and its response to cold stress. *Plant J.* 36, 652–663.
- Bah, A.M., Sun, H., Chen, F., Zhou, J., Dai, H., Zhang, G., Wu, F., 2010. Comparative proteomic analysis of *Typha angustifolia* leaf under chromium, cadmium and lead stress. *J. Hazard Mat.* 184, 191–203.
- Bareen, F., Shafiq, M., Jamil, S., 2012. Role of plant growth regulators and a saprobic fungus in enhancement of metal phytoextraction potential and stress alleviation in pearl millet. *J. Hazard. Mat.* 237–238, 186–193.
- Beere, H.M., 2005. Death versus survival: functional interaction between the apoptotic and stress-inducible heat shock protein pathways. *J. Clin. Invest.* 115, 2633–2639.
- Bona, E., Cattaneo, C., Cesaro, P., Marsano, F., Lingua, G., Cavaletto, M., Berta, G., 2010. Proteomic analysis of *Pteris vittata* fronds: two arbuscular mycorrhizal fungi differentially modulate protein expression under arsenic contamination. *Proteomics* 10, 3811–3834.
- Bravo, I., Garcès, E., Diogène, J., Fraga, S., Sampedro, N., Figueroa, R.I., 2006. Resting cysts of the toxigenic dinoflagellate genus *Alexandrium* in the recent sediments from the Western Mediterranean coast, including the first description of cysts of *A. kutnerae* and *A. peruvianum*. *Eur. J. Phycol.* 41, 3, 293–302.
- Brot, N., Weissbach, H., 2000. Peptide methionine sulfoxide reductase: Biochemistry and physiological role. *Pept. Sci.* 55, 288–296.
- Choi, Y.K., Jo, P.G., Choi, C.Y., 2008. Cadmium affects the expression of heat shock protein 90 and metallothionein mRNA in the Pacific oyster, *Crassostrea gigas*. *Comp. Biochem. Physiol. C. Toxicol. Pharmacol.* 147, 286–292.
- Clemens, S., 2006. Toxic metal accumulation, responses to exposure and mechanisms of tolerance in plants. *Biochimie* 88, 1707–1719.
- Contreras, L., Moenne, A., Guillard, F., Potin, P., Correa, J.A., 2010. Proteomic analysis and identification of copper stress-regulated proteins in the marine alga *Scytosiphon gracilis* (Phaeophyceae). *Aquat. Toxicol.* 96, 85–89.
- Dasuri, K., Zhang, L., Keller, J.N., 2013. Oxidative stress, neurodegeneration and the balance of protein degradation and protein synthesis. *Free Rad. Bio. Med.* 62, 170–185.
- Daware, V., Kesavan, S., Patil, R., Natu, A., Kumar, A., Kulkarni, M., 2012. Effects of arsenite stress on growth and proteome of *Klebsiella pneumoniae*. *J. Biotechnol.* 158, 8–16.
- Delattre, J., Beaudeau, J.L., Bonnefont-Rousselot, D., 2005. Radicaux libres et stress oxydant : aspects biologiques et pathologiques. Editions Lavoisier 548 p.
- Desimone, M., Henke, A., Wagner, E., 1996. Oxidative stress induces partial degradation of the large subunit of ribulose 1, 5- biphosphate carboxylase/oxygenase in isolated chloroplasts of Barley. *Plant Physiol.* 111, 789–796.
- Dhir, B., Sharmila, P., Pardha Saradhi, P., Sharma, S., Kumar, R., Mehta, D., 2011. Heavy metals induced physiological alterations in *Salvinia natans*. *Ecotox. Environ. Saf.* 74, 1678–1684.
- Ferrante, M., Conti, G.O., Fiore, M., Rapisarda, V., Ledda, C., 2013. Harmful algal blooms in the mediterranean sea: effects on human health. *Euromed. Biomed. J.* 8, 25–34.
- Fertouna-Bellakhal, M., Dhib, A., Fathalli, A., Bellakhal, M., Chome, N., Masseret, E., Laabir, M., Turki, S., Aleya, L. 2015. *Alexandrium pacificum* Litaker sp. nov (Group IV): Resting cyst distribution and toxin profile of vegetative cells in Bizerte Lagoon (Tunisia, Southern Mediterranean Sea). *Harmful Algae* 48, 69–82.**
- Flick, K., Kaiser, P., 2012. Protein degradation and the stress response. *Sem. Cell. Dev. Biol.* 2, 515–522.
- Führs, H., Hartwig, M., Molina, L.E.B., Heintz, D., Dorsselaer, A.V., Braun, H.P., Horst, J.W., 2008. Early manganese-toxicity response in *Vigna unguiculata* L. – a proteomic and transcriptomic study. *Proteomics* 8, 149–159.

Gao, Q., Song, L., Ni, D., Wu, L., Zhang, H., Chang, Y., 2007. cDNA cloning and mRNA expression of heat shock protein 90 gene in the haemocytes of Zhikong scallop *Chlamys farreri*. Comp. Biochem. Physiol. B Biochem. Mol. Biol. 147, 704-715.

Ge, C., Ding, Y., Wang, Z., Wan, D., Wang, Y., Shang, Q., Luo, S., 2009. Responses of wheat seedlings to Cd, Hg and trichlorobenzene stresses. J. Environ. Sci. 2, 806-813.

Gillet, S., Decottignies, P., Chardonnet, S., Le Maréchal, P. 2006. Cadmium response and redoxin targets in *Chlamydomonas reinhardtii*: a proteomic approach. Photosynth. Res. 89, 201-211.

Guillard, R.R.L., Ryther, J.H. 1962. Studies of marine planktonic diatoms, I. *Cyclotella nana* Hustedt and *Detonula confervacea* (Cleve) Gran. Can. J. Microbiol. 8, 229-239.

Herzi, F., Jean, N., Zhao, H., Mounier, S., Mabrouk, H.H., Sakka Hlaili, A., 2013. Copper and cadmium effects on growth and extracellular exudation of the marine toxic dinoflagellate *Alexandrium catenella*: 3D-fluorescence spectroscopy approach. Chemosphere 93, 1230-1239.

Herzi, F., Jean, N., Sakka Hlaili, A., Mounier, S., 2014. Three-dimensional (3D) fluorescence spectroscopy analysis of the fluorescent dissolved organic matter released by the marine toxic dinoflagellate *Alexandrium catenella* exposed to metal stress by zinc or lead. J. Phycol. 50, 665-674.

Hofmann, E., Wrench, P.M., Sharples, F.P., Hiller, R.G., Welte, W., Diederichs, K., 1996. Structural basis of light harvesting by carotenoids peridinin-chlorophyll-protein from *Amphidinium carterae*. Science 272, 1788-1791.

Hossain, Z., Makino, T., Komatsu, S., 2012. Proteomic study of β -aminobutyric acid-mediated cadmium stress alleviation in soybean. J. Proteomics 75, 4151-4164.

Ireland, E., Harding, S.J., Bonwick, G.A., Jones, M., Smith, C.J., Williams, J.H., 2004. Evaluation of heat shock protein 70 as a biomarker of environmental stress in *Fucus serratus* and *Lemna minor*. Biomarkers 9, 139-155.

Jean, N., Bogé, G., Jamet, J.-L., Richard, S., Jamet, D., 2005. Annual contribution of different plankton size classes to particulate dimethylsulfoniopropionate in a marine perturbed ecosystem. J. Marine Syst. 53, 235-247.

Jean, N., Bogé, G., Jamet, J.-L., Jamet, D., 2006. Comparison of β -dimethylsulfoniopropionate (DMSP) levels in two Mediterranean ecosystems with different trophic levels. Mar. Chem. 101, 190-202.

Jean, N., Dumont, E., Durrieu, G., Balliau, T., Jamet, J.-L., Personnic, S., Garnier, C., 2012. Protein expression from zooplankton communities in a metal contaminated NW mediterranean coastal ecosystem. Mar. Environ. Res. 80, 12-26.

Kieffer P., Dommes J., Hoffmann L., Hausman J.F. et Renaut J. 2008. Quantitative changes in protein expression of cadmium – exposed poplar plants. Proteomics 8: 2514-2530.

Kiliç, N.K., Stensballe, A., Otzen, D.E., Dönmez, G., 2010. Proteomic changes in response to chromium VI toxicity in *Pseudomonas aeruginosa*. Biores. Technol. 101, 2134-2140.

Kosová, K., Vítámvás, P., Prášil, I.T., Renaut, J., 2011. Plant proteome changes under abiotic stress. Contribution of proteomics studies to understanding plant stress response. J. Proteomics 74, 1301-1322.

Laabir, M., Collos, Y., Masseret, E., Grzebyk, D., Abadie, E., Savart, V., Sibat, M., Amzil, Z., 2013. Influence of Environmental Factors on the Paralytic Shellfish Toxin Content and Profile of *Alexandrium catenella* (Dinophyceae) Isolated from the Mediterranean Sea. Mar. Drugs 11, 1583-1601.

Laemmli, K. ,1970. Cleavage of structural proteins during the assembly of the bacteriophage T4. Nature 227, 680-685.

Leach, M.R., Cohen-Doyle, M.F., Thomas, D.Y., Williams, D.B., 2002. Localization of the lectin, ERp57 binding, and polypeptide binding sites of calnexin and calreticulin. J. Biol. Chem. 277, 29686-29697.

Leung, P.T.Y., Wang, Y., Mak, S.S.T., Ng, W.C., Leung, K.M.Y., 2011. Differential proteomic responses in hepatopancreas and adductor muscles of the green-lipped mussel *Perna viridis* to stresses induced by cadmium and hydrogen peroxide. Aquat. Toxicol. 105, 49-61.

Linares, D., Jean, N., Van Overtvelt, P., Ouidir, T., Hardouin, J., Blache, Y., Molmeret, M., 2016. The marine bacteria *Shewanella frigidimarina* NCIMB400 upregulates the type VI secretion system during early biofilm formation. Environ. Microbiol. Rep., 8, 1, 110-121.

Lovazzano, C., Serrano, C., Correa, J.A., Contreras-Porcia, L., 2013. Comparative analysis of peroxiredoxin activation in the brown macroalgae *Scytosiphon gracilis* and *Lessonia nigrescens* (Phaeophyceae) under copper stress. Physiol. Plant. 149, 378-388.

Lowry, O.H., Rosenbraugh, N.J., Farr, A.L., Randall, R.J., 1951. Protein measurement with the Folin Phenol Reagent. J. Biol. Chem. 193, 265-275.

Maksymiec, W., Wójcik, M., Krupa, Z., 2007. Variation in oxidative stress and photochemical activity in *Arabidopsis thaliana* leaves subjected to cadmium and excess copper in the presence or absence of jasmonate and ascorbate. Chemosphere 66, 421-427.

Marzano, V., Santini, S., Rossi, C., Zuccherelli, M., D'alessandro, A., Marchetti, C., Mingardi, M., Stagni, V., Barila, D., Urbani, A., 2012. Proteomic profiling of ATM kinase proficient and deficient cell lines upon blockage of proteasome activity. J. Proteomics 75, 4632-4646.

Michalak, M., Milner, R.E., Burns, K., Opas, M., 1992. Calreticulin. Bioch. J. 285, 681-692.

Miller-Morey, J.S., Van Dolah, F.M., 2004. Differential responses of stress proteins, antioxidant enzymes, and photosynthetic efficiency to physiological stresses in the Florida red tide dinoflagellate, *Karenia brevis*. *Comp. Biochem. Physiol. C Toxicol. Pharmacol.* 138, 493-505.

Mittler, R., 2002. Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci.* 7, 405-410.

Morse, D., Fritz, L., Pappenheimer, A.M., Hastings, J.W., 1989. Properties and cellular localization of a luciferin binding protein in the bioluminescence reaction of *Gonyaulax polyedra*. *J. Biolumin. Chemilumin.* 3, 79-83.

Nicolardi, V., Cai, G., Parrotta, L., Puglia, M., Bianchi, L., Bini, L., Gaggi, C., 2012. The adaptive response of lichens to mercury exposure involves changes in the photosynthetic machinery. *Environ. Poll.* 160, 1-10.

Noctor, G., Arisi, A.C.M., Jouanin, L., Christine, H.F., 1999. Photorespiratory glycine enhances glutathione accumulation in both the chloroplastic and cytosolic compartments. *J. Exp. Bot.* 50, 1157-1167.

Okamoto, O.K., Colepiccolo, P., 1998. Response of superoxide dismutase to pollutant metal stress in the marine dinoflagellate *Gonyaulax polyedra*. *Comp. Biochem. Physiol. Part C: Pharmacol. Toxicol. Endocrinol.* 119, 67-73.

Okamoto, O.K., Pinto, E., LaTorre, L.R., Bechara, E.J.H., Colepiccolo, P., 2001. Antioxidant modulation in response to metal-induced oxidative stress in algal chloroplasts. *Arch. Environ. Contam. Toxicol.* 40, 18-24.

Ozcan, S., Yildirim, V., Kaya, L., Albrecht, D., Becher, D., Hecker, M., Ozcengiz, G., 2007. *Phanerochaete chrysosporium* soluble proteome as a prelude for the analysis of heavy metal stress response. *Proteomics* 7, 1249-1260.

Page, D., Gouble, B., Valot, B., Bouchet, J.P., Callot, C., Kretzschmar, A., Causse, M., Renard, C.M., Faurobert, M., 2010. Protective proteins are differentially expressed in tomato genotypes differing for their tolerance to low-temperature storage. *Planta* 232, 483-500.

Palatnik, J.F., Valle, E.M., Carrillo, N., 1997. Oxidative stress causes ferredoxin-NADP⁺ reductase solubilization from the thylakoid membranes in methyl viologen-treated plants. *Plant Physiol.* 113, 1721-1727.

Pandey, S., Rai, R., Rai, L.C., 2012. Proteomics combines morphological, physiological and biochemical attributes to unravel the survival strategy of *Anabaena* sp. PCC7120 under arsenic stress. *J. Proteomics* 75, 921-937.

Pankovic, D., Plesnicar, M., Arsenijenic-Maksimovic, I., Petrovic, N., Sakac, Z., Kastori, R., 2000. Effects of nitrogen nutrition on photosynthesis in Cd-treated sunflower plants. *Ann. Bot.* 86, 841-847.

Péna, G., Picot, B., 1991. Métaux traces dans les sédiments d'une lagune méditerranéenne : l'étang de Thau. *Oceanol. Acta*, 14, 459-472.

Penna, A., Garcès, E., Vila, M., Giacobbe, M.G., Fraga, S., Lugliè, A., Bravo, I., Bertozzini, E., Vernesi, C., 2005. *Alexandrium catenella* (Dinophyceae), a toxic ribotype expanding in the NW Mediterranean Sea. *Mar. Biol.* 148, 1, 13-23.

Pinto, E., Sigaud-Kutner, T.C.S., Leitao, M.A.S., Okamoto, O.K., Morse, D., Colepiccolo, P., 2000. Heavy metal-induced oxidative stress in algae. *J. Phycol.* 39, 1008-1018.

Poirier, I., Jean, N., Guary, J.-C., Bertrand, M., 2009. Robustness to cell disruption of heavy metal-resistant *Pseudomonas fluorescens* BA3SM1 isolates. *Environ. Eng. Sci.* 26, 1451-1457.

Reumann, S., Weber, A.P.M., 2006. Plant peroxisomes respire in the light: Some gaps of the photorespiratory C₂ cycle have become filled-Others remain. *Biochim. Biophys. Ac. - Mol. Cell Res.* 1763, 1496-1510.

Ritter, A., Ubertini, M., Romac, S., Gaillard, F., Delage, L., Mann, A., Cock, J.M., Tonon, T., Correa, J.A., Potin, P., 2010. Copper stress proteomics highlights local adaptation of two strains of the model brown alga *Ectocarpus siliculosus*. *Proteomics* 10, 2074-2088.

Rivetta, A., Negrini, N., Cocucci, M., 1997. Involvement of Ca²⁺ calmodulin in Cd²⁺ toxicity during the early phases of radish (*Raphanus sativus* L.) seed germination. *Plant Cell Environ.* 20, 600-608.

Rodrigues, P.M., Silva, T.S., Dias, J., Jessen, F., 2012. Proteomics in aquaculture: applications and trends. *J. Proteomics* 75, 4325-4345.

Roy, H., Cannon, S., Gilson, M., 1988. Assembly of Rubisco from native subunits. *Biochim. Biophys. Ac.* 957, 323-334.

Ryu, S.Y., Hong, G.U., Kim, D.Y., Ro, J.Y., 2012. Enolase 1 and calreticulin regulate the differentiation and function of mouse mast cells. *Cell Signal* 24, 60-70.

Salvucci, M.E., Werneke, J.M., Ogren, W.L., Portis, A.R., 1987. Purification and species distribution of rubisco activase. *Plant Physiol.* 84, 930-936.

Sanders, B.M., Martin, L.S., 1991. Relationships between accumulation of a 60 kDa stress protein and scope-for-growth in *Mytilus edulis* exposed to a range of copper concentrations. *Mar. Environ. Res.* 31, 81-97.

Sigaud-Kutner, T.C.S., Pinto, E., Okamoto, O.K., Latorre, L.R., Colepiccolo, P., 2002. Changes in superoxide dismutase activity and photosynthetic pigment content during growth of marine phytoplankters in batch – cultures. *Physiol. Plantarum* 114, 566-571.

Silvestre, F., Dierick, J.F., Dumont, V., Dieu, M., Raes, M., Devos, P., 2006. Differential protein expression profiles in anterior gills of *Eriocheir sinensis* during acclimatation to cadmium. *Aquat. Toxicol.* 76, 46-58.

- Stadtman, E.R., Van Remmen, H., Richardson, A., Wehr, N.B., Levine, R.L. 2005. Methionine oxidation and aging. *Biochimica et Biophysica Acta (BBA)-Prot. Proteomics* 1703, 135–140.
- Stefan, M.I., Edelstein, S.J., Le Novère N., 2008. An allosteric model of calmodulin explains differential activation of PP2B and CaMKII. *Proc. Natl. Acad. Sci. USA* 105, 10768–10773.
- Stohs, S.J., Bagchi, D. 1995. Oxidative mechanisms in the toxicity of metal ions. *Free Rad. Biol. Med.* 18, 321–336.
- Sukalovic, V.H.T., Vuletic, M., Markovic, K., Vucinic, Z., 2011. Cell wall associated malate dehydrogenase activity from maize roots. *Plant Sci.* 181, 465–470.
- Suzuki, C.K., Rep, M., Van Dijl, J.M., Suda, K., Grivell, L.A., Schatz, G., 1997. ATP-dependent proteases that also chaperone protein biogenesis. *Trends Biochem. Sci.* 22, 118–123.
- Tessier, E., Garnier, C., Mullot, J.U., Lenoble, V., Arnaud, M., Raynaud, M., Mounier, S., 2011. Study of the spatial and historical distribution of sediment inorganic contamination in the Toulon Bay (France). *Mar. Poll. Bull.* 62, 2075 – 2086.
- Thompson, E.L., Taylor, D.A., Nair, S.V., Birch, G., Haynes, P.A., Raftos, D.A., 2011. A proteomic analysis of the effects of metal contamination on Sydney Rock Oyster (*Saccostrea glomerata*) haemolymph. *Aquat. Toxicol.* 103, 241–249.
- Thompson, E.L., Taylor, D.A., Nair, S.V., Birch, G., Haynes, P.A., Raftos, D.A., 2012. Proteomic discovery of biomarkers of metal contamination in Sydney Rock oysters (*Saccostrea glomerata*). *Aquat. Toxicol.* 109, 202–212.
- Timperio, A.M., Egidio, M.G., Zolla, L., 2008. Proteomics applied on plant abiotic stresses: role of heat shock proteins (HSP). *J. Proteomics* 71, 391–411.
- Van Assche, F.V., Clijsters, H., 1990. Effects of metals on enzyme activity in plants. *Plant Cell Environ.* 13, 195–206.
- Vannini, C., Marsoni, M., Domingo, G., Domingo, F., Biondi, S., Bracale, M., 2009. Proteomic analysis of chromate-induced modifications in *Pseudokirchneriella subcapitata*. *Chemosphere* 76, 1372–1379.
- Venn, A.A., Quinn, J., Jones, R., Bodnar, A., 2009. P-glycoprotein (multi-xenobiotic resistance) and heat shock protein gene expression in the reef coral *Montastraea franksi* in response to environmental toxicants. *Aquat. Toxicol.* 93, 188–195.
- Vido, K., Spector, D., Lagniel, G., Lopez, S., Toledano, M.B., Labarre, J., 2001. A proteome analysis of the cadmium response in *Saccharomyces cerevisiae*. *J. Biol. Chem.* 276, 8469–8474.
- Wang, D.-Z., Lin, L., Gu, H.-F., Chan, L.L., Hong, H.-S., 2008. Comparative studies on morphology, ITS sequence and protein profile of *Alexandrium tamarense* and *A. catenella* isolated from the China Sea. *Harmful Algae* 7, 1, 106–113.
- Wang, M., Wang, Y., Lin, L., Hong, H., Wang, D., 2011. Proteome profiles in medaka (*Oryzias melastigma*) liver and brain experimentally exposed to acute inorganic mercury. *Aquat. Toxicol.* 103, 129–139.
- Wingler, A., Lea, P.J., Quick, W.P., Leegood, R.C., 2000. Photorespiration: metabolic pathways and their role in stress protection. *Philos. Trans. R. Soc. Lond. Biol. Sci.* 355, 1517–1529.
- Zhang, R.G., Andersson, C.E., Savchenko, A., Skarina, T., Evdokimova, E., Beasley, S., Arrowsmith, C.H., Edwards, A.M., Joachimiak, A., Mowbray, S.L., 2003. Structure of *Escherichia coli* ribose-5-phosphate isomerase: a ubiquitous enzyme of the pentose phosphate pathway and the Calvin cycle. *Structure* 11, 31–42.
- Zhang, C., Yin, Y., Zhang, Y., Lu, Q., Wen, X., Zhu, Z., Zhang, L., Lu, C., 2012. Comparative proteomic study reveals dynamic proteome changes between superhybrid rice LYP9 and its parents at different developmental stages. *J. Plant Physiol.* 169, 387–398.
- Zhou, S., Sauvé, R., Thannhauser, T.W., 2009. Proteome changes induced by aluminium stress in tomato roots. *J. Exp. Bot.* 60, 1849–1857.
- Zou, H.-X., Pang, Q.-Y., Zhang, A.-Q., Lin, L.-D., Li, N., Yan, X.-F., 2015. Excess copper induced proteomic changes in the marine brown algae *Sargassum fusiforme*. *Ecotox. Environ. Safe.* 111, 271–280.**

Supplementary Material
[Click here to download Supplementary Material: SuppData 1.xls](#)

Supplementary Material

[Click here to download Supplementary Material: SuppData 2_.doc](#)

Highlight 1:

Soluble proteome expressed by the invasive neurotoxic dinoflagellate *Alexandrium catenella* strain ACT03 grown under control and trace metal conditions (lead or zinc at 6, 12 and 18 M) was studied thanks to two-dimensional electrophoresis (2-DE).

Highlight 2:

Zinc (6, 12 and 18 M) significantly decreased the total number of proteins constituting the soluble proteome.

Highlight 3:

Exposures to trace metal led to down-regulation of most proteins constituting the soluble proteome.

Highlight 4:

The down-regulated proteins of the soluble proteome were above all involved in the photosynthesis (RUBISCO; FNR; PCP) and in the oxidative stress response (SOD; proteasome α/β subunits).

Highlight 5:

The up-regulation of some proteins like ATP-synthase β subunit (+16.3 fold in response to contamination by lead at 12 M) suggests a proteomic adaptive response, may be contributing to the tolerance of *A. catenella* ACT03 in trace metal-contaminated ecosystems.