

# Microbes as engines of ecosystem function: When does community structure enhance predictions of ecosystem processes?

E.B. Graham, J.E. Knelman, A. Schindlbacher, S. Siciliano, M. Breulmann, A. Yannarell, J.M. Beman, G. Abell, L. Philippot, J. Prosser, et al.

## ► To cite this version:

E.B. Graham, J.E. Knelman, A. Schindlbacher, S. Siciliano, M. Breulmann, et al.. Microbes as engines of ecosystem function: When does community structure enhance predictions of ecosystem processes?. *Frontiers in Microbiology*, Frontiers Media, 2016, 7, pp.214. <10.3389/fmicb.2016.00214>. <hal-01588095>

**HAL Id: hal-01588095**

**<https://hal.archives-ouvertes.fr/hal-01588095>**

Submitted on 15 Sep 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# Microbes as Engines of Ecosystem Function: When Does Community Structure Enhance Predictions of Ecosystem Processes?

## OPEN ACCESS

### Edited by:

Gary M. King,  
Louisiana State University, USA

### Reviewed by:

Steffen Kolb,  
Landscape Biogeochemistry – Leibniz  
Centre for Agricultural Landscape  
Research, Germany  
Hongchen Jiang,  
Miami University, USA  
Kristen M. DeAngelis,  
University of Massachusetts Amherst,  
USA

### \*Correspondence:

Emily B. Graham  
emily.graham@colorado.edu

### Specialty section:

This article was submitted to  
Terrestrial Microbiology,  
a section of the journal  
Frontiers in Microbiology

**Received:** 09 December 2015

**Accepted:** 09 February 2016

**Published:** 24 February 2016

### Citation:

Graham EB, Knelman JE, Schindlbacher A, Siciliano S, Breulmann M, Yannarell A, Beman JM, Abell G, Philippot L, Prosser J, Foulquier A, Yuste JC, Glanville HC, Jones DL, Angel R, Salminen J, Newton RJ, Bürgmann H, Ingram LJ, Hamer U, Siljanen HMP, Peltoniemi K, Potthast K, Bañeras L, Hartmann M, Banerjee S, Yu R-Q, Nogaro G, Richter A, Koranda M, Castle SC, Goberna M, Song B, Chatterjee A, Nunes OC, Lopes AR, Cao Y, Kaisermann A, Hallin S, Strickland MS, Garcia-Pausas J, Barba J, Kang H, Isobe K, Papaspyrou S, Pastorelli R, Lagomarsino A, Lindström ES, Basiliko N and Nemergut DR (2016) Microbes as Engines of Ecosystem Function: When Does Community Structure Enhance Predictions of Ecosystem Processes? *Front. Microbiol.* 7:214. doi: 10.3389/fmicb.2016.00214

Emily B. Graham<sup>1,2\*</sup>, Joseph E. Knelman<sup>1,3</sup>, Andreas Schindlbacher<sup>4</sup>, Steven Siciliano<sup>5</sup>, Marc Breulmann<sup>6</sup>, Anthony Yannarell<sup>7</sup>, J. M. Beman<sup>8</sup>, Guy Abell<sup>9</sup>, Laurent Philippot<sup>10</sup>, James Prosser<sup>11</sup>, Arnaud Foulquier<sup>12</sup>, Jorge C. Yuste<sup>13</sup>, Helen C. Glanville<sup>14</sup>, Davey L. Jones<sup>14</sup>, Roey Angel<sup>15</sup>, Janne Salminen<sup>16</sup>, Ryan J. Newton<sup>17</sup>, Helmut Bürgmann<sup>18</sup>, Lachlan J. Ingram<sup>19</sup>, Ute Hamer<sup>20</sup>, Henri M. P. Siljanen<sup>21</sup>, Krista Peltoniemi<sup>22</sup>, Karin Potthast<sup>23</sup>, Lluís Bañeras<sup>24</sup>, Martin Hartmann<sup>25</sup>, Samiran Banerjee<sup>26</sup>, Ri-Qing Yu<sup>27</sup>, Geraldine Nogaro<sup>28</sup>, Andreas Richter<sup>29</sup>, Marianne Koranda<sup>30</sup>, Sarah C. Castle<sup>31</sup>, Marta Goberna<sup>32</sup>, Bongkeun Song<sup>33</sup>, Amitava Chatterjee<sup>34</sup>, Olga C. Nunes<sup>35</sup>, Ana R. Lopes<sup>35</sup>, Yiping Cao<sup>36</sup>, Aurore Kaisermann<sup>37</sup>, Sara Hallin<sup>38</sup>, Michael S. Strickland<sup>39</sup>, Jordi Garcia-Pausas<sup>40</sup>, Josep Barba<sup>41</sup>, Hojeong Kang<sup>42</sup>, Kazuo Isobe<sup>43</sup>, Sokratis Papaspyrou<sup>44</sup>, Roberta Pastorelli<sup>45</sup>, Alessandra Lagomarsino<sup>45</sup>, Eva S. Lindström<sup>46</sup>, Nathan Basiliko<sup>47</sup> and Diana R. Nemergut<sup>1,48</sup>

<sup>1</sup> Institute of Arctic and Alpine Research, University of Colorado Boulder, Boulder, CO, USA, <sup>2</sup> Biological Sciences Division, Pacific Northwest National Laboratory, Richland, WA, USA, <sup>3</sup> US Department of Energy, Joint Genome Institute, Walnut Creek, CA, USA, <sup>4</sup> Department of Forest Ecology, Federal Research and Training Centre for Forests, Bundesforschungs- und Ausbildungszentrum für Wald, Vienna, Austria, <sup>5</sup> Department of Soil Science, University of Saskatchewan, Saskatoon, SK, Canada, <sup>6</sup> Helmholtz Centre for Environmental Research – Centre for Environmental Biotechnology, Leipzig, Germany, <sup>7</sup> Department of Natural Resources and Environmental Sciences, University of Illinois at Urbana-Champaign, Urbana, IL, USA, <sup>8</sup> Life and Environmental Sciences and Sierra Nevada Research Institute, University of California – Merced, Merced, CA, USA, <sup>9</sup> School of Medicine, Flinders University, Adelaide, SA, Australia, <sup>10</sup> Institut National de la Recherche Agronomique – Agroecology, Dijon, France, <sup>11</sup> Institute of Biological and Environmental Sciences, University of Aberdeen, Aberdeen, UK, <sup>12</sup> Irstea, UR MALY, Centre de Lyon-Villeurbanne, Villeurbanne, France, <sup>13</sup> Department of Biogeography and Global Change, Museo Nacional de Ciencias Naturales, Consejo Superior de Investigaciones Científicas, Madrid, Spain, <sup>14</sup> Environment Centre Wales, Bangor University, Gwynedd, UK, <sup>15</sup> Department of Microbiology and Ecosystem Science, University of Vienna, Vienna, Austria, <sup>16</sup> Häme University of Applied Sciences, Hämeenlinna, Finland, <sup>17</sup> School of Freshwater Sciences, University of Wisconsin-Milwaukee, Milwaukee, WI, USA, <sup>18</sup> Department of Surface Waters, Eawag: Swiss Federal Institute of Aquatic Science and Technology, Kastanienbaum, Switzerland, <sup>19</sup> Centre for Carbon, Water and Food, The University of Sydney, Sydney, NSW, Australia, <sup>20</sup> Institute of Landscape Ecology, University of Münster, Münster, Germany, <sup>21</sup> Department of Environmental and Biological Sciences, University of Eastern Finland, Kuopio, Finland, <sup>22</sup> Natural Resources Institute, Vantaa, Finland, <sup>23</sup> Institute of Soil Science and Site Ecology, Technische University, Dresden, Germany, <sup>24</sup> Institute of Aquatic Ecology, Facultat de Ciències, University of Girona, Girona, Spain, <sup>25</sup> Institute for Sustainability Sciences – Agroscope, Zurich, Switzerland, <sup>26</sup> CSIRO Agriculture Flagship, Crace, ACT, Australia, <sup>27</sup> Department of Biology, University of Texas at Tyler, Tyler, TX, USA, <sup>28</sup> EDF R&D, National Hydraulics and Environmental Laboratory, Chatou, France, <sup>29</sup> Department of Microbiology and Ecosystem Science, University of Vienna, Vienna, Austria, <sup>30</sup> Division of Terrestrial Ecosystem Research, Department of Microbiology and Ecosystem Science, University of Vienna, Vienna, Austria, <sup>31</sup> Department of Ecosystem and Conservation Sciences, University of Montana, Missoula, MT, USA, <sup>32</sup> Centro de Investigación y Docencia Económicas – Consejo Superior de Investigaciones Científicas, Valencia, Spain, <sup>33</sup> Department of Biological Science, Virginia Institute of Marine Science, Gloucester Point, VA, USA, <sup>34</sup> AES School of Natural Resources Sciences, North Dakota State University, Fargo, ND, USA, <sup>35</sup> LEPABE - Laboratory for Process Engineering, Environmental, Biotechnology and Energy, Faculdade de Engenharia da Universidade do Porto, Porto, Portugal, <sup>36</sup> Southern California Coastal Water Research Project Authority, Costa Mesa, CA, USA, <sup>37</sup> UMR, Interactions Sol Plante Atmosphère, INRA Bordeaux, Villenave d'Ornon, France, <sup>38</sup> Department of Forest Mycology and Plant Pathology, Swedish University of Agricultural Sciences, Uppsala, Sweden, <sup>39</sup> Department of Biological Sciences, Virginia Polytechnic Institute, State University, Blacksburg, VA, USA, <sup>40</sup> Centre Tecnològic Forestal de Catalunya, Solsona, Spain, <sup>41</sup> Centre de Recerca Ecològica i Aplicacions Forestals, Cerdanyola del Vallès, Barcelona, Spain, <sup>42</sup> School of Civil and Environmental Engineering, Yonsei University, Seoul, South Korea, <sup>43</sup> Department of Applied Biological Chemistry, The University of Tokyo, Tokyo, Japan, <sup>44</sup> Department of Biomedicine, Biotechnology and Public Health, University of Cadiz, Puerto Real, Spain, <sup>45</sup> Research Centre for Agrobiology and Pedology, Florence, Italy, <sup>46</sup> Department of Ecology and Genetics/Limnology, Uppsala University, Uppsala, Sweden, <sup>47</sup> Vale Living with Lakes Centre and Department of Biology, Laurentian University, Sudbury, ON, Canada, <sup>48</sup> Biology Department, Duke University, Durham, NC, USA

Microorganisms are vital in mediating the earth's biogeochemical cycles; yet, despite our rapidly increasing ability to explore complex environmental microbial communities, the relationship between microbial community structure and ecosystem processes remains poorly understood. Here, we address a fundamental and unanswered question in microbial ecology: 'When do we need to understand microbial community structure to accurately predict function?' We present a statistical analysis investigating the value of environmental data and microbial community structure independently and in combination for explaining rates of carbon and nitrogen cycling processes within 82 global datasets. Environmental variables were the strongest predictors of process rates but left 44% of variation unexplained on average, suggesting the potential for microbial data to increase model accuracy. Although only 29% of our datasets were significantly improved by adding information on microbial community structure, we observed improvement in models of processes mediated by narrow phylogenetic guilds via functional gene data, and conversely, improvement in models of facultative microbial processes via community diversity metrics. Our results also suggest that microbial diversity can strengthen predictions of respiration rates beyond microbial biomass parameters, as 53% of models were improved by incorporating both sets of predictors compared to 35% by microbial biomass alone. Our analysis represents the first comprehensive analysis of research examining links between microbial community structure and ecosystem function. Taken together, our results indicate that a greater understanding of microbial communities informed by ecological principles may enhance our ability to predict ecosystem process rates relative to assessments based on environmental variables and microbial physiology.

**Keywords:** microbial diversity, functional gene, statistical modeling, microbial ecology, ecosystem processes, respiration, nitrification, denitrification

## INTRODUCTION

The links between complex environmental microbial communities and ecosystem processes remain unclear (Carney and Matson, 2005; Prosser et al., 2007; van der Heijden et al., 2008; Petersen et al., 2012; Wallenstein and Hall, 2012; Graham et al., 2014; Martiny et al., 2015), and an emerging field of research has begun to investigate the utility of microbial data for improving predictions of carbon (C) and nitrogen (N) cycling beyond estimates based solely on environmental data (Todd-Brown et al., 2012; Wieder et al., 2013; Reed et al., 2014; Powell et al., 2015). While researchers have attempted to enhance ecosystem process models by parameterizing microbial physiological properties such as drought tolerance (Manzoni et al., 2014), growth efficiency (Hagerty et al., 2014), dormancy (Wang et al., 2015), and turnover rates (Wieder et al., 2013), these models often fail to consider variation in microbial community structure that may regulate ecosystem process rates (Bouskill et al., 2012; Kaiser et al., 2014). As such, we still lack an integrated understanding of the interactions between microbial communities and ecosystem function, and a central question in ecosystem science remains: 'Under what circumstances does information on microbial communities add to our predictive power of ecosystem processes?' Addressing this question is essential not only for improving knowledge on how critical

biogeochemical cycles may respond to current and impending environmental change, but also for allowing us to identify factors that determine microbial community structure and activity in space and time.

The specific conditions in which microbial community structure – broadly defined here as information on diversity and/or abundance of taxa within a community – may improve predictions of ecosystem process rates beyond models based on environmental and physiological attributes varies with a myriad of biotic and abiotic factors (Knelman and Nemergut, 2014; Nemergut et al., 2014). Research has demonstrated global patterns in microbial communities that correlate with environmental factors such as salinity (Lozupone and Knight, 2007; Auguet et al., 2010), pH (Lauber et al., 2009), and habitat type (Dinsdale et al., 2008; Nemergut et al., 2011; Fierer et al., 2012), and links between microbial community and ecosystem processes have been observed within numerous individual studies (van der Heijden et al., 1998; Torsvik and Øvreås, 2002; Bardgett and van der Putten, 2014; Wagg et al., 2014). However, these relationships and the underlying ecological principles that generate them may vary among environments, relevant microbial traits, and processes of interest (Wallenstein and Hall, 2012; Nemergut et al., 2014). For example, stochastic assembly processes (Hubbell, 2001; Rosindell et al., 2011; Stegen et al., 2012; Nemergut et al., 2013), phenotypic plasticity

(DeWitt et al., 1998), and priority effects (Fukami, 2004; Fukami et al., 2010) can decouple microbial community structure from environmental conditions, and under such conditions, microbial community structure should be central to explaining ecosystem process rates. A unifying perspective that accounts for the dynamic relationships between microbial community structure and ecosystem function is vital for improving predictions of ecosystem process rates, yet recent work has been largely unable to decipher coherent trends in such relationships across studies (Rocca et al., 2014; Bier et al., 2015).

Thus, to assess the importance of microorganisms in explaining ecosystem processes, we used statistical modeling to evaluate the value of environmental variables and microbial community structure both alone and in conjunction for explaining rates of C and N cycling processes within global datasets. Although the literature reporting environmental variables, microbial community data, and biogeochemical processes in a single study is sparse (Rocca et al., 2014), we collected and re-analyzed 82 datasets spanning a multitude of diverse environments. We hypothesized that while the environment is a strong driver of most ecosystem processes, microbial community structure adds to our understanding of processes under certain circumstances, and we discuss patterns across ecological, biogeochemical, and phylogenetic subsets of data. Our analysis generates new insights about the relationship between microbial community structure and a variety of ecosystem processes.

## MATERIALS AND METHODS

We completed an exhaustive literature review for studies measuring environmental variables, microbial community structure, and biogeochemical processes across a range of ecosystems (Supplementary Table S1). We then constructed statistical models for three predictor sets and their additive combinations – environmental data, microbial community data, and microbial biomass measurements (Supplementary Figure S1). Because the measurement of microbial biomass (defined as microbial C and/or N content) was not a criterion for dataset selection, only 28 datasets included biomass measurements. To incorporate differences in microbial community structure into our models, we used ordination-based techniques to condense multivariate community data into vectors that we included as predictors in our models (Supplementary Methods). The Shannon diversity index of each sample was also included as a measure of alpha diversity. In total, these taxonomic metrics are referred to as ‘community diversity’ when applied to 16S rRNA genes or PLFAs and ‘functional diversity’ when applied to functional genes. Functional gene abundances were included as measures of total gene abundance per gram of soil when possible ( $n = 17$ ), although five datasets only listed gene abundances as normalized relative to 16S rRNA gene abundance.

Within each type of data, methodology to collect environmental, microbial, and process data, as well as variables collected, inevitably varied. To minimize potential error introduced by these differences, we re-analyzed data within each

dataset using a multimodel inference approach with the ‘MuMIn’ package (Barton, 2011) in R software (R Core Team, 2014) and subsequently compared results from our analyses. Multimodel inference is a broad regression-based, model-averaging statistical approach designed to reduce errors in model selection, and this method has the advantage of standardizing our approach across studies while accounting for a lack of *a priori* system-specific knowledge (Burnham and Anderson, 2002). This statistical approach has been used in other small scale studies with similar objectives (Powell et al., 2015). We validated the accuracy of MuMIn in our dataset by comparing models for several datasets with expert-built regression models, which yielded comparable results (more details available in Supplementary Methods).

Using the ‘dredge’ command in the ‘MuMIn’ package to fit and evaluate the explanatory power of all possible combinations of variables within a predictor set on a process rate, we selected a set of best fit models for each predictor set consisting of all models with a delta AICc value no more than four greater than the model with the lowest AICc value (Burnham and Anderson, 2004). We generated an averaged final model from this model set using Akaike’s weights, implemented with the ‘model.avg’ command in R (Supplementary Methods). Final models from different predictor sets were compared for statistically significant differences at a delta AICc value of four to provide conservative estimates of model improvements (Burnham and Anderson, 2004). Models were also evaluated for ecologically relevant improvement, defined by an increase in adjusted  $R^2$  value greater than 10% of the environmental model adjusted  $R^2$  value. This criterion was implemented to remove artifacts from datasets in which statistically different models, based on AICc values, yielded similar adjusted  $R^2$  values. Only models that showed both statistical and ecological improvement were considered to be improved.

Finally, model results were synthesized across studies within ecologically relevant subsets of data. We examined results within the full dataset and within biogeochemical process [C mineralization (referred to here as ‘respiration’), nitrification, denitrification, N mineralization], microbial data (PLFA, tRFLP, ARISA, DGGE, qPCR, next generation sequencing) and environment types (natural soil, sediment, agricultural soil) with sufficient replication ( $n \geq 12$ ). We report the mean adjusted  $R^2$  of models and the increase in adjusted  $R^2$  value as measures of effect size as well as the percent of models statistically improved by the addition of microbial data relative to models constructed with only environmental parameters. Differences among the explanatory power of models with environmental, microbial, or both environmental and microbial data were assessed by comparing the distribution of model adjusted  $R^2$  values within each predictor set using unpaired one-sided Mann–Whitney  $U$ -tests for non-parametric data. We also examined correlations between predictor sets to determine the extent to which environmental variables explained variation in microbial community structure and biomass and to which biomass explained variation in microbial community structure. We analyzed these relationships with redundancy analysis (RDA), utilizing the ‘ordisep’ function in the ‘vegan’ package (Oksanen



et al., 2013) in *R* to automate forward model selection, and we report the average  $R^2$  values of correlations between predictor sets as a measure of effect size (Supplementary Methods).

## RESULTS

Overall, models based on environmental variables explained significantly more variation in processes than models based on microbial community structure ( $n = 82$ , average adjusted  $R^2$  0.56 vs. 0.31, Mann–Whitney *U*,  $p < 0.0001$ , **Figure 1**). On average, models with both environmental and microbial predictors explained more variation in processes than environmental models ( $n = 82$ , average adjusted  $R^2$  0.65 vs. 0.56, Mann–Whitney *U*,  $p = 0.046$ ); however, only 29% of datasets were significantly improved by adding information on microbial community structure, by an average of 0.08 increase in adjusted  $R^2$  within all models. Data on microbial community structure from ARISA, tRFLP, qPCR, and next generation technology did not differ in their explanatory power of process rates, but DGGE and PLFA had significantly lower explanatory power than other metrics (Supplementary Figure S2A). All microbial data types showed weak correlations to environmental variables but gene abundance data displayed a trend for higher correlation ( $n = 22$ , average adjusted  $R^2$  0.36) than community ( $n = 55$ , average adjusted  $R^2$  0.20) or functional diversity ( $n = 5$ , average adjusted  $R^2$  0.21) metrics (Supplementary Figure S2B).


We also examined each process individually for those functions for which we had the most data: nitrification ( $n = 14$ ),


denitrification ( $n = 17$ ), N mineralization ( $n = 12$ ), and respiration ( $n = 26$ ). Overall, variation in N mineralization and denitrification rates was less well-explained than other processes by any predictor set (**Figure 2**). Only 17% of datasets examining N mineralization were improved by microbial community data – an average increase in adjusted  $R^2$  of 0.008 within all N mineralization datasets – and the average explanatory power of microbial community structure alone on N mineralization rates was 0.21, compared to 0.31 in the full dataset. Within N mineralization studies that assayed microbial biomass, we found that microbial biomass correlated with environmental variables ( $n = 5$ , average adjusted  $R^2$  0.58) but not with process rates; none of the N mineralization models were improved with the addition of data on biomass. No denitrification studies examined microbial biomass or PLFAs. However, despite low replication, it is notable that in three of four studies, models of denitrification were improved with the addition of 16S rRNA gene diversity data (0.13 average increase in adjusted  $R^2$ ), while only three of eleven models were improved with data on functional gene abundance (0.04 average increase in adjusted  $R^2$ ).

Rates of nitrification were well-explained by all predictor sets (**Figure 2**). Nitrification models based on microbial data yielded the highest average adjusted  $R^2$  of any process (0.38 across all data types and 0.50 if PLFA data are removed) with gene abundance ( $n = 4$ ) data showing higher adjusted  $R^2$  than community diversity data ( $n = 8$ ; 0.61 vs. 0.21, Mann–Whitney *U*,  $p = 0.008$ ). Moreover, 50% of environmental models of nitrification were improved with data on functional gene abundance or functional diversity ( $n = 6$ ), while only 30% of models were improved with data on 16S rRNA gene diversity ( $n = 6$ ), and PLFA data only improved 20% of models ( $n = 2$ ). As with N mineralization, biomass was correlated with environmental variables ( $n = 3$ , average adjusted  $R^2$  0.62) and did not add explanatory power to nitrification rates.

Respiration rates were well-explained by the environment, with an average adjusted  $R^2$  of 0.66 (**Figure 3**). Similar to the full dataset, microbial-only respiration models showed an average adjusted  $R^2$  of 0.29 and only 23% of respiration models were improved with the addition of 16S rRNA gene diversity or PLFA data (no functional gene abundance datasets existed), by an average increase in 0.06 in adjusted  $R^2$  across all respiration datasets. Interestingly, microbial biomass, when measured, improved 35% of models of respiration rates ( $n = 17$ , 0.09 average increase in adjusted  $R^2$ ), while 53% of models were improved with data on biomass and community structure in combination (0.15 average increase in adjusted  $R^2$ ). Microbial biomass was correlated to environmental variables with an average adjusted  $R^2$  of 0.56 but not to microbial community structure (average adjusted  $R^2$  0.11).

Although most of our datasets were from natural soil studies ( $n = 47$ ), we had sufficient replication of sediment ( $n = 12$ ) and agricultural soil ( $n = 12$ ) datasets to examine these independently; other environments ( $n < 12$ ) were excluded from this analysis. While statistically significant differences were not found, possibly due to the large variation in our datasets, we observed some notable trends across environments. Datasets from sediment samples were less well-explained







by environmental or microbial predictor sets than other datasets (Figure 4). Relationships between community diversity and environmental conditions were also weak in sediment systems ( $n = 8$ , average adjusted  $R^2$  0.17), and community diversity metrics provided more added value (0.12 average increase in adjusted  $R^2$ ) in sediments than other systems. Agricultural systems had high explanatory power of process by both environmental (average adjusted  $R^2$  0.59) and microbial predictors (average adjusted  $R^2$  0.38) relative to other systems, and metrics of community diversity were more strongly correlated with the environment ( $n = 7$ , average adjusted  $R^2$  0.24) and had lower added value (average adjusted  $R^2$  0.06) than in other ecosystem types.

## DISCUSSION

Despite the vast amount of variation in our datasets, our results indicate that data on microbial community structure may strengthen predictions of C and N cycling rates. Within our full dataset, environmental variables alone had greater explanatory power than either metrics of microbial community structure or microbial biomass but left 44% of variation in process rates unexplained on average (Figure 1). While we acknowledge that factors such as unmeasured variables, macroinvertebrates (Ferris et al., 2004), fungi (Talbot et al., 2008), plant communities (Hanson et al., 2000), and non-linearity could reduce the explanatory power of our models, a number of ecological dynamics can disassociate environmental conditions from

microbial communities mediating C and N cycle processes and alter the relative value of environmental and microbial data for explaining rates of ecosystem processes (Knelman and Nemergut, 2014; Nemergut et al., 2014). When such factors strongly impact microbial communities, data on microbial community structure should enhance our predictions of ecosystem processes beyond those based solely on environmental data.

We observed stronger trends in the explanatory power of microbial predictors on ecosystem process rates when we reduced noise by examining subsets of data containing a single process or environment type. However, the lack of consistent trends within the full dataset is not surprising due to a variety of ecological and methodological factors. For instance, microorganisms can enter a state of lowered activity or dormancy in response to unfavorable environmental conditions (Jones and Lennon, 2010; Lennon and Jones, 2011), and research suggests that the percent of soil microbial communities catalyzing chemical reactions in soils at a given point in time can vary widely with resource history and disturbances (Blagodatskaya and Kuzyakov, 2013). These factors also induce spatially and temporally variable relationships between microbial communities and ecosystem processes, and the same microbial community may display different relationships to ecosystem function when sampled through time. Moreover, the extent to which functional traits are phylogenetically conserved varies with the trait of interest (Martiny et al., 2013), and thus, some processes may exhibit inherently stronger correlations with taxonomic metrics than others. Although, rapidly improving transcriptomic and



proteomic sequencing technology may enhance predictions of ecosystem processes by identifying active segments of microbial communities (Abram, 2015), too few datasets presently exist to conduct a robust analysis with these data. Studies investigating the extent to which these dynamics are masked within observations of microbial community structure and function are imperative to deciphering and improving the current ability of microbial data to predict ecosystem process rates.

As well, discrepancies in methodology may also confound the relationship we observed in the full dataset between microbial communities and biogeochemical processes (Sinsabaugh et al., 2015). Microbial DNA is typically extracted from a fraction of gram of soil at a single time point and methods detect both dormant and active organisms, while corresponding process data are measured in several grams of soil over longer timeframes. Further, process rate measures are either collected as net measurements that aggregate multiple chemical reactions, potential rates that represent microbial response to substrate addition, or gross rates that trace individual chemical transformations (Bier et al., 2015). These approaches inevitably introduce data aggregation issues (Clark et al., 2011) and rely on assumptions that microbial and biogeochemical data are spatially and temporally representative. Recently, studies have attempted to reduce uncertainty associated with scalar differences by pairing spatially and temporally explicit field studies with ecological modeling approaches that interpolate data across

scales (Lira-Noriega et al., 2013; Walker and Wardle, 2014). Leveraging these approaches in conjunction requires substantial resource investment but is increasingly feasible and presents a promising avenue for maximizing the value of microbial data.

Despite inherent limitations in microbial data, our data analysis suggests that microbial data types differ in their ability to explain processes that are phylogenetically broad vs. narrow and that are products of facultative vs. obligate metabolisms (Figure 2). Nitrification is the product of an obligate chemolithotrophic metabolism, and functional markers for this process are narrowly constrained within ammonia-oxidizing bacteria and archaea that catalyze the rate-limiting step in nitrification, putatively representing a likely dataset for detecting relationships between microbial data and process rates (Schimel, 1995; Schimel and Gullede, 1998). While both functional gene abundance and community diversity metrics had similar average explanatory power within the full dataset (an effect that also did not vary among microbial technologies, Supplementary Figure S2A), models based on functional gene abundances or functional diversity explained more variation in process rates than community diversity within datasets measuring rates of nitrification (Figure 2C). Although these metrics were more strongly correlated with environmental factors than community diversity, they also provided greater explanatory power in combination with environmental variables

than community diversity metrics, which include variation in many microbial guilds external to nitrification. Notably, models based on environmental variables alone also had higher explanatory power for nitrification than for denitrification or N mineralization, suggesting that the environment may also be more strongly coupled to nitrification rates than other N cycle processes. Taken together, these results suggest that despite a relatively tight connection between the environment, microbial community structure, and process data, the microbial guilds driving nitrification rates are partially decoupled at a functional level from environmental drivers of nitrification. Importantly, although nitrification was the only narrow process in our dataset with sufficient replication to examine independently, the high explanatory power of functional genes, both alone and in combination with environmental variables, for describing nitrification rates relative to other processes also conveys the potential for the inclusion of functional metrics in improving predictions of ecosystem processes that are mediated by phylogenetically narrow functional guilds.

By contrast, denitrification is a broad process, catalyzed by diverse facultative anaerobic microorganisms, and community diversity metrics provided more added value to denitrification rates than functional gene diversity or abundances (**Figure 2B**). While we acknowledge that genetic markers may not denote all functional genes involved in a process (Jones et al., 2014; Verbaendert et al., 2014), we observed more added value from community diversity metrics than functional genes despite several studies assaying up to three genes to explain denitrification rates, as opposed to one or two functional markers measured for nitrification. For broad processes such as denitrification, representations of niche complementarity or resource availability reflected in diversity metrics (Cadotte et al., 2011) rather than functional gene abundances that do not account for interactions between individuals (Barberán et al., 2012; Gagic et al., 2015) or functional redundancy (Allison and Martiny, 2008), may be important for explaining process rates. Thus, it is plausible that denitrifying communities are decoupled from environmental variables when measured at a broad taxonomic level and that such taxonomic metrics encompass more variation in within community interactions affecting denitrification rates than functional approaches. Although the sample size of datasets measuring denitrification was small, our analysis suggests that measurements of community diversity may be more likely than functional metrics to improve predictions of ecosystem processes that are catalyzed by facultative metabolisms and/or phylogenetically broad suites of organisms.

Similarly, the ability to respire carbon is widely distributed among microorganisms, and community diversity metrics (no datasets measured functional markers) described rates of respiration better than all other processes except denitrification (**Figure 3**). In fact, environmental and microbial variables, both alone and in combination, yielded higher average explanatory power of respiration rates than all other processes we examined independently with the exception of nitrification. Importantly, while recent work has suggested that microbial biomass (Schimel and Weintraub, 2003) or physiological properties such as drought tolerance (Manzoni et al., 2014) and growth efficiency (Wieder

et al., 2013) can improve C cycling models, our results indicate that information on microbial community structure may further enhance our understanding of ecosystem C cycling. Microbial community structure and biomass appeared to explain differing portions of variation in process, as they exhibited extremely weak correlations with each other and, together, provided additive value for explaining rates of respiration. Thus, biomass and structure may jointly contribute to explaining rates of respiration as biomass may serve as a proxy for unmeasured environmental variables that regulate activity, such as soil structure (Gupta and Germida, 1988), while community structure may relate to the genetic capability of a community to respire carbon. Regardless, our results suggest that these data are at least partially independent and that measurements of microbial community structure could reduce uncertainty within Earth Systems models.

Lastly, the relationship between microbial community structure and ecosystem processes in our dataset varied by environment type (**Figure 4**). Sediment systems had weaker explanatory power of process rates by both environmental variables and microbial community structure than natural or agricultural soils. Sediment processes are influenced by both sediment and porewater chemistry (Middelburg and Levin, 2009), and these datasets may be more likely to have unmeasured variables or errors due to spatial discrepancies than other systems. In particular, dissolved oxygen concentration is a strong determinant of redox potential and ecosystem processes in sediments (Abell et al., 2011) but was only measured in three of 12 datasets. Despite the low explanatory power of sediment processes by all predictor sets, microbial community diversity added more value to explaining process rates and was more decoupled from environmental variables in sediments than in natural soil or agricultural systems, indicating that measurements of microbial community structure may improve predictions of sediment processes. Conversely, in agricultural soils, microbial community structure explained more variation in ecosystem processes but provided less improvement over environmental variables than other systems. Environmental variables were also better predictors of process in agricultural soils, and microbial community diversity was more tightly correlated with the environment than in sediments or natural soils. Agricultural systems homogenize soil structure and decrease soil organic matter (Dick, 1992; Chan and Heenan, 1996), reducing the number and complexity of microbial niches in agricultural soils. Thus, relatively coarse-scale measures of environmental conditions may more directly correlate with both community structure and ecosystem process rates than in more variable systems. Overall, these results indicate that ecosystem-specific dynamics may be crucial to understanding the value of microbial community structure for explaining ecosystem processes and emphasize the importance of future investigations into understudied biomes in enhancing our understanding of global relationships between microbial communities and ecosystem processes.

Here, we present the first extensive investigation into the relationship between environmental conditions, microbial community structure, and ecosystem function by re-analyzing 82



datasets collected from an international team of collaborators, and we demonstrate that nuanced relationships between the environment and microbial communities can influence the importance of microbial community structure for explaining ecosystem-level processes. Our analysis provides an empirical basis for future hypothesis-testing on the roles of assembly, dormancy, redundancy and phenotypic plasticity in microbial community structure and function, and despite complexity in our datasets, the trends we observed represent an encouraging step for linking microbial community data to ecosystem function. As a whole, our results indicate that a greater understanding of microbial community structure informed by ecological principles may further our ability to accurately predict rates of ecosystem processes beyond environmental variables and bulk physiological characterizations of microbial communities.

## AUTHOR CONTRIBUTIONS

All authors collected and contributed datasets for analysis as well as participated in the conceptual drafting and revision of this manuscript. EG conducted all data analysis and was the primary author in writing and revising the manuscript. DN

and JK contributed significantly in manuscript development and revision, and DN provided financial support for this work.

## FUNDING

This work was supported by NSF grant DEB-1221215 to DN, as well as grants supporting the generation of our datasets as acknowledged in their original publications and in Supplementary Table S1.

## ACKNOWLEDGMENT

We thank the USGS Powell Center 'Next Generation Microbes' working group, anonymous reviews, Brett Melbourne, and Alan Townsend for valuable feedback on this project.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fmicb.2016.00214>

## REFERENCES

- Abell, G. C., Banks, J., Ross, D. J., Keane, J. P., Robert, S. S., Revill, A. T., et al. (2011). Effects of estuarine sediment hypoxia on nitrogen fluxes and ammonia oxidizer gene transcription. *FEMS Microbiol. Ecol.* 75, 111–122. doi: 10.1111/j.1574-6941.2010.00988.x
- Abram, F. (2015). Systems-based approaches to unravel multi-species microbial community functioning. *Comput. Struct. Biotechnol. J.* 13, 24–32. doi: 10.1016/j.csbj.2014.11.009
- Allison, S. D., and Martiny, J. B. (2008). Resistance, resilience, and redundancy in microbial communities. *Proc. Natl. Acad. Sci. U.S.A.* 105, 11512–11519. doi: 10.1073/pnas.0801925105
- Auguet, J.-C., Barberan, A., and Casamayor, E. O. (2010). Global ecological patterns in uncultured Archaea. *ISME J.* 4, 182–190. doi: 10.1038/ismej.2009.109
- Barberán, A., Bates, S. T., Casamayor, E. O., and Fierer, N. (2012). Using network analysis to explore co-occurrence patterns in soil microbial communities. *ISME J.* 6, 343–351. doi: 10.1038/ismej.2011.119
- Bardgett, R. D., and van der Putten, W. H. (2014). Belowground biodiversity and ecosystem functioning. *Nature* 515, 505–511. doi: 10.1038/nature13855
- Barton, K. (2011). *MuMIn: Multi-Model Inference. R Package Version 1.0.0*. Vienna, Austria: R Foundation for Statistical Computing. Available at: <http://CRAN.R-project.org/package=MumIn>
- Bier, R. L., Bernhardt, E. S., Boot, C. M., Graham, E. B., Hall, E. K., Lennon, J. T., et al. (2015). Linking microbial community structure and microbial processes: an empirical and conceptual overview. *FEMS Microbiol. Ecol.* 91:fiv113. doi: 10.1093/femsec/fiv113
- Blagodatskaya, E., and Kuzyakov, Y. (2013). Active microorganisms in soil: critical review of estimation criteria and approaches. *Soil Biol. Biochem.* 67, 192–211. doi: 10.1016/j.soilbio.2013.08.024
- Bouskill, N. J., Tang, J., Riley, W. J., and Brodie, E. L. (2012). Trait-based representation of biological nitrification: model development, testing, and predicted community composition. *Front. Microbiol.* 3:364. doi: 10.3389/fmicb.2012.00364
- Burnham, K. P., and Anderson, D. R. (2002). *Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach*. New York, NY: Springer.
- Burnham, K. P., and Anderson, D. R. (2004). Multimodel inference understanding AIC and BIC in model selection. *Sociol. Methods Res.* 33, 261–304. doi: 10.1177/0049124104268644
- Cadotte, M. W., Carscadden, K., and Mirotchnick, N. (2011). Beyond species: functional diversity and the maintenance of ecological processes and services. *J. Appl. Ecol.* 48, 1079–1087. doi: 10.1111/j.1365-2664.2011.02048.x
- Carney, K. M., and Matson, P. A. (2005). Plant communities, soil microorganisms, and soil carbon cycling: does altering the world belowground matter to ecosystem functioning? *Ecosystems* 8, 928–940. doi: 10.1007/s10021-005-0047-0
- Chan, K., and Heenan, D. (1996). The influence of crop rotation on soil structure and soil physical properties under conventional tillage. *Soil Tillage Res.* 37, 113–125. doi: 10.1016/0167-1987(96)01008-2
- Clark, J. S., Bell, D. M., Hersh, M. H., Kwit, M. C., Moran, E., Salk, C., et al. (2011). Individual-scale variation, species-scale differences: inference needed to understand diversity. *Ecol. Lett.* 14, 1273–1287. doi: 10.1111/j.1461-0248.2011.01685.x
- DeWitt, T. J., Sih, A., and Wilson, D. S. (1998). Costs and limits of phenotypic plasticity. *Trends Ecol. Evol.* 13, 77–81. doi: 10.1016/S0169-5347(97)01274-3
- Dick, R. P. (1992). A review: long-term effects of agricultural systems on soil biochemical and microbial parameters. *Agric. Ecosyst. Environ.* 40, 25–36. doi: 10.1016/0167-8809(92)90081-L
- Dinsdale, E. A., Edwards, R. A., Hall, D., Angly, F., Breitbart, M., Brulc, J. M., et al. (2008). Functional metagenomic profiling of nine biomes. *Nature* 452, 629–632. doi: 10.1038/nature06810
- Ferris, H., Venette, R., and Scow, K. (2004). Soil management to enhance bacterivore and fungivore nematode populations and their nitrogen mineralisation function. *Appl. Soil Ecol.* 25, 19–35. doi: 10.1016/j.apsoil.2003.07.001
- Fierer, N., Leff, J. W., Adams, B. J., Nielsen, U. N., Bates, S. T., Lauber, C. L., et al. (2012). Cross-biome metagenomic analyses of soil microbial communities and their functional attributes. *Proc. Natl. Acad. Sci. U.S.A.* 109, 21390–21395. doi: 10.1073/pnas.1215210110
- Fukami, T. (2004). Assembly history interacts with ecosystem size to influence species diversity. *Ecology* 85, 3234–3242. doi: 10.1890/04-0340
- Fukami, T., Dickie, I. A., Paula Wilkie, J., Paulus, B. C., Park, D., Roberts, A., et al. (2010). Assembly history dictates ecosystem functioning: evidence from wood decomposer communities. *Ecol. Lett.* 13, 675–684. doi: 10.1111/j.1461-0248.2010.01465.x

- Gagic, V., Bartomeus, I., Jonsson, T., Taylor, A., Winqvist, C., Fischer, C., et al. (2015). Functional identity and diversity of animals predict ecosystem functioning better than species-based indices. *Proc. R. Soc. Biol. Sci. U.S.A.* 282:20142620. doi: 10.1098/rspb.2014.2620
- Graham, E. B., Wieder, W. R., Leff, J. W., Weintraub, S. R., Townsend, A. R., Cleveland, C. C., et al. (2014). Do we need to understand microbial communities to predict ecosystem function? A comparison of statistical models of nitrogen cycling processes. *Soil Biol. Biochem.* 68, 279–282. doi: 10.1016/j.soilbio.2013.08.023
- Gupta, V., and Germida, J. (1988). Distribution of microbial biomass and its activity in different soil aggregate size classes as affected by cultivation. *Soil Biol. Biochem.* 20, 777–786. doi: 10.1016/0038-0717(88)90082-X
- Hagerty, S. B., van Groenigen, K. J., Allison, S. D., Hungate, B. A., Schwartz, E., Koch, G. W., et al. (2014). Accelerated microbial turnover but constant growth efficiency with warming in soil. *Nat. Clim. Chang.* 4, 903–906. doi: 10.1038/nclimate2361
- Hanson, P., Edwards, N., Garten, C., and Andrews, J. (2000). Separating root and soil microbial contributions to soil respiration: a review of methods and observations. *Biogeochemistry* 48, 115–146. doi: 10.1023/A:1006244819642
- Hubbell, S. P. (2001). *The Unified Neutral Theory of Biodiversity and Biogeography (MPB-32)*. Princeton, NJ: Princeton University Press.
- Jones, C. M., Spor, A., Brennan, F. P., Breuil, M.-C., Bru, D., Lemanceau, P., et al. (2014). Recently identified microbial guild mediates soil N<sub>2</sub>O sink capacity. *Nat. Clim. Chang.* 4, 801–805. doi: 10.1038/nclimate2301
- Jones, S. E., and Lennon, J. T. (2010). Dormancy contributes to the maintenance of microbial diversity. *Proc. Natl. Acad. Sci. U.S.A.* 107, 5881–5886. doi: 10.1073/pnas.0912765107
- Kaiser, C., Franklin, O., Dieckmann, U., and Richter, A. (2014). Microbial community dynamics alleviate stoichiometric constraints during litter decay. *Ecol. Lett.* 17, 680–690. doi: 10.1111/ele.12269
- Knelman, J. E., and Nemergut, D. R. (2014). Changes in community assembly may shift the relationship between biodiversity and ecosystem function. *Front. Microbiol.* 5:424. doi: 10.3389/fmicb.2014.00424
- Lauber, C. L., Hamady, M., Knight, R., and Fierer, N. (2009). Pyrosequencing-based assessment of soil pH as a predictor of soil bacterial community structure at the continental scale. *Appl. Environ. Microbiol.* 75, 5111–5120. doi: 10.1128/AEM.00335-09
- Lennon, J. T., and Jones, S. E. (2011). Microbial seed banks: the ecological and evolutionary implications of dormancy. *Nat. Rev. Microbiol.* 9, 119–130. doi: 10.1038/nrmicro2504
- Lira-Noriega, A., Soberón, J., and Miller, C. P. (2013). Process-based and correlative modeling of desert mistletoe distribution: a multiscalar approach. *Ecosphere* 4:art99. doi: 10.1890/ES13-00155.1
- Lozupone, C. A., and Knight, R. (2007). Global patterns in bacterial diversity. *Proc. Natl. Acad. Sci. U.S.A.* 104, 11436–11440. doi: 10.1073/pnas.0611525104
- Manzoni, S., Schaeffer, S., Katul, G., Porporato, A., and Schimel, J. (2014). A theoretical analysis of microbial eco-physiological and diffusion limitations to carbon cycling in drying soils. *Soil Biol. Biochem.* 73, 69–83. doi: 10.1016/j.soilbio.2014.02.008
- Martiny, A. C., Treseder, K., and Pusch, G. (2013). Phylogenetic conservatism of functional traits in microorganisms. *ISME J.* 7, 830–838. doi: 10.1038/ismej.2012.160
- Martiny, J. B., Jones, S. E., Lennon, J. T., and Martiny, A. C. (2015). Microbiomes in light of traits: a phylogenetic perspective. *Science* 350:aac9323. doi: 10.1126/science.aac9323
- Middelburg, J., and Levin, L. (2009). Coastal hypoxia and sediment biogeochemistry. *Biogeosciences* 6, 3655–3706. doi: 10.5194/bgd-6-3655-2009
- Nemergut, D. R., Costello, E. K., Hamady, M., Lozupone, C., Jiang, L., Schmidt, S. K., et al. (2011). Global patterns in the biogeography of bacterial taxa. *Environ. Microbiol.* 13, 135–144. doi: 10.1111/j.1462-2920.2010.02315.x
- Nemergut, D. R., Schmidt, S. K., Fukami, T., O'Neill, S. P., Bilinski, T. M., Stanish, L. F., et al. (2013). Patterns and processes of microbial community assembly. *Microbiol. Mol. Biol. Rev.* 77, 342–356. doi: 10.1128/MMBR.00051-12
- Nemergut, D. R., Shade, A., and Vielle, C. (2014). When, where and how does microbial community composition matter? *Front. Microbiol.* 5:497. doi: 10.3389/fmicb.2014.00497
- Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R., et al. (2013). *Package 'Vegan'. Community Ecology Package, Version 2*. Available at: <http://CRAN.R-project.org/package=vegan>
- Petersen, D. G., Blazewicz, S. J., Firestone, M., Herman, D. J., Turetsky, M., and Waldrop, M. (2012). Abundance of microbial genes associated with nitrogen cycling as indices of biogeochemical process rates across a vegetation gradient in Alaska. *Environ. Microbiol.* 14, 993–1008. doi: 10.1111/j.1462-2920.2011.02679.x
- Powell, J. R., Welsh, A., and Hallin, S. (2015). Microbial functional diversity enhances predictive models linking environmental parameters to ecosystem properties. *Ecology* 96, 1985–1993. doi: 10.1890/14-1127.1
- Prosser, J. I., Bohannon, B. J., Curtis, T. P., Ellis, R. J., Firestone, M. K., Freckleton, R. P., et al. (2007). The role of ecological theory in microbial ecology. *Nat. Rev. Microbiol.* 5, 384–392. doi: 10.1038/nrmicro1643
- R Core Team (2014). *R: A Language and Environment for Statistical Computing*. Vienna: R Foundation for Statistical Computing.
- Reed, D. C., Algar, C. K., Huber, J. A., and Dick, G. J. (2014). Gene-centric approach to integrating environmental genomics and biogeochemical models. *Proc. Natl. Acad. Sci. U.S.A.* 111, 1879–1884. doi: 10.1073/pnas.1313713111
- Rocca, J. D., Hall, E. K., Lennon, J. T., Evans, S. E., Waldrop, M. P., Cotner, J. B., et al. (2014). Relationships between protein-encoding gene abundance and corresponding process are commonly assumed yet rarely observed. *ISME J.* 9, 1693–1699. doi: 10.1038/ismej.2014.252
- Rosindell, J., Hubbell, S. P., and Etienne, R. S. (2011). The unified neutral theory of biodiversity and biogeography at age ten. *Trends Ecol. Evol.* 26, 340–348. doi: 10.1016/j.tree.2011.03.024
- Schimel, J. (1995). “Ecosystem consequences of microbial diversity and community structure,” in *Arctic and Alpine Biodiversity: Patterns, Causes and Ecosystem Consequences*, ed. C. Krner (Berlin: Springer), 239–254.
- Schimel, J. P., and Gulledge, J. (1998). Microbial community structure and global trace gases. *Glob. Chang. Biol.* 4, 745–758. doi: 10.1046/j.1365-2486.1998.00195.x
- Schimel, J. P., and Weintraub, M. N. (2003). The implications of exoenzyme activity on microbial carbon and nitrogen limitation in soil: a theoretical model. *Soil Biol. Biochem.* 35, 549–563. doi: 10.1016/S0038-0717(03)00015-4
- Sinsabaugh, R. L., Shah, J. J. F., Findlay, S. G., Kuehn, K. A., and Moorhead, D. L. (2015). Scaling microbial biomass, metabolism and resource supply. *Biogeochemistry* 122, 175–190. doi: 10.1007/s10533-014-0058-z
- Stegen, J. C., Lin, X., Konopka, A. E., and Fredrickson, J. K. (2012). Stochastic and deterministic assembly processes in subsurface microbial communities. *ISME J.* 6, 1653–1664. doi: 10.1038/ismej.2012.22
- Talbot, J., Allison, S., and Treseder, K. (2008). Decomposers in disguise: mycorrhizal fungi as regulators of soil C dynamics in ecosystems under global change. *Funct. Ecol.* 22, 955–963. doi: 10.1111/j.1365-2435.2008.01402.x
- Todd-Brown, K. E., Hopkins, F. M., Kivlin, S. N., Talbot, J. M., and Allison, S. D. (2012). A framework for representing microbial decomposition in coupled climate models. *Biogeochemistry* 109, 19–33. doi: 10.1007/s10533-011-9635-6
- Torsvik, V., and Øvreås, L. (2002). Microbial diversity and function in soil: from genes to ecosystems. *Curr. Opin. Microbiol.* 5, 240–245. doi: 10.1016/S1369-5274(02)00324-7
- van der Heijden, M. G. A., Bardgett, R. D., and van Straalen, N. M. (2008). The unseen majority: soil microbes as drivers of plant diversity and productivity in terrestrial ecosystems. *Ecol. Lett.* 11, 296–310. doi: 10.1111/j.1461-0248.2007.01139.x
- van der Heijden, M. G., Klironomos, J. N., Ursic, M., Moutoglou, P., Streitwolf-Engel, R., Boller, T., et al. (1998). Mycorrhizal fungal diversity determines plant biodiversity, ecosystem variability and productivity. *Nature* 396, 69–72. doi: 10.1038/23932
- Verbaandert, I., Hoefman, S., Boeckx, P., Boon, N., and De Vos, P. (2014). Primers for overlooked nirK, qnrB, and nosZ genes of thermophilic Gram-positive denitrifiers. *FEMS Microbiol. Ecol.* 89, 162–180. doi: 10.1111/1574-6941.12346
- Wagg, C., Bender, S. F., Widmer, F., and van der Heijden, M. G. (2014). Soil biodiversity and soil community composition determine ecosystem multifunctionality. *Proc. Natl. Acad. Sci. U.S.A.* 111, 5266–5270. doi: 10.1073/pnas.1320054111
- Walker, L. R., and Wardle, D. A. (2014). Plant succession as an integrator of contrasting ecological time scales. *Trends Ecol. Evol.* 29, 504–510. doi: 10.1016/j.tree.2014.07.002

- Wallenstein, M. D., and Hall, E. K. (2012). A trait-based framework for predicting when and where microbial adaptation to climate change will affect ecosystem functioning. *Biogeochemistry* 109, 35–47. doi: 10.1007/s10533-011-9641-8
- Wang, G., Jagadamma, S., Mayes, M. A., Schadt, C. W., Steinweg, J. M., Gu, L., et al. (2015). Microbial dormancy improves development and experimental validation of ecosystem model. *ISME J.* 9, 226–237. doi: 10.1038/ismej.2014.120
- Wieder, W. R., Bonan, G. B., and Allison, S. D. (2013). Global soil carbon projections are improved by modelling microbial processes. *Nat. Clim. Chang.* 3, 909–912. doi: 10.1038/nclimate1951

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Graham, Knelman, Schindlbacher, Siciliano, Breulmann, Yannarell, Beman, Abell, Philippot, Prosser, Foulquier, Yuste, Glanville, Jones, Angel, Salminen, Newton, Bürgmann, Ingram, Hamer, Siljanen, Peltoniemi, Potthast, Bañeras, Hartmann, Banerjee, Yu, Nogaro, Richter, Koranda, Castle, Goberna, Song, Chatterjee, Nunes, Lopes, Cao, Kaisermann, Hallin, Strickland, Garcia-Pausas, Barba, Kang, Isobe, Papaspyrou, Pastorelli, Lagomarsino, Lindström, Basiliko and Nemergut. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.