**A new cheyletid mite *Metacheyletia ngaii* n. sp.
(Acariformes: Cheyletidae) from quills of *Corythaixoides*
leucogaster (Musophagidae) from Tanzania**

A.V. Bochkov, M. Skoracki

► **To cite this version:**

A.V. Bochkov, M. Skoracki. A new cheyletid mite *Metacheyletia ngaii* n. sp. (Acariformes: Cheyletidae) from quills of *Corythaixoides leucogaster* (Musophagidae) from Tanzania. *Acarologia*, 2011, 51 (1), pp.93-97. 10.1051/acarologia/20111996 . hal-01599952

HAL Id: hal-01599952

<https://hal.science/hal-01599952>

Submitted on 2 Oct 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives| 4.0
International License

ACAROLOGIA

A quarterly journal of acarology, since 1959
Publishing on all aspects of the Acari

All information:

<http://www1.montpellier.inra.fr/CBGP/acarologia/>
acarologia@supagro.inra.fr

**Acarologia is proudly non-profit,
with no page charges and free open access**

Please help us maintain this system by
encouraging your institutes to subscribe to the print version of the journal
and by sending us your high quality research on the Acari.

Subscriptions: Year 2017 (Volume 57): 380 €

<http://www1.montpellier.inra.fr/CBGP/acarologia/subscribe.php>

Previous volumes (2010-2015): 250 € / year (4 issues)

Acarologia, CBGP, CS 30016, 34988 MONTFERRIER-sur-LEZ Cedex, France

The digitalization of Acarologia papers prior to 2000 was supported by Agropolis Fondation under the reference ID 1500-024 through the « Investissements d'avenir » programme (Labex Agro: ANR-10-LABX-0001-01)

Acarologia is under **free license** and distributed under the terms of the Creative Commons-BY-NC-ND which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

A NEW CHEYLETID MITE *METACHEYLETIA* *NGAII* N. SP. (ACARIFORMES:
CHEYLETIDAE) FROM QUILLS OF *CORYTHAIXOIDES LEUCOGASTER*
(MUSOPHAGIDAE) FROM TANZANIA

Andre V. BOCHKOV^{1,2} and Maciej SKORACKI³

(Received 10 December 2010; accepted 15 February 2011; published online 30 March 2011)

¹ Zoological Institute, Russian Academy of Sciences, Universitetskaya emb. 1, 199034 St. Petersburg, Russia. andrevbochkov@gmail.com

² Museum of Zoology, University of Michigan, 1109 Geddes Ave., Ann Arbor, Michigan 48109, USA

³ Department of Animal Morphology, Adam Mickiewicz University, Faculty of Biology, Umultowska 89, 61-614 Poznan, Poland
skoracki@amu.edu.pl

ABSTRACT — *Metacheyletia ngaii* n. sp. (Acariformes: Cheyletidae) is described from quills of *Corythaixoides leucogaster* (Rüppell, 1842) (Cuculiformes: Musophagidae) from Tanzania.

KEYWORDS — Acari; Cheyletidae; *Metacheyletia*; systematics; feather quills; *Corythaixoides leucogaster*

INTRODUCTION

The genus *Metacheyletia* Fain, 1972 (Acariformes: Cheyletidae) is a single genus of the tribe Metacheyletiini (Bochkov and Fain, 2001). Mites of this genus, currently including four species, live inside the feather quills. Three species are known from various parrots (Psittaciformes: Psittacidae), *M. obesa* Fain, 1972 from *Psittacula krameri* (Scopoli, 1769) from Africa (Fain, 1972, 1980), *M. longisetosa* Atyeo *et al.*, 1984 from *Amazona finschi* (Sclater, 1864) from Mexico (Atyeo *et al.*, 1984), *M. amazonae* Bochkov & OConnor, 2003 from *Amazona amazonica* (Linnaeus, 1766) from Guyana (Bochkov and OConnor, 2003); one species, *M. degenerata* Fain & Bochkov, 2003 was described from *Serinus mozambicus* (Muller, 1776) (Passeriformes: Fringillidae) from Tanzania (Fain and Bochkov, 2003).

Few data are known on biology of these mites,

and suggestions concerning their mode of life are controversial. Atyeo *et al.* (1984) believed that these mites are predators because their movable cheliceral digits are too short to penetrate the quill wall, pierce the quill wall and reach live tissues of a host. Bochkov and Fain (2001) considered these mites parasitic based on morphological characters such as the short tarsi of legs I–III, the absence of legs IV, the short and nude eupathidia of the palpal tarsus and the relatively small sizes of the gnathosoma and suggested that *Metacheyletia* spp. use the orifices in quill walls which were made by adult syringophilids.

In this paper, we describe a new species of this genus from *Corythaixoides leucogaster* (Rüppell, 1842) (Cuculiformes: Musophagidae) from Tanzania. A key to females of all known species of the genus *Metacheyletia* is provided.

FAMILY CHEYLETIDAE

Metacheyletia ngaii n. sp.
(Figures 1 and 2)

Description — Female (holotype) — *Gnathosoma* 90 long (90 in 1 paratype) and 115 wide (120). Peritremes arch-like, each peritremal branch with 7–8 segments. Palpal femur 40 long (40) and 37 wide (35). Palpal claw with 1 basal tooth. Palpal setation: palpal femur – *dF*; palpal genu – *dG*; palpal tibia – *dTi*, *l'Ti*, *l''Ti*; palpal tarsus – smooth eupathidia *amc*, *sul*, *ul'*, *ul''*, and solenidion $\omega 1$. Subcapitular setae *n* absent, adoral setae *ao1*, *ao2*, and subcoxal setae *elcp* present. *Idiosoma* 495 long (585) and 355 wide (415),

FIGURE 2: *Metacheyleletia ngaii* sp. nov., details of holotype female. A – Peritremal branch. B – Palpal tibiotarsus in dorsal view. C – Same in ventral view. D – Tarsus I in dorsal view. E – Same in ventral view. F – Tarsus II in dorsal view. G – Same in ventral view. H – Tarsus III in dorsal view. I – Same in ventral view. J – Leg III in ventral view. Scale bars: A–I = 20 μ m; J = 50 μ m.

ovoid. Propodonal shield indistinct. No cupules observed. All idiosomal setae smooth filiform. Setae *vi* 2 times shorter than *ve*; setae *si* and *d2* 1.4–1.5 times shorter than *se* and *e2*; *c1* and *d1* about 2 times longer than *d2*; setae *c1* longer than distance between setal bases *c1* and *d1*; setae *h1* and *h2* subequal. Three pairs of pseudogenital (*ps1*–*ps3*), 2 pairs of genital (*g1* and *g2*), and 3 pairs of aggenital (*ag1*–*ag3*) setae present. Lengths of setae: *vi* 30 (35), *ve* 65 (70), *si* 50, *se* 70 (70), *c1* 130, *c2* 120 (125), *d1* 120 (115), *d2* 50 (65), *e2* 80 (75), *f2* 65 (75), *h1* 120 (115), *h2* 115 (120), *ps1*–*ps3* 23–25, *g1* and *g2* 24–25, *ag1*–*ag3* 23–27, *1a* 35 (35), *1b* 45 (55), *3a* 35 (35). Setae *4a* absent. Legs IV completely absent. Leg I–IV setation: I – tarsus with 7 setae (*ft*, *tc'*, *tc''*, *p'*, *p''*, *u'*, *u''*) + solenidion $\omega 1$, tibia with 4 setae (*d*, *l'*, *l''*, *v*) + solenidion φ , genu with 1 seta (*l'*) + solenidion σ , femur with 2 setae (*dF* and *vF*), trochanter without setae, coxal field with 1 seta (*1b*); II – tarsus with 5 setae (*tc'*, *tc''*, *p'*, *u'*, *u''*) + solenidion $\omega 1$, tibia with 4 setae (*d*, *l'*, *l''*, *v*), genu with 1 seta (*l'*), femur with 2 setae (*dF* and *vF*), trochanter and coxal field without setae; III – tarsus with 5 setae (*tc'*, *tc''*, *p'*, *u'*, *u''*), tibia with 4 setae (*d*, *l'*, *l''*, *v*), genu with 1 seta (*l'*), femur with 1 seta (*dF*), trochanter and coxal field without setae.

Male — Unknown.

Type Material — Holotype female and female paratype from *Corythaixoides leucogaster* (Rüppell, 1842) (Cuculiformes: Musophagidae), Tanzania: Tanganyika, 23 March 1960, coll. unknown.

Type deposition — All material is deposited in the Zoological Institute of the Russian Academy of Sciences, St. Petersburg, Russia.

Etymology — This new species is dedicated to the Maasai God – Ngai, the creator of everything.

Differential diagnosis. This new species is morphologically closest to *M. degenerata*. In females of both species, three pairs of aggenital setae are present and the propodonal shield is indistinct. It differs from *M. degenerata* by the following features. In females of *M. ngaii* n. sp., setae *vi* are two times shorter than *ve*, setae *d2* are two times shorter than *d1*, setae *h1* and *h2* are subequal, setae *c1*, *d1*, and *e2* are 130, 115–120, and 75–80 long, respectively. In females of *M. degenerata*, setae *vi* are 1.1–1.2 times shorter than *ve*, setae *d1* and *d2* are subequal, setae

h1 are about 1.3 times shorter than *h2*, setae *c1*, *d1*, and *e2* are 65, 60, and 50 long, respectively.

Remarks — The genus *Metacheyletia* is, probably, initially associated with parrots, because its representatives inhabit these hosts in Africa and South America. Unfortunately these mites are unknown from Australia, and records of *Metacheyletia* spp. from the Australian parrots would be very desirable as an additional prove of this hypothesis. The two species of this genus from non-parrot hosts, *M. degenerata* and *M. ngaii*, are still known exclusively from central Africa and morphologically very close to each other. Parrots are widely distributed in central Africa and, therefore, we suggest that in this region, mites of the genus *Metacheyletia* shifted on a non-parrot host from a parrot with the following dispersion and speciation on birds of different non-parrot orders.

Key to females of the genus *Metacheyletia* Fain, 1972

1. Propodonal shield weakly developed, but distinct; setae *ag3* absent (hosts – Psittacidae) 3
— Propodonal shield almost indistinct; setae *ag3* present. 2
2. Setae *vi* 2 times shorter than *ve*; setae *d2* 2 times shorter than *d1*; setae *h1* and *h2* subequal. Setae *c1*, *d1*, and *e2* 130, 115–120, and 75–80 long, respectively *M. ngaii* n. sp.
— Setae *vi* 1.1–1.2 times shorter than *ve*; setae *d1* and *d2* subequal; setae *h1* about 1.3 times shorter than *h2*. Setae *c1*, *d1*, and *e2* 65, 60, and 50 long, respectively *M. degenerata* Fain & Bochkov, 2003
3. Remnants of legs IV absent 4
— Remnants of legs IV present. *M. obesa* Fain, 1972
4. Peritremal branches with 7–8 segments each. Propodonal shield devoid of setae; length ratios of dorsal idiosomal setae: *h2*:*e2*, *f2* 2:1 and *h2*:*h1* 1.2–1.6:1, length of setae *c1* subequal to distance between bases of setae *c1*–*d1*. Setae *ve*, *si*, *se*, *c1*, *e2*, *f2* shorter than 80
..... *M. amazonae* Bochkov & OConnor, 2003
— Peritremal branches with 10–11 segments each.

Propodonal shield bearing setae *vi* and *ve*, dorsal idiosomal setae *e2*, *f2*, *h1*, and *h2* subequal in length, ratio between length of seta *c1* and distance between *c1–d1* approximately 2.8:1. Setae *ve*, *si*, *se*, *c1*, *e2*, *f2* longer than 100
.....*M. longisetosa* Atyeo, Kethley & Perez, 1984

ACKNOWLEDGEMENTS

This research was supported by a grant from the Russian Foundation for Basic Research (10-04-00160-a) to AVB. In the frames of this project visit of AVB to the Adam Mickiewicz University, Poznan, Poland was supported by the special grant from administration of this university.

REFERENCES

- Atyeo W.T., Kethley J.B., Perez T.M. 1984 — Paedomorphosis in *Metacheyletia* (Acari: Cheyletidae), with the description of a new species — J. Med. Ent., 21: 125-131.
- Bochkov A.V., Fain A. 2001 — Phylogeny and system of the Cheyletidae (Acari : Prostigmata) with special reference to their host-parasite associations — Bull. Inst. R. Sci. nat. Belg., 71: 5-36.
- Fain A., Bochkov A.V., 2003 — A new species of the genus *Metacheyletia* Fain, 1972 (Acari: Cheyletidae) parasitizing *Serinus mosambicus* (Passeriformes: Fringillidae) in Central Africa — Int. J. Acarol., 29: 119-131.
[doi:10.1080/01647950308683648](https://doi.org/10.1080/01647950308683648)
- Bochkov A.V., OConnor B.M. 2003 — New cheyletid mites (Acari, Cheyletidae) associated with birds — Acta Parasitol., 46: 265-279.
- Clements J.F. 2007 — The Clements checklist of birds of the world. 6th Edition — Ithaca, New York: Cornell University Press. pp. 864.
- Fain A. 1972 — Notes sur les acariens des famines Cheyletidae et Harpyrhynchidae producteurs de gale chez les oiseaux ou les mammiferes — Acta Zool. Pathol. Antwerp., 56: 37-60.
- Fain A. 1980 — Notes on the genera *Samsinakia* Volgin, 1965 and *Metacheyletia* Fain, 1972 (Acari: Cheyletidae) — Int. J. Acarol., 6: 103-108.
[doi:10.1080/01647958008683204](https://doi.org/10.1080/01647958008683204)
- Grandjean F. 1939 — Les segments postlarvaires de l'hysterosoma chez les oribates (Acariens) — Bull. Soc. Zool. Fr., 64: 273-284.
- Grandjean F. 1944. Observations sur les acariens de la famille des Stigmaeidae — Arch. Sci. Phys. Natur., 26: 103-131.
- Kethley J.B. 1990 — Acarina: Prostigmata (Actinedida) — In: Dindal D.L (Ed). Soil Biology Guide. New York: Wiley. p. 667-756.

COPYRIGHT

Bochkov and Skoracki. Acarologia is under free license. This open-access article is distributed under the terms of the Creative Commons-BY-NC-ND which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.