Phytoseiid mites (Acari: Phytoseiidae) of apple and almond trees in regions of western and south-western Iran

B. Asali Fayaz, M. Khanjani, F. Molavi, E.A. Ueckermann

► To cite this version:

B. Asali Fayaz, M. Khanjani, F. Molavi, E.A. Ueckermann. Phytoseiid mites (Acari: Phytoseiidae) of apple and almond trees in regions of western and south-western Iran. *Acarologia*, 2011, 51 (3), pp.371-379. 10.1051/acarologia/20112020 . hal-01600108

HAL Id: hal-01600108

<https://hal.science/hal-01600108>

Submitted on 2 Oct 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives| 4.0 International License

ACAROLOGIA

A quarterly journal of acarology, since 1959
Publishing on all aspects of the Acari

All information:

<http://www1.montpellier.inra.fr/CBGP/acarologia/>
acarologia@supagro.inra.fr

**Acarologia is proudly non-profit,
with no page charges and free open access**

Please help us maintain this system by
encouraging your institutes to subscribe to the print version of the journal
and by sending us your high quality research on the Acari.

Subscriptions: Year 2017 (Volume 57): 380 €

<http://www1.montpellier.inra.fr/CBGP/acarologia/subscribe.php>

Previous volumes (2010-2015): 250 € / year (4 issues)

Acarologia, CBGP, CS 30016, 34988 MONTFERRIER-sur-LEZ Cedex, France

The digitalization of Acarologia papers prior to 2000 was supported by Agropolis Fondation under the reference ID 1500-024 through the « Investissements d'avenir » programme (Labex Agro: ANR-10-LABX-0001-01)

Acarologia is under **free license** and distributed under the terms of the Creative Commons-BY-NC-ND which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

PHYTOSEIID MITES (ACARI: PHYTOSEIIDAE) OF APPLE AND ALMOND TREES IN REGIONS OF WESTERN AND SOUTH-WESTERN IRAN

Bahman ASALI FAYAZ¹, Mohammad KHANJANI^{1*}, Fatemeh MOLAVI¹,
Edward A. UECKERMANN^{2,3}

(Received 28 February 2011; accepted 09 May 2011; published online 23 September 2011)

¹ Department of Plant Protection, College of Agriculture, Bu Ali-Sina University, Hamadan, I. R. Iran.
mkhanjani@gmail.com (*corresponding author); basalifayaz@gmail.com

² ARC-Plant Protection Research Institute, Private Bag X134, Queenswood, Pretoria, 0121 South Africa. UeckermannE@arc.agric.za

³ School of Environmental Sciences and Development, North-West University, Potchefstroom Campus 2520, South Africa

ABSTRACT — Eight phytoseiid species belonging to three genera were collected in apple and almond orchards from the Provinces of Hamedan, Kurdistan and Chahar Mahal va Bakhtiari, in western and south-western Iran. *Typhlodromus (Anthoseius) bagdasarjani* Wainstein and Arutunjan, 1967 was the most abundant species, being collected from both plant species in the three provinces.

KEYWORDS — Acari; predatory mites; Phytoseiidae; apple; almond; Iran

INTRODUCTION

In the western and south-western areas of Iran, apples, *Malus domestica* L. (Rosaceae), are cultivated on more than 2,000 ha, with a total production of over 153,000 t, whereas almonds, *Prunus dulcis* (Mill.) D. A. Webb (Rosaceae), are cultivated on almost 14,000 ha with a total production of almost 13,000 t (Anonymous, 2009). Members of the phytophagous mite families Eriophyidae and Tetranychidae are important pests of those crops in Iranian orchards (Khanjani and Hadad Irani Nejad, 2006). Species of the mite family Phytoseiidae are important predators of eriophyids and tetranychids. More than 20 phytoseiid species are commercially reared for use as biological control agents of mites and small insects harmful to greenhouse and field crops worldwide (Chant and McMurtry, 2007; Gerson *et al.*,

2003). Moraes *et al.* (2004) mentioned 39 phytoseiid species described and recorded from apple trees but none from almonds. Daneshvar and Denmark (1982) described *T. (A.) iranensis* and Ueckermann *et al.* (2009) redescribed *Typhlodromus (Anthoseius) khosrovensis* Arutunjan from apple trees in north-west Iran. In this study, the results of a survey conducted in apple and almond orchards during 2008–2010 in 3 provinces in western and south-western Iran are given.

MATERIALS AND METHODS

This study was carried out in 8 commercial apple orchards in the Provinces of Hamedan and Kurdistan and in 7 commercial almond orchards in the Province of Chahar Mahale va Bakhtiari, in 2008–

2010. Phytoseiid mites were collected by beating or shaking shoots over a white tray or by extracting them from samples of soil and fallen fruits in Berlese funnels. The mites were preserved in 70 % ethanol and then mounted on slides in Hoyer's medium (Walter and Krantz, 2009) for later examination under a Nomarski Interference Contrast Olympus microscope.

All measurements are given in micrometers with the means followed by the respective ranges in parentheses. A key to the species found in this survey is given. The classification systems used follows that of Chant and McMurtry (2007). The dorsal and ventral setal nomenclature is that of Rowell *et al.* (1978) and Chant and Yoshida-Shaul (1991). Idiosomal setal patterns are given according to Chant and Yoshida-Shaul (1992). All specimens are deposited in the Collection of the Acarology Laboratory, University of Bu-Ali Sina, Hamedan, Iran.

RESULTS

Typhlodrominae

Typhlodromus (Anthoseius) bagdasarjani Wainstein and Arutunjan

Typhlodromus bagdasarjani Wainstein and Arutunjan, 1967: 1765 *Typhlodromus (Anthoseius) bagdasarjani*, Moraes *et al.*, 2004: 311

Remarks — Measurements of setae agree with those included in the original description of this species.

Female

Dorsum — Dorsal setal pattern: 12A:8A. Tectum convex, 30 (26 – 33) wide. Dorsal shield reticulated, 362 (355 – 368) long, width at the widest level of podonotal region of the dorsal shield 174 (173 – 175); five pairs of solenostomes. Measurements of dorsal setae as follows: j1 25 (24 – 25), j3 32 (31 – 33), j4 20 (19 – 20), j5 19 (18 – 19), j6 22 (21 – 23), J2 28 (26 – 30), J5 11 (10 – 11), z2 23 (20 – 25), z3 28 (27 – 28), z4 29 (28 – 30), z5 19 (18 – 20), Z4 54 (52 – 55), Z5 69 (65 – 73); s4 32 (28 – 35); s6 38 (35 – 40), S2 40 (35 – 45); S4 37 (33 – 40); S5 30 (27 – 33); r3 31 (28 – 33);

R1 27 (24 – 30). All setae smooth, except Z4 and Z5, serrated.

Venter — Setae set on venter of gnathosoma of the following lengths: h1 23 (22 – 23), h2 21 (20 – 21), h3 18 (16 – 20) and palp coxal seta 26 (25 – 26). Length of tritosternum 90 – 93 (92). Sternal shield smooth and with two pairs of setae (St1 and St2) and two pairs of lyrifissures; setae St3 and St4 each set on separated platelets. Length of sternal setae as follows: St1 33 (29 – 35), St2 31 (28 – 33), St3 31 (28 – 31), St4 30 (28 – 31), St5 27 (23 – 31). Genital shield 113 (110 – 115) long and 75 (72 – 78) at the widest level. Ventrianal shield without pre-anal pores and with a few transverse lines, 110 – 120 long and 80 – 90 wide at level setae ZV2. Setae JV5 50 (48 – 53).

Chelicera — Movable digit 25, toothless; fixed digit 27, with four teeth and a pilus dentilis.

Peritreme — Extending to level between setae z3-4, 106 (100 – 112) long.

Spermatheca — Calyx 14 (11 – 17) long and 10 (9 – 10) wide, bell-shaped.

Legs — Length of legs I-IV as follows: leg I 305 (301 – 308), leg II 264 (261 – 267), leg III 265 (263 – 266) and leg IV 351 (347 – 355). Basitarsus IV with one knobbed macroseta (St) 48 (45 – 49).

Specimens examined — Hamedan Province – Heydareh village (34° 48' N; 48° 28' E), Hamedan (Apr. 2009) (n = 2), Hamedan (Oct. 2009) (n = 1), Ghorveh Dar Jazin (Oct. 2009) (n = 4), Kurdistan Province – Kamyaran vicinity (Nov. 2009) (n = 4), Chenu village, (Nov. 2009) (n = 2). All on apple trees. Chahar Mahal va Bakhtiari Province – Emamieh (32° 32' 11" N; 50° 8' 50" E) (Jul. and Oct. 2009), Hureh (Jul. 2008); Cham Heidar (32° 30' 40" N; 51° 00' 27" E) (Sept. 2009) infested with eriophyid mites; Cham Khoram (32° 28' 01" N; 50° 57' 15" E), (Aug. and Sept. 2009); Shoorab (32° 29' 23" N; 50° 57' 03" E) (Jul. 2009) Cham Ali (32° 30' 40" N; 50° 53' 30" E) (Oct. 2008); Saman (32° 29' 48" N; 51° 00' 27" E) (Jul. 2009). All on almond trees (n = 31).

Typhlodromus (Anthoseius) iranensis (Denmark and Daneshvar)

Amblydromella iranensis Denmark and Daneshvar in Daneshvar and Denmark, 1982: 8 *Typhlodromus (An-*

thoseius) *iranensis*, Moraes *et al.*, 2004: 332

Note — This species is suspected to be junior synonym of *T. (A.) kazachstanicus* Wainstein, 1958 (Faraji *et al.*, 2007).

Remarks — Measurements of setae agree with those included in the original description of this species.

Female

Dorsum — Dorsal setal pattern: 12A:8A. Tectum convex, 32 (30 – 34) wide. Dorsal shield reticulated, 370 (370) long, width at the widest level of podonotal region of the dorsal shield 171 (166 – 175); three pairs of solenostomes. Length of dorsal setae as follows: j1 21 (20 – 21), j3 31 (30 – 31), j4 20 (19 – 20), j5 21 (20 – 21), j6 24 (23 – 24), J2 28 (27 – 28), J5 4 (3 – 4), z2 27 (26 – 27), z3 29 (28 – 30), z4 31 (30 – 31), z5 21 (20 – 21), Z4 48 (46 – 49), Z5 59 (57 – 60); s4 32 (31 – 32); s6 36 (33 – 38), S2 40 (38 – 41); S4 36 (35 – 38); S5 29 (27 – 30); r3 31 (30 – 31); R1 32 (30 – 33). All dorsal setae smooth, except Z4 and Z5, serrated.

Venter — Setae set on venter of gnathosoma of the following lengths: h1 25 (24 – 25), h2 25 (24 – 25), h3 23 (20 – 25) and palp coxal seta 29 (28 – 30). Tritosternum bifid and 94 (92 – 95) long; Sternal shield smooth and with two pairs of setae (St1 and St2) and two pairs of lyrifissures; setae St3 and St4 each set on separated platelets. Length of sternal setae as follows: St1 31 (30 – 32), St2 31 (30 – 31), St3 34 (33 – 35), St4 31 (30 – 31), St5 33 (31 – 34). Genital shield 140 (140) long; 75 (75 – 79) at the widest level. Ventrianal shield with a few transverse lines, 121 (116 – 125) long and 81 (80 – 82) wide at level of setae ZV2; no pre-anal pores. Setae JV5 50 (50).

Chelicera — Movable digit 28 (27 – 28), with one tooth; fixed digit 29 (28 – 30), with three teeth and a pilus dentilis.

Peritreme — Apex of peritreme reaches level between setae s4 and z4.

Spermatheca — Calyx 17 (15 – 18) and 13 (10 – 15) wide, fundibular; atrium nodular.

Legs — Length of legs I–IV as follow: leg I 313 (300 – 325), leg II 260 (260), leg III 274 (267 – 280)

and leg IV 354 (345 – 363). Basitarsus IV with one knobbed macroseta (St) 53 (51 – 54).

Specimens examined — Hamedan Province – Qorveh Dar Jazin, Razan vicinity (35° 21' N; 49° 06' E), (Oct. 2009), (n = 2), on apple trees.

***Typhlodromus (Anthoseius) khosrovensis* Arutunjan**

Typhlodromus khosrovensis Arutunjan, 1971: 306 *Typhlodromus (Anthoseius) khosrovensis*, Moraes *et al.*, 2004 : 333 Redescription: Ueckermann *et al.*, 2009

Remarks — The specimens examined are very similar to the original description of the species.

Female

Dorsum — Dorsal setal pattern: 12A:8A. Tectum convex, 32 (31 – 33) wide. Dorsal shield reticulated, 334 (330 – 338) and width at the widest level of podonotal region of the dorsal shield 158 (155 – 160); four pairs of solenostomes. Length of dorsal setae as follows: j1 22 (21 – 23), j3 31 (30 – 31), j4 19 (18 – 19), j5 20 (19 – 20), j6 24 (22 – 25), J2 26 (25 – 27), J5 4 (3 – 4), z2 24 (23 – 24), z3 30 (29 – 30), z4 29 (29), z5 20 (19 – 20), Z4 41 (39 – 42), Z5 55 (51 – 59); s4 33 (32 – 34); s6 36 (35 – 36), S2 38 (35 – 40); S4 33 (32 – 33); S5 17 (15 – 19); r3 28 (27 – 28); R1 29 (28 – 30). Setae smooth except Z4 and Z5, serrated.

Venter — Setae set on venter of gnathosoma of the following lengths: h1 22 (21 – 22), h2 21 (18 – 23), h3 22 (20 – 23) and palp coxal seta 26 (24 – 27). Tritosternum bifid, 79 (75 – 82) long; Sternal shield smooth with two pairs of setae (St1 and St2) and two pairs of lyrifissures; posterior margin with a median lobe; setae St3 and St4 each set on separated platelets. Length of sternal setae as follows: St1 29 (27 – 30), St2 30 (30), St3 29 (27 – 31), St4 27 (25 – 28), St5 27 (25 – 28). Genital shield 120 (118 – 121) long and 70 (70) wide at level of setae St5. Ventrianal shield with a few transverse lines, 112 (107 – 117) long and 80 (80) wide at level of ZV2, with no pre-anal pores. Setae (JV5) 49 (48 – 50).

Chelicera — Movable cheliceral digit 26 (25 – 26), with one tooth; fixed digit 27 (26 – 27), with four teeth and a pilus dentilis.

Peritreme — Apex of peritreme extending to level of z4.

Spermatheca — Calyx 12 (11 – 13) long and 9 (8 – 9) wide, cup or bell-shaped; atrium incorporated in calyx.

Legs — Length of legs as follows: Leg I 305 (300 – 310), leg II 260 (255 – 265), leg III 257 (255 – 260) and leg IV 234 (230 – 237). Basitarsus IV with a knobbed macroseta (St) 51 (50 – 51).

Specimens examined — Hamedan Province Heidareh village (34° 48' N; 48° 28' E), Hamedan, (Sept. 2009) (n = 3), on apple trees (n = 3).

Paraseiulus talbii (Athias-Henriot)

Typhlodromus talbii Athias-Henriot, 1960: 75 *Paraseiulus talbii*, Abbasova, 1972: 11; Moraes *et al.*, 2004: 301

Remarks — The specimens examined are very similar to the original description of the species.

Female

Dorsum — Dorsal setal pattern: 13A: 9A. Tectum convex, 30 (26 – 33) wide. Dorsal shield reticulated, 381 (367 – 395) long; width at the widest level of podonotal region of the dorsal shield 203 (203); three pairs of solenostomes. Length of dorsal setae as follows: j1 20 (17 – 23), j3 29 (28 – 30), j4 27 (26 – 27), j5 29 (28 – 29), j6 35 (33 – 37), J2 38 (37 – 38), J5 12 (12), z2 32 (30 – 33), z3 34 (32 – 36), z4 36 (36), z5 28 (27 – 28), z6 31 (31), Z3 39 (38 – 40), Z4 43 (42 – 43), Z5 45 (44 – 45), s4 44 (43 – 45); s6 35 (34 – 35), S2 41 (40 – 42); S4 36 (35 – 36), S5 40 (39 – 40); r3 32 (30 – 34); R1 34 (33 – 35). All dorsal setae smooth, except Z4 and Z5, serrated.

Venter — Setae set on venter of gnathosoma of the following lengths: h1 20 (20), h2 21 (20 – 21), h3 16 (15 – 16) and palp coxal seta 26 (25 – 26). Tritosternum bifid and 79 (79) long; Sternal shield smooth and with two pairs of setae (St1 and St2) and 2 pairs of lyrifissures; setae St3 set on unsclerotized cuticle and setae St4 setae on platelets. Length of sternal setae St1 29 (28 – 30), St2 27 (26 – 28), St3 29 (28 – 30), St4 29 (28 – 29), St5 29 (28 – 29). Genital shield 150 (150) long and 68 (68) wide at level of

setae St5. Ventrianal shield with a few transverse lines, 140 (140) long, and 62 (62) at level of anal opening, with no pre-anal pores and two pairs of pre-anal setae JV1 26 (25 – 26), ZV2 28 (27 – 28), one pair of para-anal seta 20 (20), post anal setae 17 (17). Seta JV5 47 (45 – 48).

Peritreme — Apex of peritreme extending to level between setae j3-z2.

Spermatheca — Calyx 16 (15 – 17) long, 10 (8 – 11) wide and saccular.

Legs — Length of legs I-IV as follows: leg I 353 (350 – 355), leg II 273 (270 – 275), leg III 281 (279 – 282) and leg IV 373 (365 – 380). Basitarsus IV with one macroseta (St) 31 (30 – 31).

Specimens examined — Chahar Mahal va Bakhtiari Province – Cham Khoram (32° 28' 01" N; 50° 57' 15" E) (Agu. 2009), Cham Heidar (32° 30' 40" N; 51° 00' 27" E) (Sept. 2009) and Saman (Jul. 2008). All seven specimens were collected from foliage of almond trees infested with *Tetranychus urticae* Koch and eriophyid mites.

AMBLYSEIINAE

Neoseiulus agrestis (Karg)

Typhlodromus agrestis Karg, 1960: 449 *Neoseiulus agrestis*, Moraes *et al.*, 2004: 98

Remarks – The specimens examined are very similar to the original description of the species.

Female

Dorsum — Dorsal setal pattern: 10A: 9B. Tectum convex, 40 (40) wide. Dorsal shield reticulated and 368 (360 – 375) long, width at the widest level of podonotal region of the dorsal shield 193 (190 – 195); four pairs of solenostomes. Measurements of dorsal setae as follows: j1 14 (12 – 15); j3 20 (19 – 21); j4 18 (17 – 18); j5 18 (17 – 18); j6 21 (20 – 22), J2 24 (23 – 25); J5 13 (12 – 13); z2 18 (17 – 18); z4 21 (20 – 22); z5 18 (17 – 18); Z1 21 (18 – 23), Z4 25 (24 – 26), Z5 37 (34 – 40); s4 24 (23 – 25); S2 23 (22 – 23); S4 26 (24 – 27); S5 21 (21); r3 21 (20 – 21); R1 21 (20 – 21). All dorsal setae smooth.

Venter — Setae set on venter of gnathosoma of the following lengths: h1 25 (24 – 25), h2 24 (23 – 24),

h3 25 (24 – 25) and palp coxa setae 26 (25 – 27). Tritosternum bifid and 89 (88 – 90) long; Sternal shield smooth and with three pairs of setae (St1, St2 and St3) and two pairs of lyrifissures. Lengths of sternal setae as follows: St1 28 (27 – 28), St2 25 (24 – 25), St3 23 (22 – 24), St4 26 (25 – 26), St5 24 (22 – 26). Sternal setae with two pairs of lyrifissures. Genital shield 123 (120 – 125) long; 74 (70 – 78) wide at level of setae St5. Ventrianal shield with a few transverse lines, 130 (128 – 131) long, and 99 (98 – 100) wide at level of setae ZV2, with a pair of pre-anal pores posteromedially to seta JV2. Setae JV5 34 (33 – 35).

Chelicera — Movable digit 30, with one tooth; fixed digit 34 (32 – 35) three teeth and a pilus dentilis.

Peritreme — Apex of peritreme extending to level seta j1.

Spermatheca — Calyx 29 (27 – 30) long and 13 (12 – 13) wide, tubular and flaring towards vesicle.

Legs — Length of legs I–IV as follows: leg I 355 (350 – 360), leg II 259 (258 – 260), leg III 244 (238 – 250) and leg IV 253 (345 – 360). Basitarsus IV with one pointed macroseta (St) 51 (50 – 51).

Specimens examined — Hamedan Province Moejin village, Marianaj vicinity, (Apr. 2010) (n = 2), on apple trees.

Neoseiulus bicaudus (Wainstein)

Amblyseius bicaudus Wainstein, 1962: 146 *Neoseiulus bicaudus*, Congdon, 2002: 23; Moraes *et al.*, 2004: 108

Remarks — The specimens examined are very similar to the original description of the species. However, 7 pairs of solenostomes are observed on the dorsal shield of the specimens examined but only 6 pairs are mentioned in the original description.

Female

Dorsum — Dorsal setal pattern: 10A: 9B. Tectum convex, 44 (43 – 44) wide. Dorsal shield reticulated, 404 (398 – 410) long; width at the widest level of podonotal region of the dorsal shield 193 (185 – 200) at level of setae R1; seven pairs of solenostomes. Length of setae as follows: j1 22 (21 – 23), j3 32 (30

– 33), j4 16 (14 – 17), j5 16 (15 – 16), j6 18 (16 – 20), J2 17 (16 – 18), J5 14 (13 – 15), z2 27 (25 – 30), z4 21 (21 – 22), z5 16 (16), Z1 22 (20 – 24), Z4 41 (40 – 42), Z5 85 (80 – 89), s4 34 (33 – 35), S2 35 (34 – 36), S4 37 (36 – 38), S5 42 (40 – 44); r3 33 (30 – 35); R1 30 (29 – 31), both of setae r3 and R1 on soft integument. All dorsal setae smooth except S4, S5, Z4 and Z5, serrated.

Venter — Setae set on venter of gnathosoma of the following lengths: h1 22 (21 – 23), h2 21 (20 – 22), h3 23 (21 – 25) and palp coxal seta 27 (26 – 27). Length of tritosternum 93 (92 – 94). Sternal shield reticulated, bears three pairs of setae (St1, St2 and St3) and two pairs of lyrifissures. Length of sternal setae St1 27 (24 – 30), St2 25 (23 – 26), St3 24 (22 – 25), St4 28 (26 – 30), St5 23 (20 – 25). Seta St4 set on platelet. Genital shield 124 (123 – 125) long and 81 (75 – 87) at level of St5. Ventrianal shield reticulated, 135 (130 – 140) long, 106 (105 – 107) wide at level of setae ZV2, one pair of pre-anal pores posterior to JV2 and distance between their bases 41 (41), three pairs of pre-anal setae JV1 16 (14 – 18), ZV2 21 (20 – 21) and JV2 17 (15 – 19), one pair of para-anal setae 18 (17 – 18), post anal seta 21 (20 – 21). Seta JV5 54 (53 – 55), serrated.

Chelicera — Movable digit 33 (30 – 35), with one tooth; fixed digit 37 (35 – 40) long, with six teeth and a pilus dentilis.

Peritreme — Apex of peritreme extending to level between setae j1 and j3.

Spermatheca — Calyx 11 (9 – 12) long and 9 (7 – 10) wide, bowl-shaped with short neck between the latter and the nodular atrium.

Legs — Length of legs I–IV as follows: leg I 350 (340 – 360), leg II 271 (263 – 280), leg III 277 (265 – 290) and leg IV 355 (345 – 360). Basitarsus IV with a pointed macroseta, 75 (71 – 78) long.

Specimens examined — Chahar Mahal va Bakhtiari province – Cham Khoram (32° 28' 01" N 50° 57' 15" E) (Jun. 2010) (n = 2), Saman (32° 30' 40" N 51° 00' 27" E) (Oct. 2008) (n = 1), Cham Heidar (32° 30' 0" N 50° 53' 51" E) (Aug. 2009) (n = 2). All on almond trees.

***Neoseiulus sugonjaevi*
(Wainstein and Abbasova)**

Amblyseius sugonjaevi Wainstein and Abbasova, 1974: 796 *Neoseiulus sugonjaevi*, Moraes *et al.*, 2004: 145

Remarks — The specimens examined are very similar to the original description of the species.

Female

Dorsum — Dorsal setal pattern: 10A: 9B. Tectum convex, 38 (36 – 40) wide. Dorsal shield reticulated, 359 (355 – 363) long; width at the widest level of podonotal region of the dorsal shield 184 (170 – 198); three pairs of solenostomes. Length of setae as follows: j1 20 (19 – 20), j3 26 (25 – 27), j4 20 (20), j5 20 (20), j6 20 (19 – 20), J2 26 (25 – 27), J5 12 (11 – 12), z2 23 (22 – 23), z4 26 (26), z5 21 (20 – 21), Z1 25 (25), Z4 32 (31 – 32), Z5 43 (42 – 43); s4 28 (28); S2 26 (25 – 27); S4 26 (25 – 26); S5 23 (21 – 25); r3 21 (21); R1 22 (21 – 22). All dorsal setae smooth, except Z5, serrated.

Venter — Setae set on venter of gnathosoma of the following lengths: h1 23 (21 – 25), h2 23 (20 – 25), h3 24 (22 – 25) and palp coxal seta 26 (25 – 26). Tri-tosternum 88 (86 – 90) long; Sternal shield smooth and with three pairs of setae (St1, St2 and St3) and two pairs of lyrifissures. Length of sternal setae St1 27 (26 – 28), St2 26 (24 – 27), St3 26 (26), St4 26 (24 – 27), St5 25 (23 – 27). Genital shield 113 (112 – 115) long and 75 (74 – 75) wide at level of setae St5. Ventrianal shield with longitudinal lines, 123 (115 – 130) long and 103 (100 – 106) wide at level of setae ZV2; with a pair of pre-anal pores at posterior of setae JV2. Seta JV5 39 (37 – 40).

Chelicera — Movable digit 31 (30 – 33), with one tooth; fixed digit 35, with four teeth and a pilus dentilis.

Peritreme — Apex of peritreme extending to level of seta j1.

Spermatheca — Calyx 22 (20 – 23) long and 9 (7 – 10) wide, distal half of calyx slightly bulged and proximal half flared.

Legs — Length of legs I-IV as follows: leg I 335 (325 – 345), leg II 248 (240 – 255), leg III 250 (240 – 260) and leg IV 351 (341 – 360). Basitarsus IV with

one pointed macroseta (St) 59 (56 – 62) and genu IV with one pointed macroseta (Sge) 25 (25).

Specimens examined — Hamedan Province – Pole Shekasteh region, between Toyserkan and Asad Abad vicinities, (Oct. 2009) (n = 2). Kurdistan Province – Naran village, Sanandaj, (Nov. 2009) (n = 1) and Chenu village, Sanandaj, (Nov. 2009) (n = 1). All from soil and litter under apple tree.

***Neoseiulus zwoelferi* (Dosse)**

Typhlodromus zwoelferi Dosse, 1957: 301 *Neoseiulus zwoelferi*, Moraes *et al.*, 2004: 152.

Remarks — The specimens examined are very similar to the original description of the species.

Female

Dorsum — Dorsal setal pattern: 10A: 9B. Tectum convex, 43 (40 – 45) wide. Dorsal shield reticulated 408 (400 – 415) long, width at the widest level of podonotal region of the dorsal shield 209 (207 – 210); seven pairs of solenostomes. Length of dorsal setae as follows: j1 20 (19 – 20), j3 26 (25 – 26), j4 18 (18), j5 18 (18), j6 21 (21), J2 24 (22 – 25), J5 11 (10 – 11), z2 25 (24 – 25), z4 21 (20 – 21), z5 19 (19), Z1 23 (22 – 23), Z4 35 (33 – 36), Z5 50 (49 – 50); s4 31 (30 – 32); S2 28 (26 – 30); S4 25 (24 – 25); S5 24 (24); r3 24 (24); R1 25 (25). All dorsal setae smooth, except Z5, serrated.

Venter — Setae set on venter of gnathosoma of the following lengths: h1 26 (25 – 26), h2 26 (25 – 27), h3 26 (24 – 27) and palp coxal seta 30 (29 – 30). Tri-tosternum with two lacinia, 97 (95 – 98) long; Sternal shield smooth and with three pairs of setae (St1, St2 and St3) and two pairs of lyrifissures; setae St4 set on platelets. Length of sternal setae St1 35 (35), St2 30 (30), St3 30 (29 – 31), St4 33 (33), St5 30 (28 – 31). Genital shield 127 (120 – 135) long and 83 (80 – 85) wide at level of setae St5. Ventrianal shield with a few transverse lines, 141 (138 – 143) long and 116 (112 – 120) wide at level of setae ZV2; one pair of pre-anal pores posterior to setae JV2. Setae JV5 47 (43 – 50).

Chelicera — Movable digit 32 (31 – 33) long, toothless, fixed digit 35, with four teeth and a pilus dentilis.

FIGURE 1: Relative abundance of specimens of phytoseiid mites from apple and almond orchards in the Hamedan, Kurdistan and Chahar Mahale va Bakhtiari Provinces of Iran, based on samples collected in 2008-2010.

Peritreme — Apex of peritreme extending to near seta j1.

Spermatheca — Calyx 22 (21 – 23) long and 14 (12 – 15) wide, with a slender neck that flares towards vesicle.

Legs — Length of legs I-IV as follows: leg I 299 (286 – 290), leg II 299 (295 – 303), leg III 296 (292 – 300) and leg IV 418 (415 – 420). Basitarsus IV with one pointed macroseta (St) 71 (66 – 75); genu IV also with one pointed macroseta (Sge) 31 (30 – 31).

Specimens examined — Hamedan Province – Heydareh village (34° 48' N; 48° 28' E), Hamedan, (Nov. 2009) (n = 1), from soil and litter under apple trees. Kurdistan Province – Chenu village, Sanandaj, (Nov. 2009) (n = 4), from apple trees (n = 5).

Key to phytoseiid species found in this survey (females).

1. Setae z3 and s6 absent (*Amblyseiinae-Neoseiulus*)... 2
— Both setae z3 and s6 present (*Typhlodrominae*)... 5
2. Seta Z5 longer than 80 μ m... *N. bicaudus* Wainstein, 1962
— Seta Z5 shorter than 80 μ m... 3

3. Dorsal shield with 7 pairs of solenostomes... *N. zwoelferi* Dosse, 1957
— Dorsal shield with 3-4 pairs of solenostomes... 4

4. Dorsal shield with 3 pairs of solenostomes... *N. sugonjaevi* Wainstein and Abbasova, 1974
— Dorsal shield with 4 pairs solenostomes... *N. agrestis* Karg, 1960

5. Dorsal setae z6 and Z3 present... *Paraseiulus talbii* Athias-Henriot, 1960
— Dorsal setae z6 and Z3 absent... 6

6. Movable digit of chelicera without tooth and 5 pairs of solenostomes on dorsal shield... *T.(A.) bagdasarjani* Wainstein and Arutunjan, 1967
— Movable digit of chelicera with 1 tooth and 3-4 pairs of solenostomes on dorsal shield... 7

7. Dorsal shield with 3 pairs of solenostomes and $Z4/S5 \approx 1.6$... *T.(A.) iraniensis* Denmark and Daneshvar, 1982
— Dorsal shield with 4 pairs of solenostomes and $Z4/S5 \approx 2.4$... *T.(A.) khosrovensis* Arutunjan, 1971

DISCUSSION

This survey showed that *Typhlodromus (Anthoseius) bagdasarjani* was the dominant of the eight species collected in apple and almond orchards in the three provinces (Figure 1). The following step will be to evaluate these predators specially the dominant species for its effectiveness against important pest mite species of the families Tetranychidae and Eriophyidae in Iran. The aim is to reduce chemical control.

ACKNOWLEDGEMENTS

We wish to thank Prof G. J. De Moraes (Depto. Entomologia, Fitopatologia e Zoologia Agrícola, Universidade de São Paulo/ Escola Superior de Agricultura "Luiz de Queiroz", 13418-900 Piracicaba-SP, Brazil) for supplying us with literature.

REFERENCES

- Abbasova E.D. 1972 — Phytoseiid mites (Parasitiformes: Phytoseiidae) of Azerbaijan — Avtoreferat Dissertatsii na Soiskanie Uchenoy Stepeni Kandidata Biologicheskikh Nauk - Akademiya Nauk Azerbaydzhanskoy SSR, Institut Zoologii, Baku, Azerbaijan, 34 pp. [in Russian].
- Anonymous. 2009 — The result of sampling design of Horticultural productions of Iran — Ministry of Agricultural Jihad: Publisher. pp. 95. [In Persian]
- Arutunjan E.S. 1971 — New species of the genus *Typhlodromus* Scheuten, 1857 (Parasitiformes, Phytoseiidae) — Dokl. Akad. Nauk. Armyanskoy SSR, Armenia, 52: 305-308 [in Russian].
- Athias-Henriot C. 1960 — Phytoseiidae et Aceosejidae (Acarina: Gamasina) d' Algérie. IV. Genre *Typhlodromus* Scheuten, 1857 — Bull. Soc. Hist. Nat. Afrique du Nord, 51: 62-107.
- Chant D.A., McMurtry J.A. 2007 — Illustrated keys and diagnoses for the genera and subgenera of the Phytoseiidae of the world (Acari: Mesostigmata) — Indira Publishing House, West Bloomfield, Michigan, USA. pp. 220.
- Chant D.A., Yoshida-Shaul E. 1991 — Adult ventral setal patterns in the family Phytoseiidae (Acari: Gamasina) — Int. J. Acarol., USA, 17: 187-199.
- Chant D.A., Yoshida-Shaul E. 1992 — Adult idiosomal setal patterns in the family Phytoseiidae (Acari: Gamasina) — Int. J. Acarol., 18: 177-193.
- Congdon B.D. 2002 — The family Phytoseiidae (Acari) in western Washington State with descriptions of three new species — Int. J. Acarol., USA, 28(1): 3-27.
- Daneshvar H., Denmark H.A. 1982 — Phytoseiids of Iran (Acarina: Phytoseiidae) — Int. J. Acarol., USA, 8: 3-14.
- Dosse G. 1957 — Morphologie und biologie von *Typhlodromus zwoelferi* n. sp. (Acar., Phytoseiidae) — Z. Angew. Entomol., Germany, 41(2-3): 301-311.
- Faraji F., Hajizadeh J., Ueckermann E.A. Kamali, K., McMurtry J.A. 2007 — Two new records for Iranian phytoseiid mites with synonymy and keys to the species of *Typhloseiulus* Chant and McMurtry and Phytoseiidae in Iran (Acari: Mesostigmata) — Int. J. Acarol., 33: 231-239.
- Gerson U., Smiley R. L., Ochoa R. 2003 — Mites (Acari) for Pest Control — Blackwell Science Ltd: Publisher, UK. pp. 539.
- Karg W. 1960 — Zur Kenntnis der Typhlodromiden (Acarina: Parasitiformes) aus Ackerund Grünlandboden — Z. Angew. Entomol., Germany, 47: 440-452.
- Khanjani M., Hadad Irani Nejad K. 2006 — Injurious mites of Agricultural crops in Iran — Bu-Ali Sina University: Publisher. pp. 515. [In persian].
- Moraes G.J. De., McMurtry J.A., Denmark H., Campos C.B. 2004 — A revised catalog of the mite family Phytoseiidae — Zootaxa. 434:1-494.
- Rowell H.J., Chant D.A., Hansell R.I.C. 1978 — The determination of setal homologies and setal patterns on the dorsal shield in the family Phytoseiidae (Acarina: Mesostigmata) — Can. Entomol., 110: 859-876.
- Ueckermann E.A., Jalaieian M., Saboori A., Seyedoslami H. 2009 — Re-description of *Typhlodromus (Anthoseius) khosrovensis* Arutunjan, first record for Iran (Acari: Phytoseiidae) — Acarologia, XLIX, 1-2: 23-27.
- Wainstein B.A. 1958 — New species of mites of the genus *Typhlodromus* (Parasitiformes: Phytoseiidae) from Georgia — Soobshcheniya Akad. Nauk Gruzinskoy SSR, 21(2): 201-207 [in Russian].
- Wainstein B.A. 1962 — Some new predatory mites of the family Phytoseiidae (Parasitiformes) of the USSR fauna — Entomologicheskoe Obozrenie, Russia, 41, 230-240 [in Russian]; Entomol. Rev., Russia, 41: 139-146 [English translation].
- Wainstein B.A., Abbasova E. D. 1974 — Two new species of the genus *Amblyseius* (Parasitiformes: Phytoseiidae) from Azerbaijan [in Russian] — Zool. Zh., Russia, 53: 796-798.
- Wainstein B.A., Arutunjan E.S. 1967 — New species of predaceous mites of the genera *Typhlodromus* Scheuten and *Paraseiulus* Muma (Parasitiformes, Phytoseiidae) [in Russian] — Zool. Zh., Russia, 46: 1764-1770.

Walter D., Krantz G. W. 2009 — Collection, rearing and preparing specimens. *In*: Krantz G.W., Walter D. [Eds.]. A Manual of Acarology. 3rd ed. — Oregon State University Book Stores, Inc., Corvallis, Oregon. 83-96.

COPYRIGHT

Asali Fayaz et al.. Acarologia is under free license. This open-access article is distributed under the terms of the Creative Commons-BY-NC-ND which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.