Effects of a high FODMAPS diet on visceral sensitivity: Involvement of advanced glycation end products and colonic mast cells

Jasper Kamphuis, Pauline Riviere, Sophie Yvon, Valérie Tondereau, Hélène Eutamène, Vassilia Theodorou

► To cite this version:

Jasper Kamphuis, Pauline Riviere, Sophie Yvon, Valérie Tondereau, Hélène Eutamène, et al.. Effects of a high FODMAPS diet on visceral sensitivity: Involvement of advanced glycation end products and colonic mast cells. 2. Federation of Neurogastroenterology and Motility Meeting, Aug 2016, San Francisco, California, United States. 116 p., 10.1111/nmo.12881 . hal-01602901

HAL Id: hal-01602901

<https://hal.science/hal-01602901>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution| 4.0 International License

291

Intestinal dysfunctions induced by intrauterine growth retardation are associated with altered autophagy in the enteric nervous system

E. OLEYNIKOVA*, P. AUBERT*, H. RIZVI*, C. LE BERRE-SCOUL*, P. A. GROHARD*, J. CHEVALIER*, J.-P. SEGAIN†, G. LE DREAN†, M. NEUNLIST* and H. BOUDIN*

*INSERM UMR913, Institut des Maladies de l'Appareil Digestif, University of Nantes, Nantes, France and †Inra Phan, Institut des Maladies de l'Appareil Digestif, University of Nantes, Nantes, France

Objective: The intra uterine growth retardation (IUGR) is characterized by a low birth body weight (<2.5 kg) and represents 4% of births. IUGR has long term consequences on health in adulthood, with an elevated risk to develop metabolic diseases and hypertension. In this study, we examined whether IUGR alters gastrointestinal (GI) and enteric nervous system (ENS) functions during life.

Methods: Pregnant rat dams were fed with a normal protein diet (20% protein; control group) or with an isocaloric low protein diet (8% protein; IUGR group) until weaning. The study was performed on the progeny at post-natal day 35 (P35). Digestive functions (motility and permeability) and their response to a stress, the water avoidance stress (WAS), were analyzed *in vivo* and *ex vivo*. Changes in ENS phenotype were evaluated in whole mounts and in primary culture of ENS from control or IUGR group.

Results: No difference was observed between control and IUGR rats for motility and permeability under basal conditions both *in vivo* and *ex vivo*. By contrast, following WAS, paracellular permeability in the distal colon was increased in control but not in IUGR group. Similarly, following WAS, fecal pellet output was increased in control but not in IUGR group. Analysis of ENS revealed a reduction of nitrergic neurons in the ileum of RCIU rats compared to control animals. In addition, autophagy, a key cellular pathway during stress condition, was decreased in the nitrergic, but not cholinergic, neurons in the distal colon of IUGR rats. Moreover, using a coculture system of enteric neurons and glia, we found that IUGR neurons were less resistant than control neurons to cell death induced by glial deprivation. In this model, pharmacological blockade of autophagy indicated that survival of IUGR neurons critically required autophagy while control neurons did not.

Conclusions: Our study suggests that early life stress, such as IUGR, can alter the capacity of the GI tract to respond to an environmental stress later in life, in part via alterations of ENS functions. These ENS dysfunctions might involve impaired autophagic pathways. Funding: INSERM, Region Pays de la Loire (Parimad), Fondation LCL, Fondation SantéDige.

292

Diagnostic yield of 24 h esophageal manometry for non-cardiac chest pain

M. BARRET, T. V. K. HERREGODS, J. M. OORS, A. J. P. M. SMOUT and A. J. BREDENOORD

Department of Gastroenterology and Hepatology, Academic Medical Center, The Netherlands

Background and aim: Non-cardiac chest pain (NCCP) can be caused by reflux or esophageal motor disorders. With the diagnostic improvements brought by pH-impedance monitoring and high-resolution manometry

(HRM), the contribution of ambulatory 24 h manometry in the setting of NCCP is uncertain. Our aim was to assess the diagnostic yield of ambulatory 24 h manometry in this patient population.

Methods: Twenty-four-hour ambulatory pressure-pH-impedance monitoring off proton pump inhibitors was used to study all patients who were referred to our tertiary hospital for NCCP between 2011 and 2015. Patients had a preliminary work-up including cardiological assessment, upper gastro-intestinal endoscopy and esophageal HRM. Diagnostic measurements were independently analyzed by two physicians, and in ambiguous cases a consensus had to be reached. Symptom association probability (SAP) was calculated for both reflux and esophageal spasm using a 2-min time window.

Results: Of the 334 patients selected, 59 (24 males) met the inclusion criteria. Retrosternal pain was present in all patients. However, heartburn, regurgitation, dysphagia, nausea or vomiting, belching and weight loss were also reported, by 9 (15.3%), 15 (25.4%), 14 (23.7%), 2 (3.4%), 4 (6.8%) and 9 (15.3%) patients, respectively. Nine (15.2%) patients had signs of gastroesophageal reflux on upper gastrointestinal endoscopy. Twenty-nine (49.1%) patients had a final diagnosis of functional chest pain, 22 (37%) of gastroesophageal reflux disease, 4 (6.8%) of esophageal spasm, and 4 (6.8%) had other diagnoses. Esophageal spasm was diagnosed in all patients using ambulatory manometry. HRM, using the Chicago Classification v3.0 criteria alone, did not identify any of these patients. However, taking into account suggestive abnormalities on HRM, such as rapid or repetitive contractions, HRM had a sensitivity of 75% and a specificity of 94.5% for the diagnosis of distal esophageal spasms. Eighteen (30.5%) patients did not experience chest pain during the 24 h recording.

Conclusion: Ambulatory 24 h manometry has a low diagnostic yield for esophageal hypertensive peristaltic disorders in non-cardiac chest pain patients. However, it remains the best currently available diagnostic modality. This is particularly relevant if a definitive diagnosis is needed before invasive treatment.

293

Pain modulation in youth with functional gastrointestinal disorders (FGID) may be normal: a role for baseline norepinephrine?

G. CHELIMSKY*,†, C. HILLARD*, C. WELZIG*, S. MARCHAND‡, L. CONANT* and T. CHELIMSKY*,†

*Medical College of Wisconsin, Milwaukee, WI, USA, †Children's Hospitals of Wisconsin, Milwaukee, WI, USA and ‡Université de Sherbrooke, Sherbrooke, QC, Canada

Background: Children with FGID report pain complaints outside the gastrointestinal tract, suggesting abnormal Diffuse Noxious Inhibitory Control (DNIC). Catecholamines impact pain modulation in animals and are a logical target of investigation in humans.

Hypothesis: DNIC is reduced in children with FGID and correlates with functional disability. The DNIC response reflects circulating norepinephrine levels.

Methods: In this prospective IRB approved study, we compared DNIC (Chalaye *et al.*) in ascending and descending immersion of the arm up to the shoulder in 12 °C cold water, with a 45 min rest in between. Each of 4 segments (fingers, wrist, forearm and shoulder) was immersed for 2 min in cold water with a rest period of 5 min between segments, and pain reported by the subject on a 10-point numeric rating scale every 15 s

during the immersion. DNIC is calculated by subtracting finger and wrist immersion during the ascending period (DNIC not activated) from the same report during the descending period (full DNIC activation). Blood was obtained through an IV at baseline, at end of ascending, prior to descending and after descending is complete. Functional Disability Inventory (FDI) measured functional disability.

Results: We enrolled 11 subjects with FGID (1 male; mean age 15.1 years, range 13–17 years). FDI ranged from 4 to 28 with a median of 22 points [±SEM? N=?]. DNIC could not be calculated in 2 subjects because they reported no significant pain (<2 in NRS) when immersing their hand in cold water. The other 10 FGID subjects had a normal DNIC activation of a median of 3.1 points with a range of 2.8–10. Catecholamines and present DNIC were available in 7 subjects. In those subjects, DNIC activation correlated highly with baseline norepinephrine value ($r = 0.97$; $p < 0.01$). The FDI did not correlate norepinephrine ($r = 0.36$; $p = 0.42$).

Conclusions: DNIC is preserved in youth with FGID. Baseline norepinephrine correlated with DNIC activation, suggesting that norepinephrine levels directly regulate pain modulation.

Supported by a Digestive Disease Center grant and Advancing Healthier Wisconsin 5520298.

294

Low bioenergetics in functional disorders parallel disability score

T. CHELIMSKY*,†, P. SIMPSON*,†, L. ZHANG*,†, J. BANDA*,†, D. DIMOCK*,†, S. KOMAS* and G. CHELIMSKY*,†

*Medical College of Wisconsin, Milwaukee, WI, USA and †Children's Hospitals of Wisconsin, Milwaukee, WI, USA

Background: We hypothesized that patients with functional disorders (FD e.g. chronic migraine, functional gastrointestinal disorders, chronic fatigue syndrome, etc.,) show impaired mitochondrial bioenergetics.

Methods: We compared blood from youth with an FD with carefully screened healthy controls (HC) with no functional disorder or chronic medical condition. The Oxygen Consumption Rate and Extracellular Acidification Rate (ECAR) measurements utilized the Seahorse XF96 Extracellular Flux Analyzer (North Billerica, MA), in unbuffered/serum free RPMI assay media supplemented with 1 mM pyruvate. Peripheral Blood Mononuclear Cells (PBMCs) were seeded in a PS V7 cell culture plate at a density of 3.25×10^5 cells per well, then placed in a non-CO₂ incubator for 1 h. Oligomycin (1 µg/mL), Carbonyl cyanide-4-phenyl-hydrazone (FCCP; 1 µM) and Antimycin A (10 µM) determined the mitochondrial parameters. A Mann-Whitney test compared skewed variables, and Fisher's exact test dichotomous variables. Regression tree tested predictors (among age, gender, basal, ECAR and SRC) of functional outcome (Functional Disability Inventory, FDI) optimized by least absolute deviation and 10% leave out samples for cross validation.

Results: 45 subjects (36 female) with a median (range) age of 16 years (10, 20), did not differ by age or gender between 15 FD and 30 HC FD subjects. FD subjects demonstrated lower resting mitochondrial function (basal respiration BR: FD 33.7 [13.3, 96.8] pmol/min units for all values, HC 56.0 [27.1, 171.1] $p = 0.002$), and lower reserve energy (spare respiratory capacity – SRC: FD 68.4 [5.2, 264.7]; HC 118.0 [32.6, 377.3] $p = 0.016$). BR correlated with SRC ($p < 0.0001$). Interestingly, of all