Genetic structure of *Hylesia metabus* (Lepidoptera: Saturniidae), moths responsible for lepidopterism outbreaks in French Guiana and Venezuela.

Marina Ciminera, Alain Roques, Niklas Tysklind, Marie-Anne Auger-Rozenberg

► To cite this version:

Marina Ciminera, Alain Roques, Niklas Tysklind, Marie-Anne Auger-Rozenberg. Genetic structure of *Hylesia metabus* (Lepidoptera: Saturniidae), moths responsible for lepidopterism outbreaks in French Guiana and Venezuela.. ICE 2016; XXV International Congress of Entomology, Sep 2016, Orlando, United States. 2016, ICE 2016; XXV International Congress of Entomology. Program book. 10.1603/ICE.2016.113657 . hal-01607289

HAL Id: hal-01607289

<https://hal.science/hal-01607289>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution - ShareAlike| 4.0 International License

Genetic structure of *Hylesia metabus* (Lepidoptera: Saturniidae): Moths responsible for lepidopteran outbreaks in French Guiana and Venezuela

Marina Ciminera , National Center for Scientific Research, Kourou, French Guiana

Alain Roques , UR633 Zoologie Forestière, INRA, Orléans, France

Niklas Tysklind , INRA, Kourou, French Guiana

Marie-Anne Auger-Rozenberg , UR633 Zoologie Forestière, INRA, Orléans, France

doi: 10.1603/ICE.2016.113657

Introduction: The genus *Hylesia* Hübner (Lepidoptera Saturniidae) involves moths of human health importance in South America, inducing epidemic outbreaks of lepidopterism, a puriginous dermatitis caused by the urticating properties of the females' abdominal hairs. The genus includes about 110 species in South America, several species having been clearly identified as responsible for health problems. The species *H. metabus* is considered as widespread at low elevation in South America. Furthermore, outbreaks of *H. metabus* have been reported in Atlantic coastal countries such as Venezuela, Guyana, Suriname and French Guiana since the last century. In spite of their sanitary and economics importance these moths have been poorly studied, and their ecology and systematics remain unclear.

Methods: In order to evaluate genetic differentiation among populations of *Hylesia metabus*, specific microsatellite primers have been developed on moths sampled during outbreak events. Moths from 20 sampling locations throughout the outbreak areas in French Guiana and Venezuela were analyzed at 7 microsatellite loci.

Results/Conclusion: We found significant genetic population structure as evidenced by Bayesian-based and Multivariate analysis. Our data support the hypothesis of multiple highly differentiated populations . Interestingly only few of them seem to be involved in lepidopterism events occurring in French Guiana and in Venezuela. These findings emphasize the utility of molecular approaches for identifying population structure and connectivity. Such approaches constitute useful tools for understanding outbreak patterns.