

# On the domain of implicit functions in a projective limit setting without additional norm estimates

Jean-Pierre Magnot

► To cite this version:

Jean-Pierre Magnot. On the domain of implicit functions in a projective limit setting without additional norm estimates. *Demonstratio Mathematica*, De Gruyter, 2020, 53 (1), pp.112-120. hal-01626434

HAL Id: hal-01626434

<https://hal.archives-ouvertes.fr/hal-01626434>

Submitted on 30 Oct 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# ON THE DOMAIN OF IMPLICIT FUNCTIONS IN A PROJECTIVE LIMIT SETTING WITHOUT ADDITIONAL NORM ESTIMATES

JEAN-PIERRE MAGNOT

ABSTRACT. We examine how implicit functions on ILB-Fréchet spaces can be obtained without metric or norm estimates which are classically assumed. We obtain implicit functions defined on a domain  $D$  which is not necessarily open, but which contains the unit open ball of a Banach space. The corresponding inverse functions theorem is obtained, and we finish with an open question on the adequate (generalized) notion of differentiation, needed for the corresponding version of the Fröbenius theorem.

*Keywords: implicit functions; ILB spaces*  
*MSC (2010): 58C15*

## INTRODUCTION

Classical inverse functions theorems, implicit functions theorems and Fröbenius theorems on Banach spaces are known to be equivalent. There exists numerous extensions to setting on Fréchet or locally convex spaces, and to our knowledge almost all proofs are based on a contraction principle. In order to obtain in the proofs a mapping which is contracting, one needs to assume conditions which are not automatically fulfilled by a mapping on Fréchet spaces, but which are automatically locally fulfilled (on an open set) by a sufficiently regular mapping on Banach spaces. For classical statements, one can see [1, 3, 4, 5, 6, 12, 13].

We analyze here how a very classical proof of the implicit function theorem can be adapted on a ILB setting, more precisely on a slightly more general framework, that is when Fréchet spaces  $E_\infty$  considered are projective limits of a sequence of Banach spaces  $(E_i)$  without assumption of density for the inclusion  $E_{i+1} \subset E_i$ , and when the functions  $f_\infty$  on Fréchet spaces are restrictions of bounded functions  $f_i$  on the sequence of Banach space. This is what one may call order 0 maps, by analogy with the order of differential operators. Then we get (Theorem 1.6) an implicit function which is defined on a domain  $D$ , which is not a priori open in the Fréchet topology, but which contains the open ball of a Banach space. This results can be adapted to some functions  $f_\infty$  for which there does not exist any extension to a Banach space. These functions have to be controlled by a family of injective maps  $\{\Phi_x\}$ , which explains the terminology "tame" (Theorem 1.9). As a special case of applications, we recover the maps  $f_\infty$  which extend to maps  $E_i \rightarrow E_{i-r}$  (called order  $r$  maps)

We have to remark that the domain  $D$  can be very small. This is the reason why regularity results on implicit functions cannot be stated: differentiability, in a classical sense, requires open domains or at least manifolds. This leads to natural

questions for the adequate setting for analysis beyond the Banach setting. Even if not open, following the same motivations as the ones of Kriegl and Michor in [8] when they consider smoothness on non open domains, the domain  $D$  may inherit some kind of generalized setting for differential calculus, such as diffeologies [7] which are used in [10]. This question is left open, because out of the scope of this work: the most adapted (generalized) framework for the extension of the regularity (i.e. differentiability) has to be determined.

We then give consequences for an inverse functions theorem, which can be stated with the same restrictions as before on the nature of the domain  $D$ , and with an obstruction to follow the classical proof of the Fröbenius theorem from [13] where differentiation on  $D$  is explicitly needed. Finally, in last section, we show how this theorem can describe a Banach Lie group of a topological group arising in the ILH setting.

### 1. IMPLICIT FUNCTIONS FROM BANACH SPACES TO PROJECTIVE LIMITS

Let  $(E, \|\cdot\|_E)$  and  $(F, \|\cdot\|_F)$  be two Banach spaces. The Banach space  $E \times F$  is endowed with the norm  $\|(x, y)\|_{E \times F} = \max\{\|x\|_E, \|y\|_F\}$ . We note by  $D_1$  and  $D_2$  the (Fréchet) differential with respect to the variables in  $E$  and  $F$  respectively. Let us first give the statement and a proof of a classical implicit function theorem on Banach spaces, for the sake of extracting key features for generalization. For this, let  $U$  be an open neighborhood of  $O$  in  $E$ , let  $V$  be an open neighborhood of  $0$  in  $F$ , and let

$$(1) \quad f : U \times V \rightarrow F$$

be a  $C^r$ -function ( $r \geq 1$ ) in the Fréchet sense, such that

$$(2) \quad f(0, 0) = 0$$

and

$$(3) \quad D_2 f(0; 0) = Id_F.$$

**Theorem 1.1.** *There exists a constant  $c > 0$  such that, on the open ball  $B(0, c) \subset E$ , there is a unique map*

$$u : B(0, c) \rightarrow V$$

such that

$$(4) \quad \forall x \in B(0, c), \quad f(x, u(x)) = 0.$$

Let us remark that regularity of the function  $u$  is ignored for it is not necessary for next developments. We now divide the main arguments of the classical direct proof of this theorem into three lemmas.

**Lemma 1.2.** *There exists  $c_0 > 0$  and  $K > 0$  such that*

$$\|(x, y)\|_{E \times F} < c_0 \Rightarrow \|D_1 f(x, y)\|_{L(E, F)} < K.$$

*Proof.* Since  $f$  is  $C^1$ , there exists a neighborhood of  $(0, 0) \in E \times F$  such that  $\|D_1 f(x, y)\|_{L(E, F)}$  is bounded.  $\square$

**Lemma 1.3.** *There exists  $c_1 > 0$  such that*

$$(5) \quad \|(x, y)\|_{E \times F} < c_1 \Rightarrow \|D_2 f(x, y) - Id_F\|_{L(F)} < \frac{1}{2}.$$

*Proof.* The map  $f$  is of class  $C^r$ , with  $r \geq 1$ , so that, the map  $D_2f : (x, y) \in U \times V \mapsto D_2f(x, y)(\cdot) \in L(F)$  is of class  $C^{r-1}$ , and in particular of class  $C^0$ . By the way,

$$\exists c_1 > 0, \|(x, y)\|_{E \times F} < c_1 \Rightarrow \|D_2f(x, y) - D_2f(0; 0)\|_{L(F)} < \frac{1}{2}.$$

□

**Lemma 1.4.** *Let  $c_1$  be the constant of Lemma 1.3. There exists  $c_2 > 0$  such that*

$$(6) \quad \|x\|_E < c_2 \Rightarrow \|f(x, 0)\|_F < \frac{c_1}{4}.$$

*Proof.* The map  $f$  is of class  $C^r$ , with  $r \geq 1$ , so that it is in particular of class  $C^0$ . By the way, there exists a constant  $c_2$  such that

$$\|x - 0\|_E < c_2 \Rightarrow \|f(x, 0) - f(0, 0)\|_F < \frac{c_1}{4}.$$

□

**Lemma 1.5.** *Let  $c = \min\{c_0, c_1, c_2, 1\}$ . Let  $x$  such that*

$$\|x\| < c.$$

*Then the sequence  $(u_n)_{\mathbb{N}} \in F^{\mathbb{N}}$ , defined by induction by*

$$(7) \quad \begin{cases} u_0 = 0 \\ \forall n \in \mathbb{N}, u_{n+1} = u_n - f(x, u_n) \end{cases}$$

*is well-defined and converges to  $u(x) \in V$ .*

*Proof.* Let us assume  $x$  fixed. Let  $g(x, y) = y - f(x, y)$ . By the way,  $u_n = g^n(u_0)$ . Applying Lemma 1.3, let  $(y, y') \in F^2$  such that both  $(x, y)$  and  $(x, y')$  are in  $B(0, c) \subset B(0, c_1)$ .

$$f(x, y) - f(x, y') = \int_0^1 D_2f(x, ty + (1-t)y') \cdot (y' - y) dt$$

By the way

$$\begin{aligned} \|g(x, y) - g(x, y')\|_F &\leq \int_0^1 \|D_2g(x, ty + (1-t)y') \cdot (y' - y)\|_F dt \\ &\leq \int_0^1 \|D_2f(x, ty + (1-t)y') - Id_F\|_{L(F)} \cdot \|y' - y\|_F dt \\ &\leq \frac{\|y' - y\|_F}{2} \end{aligned}$$

By the way,  $g$  is  $\frac{1}{2}$ -Lipschitz. Thus, applying Lemma 1.4, we obtain by induction:

$$\|u_1 - u_0\|_F < \frac{c_1}{4} \Rightarrow \|u_{n+1} - u_n\|_F < \frac{c_1}{2^{n+2}}.$$

and

$$\forall n \in \mathbb{N}, \|u_n\|_F \leq \frac{c_1(2^n - 1)}{2^{n+2}} < c_1.$$

Hence  $(u_n)$  is converging to  $u(x)$ , which is in  $V$ .

□

*Proof of Theorem 1.1.* By Lemma 1.5 the function  $x \mapsto u(x)$  exists for  $\|x\|_E < c$ .

□

We now adapt these results to the following setting. Let  $(E_i)_{i \in \mathbb{N}}$  and  $(F_i)_{i \in \mathbb{N}}$  be two decreasing sequences of Banach spaces, i.e.  $\forall i \in \mathbb{N}$ , we have  $E_{i+1} \subset E_i$  and  $F_{i+1} \subset F_i$ , with continuous inclusion maps. We then consider  $U_0$  and  $V_0$  two open neighborhoods of 0 in  $E_0$  and  $F_0$  respectively, and a function  $f_0$ , of class  $C^r$  with the same properties as in equations (1,2,3). Let us now define, for  $i \in \mathbb{N}$ ,  $U_i = U_0 \cap E_i$  and  $V_i = V_0 \cap F_i$ , and let us assume that  $f_0$  restricts to  $C^r$ -maps

$$(8) \quad f_i : U_i \times V_i \rightarrow F_i.$$

Let  $E_\infty = \varprojlim \{E_i; i \in \mathbb{N}\}$ , let  $F_\infty = \varprojlim \{F_i; i \in \mathbb{N}\}$  and let  $f_\infty = \varprojlim \{f_i; i \in \mathbb{N}\}$ .

**Theorem 1.6.** *There exists a non-empty domain  $D_\infty \subset U_\infty$ , possibly non-open in  $U_\infty$ , and a function  $u_\infty : D_\infty \rightarrow V_\infty$  such that,*

$$\forall x \in D_\infty, f_\infty(x, u_\infty(x)) = 0,$$

and such that  $D_\infty$  contains the unit ball of the Banach space  $B_{f_\infty} \subset E_\infty$  defined as the domain of the norm

$$\|x\|_{f_\infty} = \sup_{i \in \mathbb{N}} \left\{ \frac{\|x\|_{E_i}}{c_i} \right\}.$$

*Proof.* Let  $i \in \mathbb{N}$ . We now consider a maximal domain  $D_i \subset U_i$  where there exists an unique function  $u_i$  such that

$$\forall x \in D_i, f_i(x, u_i(x)) = 0.$$

This domain is non empty since it contains  $0 \in E_i$  and, applying Theorem 1.1, there exists a constant  $c_i > 0$  such that

$$\|x\|_{E_i} < c_i \Rightarrow x \in D_i.$$

By the way, any maximal  $D_i$  is an open neighborhood of 0. By the way, setting  $D_\infty$  as the intersection of such a family  $(D_i)$ , we get that  $D_\infty$  contains  $0 \in E_\infty$ . Of course,  $D_\infty$  is not a priori open in the projective limit topology. However, let

$$B_{f_\infty} = \left\{ x \in E_\infty \mid \sup_{i \in \mathbb{N}} \left\{ \frac{\|x\|_{E_i}}{c_i} \right\} < +\infty \right\}.$$

This space is a Banach space for the norm

$$\|\cdot\|_{f_\infty} = \sup_{i \in \mathbb{N}} \left\{ \frac{\|\cdot\|_{E_i}}{c_i} \right\}.$$

Since

$$\|x\|_{f_\infty} < 1 \Leftrightarrow \forall i \in \mathbb{N}, \|x\|_{E_i} < c_i,$$

we get that the open ball of radius 1 in  $B_{f_\infty}$  is a subset of  $D_\infty$ , which ends the proof.  $\square$

Let us now extend it to a class of functions that we call **tame**. For this, we define the sequences  $(E_i)_{i \in \mathbb{N}}$  and  $(F_i)_{i \in \mathbb{N}}$  as before, as well as  $E_\infty$  and  $F_\infty$ . We also define a similar sequence  $(G_i)_{i \in \mathbb{N}}$  of Banach spaces and  $G_\infty$  the projective limit of this family.

**Definition 1.7.** *Let  $U_0 \times V_0$  be an open neighborhood of 0 in  $E_0 \times F_0$  and let  $U_\infty = U_0 \cap E_\infty$  and  $V_\infty = V_0 \cap F_\infty$ . Let us fix  $\{\Phi_x\}_{x \in U_\infty}$  be a family of injective maps from  $G_\infty$  to  $F_\infty$ . A map*

$$f : U_\infty \times V_\infty \rightarrow G_\infty$$

is  $\Phi$ -tame if and only if

$$f_\infty = \Phi_x \circ f : U_\infty \times V_\infty \rightarrow V_\infty$$

extends to  $C^r$ -maps ( $r \geq 1$ )

$$f_i : U_i \times V_i \rightarrow V_i$$

**Example 1.8.** If there exists a linear isomorphism  $A : E_0 \rightarrow E_1$  which restricts to isomorphisms  $E_i \rightarrow E_{i+1}$ , setting  $\Phi_x = A^r$ , the family of tame maps are exactly the family of functions  $f_\infty$  which extend to  $C^r$ -maps  $E_i \rightarrow E_{i-r}$ .

**Theorem 1.9.** Let  $f$  be a  $\Phi$ -tame map, such that,  $\forall i \in \mathbb{N}$ ,

$$D_2 f_i(0, 0) = Id_{E_i}.$$

Then there exists a non-empty domain  $D_\infty \subset U_\infty$ , possibly non-open in  $U_\infty$ , and a function  $u_\infty : D_\infty \rightarrow V_\infty$  such that,

$$\forall x \in D_\infty, f(x, u_\infty(x)) = 0,$$

Moreover, there exists a sequence of positive real numbers  $(c_i)$  such that  $D_\infty$  contains the unit ball of the Banach space  $B_{f, \Phi} \subset E_\infty$  defined as the domain of the norm

$$\|x\|_{f, \Phi} = \sup_{i \in \mathbb{N}} \left\{ \frac{\|x\|_{E_i}}{c_i} \right\}.$$

*Proof.* We apply Theorem 1.6 to  $\Phi \circ f$ . Then, since  $\forall x, \Phi_x$  is injective,  $\Phi_x \circ f(x, \cdot) = 0 \Leftrightarrow f(x, \cdot) = 0$ .  $\square$

## 2. TENTATIVES FOR INVERSE FUNCTIONS AND FROBENIUS THEOREM

**2.1. "Local" inverse theorem.** Let  $(E_i)_{i \in \mathbb{N}}$  be an decreasing sequence of Banach spaces, i.e.  $\forall i \in \mathbb{N}$ , we have  $E_{i+1} \subset E_i$  with continuous inclusion maps and let  $E_\infty$  be the projective limit of the family  $(E_i)_{i \in \mathbb{N}}$ . Let  $U_0$  be an open neighborhood of 0 in  $E_0$ , and define for  $i \in \mathbb{N} \cup \{\infty\}$ ,  $U_i = U_0 \cap E_i$ . Let  $V_0$  be an open neighborhood of 0 in  $E_0$ , and define for  $i \in \mathbb{N} \cup \{\infty\}$ ,  $V_i = V_0 \cap E_i$ . Let  $f_\infty : U_\infty \rightarrow V_\infty$  be a  $C^r$ -map ( $r \geq 1$ ) such that  $f(0) = 0$ , which extends to  $C^r$ -maps  $f_i : U_i \rightarrow V_i$  and such that  $Df_i(0) = Id_{E_i}$ .

**Theorem 2.1.** There exists a domain  $D \subset U_\infty$ , which contains the open unit ball of a Banach space  $B_{f_\infty} \subset E_\infty$ , with norm defined by a sequence  $(k_i)$  of positive numbers by

$$\|\cdot\|_{f_\infty} = \sup_{i \in \mathbb{N}} \left\{ \frac{\|\cdot\|_{E_i}}{k_i} \right\}$$

such that  $f_\infty|_D$  is a bijection  $D \subset U_\infty \rightarrow f_\infty(D) \subset V_\infty$ .

*Proof.* We apply Theorem 1.6 to  $g(x, y) = x - f_\infty(y)$ , for  $(x, y) \in V_\infty \times U_\infty$ . Indeed, we define a  $C^r$ -map  $g : V_\infty \times U_\infty \rightarrow E_\infty$  which extends to the maps

$$g_i : (x, y) \in V_i \times U_i \mapsto x - f_i(y) \in E_i.$$

We have that  $D_2 g_i(0; 0) = Df_i(0) = Id_{E_i}$ , so that there exists a domain  $D_\infty \subset V_\infty$ , and a sequence  $(c_i)$  of positive real numbers such that  $D_\infty$  contains the unit open ball of the Banach space  $B_{g_\infty} \subset E_\infty$  with norm  $\|\cdot\|_{g_\infty} = \sup_{i \in \mathbb{N}} \left\{ \frac{\|\cdot\|_{E_i}}{c_i} \right\}$  and a function  $u_\infty : D_\infty \rightarrow U_\infty$  such that

$$\forall x \in D_\infty, x - f_\infty(u_\infty(x)) = 0.$$

We set  $D = u_\infty(D_\infty)$ . Since each  $f_i$  is a  $C^0$ -map, there exists a sequence  $(k_i)$  of positive numbers such that

$$\|x\|_{E_i} < k_i \Rightarrow \|f_i(x)\|_{E_i} < c_i.$$

By the way,  $\forall x \in U_\infty$ ,

$$\sup_{i \in \mathbb{N}} \left\{ \frac{\|x\|_{E_i}}{k_i} \right\} < 1 \Rightarrow \sup_{i \in \mathbb{N}} \left\{ \frac{\|f_\infty(x)\|_{E_i}}{c_i} \right\} < 1 \Rightarrow f_\infty(x) \in D_\infty \Rightarrow x = u_\infty \circ f_\infty(x) \in D.$$

□

**2.2. An obstruction for a Frobenius theorem.** A setting for an adapted Frobenius theorem would be the following: *Let*

$$f_i : O_i \rightarrow L(E_i, F_i), \quad i \in -\mathbb{N}$$

*be a collection of smooth maps satisfying the following condition:*

$$i > j \Rightarrow f_j|_{O_i} \text{ restricts as a linear map to } f_i$$

*and such that,*

$$\begin{aligned} & \forall (x, y) \in O_i, \forall a, b \in E_i \\ & (D_1 f_i(x, y)(a))(b) + (D_2 f_i(x, y))(f_i(x, y)(a))(b) = \\ & (D_1 f_i(x, y)(b))(a) + (D_2 f_i(x, y))(f_i(x, y)(b))(a) \end{aligned}$$

*(this condition is the analogous of the Frobenius condition in a Banach setting, that we call the ILH Frobenius condition).*

*Then,  $\forall (x_0, y_0) \in O_\infty$ , there exists Frölicher space  $D$  that contains  $(x_0, y_0)$  and a smooth map  $J : D \rightarrow \mathbf{F}$  such that (conditions linked to differentiability of  $J$ )*

Let us now try to adapt the classical proof in e.g. [13], with the help of theorem 1.6. We can assume with no restriction that  $f(0; 0) = 0$ . We consider

$$G_i = C_b^1([0, 1], F_i) = \{\gamma \in C^1([0, 1], F_i) | \gamma(0) = 0\}$$

and

$$H_i = C^0([0, 1], F_i),$$

endowed with their usual topologies. Obviously, if  $i < j$ , the injections  $G_j \subset G_i$  and  $H_j \subset H_i$  are continuous.

Let us consider  $B_0$  an open ball of  $E_0$  centered in  $x_0$ ,  $B'_0$  an open ball of  $F_0$  centered in  $y_0$ ,  $B''_0$  an open ball of  $G_0$  centered in 0. We set  $B_i = B_0 \cap E_i$ ,  $B'_i = B'_0 \cap F_i$  and  $B''_i = B''_0 \cap G_i$ . Then, we define, for  $i \in \mathbb{N} \cup \{\infty\}$ ,

$$g_i : B_i \times B'_i \times B''_i \rightarrow H_i$$

$$g(x, y, \gamma)(t) = \mathfrak{D}\gamma dt(t) - f_i(t(x - x_0) + x_0, y + \gamma(t)).(x - x_0).$$

We then apply Theorem 1.6 to  $\int_0^{(\cdot)} \circ g_\infty : B_\infty \times B'_\infty \times B''_\infty \subset (E_\infty \times E_\infty) \times G_\infty \rightarrow G_\infty$ . There exists a domain  $D_\infty$  such that we can define the function  $\alpha_\infty$  as the unique function such that

$$\begin{cases} \alpha_\infty(x_0) = 0 \\ g_i(x, y, \alpha_\infty(x, y)) = 0, \quad \forall (x, y) \in D_\infty \end{cases}.$$

Since we set  $J(x, y) = y + \alpha_\infty(x, y)(1)$  Uniqueness follows from Theorem 1.6.

**Open problem:** We are now facing a theoretical impossibility. Classical theory of differentiation is valid for functions on open domains. We need here to consider

$D_1J$ , which is here defined on  $D_\infty$  which is not a priori open. There exists numerous extensions of the classical theory of differentiation, one of them is used in [10] based on [7]. Which one is better for this setting?

### 3. AN APPLICATION OF THE IMPLICIT FUNCTIONS THEOREM ON $\mathcal{L}_\infty$

We consider here a sequence of Banach spaces  $(E_i)$  as before, and we assume also that  $\forall i, E_{i+1}$  is dense in  $E_i$ . Following [2, 12], we consider the set of linear maps  $E_\infty \rightarrow E_\infty$  which extend to bounded linear maps  $E_i \rightarrow E_i$ . Let us note it as  $\mathcal{L}_\infty$ , and  $G\mathcal{L}_\infty = \bigcap_{i \in \mathbb{N}} GL(E_i)$  is a group known as a topological group [2], and [12] quotes “natural differentiation rules” that are identified in [9, 11] as generating a smooth Lie group for generalized differentiation on Frölicher or diffeological spaces. Let  $i \in \mathbb{N}$ , we define

$$\mathcal{L}_i = \left\{ a \in L(E_i) \mid a|_{E_\infty} \in \mathcal{L}_\infty \right\}.$$

We equip these spaces with the norms

$$\|a\|_i = \max \left\{ \|a\|_{L(E_{i-r})} \mid 0 \leq r \leq i \right\}.$$

We apply Theorem 1.6 to the map

$$f_\infty : (a, b) \in \mathcal{L}_\infty^2 \mapsto (Id + a)(Id + b) - Id,$$

for the sequence of Banach spaces  $(\mathcal{L}_i)$  with projective limit  $\mathcal{L}_\infty$ . We already know that the maximal domain  $D_\infty$  of the implicit function obtained will be

$$D_\infty \supset \left\{ a \in \mathcal{L}_\infty \mid Id + a \in G\mathcal{L}_\infty \right\}$$

and the implicit function will be

$$u_\infty : a \in D_\infty \mapsto (Id + a)^{-1} - Id$$

where  $(Id + a)^{-1}$  is the left inverse of  $Id + a$ . But the main question about  $G\mathcal{L}_\infty$  is the most adequate structure for it: it behaves like a Lie group [12, 9], but does not carry a priori charts which allows us only to consider it as a topological group [2]. Applying Theorem 1.6, there exists a Banach subspace  $B$  of  $\mathcal{L}_\infty$  defined by the norm

$$\|a\| = \sup_{i \in \mathbb{N}} \left\{ \frac{\|a\|_i}{c_i} \right\}.$$

But we easily show that each  $L_i$  is a Banach algebra, so that,  $c_i = 1$  since its group of the units contains the open ball of radius 1 centred at  $Id$ . By the way,

$$\|a\| = \sup_{i \in \mathbb{N}} \{ \|a\|_{L(E_i)} \},$$

and  $B$  is a Banach algebra, with group of the units  $GL(B) \subset G\mathcal{L}_\infty$  which is a Banach Lie group.

We finish with the special case when  $(E_i)$  is a ILH sequence (i.e. a sequence of Hilbert spaces with bounded and dense inclusion, see [12]) and when there exists a self-adjoint, positive (unbounded) operator  $Q$  such that

$$(Q^i a, b)_{E_0} = (a, b)_{E_i}.$$

In this case, there exists  $(e_k)_{k \in \mathbb{N}}$  an orthonormal base in  $E_0$  of eigenvectors of  $Q$  in  $E_\infty$  with is also orthogonal in  $E_i$ . In this case, the orthogonal projections

$$a \mapsto (e_k, a)_{E_0} e_k$$



restrict to operators in  $B$ , which shows that  $B$  is an infinite dimensional Banach algebra.

**Open question:** There is a natural right action of  $GL(B)$  on  $GL_\infty$  by composition. What is the structure of  $GL_\infty/GL(B)$ ?

#### REFERENCES

- [1] Dieudonné, J.; *Foundations of modern analysis* Academic Press (1960)
- [2] Dodson, C.T.J.; Galanis, G. Vassiliou, E.; *Geometry in a Fréchet Context: A Projective Limit Approach* london Math. Soc. Lect . Note Series **428** (2016)
- [3] Glöckner, H., Implicit functions from topological vector spaces to Banach spaces , *Israel J. Math.* **155** , 205–252 (2006).
- [4] Hamilton, R.S.; The inverse function theorem of Nash and Moser; *Bull. Amer. Math Soc. (NS)* **7** (1984) 65-222
- [5] Hogbe-Nlend, H.; *Théorie des bornologies et applications* Lect. Notes in Math. **273** (1971)
- [6] Krantz, S. G.; Parks, H.; *The Implicit Function Theorem*, Birkhauser, Boston (2002)
- [7] Iglesias-Zemmour, P.; *Diffeology* Mathematical Surveys and Monographs **185** (2013)
- [8] Kriegl, A.; Michor, P.W.; *The convenient setting for global analysis* Math. surveys and monographs **53**, American Mathematical society, Providence, USA. (2000)
- [9] Magnot, J-P., Ambrose-Singer theorem on diffeological bundles and complete integrability of the KP equation, *Int. J. of Geom. Meth. in Mod. Phys.*, **10** no9, articleID 1350043, 31 p. (2013)
- [10] Magnot, J-P.; Cauchy diffeology, numerical methods and implicit function theorem [arXiv:1607.02636](https://arxiv.org/abs/1607.02636)
- [11] Magnot, J-P.; Watts, J. The diffeology of Milnor’s classifying space *Top. Appl.* **232** 189-213 (2017)
- [12] Omori, H.; *Infinite dimensional Lie groups* AMS translations of mathematical monographs **158** (1997)
- [13] Penot, J-P.; Sur le théorème de Frobenius *Bull. SMF* **98** 47-80 (1970)

LAREMA, UNIVERSITÉ D'ANGERS, 2 BD LAVOISIER , 49045 ANGERS CEDEX 1, FRANCE AND  
 LYCÉE JEANNE D'ARC, 40 AVENUE DE GRANDE BRETAGNE, 63000 CLERMONT-FERRAND, FRANCE  
*E-mail address:* [jean-pierr.magnot@ac-clermont.fr](mailto:jean-pierr.magnot@ac-clermont.fr)