The essential role of charge-shift bonding in hypervalent prototype XeF₂

Benoît Braïda, Philippe C. Hiberty

► To cite this version:

Benoît Braïda, Philippe C. Hiberty. The essential role of charge-shift bonding in hypervalent prototype XeF₂. Nature Chemistry, 2013, 5 (5), pp.417–422. 10.1038/nchem.1619 . hal-01627883

HAL Id: hal-01627883

<https://hal.science/hal-01627883>

Submitted on 9 Nov 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The essential role of charge-shift bonding in hypervalent prototype XeF_2

Benoît Braïda^{1*} and Philippe C. Hiberty^{2*}

Hypervalency in XeF_2 and isoelectronic complexes is generally understood in terms of the Rundle-Pimentel model (which invokes a three-centre/four-electron molecular system) or its valence bond version as proposed by Coulson, which replaced the old expanded octet model of Pauling. However, the Rundle-Pimentel model is not always successful in describing such complexes and has been shown to be oversimplified. Here using *ab initio* valence bond theory coupled to quantum Monte Carlo methods, we show that the Rundle-Pimentel model is insufficient by itself in accounting for the great stability of XeF_2 , and that charge-shift bonding, wherein the large covalent-ionic interaction energy has the dominant role, is a major stabilizing factor. The energetic contribution of the old expanded octet model is also quantified and shown to be marginal. Generalizing to isoelectronic systems such as ClF_3 , SF_4 , PCl_5 and others, it is suggested that charge-shift bonding is necessary, in association with the Rundle-Pimentel model, for hypervalent analogues of XeF_2 to be strongly bonded.

Noble gas-containing compounds were long considered unstable because of their violation of the octet rule. However, they began to attract a lot of attention soon after the discovery of XePtF_6 by Bartlett¹, which was shortly followed by the synthesis of XeF_2 (ref. 2). This latter compound, the smallest product of noble gas chemistry, is also the prototype for a wide category of isoelectronic hypervalent molecules such as XeCl_2 , KrF_2 , RnF_2 , ClF_3 , SF_4 and PCl_5 , among others. It is therefore of primary importance to fully understand the nature of bonding in this prototype molecule.

The first tentative explanation for the hypervalency of P and S atoms was proposed by Pauling in terms of an expanded octet model, through promotion of electrons into vacant high-lying d orbitals, leading to sp^3d hybridization³. However, it has been shown by many researchers^{4–9} that, even if d orbitals are necessary to provide quantitative bond energies in hypervalent species, these orbitals have occupancies of only 0.3 electrons at most¹⁰, and therefore do not act primarily as valence orbitals but instead as polarization functions or as acceptor orbitals for back-donation from the ligands. Accordingly, the energetic contributions of sp^3d hybridization to hypervalent bonding are expected to be small (amazingly, to our knowledge, such quantities have never been estimated, even approximately).

It was soon realized that the explanation for hypervalency needed an alternative to the expanded octet model. In 1951, Pimentel and Rundle presented a simple and lucid model of hypervalency that does not require d -orbital participation^{11,12}. The model is based on a general three-centre–four-electron (3c–4e) molecular system, comprising three atoms or fragments that each contribute a single atomic orbital, from which one can construct a set of three molecular orbitals (MOs) of bonding, non-bonding and antibonding character, respectively. In the case of XeF_2 , three pure p_z orbitals (or s orbitals in the H_3^- case) combine to form the set of MOs φ_1 – φ_3 (Fig. 1), of which only the first two are occupied, giving a net bond order of 0.5. The Rundle–Pimentel model was immediately accepted by a wide number of chemists and served to rationalize and predict many structures.

However, Hoffmann and colleagues showed, in a systematic theoretical study of electron-rich X_3 systems, that the simple Rundle–Pimentel model is oversimplified, as it does not take s,p orbital mixing into account¹³. As this s,p mixing gives some antibonding character to the φ_2 MO in Fig. 1, some 3c–4e systems become unstable when this mixing is strong. However, even this addition to the Rundle–Pimentel model fails to explain why the simplest 3c–4e system, the H_3^- anion, is unstable, but isoelectronic F_3^- is stable. Clearly then, for the Rundle–Pimentel model to be entirely successful, there must be other factors yet to be taken into account.

One variant of the Rundle–Pimentel model is the resonance valence bond (VB) model, proposed by Coulson¹⁴. In this model, the electronic structure of XeF_2 is described as a resonating combination of a few VB structures:

where the first two VB structures display a covalent bond between one fluorine atom and a central Xe cation ($\bullet$ symbolizes a single electron), while the third structure is fully ionic. This model clearly shows that an electron must be transferred from the central atom to the ligands for hypervalency to take place. This stresses the importance of a low ionization potential for the central atom, associated with a strong electron affinity of the ligands.

Coulson demonstrated the close connection between the 3c–4e MO model and the VB resonance model¹⁴ by expanding the single-determinant MO wavefunction in terms of atomic-orbital determinants, which are then regrouped as VB structures (Supplementary Section S1). The result of the transformation (equation (2)), shows that the MO wavefunction is equivalent to a six-structure VB function.

$$\begin{aligned} \Psi(\varphi_1^2 \varphi_2^2) = & \text{F} \bullet - \bullet \text{Xe}^+ \text{F}^- + \text{F}^- \text{Xe}^+ \bullet - \bullet \text{F} + \text{F}^- \text{Xe}^{2+} \text{F}^- \\ & + \frac{1}{2}[\text{F}^- \text{XeF}^+ + \text{F}^+ \text{XeF}^- + \sqrt{2} \bullet \text{FXeF} \bullet] \end{aligned} \quad (2)$$

Even if, as we shall see, the coefficients of the three last structures are largely overestimated in the MO–VB expansion, the connection

¹UPMC Université Paris 06, CNRS UMR 7616, Laboratoire de Chimie Théorique, C. 137, 4 Place Jussieu, 75252 Paris Cedex 05, France, ²Laboratoire de Chimie Physique, UMR CNRS 8000, Université de Paris Sud, 91405 Orsay Cédex, France. *e-mail: braïda@lct.jussieu.fr; philippe.hiberty@u-psud.fr

Figure 1 | The Rundle-Pimentel orbital model for 3c-4e hypervalent complexes. Each centre contributes a single atomic orbital from which one constructs three molecular orbitals, one each of bonding, non-bonding and antibonding character. Only the bonding and non-bonding orbitals are occupied, thus allowing the complex to be stable.

between the two models is still apparent, and all the more so as both explain hypervalency without the need for *d*-orbital participation.

The Coulson-Rundle-Pimentel (CRP) model therefore provides a plausible explanation for hypervalency of heavy noble gases, which have a low ionization potential, associated with light halogens, which are strongly electronegative. However, it still does not explain why F_3^- is stable but H_3^- is not. Moreover, the CRP model does not tell us anything about one important feature, the strength of the hypervalent bonds. Are they strong or weak? In fact, they can be amazingly strong, as in the XeF_2 prototype, which has an atomization energy as large as 2.78 eV (64.1 kcal mol⁻¹)¹⁵, quite a large value if one recalls that the octet rule would predict noble gases to be inert. The question becomes even more intriguing if one tries to estimate the stabilities of the VB structures in equation (1) relative to separate atoms, using experimentally known quantities^{16,17}. For example, the bond dissociation energy D_0 of the two first VB structures, $\text{F} \bullet - \bullet \text{Xe}^+ \text{F}^-$ or $\text{F}^- \text{Xe}^+ \bullet - \bullet \text{F}$, may be estimated by summing the reaction energies of the thermodynamic cycle (for details see Supplementary Section S2):

Note that the purely covalent bond (not the full bond) between Xe^+ and F in equation (5) is assumed to be repulsive by analogy with the F_2 covalent bond^{18,19}, a hypothesis that will be verified in the following. Thus, summing equations (3) to (6) predicts the VB structures $\text{F} \bullet - \bullet \text{Xe}^+ \text{F}^-$ and $\text{F}^- \text{Xe}^+ \bullet - \bullet \text{F}$ to be definitely unstable relative to $\text{Xe} + 2\text{F} \bullet$, by at least 1.45 eV (33.4 kcal mol⁻¹). A similar thermodynamic cycle would define the third (fully ionic) VB structure, $\text{F}^- \text{Xe}^{2+} \text{F}^-$, as also largely unbound.

It follows that the large atomization energy of XeF_2 cannot come from the bonding energies of any of the individual VB structures, but solely from an exceptionally large resonance energy arising from their mixing. Such bonds, in which the covalent-ionic resonance energy is the major cause for bonding, have been termed ‘charge-shift bonds’^{21,19}, and a typical example is found in the F_2 molecule where both the covalent and ionic structures are individually unbound but their mixing provides a final bonding energy of 38.3 kcal mol⁻¹ (ref. 20). A further clue suggestive of charge-shift bonding is that this bonding mode is often present when the bonded atoms bear lone pairs, which is the case in most hypervalent compounds with S, P, Cl or a noble gas as the central atom and

halogens as ligands. Thus, as can be seen, there are some indications that charge-shift bonding might be an essential feature of hypervalency, and perhaps its root cause in many compounds. This prompted us to perform a thorough study of the nature of bonding in XeF_2 , as a prototype of many hypervalent compounds. This was done in the framework of the CRP model (equation (1)) using *ab initio* VB methods.

Results and discussion

The present calculations use either the breathing-orbital VB method (BOVB)²¹, or VB theory coupled to quantum Monte Carlo (QMC) methods^{22,23} (see Methods). Two levels of the VB-QMC method have been used: the VB-variational Monte Carlo method (VB-VMC), which allows the calculation of diabatic energies for individual VB structures, and the VB-diffusion Monte Carlo method (VB-DMC), which lacks this latter capability but is more accurate regarding the multistate ground state. A basis set of polarized triple-zeta quality is used throughout, with relativistic pseudopotentials, referred to as ps-VTZ.

Computational background. Since its synthesis² in 1962, the XeF_2 molecule has been the subject of a number of *ab initio* theoretical studies^{13,24–31}. These studies show a consensus that electron correlation effects are essential^{25–30}. A very accurate study was performed by Dixon and colleagues²⁷ using a coupled-cluster (CCSD(T)) method with estimation of the complete basis set limit and corrections from core-valence excitations and relativistic effects. They found an atomization energy of 2.61 eV (60.26 kcal mol⁻¹), close to experimental D_0 values^{16,32,33}. They also computed a bonding energy D_e of 1.81 eV (41.74 kcal mol⁻¹) for XeF^+ , again close to the experimental value of 1.95 ± 0.16 eV obtained from collision-induced dissociation studies³⁴. D_e values can be obtained from D_0 values by adding a zero point energy of 2.16 kcal mol⁻¹ for XeF_2 and 0.92 kcal mol⁻¹ for XeF^+ (ref. 27). However, the question of the nature of the bonds in these molecules is still open, hence our motivation to use VB theory to obtain more insight.

If one ignores the lone pairs of the Xe and F atoms, which play no direct role in axial bonds, XeF_2 can be considered a 3c-4e system isoelectronic to H_3^- and isovalent to F_3^- . Now, recalling that H_3^- is an unstable transition state but F_3^- is stable³⁵, it is interesting to note that the corresponding normal-valent compounds, F_2 and H_2 , display correspondingly different bonding features: the H-H bond is of the classical covalent type and the F-F bond is a typical charge-shift bond. Therefore, it appears to be of primary importance to study the FXe^+ normal-valent compound before studying XeF_2 .

The nature of the two-electron bond in normal-valent F-Xe^+ . The VB description of FXe^+ involves three VB structures (Fig. 2). The weights of each structure are displayed below the corresponding cartoons, as calculated at the BOVB and VB-VMC levels. At both levels, structure 1 (which represents a purely covalent bond between F and Xe^+) is the major one. Of the two ionic structures, 2 is the least important; and displays only one

1	2	3
0.598	0.097	0.305
0.545	0.053	0.402

Figure 2 | The three VB structures of F-Xe^+ and their weights as calculated by the BOVB (upper line) and VB-VMC (lower line) methods, in the ps-VTZ basis set. The hybridized atomic orbitals that are represented are mainly p_z orbitals with some minor *s* components and contributions of polarization functions.

Table 1 | Energies of F-Xe⁺ relative to the separate atoms, and resonance energies arising from covalent-ionic mixing.

	BOVB/ps-VTZ (kcal mol ⁻¹)	VB-VMC/ps-VTZ (kcal mol ⁻¹)	CCSD(T)/ps-VTZ (kcal mol ⁻¹)	CCSD(T)/CBS (kcal mol ⁻¹)
Structure 1 alone	27.6	38.6	–	–
Full ground state*	–40.7	–31.1	–36.9	–42.6 [‡]
RE [†]	68.3	69.7	–	–

*For BOVB and VB-VMC calculations, this corresponds to the wavefunction including all structures **1–3**.

[†]Resonance energies (RE) due to mixing of **1** with **2** and **3**.

[‡]From ref. 27.

ionized atom. In contrast, structure **3**, in which the xenon atom is doubly ionized, has a remarkably large weight (0.31–0.40).

The importance of structure **3** is also revealed by the bonding energies of FXe⁺, as calculated with structure **1** alone or with the full set of structures **1–3** (Table 1). At both VB levels, covalent structure **1** is unbound, as expected from analogy with the F₂ case. It therefore requires a considerable resonance energy, due to the mixing of structure **1** with ionic structures **2** and especially **3**, to achieve a reasonable bonding energy (41 kcal mol⁻¹ at the BOVB level versus 42.6 kcal mol⁻¹ at the CCSD(T)/CBS level²⁷). This resonance energy amounts to 68–70 kcal mol⁻¹ and is thus significantly larger than the bonding energy itself. As such, the F–Xe⁺ bond is a typical charge-shift bond, owing all its bonding energy to a very large resonance energy due to covalent–ionic mixing.

The VB description of XeF₂. The VB description of the active space of XeF₂ involves the six VB structures of equation (2), from which we can eliminate F[–]XeF⁺ + F⁺XeF[–] (presumably very high in energy). Now, for the sake of better elucidating the role of the 5d_{z²} orbitals of Xe, we will split the four remaining VB structures into **4–7** (where the active electrons of Xe are in the axial p_z orbital) and **8–11** (where at least one covalent bond involves the 5d_{z²} orbital of Xe) with optimal contribution from 5s components (called 5d_{z²} in the following). The combination of **4** and **8** (or **5** and **9**) is therefore equivalent to a unique VB structure that displays a single bond to one F atom by means of a central orbital, which is a p_z orbital polarized by the 5d_{z²} orbital. Structures **10** and **11** are fundamentally different. Each displays two Xe–F bonds and characterizes the direct participation of 5d_{z²} to bonding, following the sp³d hybridization scheme of Pauling³. Accordingly, the contribution of structures **10** and **11**, in terms of weights or in terms of stabilization energy, is a direct measure of the importance of the expanded octet model.

The calculated relative energies of XeF₂ with respect to its separate atoms are presented in Table 2. At the VB-VMC/ps-VTZ level, the calculated atomization energy is 48.7 kcal mol⁻¹, better than the CCSD(T) value in the same basis set, but somewhat smaller than the experimental and CBS extrapolated values, reflecting the necessity of very large basis sets to obtain accurate energetics for this molecule. The VB-DMC level corrects for most of the remaining basis set deficiencies and correlation effects missing in the former wavefunction, and provides an atomization energy in excellent agreement

Table 2 | Atomization energies of XeF₂, as calculated by VB-VMC, VB-DMC and CCSD(T) methods.

Computational method	Relative energy (kcal mol ⁻¹)*
VB-VMC/ps-VTZ	48.7
CCSD(T)/ps-VTZ	40.1
VB-DMC/ps-VTZ	60.6
CCSD(T)/CBS	63.7
Experiment	62.2 [†] , 63.4 [‡]

*Energies relative to the separate atoms.

[†]Ref. 32.

[‡]Ref. 33.

with the experimental value, showing that the set of VB structures **4–11** captures the essential physics of the ground state of XeF₂.

How much does sp³d hybridization contribute to bonding in XeF₂? The weights of structures **4–11** of XeF₂, as calculated at the VB-VMC/ps-VTZ level, are displayed in Fig. 3 (under their respective structures). It is apparent that the three VB structures of the Coulson–Rundle–Pimentel model (equation (1)) are largely dominant. These are covalent structures **4** + **8** and **5** + **9**, and ionic structure **6**, which together represent 81% of the electronic state. In contrast, diradical structure **7** has a weight of only 0.078 and may look amazingly marginal if one compares XeF₂ to the π-system of ozone, another 3c–4e system, where the corresponding •O–Ö–O• structure predominates over O=O⁺–O[–] (for an explanation see Supplementary Section S3). Finally, structures **10** and **11** together contribute only 11.2% to the wavefunction, thereby demonstrating the marginal contribution of sp³d hybridization. The stabilization energy contributed by structures **10** and **11** can also be estimated by comparing the atomization energies calculated with and without inclusion of **10** and **11** in the VB set.

Table 3 displays the energies of some individual VB structures (or combinations of them) for XeF₂ relative to its separate atoms. A comparison of entries 5 and 6 shows that the effect of removing **10** and **11** from the wavefunction results in an energy increase of only 7.2 kcal mol⁻¹, a value that can be considered a quantitative measure of the contribution of sp³d hybridization to the stability of XeF₂.

What is the contribution of charge-shift bonding to the hypervalency of XeF₂? The energies of the diabatic states, displayed in Table 3, can also be used to estimate the importance of resonance energies for the stability of XeF₂. As has been stated, combination (**4** + **8**) represents the complete diabatic wavefunction for the covalent VB structure F•–•Xe⁺F[–], the first term of the CRP model in equation (1). As anticipated on the basis of experimental quantities, this VB structure is unbound, and so is ionic VB structure **6**, or F[–]Xe²⁺F[–], the third term in equation (1). On the other hand, the combination **4** + **5** + **8** + **9** (entry 3 in Table 3) shows much lower energies than either (**4** + **8**) or (**5** + **9**), by as much as 82.9 kcal mol⁻¹, meaning that mixing the two covalent structures is exceptionally stabilizing. This deserves some comment.

Any symmetrical hypercoordinated species, say [A–B–A'], can be considered as an intermediate or transition state in the exchange reaction AB + A' → A + BA', where AB and BA' are normal-valent species. Generally, when AB is bound by a classical covalent bond, the so-called transition state resonance energy (TSRE) arising from mixing between the two VB structures A–B/A' and A/B–A' in [A–B–A'] is about one-half of the AB bonding energy or less³⁶. However, much larger resonance energies can be found if AB is bound by a charge-shift bond. For example, the TSRE is 41 kcal mol⁻¹ in H₃[–], less than half the bonding energy of H₂, but it is 38.9 kcal mol⁻¹ in F₃[–], slightly larger than the bonding energy of F₂ (ref. 35). Thus, a large TSRE in the hypercoordinated complex originates from the charge-shift character of the bond in

Figure 3 | VB structures of XeF₂ and their weights, as calculated at the VB-VMC/ps-VTZ level. The central atomic orbital is a pure 5p_z orbital in **4–7** and an optimized mixture of 5s and 5d₂₂ in **8** and **9**. **10** and **11** each display two simultaneous covalent bonds, one by means of the p_z atomic orbital and the other by means of the 5s–5d₂₂ one. The set of structures **4–9** corresponds to the VB version of the Rundle–Pimentel model, in which the d orbitals of Xe only have a role of polarization. Structures **10** and **11** characterize the sp³d expanded octet model.

the normal-valent compound. Where does XeF₂ fit with respect to these two typical cases? Clearly, XeF₂ falls on the same side as F₃[−], and even beyond, because the resonance energy between F••Xe⁺F[−] and F[−]Xe⁺••F, 82.9 kcal mol^{−1}, is about twice as large as the F–Xe⁺ bonding energy. This large resonance energy is connected to the charge-shift character of the F–Xe⁺ bond, and indeed this is already a strong argument in support of the role of charge-shift bonding in the stability of XeF₂. However, this very large resonance energy is still insufficient to make XeF₂ stable. About 18.6 kcal mol^{−1} can still be gained by adding diradical structure **7**, making the XeF₂ complex nearly isoenergetic with the separate atoms. On the other hand, it is the covalent–ionic mixing, obtained by also adding structure **6**, that definitely makes the system stable by contributing a resonance energy of 44.3 kcal mol^{−1} (entry 5 versus entry 4 in Table 3). At this level, XeF₂ is stable by 41.5 kcal mol^{−1} relative to its separate atoms, close to the final value of 48.7 kcal mol^{−1} obtained by adding sp³d structures **10** and **11**. Such systems, where stability is entirely due to covalent–ionic mixing, typically belong to the category of charge-shift-bonded molecules.

Comparison with other rare-gas-fluorine complexes

Coulson has already pointed out the importance of a low ionization potential for the central atom in compounds of the XeF₂ type¹⁴. As a complement to Coulson’s statement, our study emphasizes the contribution of the ionic structure F[−]Xe²⁺F[−] to the ground state of XeF₂, and the concomitant necessity for a low second ionization potential for the central atom. This feature is also a necessary condition for charge-shift bonding to take place in XeF₂, a property that can already be traced to the large resonance energy arising from the mixing of the Xe⁺••F and Xe²⁺F[−] structures in the normal-valent compound. For the sake of comparison, Table 4 reports the first and second ionization potentials of some rare gases Rg (Rg=Ne–Xe), the resonance energies in the [RgF]⁺ normal-valent molecules, as well as the experimental atomization energies of RgF₂. Not only does the rare-gas first ionization potential increase from Xe to Ne, but the second ionization

Table 4 | Ionization potentials (IPs) of rare gas Rg, resonance energies of normal-valent [RgF]⁺ and atomization energies of hypercoordinated complex RgF₂ (Rg = Ne–Xe).

[RgF] ⁺	1st IP of Rg* (eV)	2nd IP of Rg* (eV)	RE ^{†,‡} of [RgF] ⁺ (eV)	Stability of RgF ₂ (eV)
[NeF] ⁺	21.6	41.0	9.7	Unstable
[ArF] ⁺	15.8	27.6	37.3	Unstable
[KrF] ⁺	14.0	24.4	44.5	1.02 [§]
[XeF] ⁺	12.1	21.2	64.0	2.78

*Experimental values from Ref. 16.

[†]Defined as RE = E(Rg⁺••F) – E(Rg⁺••F ↔ Rg²⁺F[−]).

[‡]Calculated at the VB-VMC/ps-VTZ level.

[§]Ref. 49.

^{||}Ref. 15.

potential increases even faster in the same series (up to 41.0 eV for Ne). Meanwhile, the resonance energies in [RgF]⁺ gradually decrease from Xe to Ne, finally being almost negligible in [NeF]⁺. Quite in harmony with these trends, XeF₂ is strongly stable, KrF₂ is weakly stable, and ArF₂ and NeF₂ are unstable.

Conclusion and perspectives

The XeF₂ molecule, a classical prototype of hypervalent compounds, has been studied by using a VB-QMC method to gain detailed insight into the root causes for its amazing stability relative to its separate atoms. One side result of this study is a quantitative measure of the contribution of the old expanded octet model, which was thought long ago to explain hypervalency through sp³d hybridization. It is shown that the VB structures corresponding to this model account for 11.2% of the wavefunction, and bring a stabilization energy of only 7.2 kcal mol^{−1}, much less than the total bonding energy. These results demonstrate, in a more quantitative way than has been done before, the definite superiority of the Rundle–Pimentel model over the expanded octet proposal.

In the VB framework, XeF₂ is mainly described in terms of three VB structures, two covalent and one ionic, in accordance with the VB version of the Rundle–Pimentel model. However, none of these structures is bonding by itself, as they all lie much higher in energy than the dissociation limit. The mixing between the two covalent structures generates a resonance energy that is unexpectedly large (82.9 kcal mol^{−1}). This large resonance energy in XeF₂ arises from the charge-shift character of the corresponding normal-valent compound, FXe⁺, which displays a considerable covalent–ionic resonance energy, larger than the bonding energy itself, and therefore qualifies as a typical charge-shift bond. Thus, the charge-shift character of FXe⁺ carries over to XeF₂ and makes

Table 3 | Energies of some diabatic states of XeF₂, as calculated at the VB-VMC/ps-VTZ level.

Set of VB structures	Relative energy (kcal mol ^{−1})*
4 + 8 or 5 + 9	104.3
6	78.8
4 + 5 + 8 + 9	21.4
4 + 5 + 7 + 8 + 9	2.8
4 + 5 + 6 + 7 + 8 + 9	−41.5
4 + 5 + 6 + 7 + 8 + 9 + 10 + 11	−48.7

*Energies relative to the separate atoms.

the combination of the covalent structures much lower in energy than either of them taken separately.

Another expression of charge-shift bonding in XeF₂ can be seen in the large resonance energy (44.3 kcal mol⁻¹) arising from the mixing of the ionic structure with the combination of covalent structures. This covalent-ionic resonance energy is close to the total bonding energy at this level of calculation, which is the very definition of a charge-shift-bonded system. Thus, charge-shift bonding is a key feature of hypervalency in XeF₂, in agreement with an early proposal by Shaik³⁷, and without the large resonance energies that are attached to this type of bonding the molecule would be unstable. This means that the complete explanation for the strong stability of XeF₂ is a combination of the Rundle–Pimentel model with the occurrence of charge-shift bonding. Incidentally, our model highlights the importance of the diionic structure F⁻Xe²⁺F⁻ (often omitted in papers or textbooks), which must have a significant weight and, overall, be low enough in energy to efficiently mix with the VB structures F••Xe⁺F⁻ and F⁻Xe⁺••F. This necessity for a low second ionization potential reinforces the reason why Ne and Ar cannot be hypervalent, and why XeF₂ has a greater atomization energy than KrF₂.

Of course, the above discussed model is restricted to atoms that are prone to charge-shift bonding, that is, those from the right-hand side of the periodic table or bearing lone pairs, a necessary condition for charge-shift bonding to take place^{18,19}. For atoms from the left-hand side, another hypervalency model already exists, the VB state correlation diagrams, published long ago by Shaik and colleagues³⁸. However, the above model covers complexes of rare gas or isoelectronic groups, linked to electronegative atoms like halogens, and provides a simple explanation as to why some hypercoordinated species are stable but others are not. Thus, replacing Xe in XeF₂ by isoelectronic groups such as Kr, Rn, ClF or SF₂, one obtains KrF₂, RnF₂, ClF₃ and SF₄, while with the PCl₃ fragment and Cl ligands one obtains PCl₅, and so on; these are known hypervalent species in which we predict that charge-shift bonding is important. More *ab initio* VB studies are on their way to confirming the latter predictions and to generalize the Rundle–Pimentel/charge-shift model of hypervalency.

Methods

A many-electron system wavefunction Ψ in VB theory is expressed as a linear combination of Heitler–London–Slater–Pauling (HLSP) functions, Φ_K in equation (7):

$$\Psi = \sum_K C_K \Phi_K \quad (7)$$

where Φ_K corresponds to ‘classical’ VB structures and C_K are structural coefficients. The weights of the VB structures are defined by the Coulson–Chirgwin formula³⁹ (equation (8)), which is the equivalent of a Mulliken population analysis in VB theory:

$$W_K = C_K^2 + \sum_{L \neq K} C_K C_L \langle \Phi_K | \Phi_L \rangle \quad (8)$$

where $\langle \Phi_K | \Phi_L \rangle$ is the overlap integral of two VB structures. An important feature of our VB calculations is that all the active orbitals, here the ones that are involved in the axial F–Xe bonds, are strictly localized on a single atom, as in the classical VB method, so as to ensure a clear correspondence between the mathematical expressions of the VB structures and their physical meaning, ionic or covalent.

There are several computational approaches for VB theory at the *ab initio* level⁴⁰. In the VB self-consistent-field (VB–SCF) procedure⁴¹, both the VB orbitals and structural coefficients are optimized simultaneously to minimize the total energy. The BOVB²¹ method improves the accuracy of VB–SCF without increasing the number of VB structures Φ_K . This is done by allowing each VB structure to have its own specific set of orbitals during the optimization process such that they can be different from one VB structure to the other. In this manner, the orbitals can fluctuate in size and shape so as to fit the instantaneous charges of the atoms on which these orbitals are located.

The BOVB method has several levels of sophistication²¹. Here we chose the ‘SD–BOVB’ level, whereby each doubly occupied active orbital is split into a pair of singly occupied orbitals accommodating a spin pair, so as to account for the radial correlation of the electrons involved in the lone pair. Moreover, in this option

the orbitals that do not belong to the active space are allowed to delocalize. The BOVB calculations were carried out with the Xiamen Valence Bond (XMVB) program^{42–44}.

Very recently, a mixed VB–QMC method was proposed that managed to provide high accuracy while keeping the full interpretative capabilities of classical VB methods²². In essence, the VB–QMC wavefunction we used consists of a VB–SCF determinant part multiplied by a Jastrow function, which is included to account for electronic correlation. In this study, the bond atomization energies were computed by optimizing Jastrow parameters, VB structure coefficients and all orbitals simultaneously, using the VMC optimization algorithm on the multistructure wavefunction and separate atoms, respectively. These VB–VMC wavefunctions are then projected onto the ground state using the DMC algorithm under the fixed-node approximation, a procedure that recovers most of the remaining correlation effects that are missing at the VB–VMC level, while at the same time compensating for basis set deficiencies, and usually provides very accurate energy differences^{22,23}. This method is referred to here as VB–DMC. This latter method is more accurate for the ground state energy, but does not lend itself to the calculation of diabatic energies for individual VB structures, and cannot therefore be used for calculating resonance energies. For this purpose, VB–VMC must be used. For VB–VMC calculations of diabatic states, orbitals that are optimal for each separate combination of structures have been taken from an initial VB–SCF calculation, and then Jastrow parameters and VB structure coefficients are optimized at the VMC level. For all VB–QMC calculations, we used a systematically convergent triple-zeta polarized basis set from Burkatzki and colleagues⁴⁵, together with its corresponding pseudopotentials for both Xe and F atoms. (Basis sets and corresponding pseudopotentials are available online at <http://burkatzki.com/pseudos/index.2.html>.) For Xe and Kr atoms, to ensure enough flexibility for the *d* shell set of basis functions, Burkatzki *et al.* TZP polarization functions were replaced by the *d* set of basis functions taken from the cc-pvqz-PP basis set of Peterson and colleagues⁴⁶. The so-constructed basis set is referred to as ps-VTZ in this work. All geometries were optimized at the CCSD(T) level using the latter basis sets with the GAUSSIAN09 program⁴⁷ (for details see Supplementary Information). All QMC calculations were carried out using the CHAMP program⁴⁸.

References

- Bartlett, N. Xenon hexafluoroplatinate (V) Xe⁺PtF₆⁻. *Proc. Chem. Soc. Lond.* 218 (1962).
- Hope, R., Dähne, W., Rödder, K. M. & Mattauch, H. Fluorierung von Xenon. *Angew. Chem.* **74**, 903 (1962).
- Pauling, L. *The Nature of the Chemical Bond* 2nd edn, 145 (Cornell Univ. Press, 1940).
- Rundle, R. E. Probable structure of XeF₄ and XeF₂. *J. Am. Chem. Soc.* **85**, 112–113 (1963).
- Pitzer, K. S. Bonding in xenon fluorides and halogen fluorides. *Science* **139**, 414 (1963).
- Kutzelnigg, W. Chemical bonding in higher main group elements. *Angew. Chem. Int. Ed.* **23**, 272–295 (1984).
- Malm, J. G., Selig, H., Jortner, J. & Rice, S. A. The chemistry of xenon. *Chem. Rev.* **65**, 199–236 (1965).
- Reed, A. E. & Schleyer, P. v. R. Chemical bonding in hypervalent molecules. The dominance of ionic bonding and negative hyperconjugation over *d*-orbital participation. *J. Am. Chem. Soc.* **112**, 1434–1445 (1990).
- Musher, J. I. The chemistry of hypervalent molecules. *Angew. Chem. Int. Ed.* **8**, 54–68 (1969).
- Reed, A. E. & Weinhold, F. On the role of *d* orbitals in SF₆. *J. Am. Chem. Soc.* **108**, 3586–3593 (1986).
- Hach, R. J. & Rundle, R. E. The structure of tetramethylammonium pentafluoride. *J. Am. Chem. Soc.* **73**, 4321–4324 (1951).
- Pimentel, G. C. The bonding of trihalide and bifluoride ions by the molecular orbital method. *J. Chem. Phys.* **19**, 446–448 (1951).
- Munzarova, M. L. & Hoffmann, R. Electron-rich three-center bonding: role of *s,p* interactions across the *p*-block. *J. Am. Chem. Soc.* **124**, 4787–4795 (2002).
- Coulson, C. A. The nature of the bonding in xenon fluorides and related molecules. *J. Chem. Soc.* 1442–1454 (1964).
- Pepkin, V. I., Lebedev, Y. A. & Apin, A. Y. Enthalpy of formation of xenon difluoride. *Fiz. Zh. Khim.* **43**, 1564 (1969).
- Weast, R. C. (ed.) *Handbook of Chemistry and Physics* 70th edn (CRC Press, 1989).
- Bürger, H., Kuna, R., Ma, S., Breidung, J. & Thiel, W. The vibrational spectra of krypton and xenon difluorides—high-resolution infrared studies and *ab initio* calculations. *J. Chem. Phys.* **101**, 1–14 (1994).
- Shaik, S., Danovich, D., Silvi, B., Lauvergnat, D. & Hiberty, P. C. Charge-shift bonding—a class of electron-pair bonds that emerges from valence bond theory and is supported by the electron localization function approach. *Chem. Eur. J.* **11**, 6358–6371 (2005).

19. Shaik, S., Danovich, D., Wu, W. & Hiberty, P. C. Charge-shift bonding and its manifestations in chemistry. *Nature Chem.* **1**, 443–449 (2009).
20. Huber, K. P. & Herzberg, G. *Molecular Spectra and Molecular Structures* Vol. 4 (van Nostrand Reinhold, 1979).
21. Hiberty, P. C. & Shaik, S. Breathing-orbital valence bond method—a modern valence bond method that includes dynamic correlation. *Theor. Chem. Acc.* **108**, 255–272 (2002).
22. Braida, B., Toulouse, J., Caffarel, M. & Umrigar, C. J. Quantum Monte Carlo with Jastrow-valence-bond wave functions. *J. Chem. Phys.* **134**, 084108 (2011).
23. Petruzielo, F. R., Toulouse, J. & Umrigar, C. J. Approaching chemical accuracy with quantum Monte Carlo. *J. Chem. Phys.* **136**, 124116 (2012).
24. Bagus, P. S., Liu, B., Liskow, D. H. & Schaefer, H. F. Electron correlation and the reality of xenon difluoride. *J. Am. Chem. Soc.* **97**, 7216–7219 (1975).
25. Styszynski, J., Cao, X., Mali, G. L. & Visscher, L. Relativistic all-electron Dirac-Fock-Breit calculations on xenon fluorides (XeF_n , $n = 1, 2, 4, 6$). *J. Comput. Chem.* **18**, 601–608 (1997).
26. Liao, M. S. & Zhang, Q. E. Chemical bonding in XeF_2 , XeF_4 , KrF_2 , KrF_4 , RnF_2 , XeCl_2 , and XeBr_2 from the gas phase to the solid state. *J. Phys. Chem. A* **102**, 10647–10654 (1998).
27. Dixon, D. A., de Jong, W. A., Peterson, K. A., Christe, K. O. & Schrobilgen, G. Heats of formation of xenon fluorides and the fluxionality of XeF_6 from high level electronic structure calculations. *J. Am. Chem. Soc.* **127**, 8627–8634 (2005).
28. Haiduke, R. L. A., Filho, H. P. M. & da Silva, A. B. F. A theoretical study on the XeF_2 molecule. *Chem. Phys.* **348**, 89–96 (2008).
29. Perpointner, M. & Cederbaum, L. S. Effect of relativity on the ionization spectra of the xenon fluorides XeF_n ($n = 2, 4, 6$). *J. Chem. Phys.* **122**, 214302 (2005).
30. Kraupp, M., van Wüllen, Ch., Franke, R., Schmitz, F. & Kutzelnigg, W. The Structure of XeF_6 and of compounds isoelectronic with it. A challenge to computational chemistry and to the qualitative theory of the chemical bond. *J. Am. Chem. Soc.* **118**, 11939–11950 (1996).
31. Kurzydowski, D., Zaleski-Ejgierd, P., Grochala, W. & Hoffmann, R. Freezing in resonance structures for better packing: XeF_2 becomes $(\text{XeF}^+)(\text{F}^-)$ at large compression. *Inorg. Chem.* **50**, 3832–3840 (2011).
32. Weinstock, B., Weaver, E. E. & Knop, C. P. The xenon–fluorine system. *Inorg. Chem.* **5**, 2189–2203 (1966).
33. Berkowitz, J., Chupka, W. A., Guyon, P. M., Holloway, J. H. & Spohr, R. Photoionization mass spectrometric study of XeF_2 , XeF_4 , and XeF_6 . *J. Phys. Chem.* **75**, 1461–1471 (1971).
34. Krouse, I. H. & Wenthold, P. G. Bond dissociation energy and Lewis acidity of the xenon fluoride cation. *Inorg. Chem.* **42**, 4293–4298 (2003).
35. Braida, B. & Hiberty, P. C. What makes the trifluoride anion F_3^- so special? A breathing-orbital valence bond *ab initio* study. *J. Am. Chem. Soc.* **126**, 14890–14898 (2004).
36. Shaik, S. & Hiberty, P. C. *A Chemist's Guide to Valence Bond Theory* 142 (Wiley, 2008).
37. Shaik, S. S. in *An Encomium to Linus Pauling. Molecules in Natural Science and Medicine* (eds Maksic, Z. B. & Maksic, M. E.) 253–266 (Ellis Horwood, 1991).
38. Shaik, S. & Shurki, A. VB diagrams and chemical reactivity. *Angew. Chem. Int. Ed.* **38**, 586–625 (1999).
39. Chirgwin, H. B. & Coulson, C. A. The electronic structure of conjugated systems. VI. *Proc. R. Soc. Lond. A* **201**, 196–209 (1950).
40. Hiberty, P. C. & Shaik, S. A survey of recent developments in *ab initio* valence bond theory. *J. Comput. Chem.* **28**, 137–151 (2007).
41. van Lenthe, J. H. & Balint-Kurti, G. G. The valence-bond self-consistent field method (VB-SCF): theory and test calculations. *J. Chem. Phys.* **78**, 5699–5713 (1983).
42. Song, L., Wu, W., Mo, Y. & Zhang, Q. *XMVB: an ab initio non-orthogonal valence bond program* (Xiamen University, 2003).
43. Song, L., Mo, Y., Zhang, Q. & Wu, W. XMVB: a program for *ab initio* nonorthogonal valence bond computations. *J. Comput. Chem.* **26**, 514–521 (2005).
44. Song, L., Song, J., Mo, Y. & Wu, W. An efficient algorithm for energy gradients and orbital optimization in valence bond theory. *J. Comput. Chem.* **30**, 399–406 (2009).
45. Burkatzki, M., Filippi, C. & Dolg, M. Energy-consistent pseudopotentials for quantum Monte Carlo calculations. *J. Chem. Phys.* **126**, 234105 (2007).
46. Peterson, K. A., Figgen, D., Goll, E., Stoll, H. & Dolg, M. Systematically convergent basis sets with relativistic pseudopotentials. II. Small-core pseudopotentials and correlation consistent basis sets for the post-*d* group 16–18 elements. *J. Chem. Phys.* **119**, 11113–11123 (2003).
47. Frisch, M. J. *et al. Gaussian 09, Revision A.02* (Gaussian, 2009).
48. Umrigar, C. J., Filippi, C. & Toulouse, J. CHAMP, a Quantum Monte Carlo program *ab initio* program; available at <http://pages.physics.cornell.edu/~cyrus/champ.html>
49. Bartlett, N. & Sladky, F. O. *Comprehensive Inorganic Chemistry* Vol. 1 (eds Blair, J. C. & Emeleus, H. J.) Ch. 6 (Pergamon, 1973).

Acknowledgements

The authors thank W. Wu and C.J. Umrigar for respectively making their XMVB and CHAMP code available to us.

Author contributions

Both authors designed the project. B.B. performed all the calculations. Both authors analysed the data. P.C.H. wrote the manuscript and the Supplementary Information. Both authors discussed the results and commented on the manuscript.

Competing financial interests

The authors declare no competing financial interests.