Aggregating exposures & cumulating risk for semivolatile organic compounds: A review

Maud Pelletier, Nathalie Bonvallot, Philippe Glorennec

► To cite this version:

Maud Pelletier, Nathalie Bonvallot, Philippe Glorennec. Aggregating exposures & cumulating risk for semivolatile organic compounds: A review. *Environmental Research*, 2017, 158, pp.649-659. 10.1016/j.envres.2017.06.022 . hal-01635759

HAL Id: hal-01635759

<https://hal.science/hal-01635759>

Submitted on 15 Nov 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Aggregating exposures & cumulating risk for Semivolatile Organic

Compounds: A review

Maud Pelletier^{1, 2}, Nathalie Bonvallot^{1, 2} Philippe Glorennec^{1, 2*}

¹ EHESP-School of Public Health, Sorbonne Paris Cité, Rennes, France

² INSERM-U1085, Irset-Research Institute for Environmental and Occupational Health,
Rennes, France

*Corresponding author:

INSERM-U1085, Irset-Research Institute for Environmental and Occupational Health,
Rennes, France. EHESP-School of Public Health, Sorbonne Paris Cité, Rennes, France.

Tel.: +33 (0) 299 022 680.

Email: philippe.glorennec@ehesp.fr

ABSTRACT

Increasingly, health risk assessment is addressing multiple pathway exposures to multiple contaminants. We reviewed aggregated exposure and cumulative risk approaches for contemporary and ubiquitous semivolatile organic compounds (SVOC). We identified 22 studies aggregating exposure pathways, and 31 cumulating risk. Exposure aggregation is based on the addition of pathway-specific doses, using kinetic modeling where it exists, and classic external dose equations otherwise. In most cases, exposure is dominated by a single route or source of exposure - mainly the oral pathway - via dietary or non-dietary exposure. Preferential routes and sources of exposure are influenced by SVOC physical-chemical properties such as vapor pressure. The cumulative risk approach for contaminants is based on dose addition. Simple sum of hazard quotient (Hazard Index: HI) is the most commonly used cumulative risk assessment approach, while Relative Potency Factor (RPF) appeared to be the best suited – although this calls for a level of toxicological information that limits the number of compounds that can be studied simultaneously. Where both were performed, moving from HI to more refined approach produced similar results. In conclusion, both approaches - exposure aggregation and cumulative risk - rely on simple assumptions. Nevertheless, they allow uncertainty to be reduced, in comparison with source-by-source or chemical-by-chemical approaches.

KEY WORDS

Mixture; contaminants; environmental exposure; health; exposure pathway; cumulative risk assessment; chemical.

INTRODUCTION

People are continuously and increasingly exposed to a multitude of organic chemicals (NHANES 2015) from various sources (e.g. food, dust, cosmetics and personal care products (C&PCPs), textiles and materials) and media (e.g. air, water and soil), and by different routes of exposure such as inhalation, ingestion or dermal contact. Organic chemicals include a high number of compounds having various physical-chemical properties, and can be classified as volatile, semivolatile or non-volatile compounds. Semivolatile organic compounds (SVOCs) are defined as having a boiling point temperature of between 240 –and 400 °C (NF ISO 16000-6, 2006). This group includes a high number of organic molecules from different chemical families (e.g. phthalates, bisphenols, polycyclic aromatic hydrocarbons (PAHs), organophosphorus (OPs), organochlorines (OCs), synthetic musks, polychlorinated biphenyls (PCBs), polybromodiphenylethers (PBDEs)). The scientific community's growing interest in studying exposure to SVOCs is motivated by a rise in their use in consumer products as well as by improved analytical techniques that have shown their ubiquity, for example, in dwellings (Rudel et al. 2003; Weschler and Nazaroff 2008). Moreover most are reprotoxic (Peretz et al. 2014; Kay et al. 2014), neurotoxic (Muñoz-Quezada et al. 2013; Costa et al. 2014), or carcinogenic compounds (Armstrong et al. 2004; IARC 2015a, 2015b), and have been found in human biological fluids (blood and urine).

When evaluating the impact of SVOC exposure on human health, exposures may be assessed using either external or internal doses. External dose is estimated from contamination data and human parameters such as body weight, inhalation and dust ingestion rate... Using a pharmacokinetic model or absorption factors, it can be converted to internal dose. Internal dose is preferentially assessed by biomonitoring. Biomonitoring data directly reflect internal aggregate exposures and could, with back calculation using pharmacokinetics models, inform as to the external dose attributable to each exposure pathway. However, using biomonitoring

data does not inform on the source of exposure that is so useful for prevention strategies. Once exposure is assessed, the risk may be assessed by comparing exposures to a toxicological reference value. This value may be expressed as an external dose for a unique route of exposure, i.e. a reference dose or acceptable daily intake for ingestion (or a reference concentration for inhalation). They are usually estimated from an indicator of the dose-response relationship in laboratory mammals to which uncertainty factors are applied to take into account inter/intra-species differences (Dybing et al. 2002). For some chemicals, internal reference dose are proposed such as biomonitoring equivalent which are estimates of the concentration of the substance (or a metabolite) in blood or urine that is consistent with the reference dose (Hays and Aylward 2009).

In the case of indoor SVOCs, exposure occurs via different sources and media and concerns many compounds simultaneously. Multiple exposure to mixtures is now widely recognized and addressed, notably by international and national institutions in charge of risk assessment. In the WHO/IPCS Framework for Risk Assessment of Combined Exposures to Multiple Chemicals, aggregate exposure is defined as exposure to the same substance from multiple sources, via multiple pathways, and routes (Meek et al. 2011). Cumulative risk assessment linked to co-exposure to different substances is also recommended when feasible (Meek et al. 2011).

The objective of this study is to review the methods employed in the literature regarding (1) exposure aggregation to an SVOC excluding biomonitoring and (2) cumulative risk assessment for a mixture of SVOCs.

METHODS

The search covered the period from 1985 until 2016. Publications were selected in line with the following criteria: addressing SVOC human exposure, aggregating at least two routes of exposure for one SVOC or assessing cumulative risk to at least two SVOCs. Because they do not provide methods for the aggregation of several routes of exposure; publications that considered exposures to an SVOC from different sources (e.g. foodstuff) but via a single exposure route (e.g. ingestion) were discarded, even though aggregation was the mentioned terminology. Using the Web Of KnowledgeTM website (Thomson Reuters, www.webofknowledge.com), publications were selected primarily by searching in the field “topic” (title, abstract and key words): ((exposure AND (assessment OR evaluation) AND (cumulat* OR aggregat*)) AND (chemicals OR products OR substances OR compounds OR pollutants) AND health) OR (PBPK AND (SVOC OR pesticide OR phthalate OR flame retardant OR plasticizer OR PCB OR musk OR PAH OR organochlor* OR organophosph* OR pyrethroid) NOT (rat* OR mice OR mouse OR bear). Secondly, selected publications were screened on the basis of the subject-title match. Thirdly, those publications remaining were subjected to an exhaustive abstract reading in order to remove those which did not address the research subject and to separate publications performing exposure aggregation from those dealing with cumulative risk assessment (CRA). Publications identified in the reference list of others were added. Since the objective was to review methods, publications’ results are presented only where they allow the comparison of different methods.

Regarding publications that aggregated exposure, the following items were retrieved: references, chemical class, exposure characteristics (source, media, and route of exposure), exposure modeling (type of approach: deterministic or probabilistic, type of aggregated dose, and the doses aggregation level: external or internal), and whether comparison with biomonitoring data was made. Regarding publications that cumulated risks, the following

106 items were retrieved: references, number of chemical cumulated, SVOC selection criteria
107 (related to occurrence and toxicity), CRA characteristics (grouping by chemical class or not,
108 the dose calculation basis, and the method employed), and toxicity indicators characteristics
109 (type, source, point of departure, and type of toxicological endpoint).

RESULTS AND DISCUSSION

This search identified 22 publications that carried out an exposure aggregation from at least two routes of exposure to an SVOC and 31 publications that carried out a CRA to at least two SVOCs.

1. Aggregate exposure assessment

Table 1: Publications performing exposure aggregation on semivolatile organic compounds.

[Table 1]

Twenty-two publications carrying out SVOC exposure aggregation were identified, and these are presented in Table 1. This may seem few, given the high number of compounds concerned, yet assessments are often focused on exposure to a compound from various sources but via a single exposure route. The oldest dates back to 1998 and from 2010 to 2016, one to five publications per year were retrieved from the literature.

Seven publications aggregated exposure to OPs, six to phthalates, four to pyrethroids, three to PAHs, three to PBDEs, three to phenols, two to PCBs, two to parabens, one to musks, one to OCs, one to polychlorinated naphthalenes (PCNs), one to perfluoroalkyl acids (PFAAs) and one to replacement flame retardants (RFRs). Residential exposure was studied most (18 publications), addressing exposure to air (n=16), dust (n=13), surfaces (n=7), soil (n=6) and dental sealing (n=1). Fourteen of the publications addressed dietary exposure: all were interested in solid food exposure, four took drinking water into account and three were also interested in breast milk exposure. Three publications addressed occupational source with exposures to air (n=2) or to air and soil (n=1). Two publications addressed exposure to C&PCPs.

Human exposure to these various sources and media could occur via three pathways: inhalation, ingestion or dermal contact. Of the 22 publications selected, a majority (n=14) aggregated exposure by three pathways, whereas the rest did so by two pathways. As an example, Gosens et al. (2014) aggregated exposure to four parabens contained in C&PCPs by dermal and ingestion pathways. The reason put forward for not studying exposure by inhalation was that none of the investigated C&PCPs applied by spray contained parabens. Another reason for not studying a route of exposure is negligible absorption, in comparison with the other pathway. For instance, Wilson et al. (2003) assessed the aggregated exposure by ingestion and inhalation of 37 SVOCs present in several media (indoor and outdoor air, indoor dust, soil and food) rather than by dermal absorption because this was considered negligible in comparison. Because of recent advances in dermal exposure importance (Weschler and Nazaroff 2012) and modeling (Gong et al. 2014; Shi and Zhao 2014; Morrison et al. 2016), this pathway is expected to be more systematically considered in the future. Indeed human experiments with exposure to phthalates have proven this pathway to be significant for some SVOCs (Weschler et al. 2015).

Aggregation of exposure was always conducted by estimating exposure for each route individually, then summing them. These exposures were often assessed by employing classic external dose equations (e.g. U.S. EPA, 1992) combining human parameters (such as weight, body surface area, inhalation rate, amount of ingested dust and frequency of use) with media or product contamination (e.g. SVOC concentration in the media or product). Thirteen publications have used this method. An alternative could be the employment of pharmacokinetic (PK) and/or pharmacodynamics (PD) models to aggregate exposure to individual SVOCs by several routes, and provide an absorbed dose. The physiologically-based pharmacokinetic model (PBPK), describes the physiological, biochemical, and physical-chemical processes governing the absorption, distribution, metabolism, and excretion

of a xenobiotic in the body. It consists of a series of compartments receiving the xenobiotic via systemic circulation. These models could also incorporate a pharmacodynamic dimension (PBPK/PD) relating to processes leading to a critical health outcome (e.g. acetylcholinesterase inhibition). PK and PD models require the same parameters as for external dose equations, with the addition of physiological, physiochemical (e.g. partition coefficient) and biochemical (e.g. metabolic rate constant) parameters. Eight publications have used PK models: four used a PBPK model, two a one-compartment PK model and one a PBPK/PD model. The four PBPK models were developed for three different chemical families (PAHs, pyrethroids and OPs), the two one-compartment PK models for PBDE compounds and for PAH compounds, whereas the PBPK/PD model was only developed for OP compounds – chlorpyrifos, to be precise. When PK and/or PD models were available these appeared to be the preferred method for assessment of overall exposure to an SVOC. However, because these are available only for certain specific SVOCs, and because their construction entails high consumption of both data and resources, they are not systematically used. In order to estimate and aggregate exposures to an SVOC, some publications have also used specific exposure models, which include PK modeling components, such as the Stochastic Human Exposure and Dose Simulation model for multimedia (SHEDS-Multimedia) which was developed by the U.S. EPA's Office of Research and Development to simulate aggregate exposure via dietary and residential routes (inhalation and dermal contact) to a variety of environmental chemicals such as pesticides, metals and persistent organic pollutants (<https://www.epa.gov/chemical-research/stochastic-human-exposure-and-dose-simulation-sheds-estimate-human-exposure>). Three publications used SHEDS-Multimedia to assess exposure to pyrethroid and OP compounds. Another specific model is the Probabilistic Aggregate Consumer Exposure Model (PACEM), first used by Gosens et al. (2014) to facilitate aggregate exposure assessment of parabens compounds present in C&PCPs. However, though these are more

adaptable to various SVOCs than are PK and/or PD models, they could be limited by the use of predefined scenarios (e.g. residential pesticide use scenarios using SHEDS-Multimedia, such as indoor flying insect killer, pet treatment, vegetable patch, etc.) or data (e.g. Dutch C&PCPs use data for PACEM). Compared to classical external dose modeling, most of the time achieved with a deterministic approach, these models use distributions for the human parameters inputs and contamination data in order to allow a probabilistic approach.

Exposure assessment to a contaminant could be expressed as an external or internal dose, the latter reflecting contaminant uptake after crossing absorption barriers. Only Wilson et al. (2003) and Roosens et al. (2010) expressed results as external doses for each route of exposure. Fifteen other publications estimated internal doses for each route of exposure, and five estimated internal dose only for the dermal pathway. Of these, nine publications used a PK approach to convert external exposure into organ, tissue or fluid concentrations. Eleven other publications converted external exposure into internal exposure by applying absorption factors retrieved from the literature or considered a default value of 100%. For the dermal pathway the authors used dermal absorption fraction (Shurdut et al. 1998; Duggan et al. 2003; Linares et al. 2010; Gosens et al. 2014) or a transdermal permeability coefficient (Beko et al. 2013). Transdermal permeability coefficients are more precise, because they take into account partition through the different layers of the skin to the dermal capillaries, and these were recently available in the literature for several SVOCs (Weschler and Nazaroff 2012). This explains why their use has increased since 2013 (in comparison with 1998) for dermal absorption factors. Internal dose appears as the logical metric when considering aggregation of exposure. The most sophisticated approach is PBPK modeling, followed by PK modeling, use of absorption factor, or default 100% absorption. Using external doses (most often observed for inhalation and ingestion) actually consists in considering that there is no

206 differential absorption between exposure routes. This should at least deserve an uncertainty
207 analysis with reported values for bioavailability.

208 Both external and internal exposure assessments require input parameters (such as human
209 parameters, media or product contamination data and specific parameters of SVOCs). With
210 regard to these parameters, single values could be used to conduct a deterministic approach. A
211 conservative choice of parameters is considered as a lower tier approach by several authors
212 (Delmaar and van Engelen 2006; Sarigiannis et al. 2013), and was indeed employed in ten
213 publications. A higher tier approach could be conducted by probabilistic analyses with Monte
214 Carlo simulation by introducing distributions rather than single values, and this was employed
215 by twelve authors. Both approaches were used and compared by Gosens et al. (2014) to
216 estimate aggregate exposure, by ingestion and dermal contact, to four parabens. Although
217 exposure assessment results were comparable in both approaches, internal exposures
218 estimated by the higher tier approach were always below those estimated by the lower tier
219 approach for each paraben. This result confirmed that a deterministic approach gave a good
220 conservative starting point for exposure assessment but that a probabilistic approach gave a
221 more realistic and more informative exposure assessment where enough data were available
222 (by, for example, allowing estimation of the proportion of population above the reference
223 dose, and enabling sensitivity analysis).

224 Most of the studies (n=16) aggregated exposures for an average individual using
225 representative data from a population, e.g. a specific age group. Six studies aggregated
226 exposures for specific individuals with their own data.

227 Most studies (n=16) compared their estimations of aggregate exposure to human
228 biomonitoring data (e.g. NHANES), and eleven of these found consistent results. A reason for

the discrepancies could be for instance the non-inclusion of a major pathway, such as diet for phthalates (Beko et al. 2013).

It is noteworthy that the authors discussed the contribution made to exposure by each pathway, source or media. The major route of exposure depends on: the studied compounds; the conditions of exposure to these compounds (e.g. residential or occupational), the routes of exposure assessed, and whether the population of interest included children more susceptible to dust ingestion (Beamer et al. 2012; Wason et al. 2012; Gaspar et al. 2014; Mitro et al. 2016). Some authors investigated influent parameters such as SVOC physical-chemical parameters with molecular weight (Beko et al. 2013), and vapor pressure (Wilson et al. 2003), which are related to their partitioning among various media (e.g. indoor air, dust), leading to a preferential route of exposure. As an example, Beko et al. (2013) found the dermal pathway to be the major route of exposure for two phthalates (DnBP and DiBP). Xue et al. (2014) studied exposure to seven pyrethroids when using, or not using, pesticide at home. These authors found ingestion to be the major route of exposure in both cases, with dietary exposure being the most contributive for the population not using pesticides, and non-dietary exposure for the population that does use them. Zartarian et al. (2012) found the same results regarding children's aggregate (dietary and non-dietary) residential exposure to permethrin. The circumstance of exposure also has an influence. For example Wei et al. (2013) studied global exposure to a pyrethroid during and after aircraft disinfection treatments, finding inhalation to be the major route of exposure during the pre-flight spray, and dermal pathway to be the major route of exposure for the residual treatment scenario. The dominant pathway may change depending on exposure level. For example Trudel et al. (2011) found ingestion of food to be the dominant pathway for several PBDEs up to about the 60th quantile of the total dose, while oral and dermal uptake of dust dominated above the 60th quantile, because of a larger variability of concentration and dust ingestion rate. Because few studies (n=6) addressed

global exposure to SVOCs via ingestion, inhalation and dermal contact to at least food, air, dust or soil and objects (such as toys, clothing or indoor surfaces), it is difficult to determine a general dominant pathway trend in global exposure assessment to SVOC. However, as observed by Wilson et al. (2003), on the basis of 54 SVOCs, ingestion was found to be the main route of exposure for the less volatile compounds, whereas inhalation was the most significant for the more volatile compounds.

To summarize, choice of method depends strongly on research objectives. A PK and/or PD model seemed preferential when assessing total exposure to an SVOC. However, because they are available only for certain SVOCs, and because their construction involves high data and resource consumption, a lower tier approach - such as employing external dose equation - appeared to be more adequate (at least as a first step) and provided consistent estimations. Inclusion of an uncertainty analysis in exposure assessment appears essential in this situation.

Lastly, even where exposure to an SVOC is governed by a major route, aggregating exposure provides useful information regarding exposure characteristics such as influent media, or variation between compounds of the same chemical family.. Aggregation of all routes of exposure to an SVOC appears essential, especially when performing a risk assessment in a second step.

2. Cumulative Risk Assessment

Table 2: Publications performing Cumulative Risk Assessment on semivolatile organic compounds.

[Table 2]

Thirty-one publications conducting SVOC CRA were identified for inclusion in this review, and these are presented in Table 2. The oldest dates back to 2003, and since 2009, from one to seven publications per year were retrieved from the literature.

For each study, SVOCs were included in a CRA on the basis of their occurrence and/or a similar pattern of toxicity. Firstly, regarding occurrence, dietary contamination was the most used criterion (n=16), followed by the occurrence of SVOC metabolites in humans (n=8). In addition, two studies chose on the basis of their presence in industrial emissions, three on their presence in the residential environment, one on their presence in drinking water, one on their presence in the water, sediment and biota of a specific river and lastly, four studies provided no explanation regarding occurrence for the selection of compounds. Secondly, regarding toxicity patterns for compound selection, having an anti-androgenic mechanism of action (n=11), or inhibiting the acetylcholinesterase (AChE) mechanism (n=11) were the most studied effects. In addition, sodium channel modulators, toxic effects on the nervous system, toxic effects on the liver, the reduction of hepatic retinoid, and respiratory irritation were also studied. Finally, five publications based their compound selection on multiple toxic effects. Chemical classes investigated were: carbamates (CBs), OPs, OCs, PAHs, parabens, PBDEs, PCBs, phenols, phthalates, polyfluoroalkylated substances (PFASs) and pyrethroids. In fact, 12 publications studied OP compounds, 11 phthalate compounds, 7 carbamate compounds and 6 pyrethroid compounds. The scientific community's growing interest in OPs and phthalates is driven by suspicion of toxic health effects, such as the anti-androgenic effects of some phthalates on the male reproductive system (Joensen et al. 2012), or the neurotoxic effects of some OPs (Miles et al. 1998), their ubiquitous occurrence in various media (Rudel et al. 2003; Blanchard et al. 2014), and the acknowledged presence of these compounds and their metabolites in human biological fluids (NHANES 2015), as well as - specifically for OPs - the availability of relative potency factors (RPF) published by governmental agencies

(U.S. EPA 2006). Finally the review reveals that a majority of the publications (n=26) performed CRA for SVOCs coming from the same chemical class and only five encompassed compounds from different families.

A CRA can be performed using several methods that have been developed and used for decades now, notably for regulatory purposes on polluted sites (U.S. EPA 1989). Most of these are based on the following additivity assumptions, notably with the “Incremental Lifetime Cancer Risk”, a sum of cancer risks commonly used in regulatory risk assessment. “Dose additivity” assumes that each compound of a mixture is supposed to contribute to the overall observed effect in proportion to its concentration (dose), with the assumption that “toxicants in a mixture act upon similar biological systems and contribute to a common response in proportion to their respective potencies” (Calabrese 1990). On the other hand, “response additivity” assumes that each compound of a mixture is supposed to contribute to the overall observed effect in proportion to its effect (response), with the assumption that “toxicants in a mixture act upon dissimilar biological systems and act independently” (Calabrese 1990). This notion does not take into account the possibility of such toxicological interaction as antagonism or potentiation and could lead to biased estimation of the risk. For example, Boon et al. (2008) have highlighted an overestimation when conducting a CRA to OPs and carbamates exposures. Even though these are known for their common AChE inhibiting activity, their specific mechanisms of action are different - indeed AChE receptor binding is mostly irreversible regarding OPs whereas it is reversible regarding carbamates. However, toxicological interaction (i.e. antagonism or potentiation) is mostly based on the saturation phenomena of metabolic systems, which can be considered as never being achieved regarding low, even sometimes very low (e.g. up to ng/m^3 in indoor air regarding several SVOCs) (Blanchard et al. 2014) compound concentrations in the exposure media (ATSDR 2004). Nevertheless, an empirical analysis of low-dose synergy, by Boobis et al. (2011)

revealed results of low magnitude (1.5 to 3.5). Furthermore, the principal purpose of risk assessments is to help regulatory or policy-making organizations make decisions and to develop operative strategies regarding the control of human exposure to environmental, residential and occupational pollutants, leading to consideration of the additivity assumptions as operational.

Several CRA methods exist and the most commonly used in the literature are, from the lowest to the highest tier, the Hazard Index (HI) method, the Point of Departure Index (PODI) and the RPF approaches, described extensively elsewhere (U.S. EPA 2000; Sarigiannis and Hansen 2012; Fournier et al. 2014a). Briefly, the HI method is based on the sum of hazard quotients (HQ) for substances affecting the same target organ or system. For each substance involved, the HQ is calculated by dividing exposure by reference dose (i.e. the level at which no adverse effects are expected in humans) (U.S. EPA 2000). This method was used in 20 publications. Although it provides a transparent and comprehensive result (HI<1 indicates that there is little likelihood that an adverse effect might be observed from chronic exposure) this method is also a lower tier approach, because the reference doses are not necessarily constructed with regard to the same target organ, and can sometimes be imprecise with regard to the compounds' mechanism of action. However, it does allow the study of compounds from different chemical classes, but affecting the same target organ or system. Four publications are concerned in this review; all the others studied compounds from a single class (n=16). In order to quantify interaction effects in a mixture, Ragas et al. (2011) followed the interaction-based HI approach developed by the U.S. EPA (2000). This method assumes that interactions among compounds in a complex mixture are dependent on those of all possible binary combinations of the individual mixture components. The HI is multiplied by a weight factor reflecting the significance of these interactions. The authors observed interaction effects in two groups (with, for instance, four and two pesticides respectively) and

found that the interaction-based HI was always higher than the conventional one. However, because of weak scientific evidence regarding the calculation procedure, and a weak empirical basis regarding parameter values, the authors have warned that these interaction-based HIs should be used with caution (Ragas et al. 2011).

The PODI approach differs from the HI in that the sum of exposures to each compound is expressed as a fraction of its respective and comparable point of departure (POD) for effects of toxicological relevance (for example, a benchmark dose) rather than a fraction of the reference dose – inducing less uncertainty. The reciprocal of the PODI is the combined margin of exposure (MOE) based on the sum of individual MOEs. For each compound involved, the MOE is calculated by dividing its POD to human exposure (EFSA 2008). Three of the publications used a combined MOE approach.

The RPF approach assumes similarity of mechanisms of action between individual compounds and uses toxicity data for an index compound to normalize the potencies of each other compound in a mixture. Usually the potencies are POD derived from dose-response curves. The combined toxicity of the mixture corresponds to the sum of the potency normalized doses to yield total equivalent exposure, expressed as index compound equivalents, which is then compared to the reference dose of the index compound (EFSA 2008). This method was used in 13 publications. Seven publications studied compounds from the same chemical class - mostly OPs (n=4) - whereas the others studied compounds from different classes (n=6). Four of them applied a RPF approach class by class, whereas Caldas et al. (2006) and Blaznik et al. (2015) performed a RPF approach to OP and CB compounds having the same mechanism of action.

Several publications compared CRA methods. Three publications used both a RPF and a combined MOE approach. One of these is interested in cumulating risks linked to exposure to

five PCBs, and although the authors found similar results with respect to estimates of median MOE values, they pointed out that the advantage of the RPF approach is that it is more easily implemented in practice, and accounts quantitatively for both variability and the various uncertainties involved (Kalantari et al. 2013). Furthermore, RPF data published by the U.S. EPA are available for the OPs (U.S. EPA 2006), CBs (U.S. EPA 2007), and pyrethroids (U.S. EPA 2011). Benson et al. (2009) and Beamer et al. (2012) found similar results when using both a RPF and an HI approach. Beamer et al. (2012) did not assess the potential added value of a RPF approach to an HI approach, yet Benson et al. (2009) suggested using the less complex HI approach for future CRA to simultaneous phthalates.

The reference doses employed by the authors could differ for a single SVOC according to: the toxicological endpoint studied and the type of POD chosen (no observed adverse effect level (NOAEL), lowest observed adverse effect level (LOAEL) or Benchmark Dose (BMD)); the value of the uncertainty factor applied, and the date of construction, depending on the organization producing these values (e.g. acceptable/tolerable daily intakes (A/TDI) from EFSA or WHO, and reference doses (RfD) from U.S. EPA). Dewalque et al. (2014) and Gao et al. (2016) both calculated HI to several phthalates using two different reference values, (1) TDI from EFSA based on effects on reproduction and development and (2) reference dose from Kortenkamp and Faust (2010) based on anti-androgenic effects. Each author found different results, attesting to the fact that cumulative assessment is dependent on the reference value taken into account.

Dose calculation for exposure estimation was based on contamination data alone in 21 publications and on both contamination and biomonitoring data in 9. Only Beko et al. (2013) estimated daily intakes from both media phthalates compound contaminations (with the exception of foodstuffs) and urinary concentrations of their metabolites before conducting a

398 CRA using an HI approach. The authors found an HI from urinary concentrations to be twice
399 as high as from media contaminations.

400 Following a tiered approach when conducting a CRA is now recognized by institutions such
401 as the EFSA's Scientific Panel on Plant Protection products and their Residues (PPR) which
402 recommends the use of tiered approaches when assessing risks of chronic exposure to
403 pesticide residues (EFSA 2008). Several authors of this review have confirmed the pertinence
404 of this recommendation for conducting a CRA on SVOC exposure. A lower tier method (such
405 as the HI approach) could be used as a first step, with recourse to a higher tier method (such
406 as the RPF approach) being justified where the outcome suggests further refinement is called
407 for.

CONCLUSIONS AND PERSPECTIVES

Aggregate exposures and cumulative risks need to be taken into consideration in addressing total SVOC exposure and the overall risk they pose. In particular, the dermal pathway has proven to be significant in the recent years. This literature review has revealed that relatively simple approaches based on the sum of exposures, doses or effects can be useful, at least in a first tier. However, these first-tier approaches carry their own uncertainties and should be accompanied by an uncertainty analysis of simplifying assumptions. It is however important to keep in mind that not choosing to aggregate exposures or cumulate risks creates more uncertainty because of the implicit assumption of independence of exposures and effects.

This review has highlighted that, because they consider differential absorption rates related to the different routes of exposure, as well as metabolism phenomena, pharmacokinetic models are useful tools for aggregating exposure. Because these are currently available only for certain SVOCs, there is a need for development of generic models for those SVOCs sharing common toxicological effects (e.g. a PBPK model for reprotoxic SVOC). They could also be used for reverse dosimetry using biomonitoring data.

When assessing cumulative risks to several compounds (e.g. in order to build a RPF), a compromise needs to be reached between the exhaustiveness of the compounds list and the required precision of the toxicity estimates for each substance involved in the mixture. A great progress towards putting cumulative risk assessment into practice would be to generate comparable toxicity data in order to build RPF. This may be achieved, for instance, with the use of standardized high-throughput assays as developed last years by the US EPA (Kavlock et al. 2012), associated with the development of omics strategies (i.e. proteomics or transcriptomics). Recent works in these topics are very promising (Hannas et al. 2011;

431 Thomas et al. 2011, 2012; Van Ravenzwaay et al. 2014; Chauhan et al. 2016; Labib et al.
432 2016).

433 Both PBPK building and toxicity testing is time and resources consuming and a proposal
434 would be to focus efforts on ubiquitous compounds sharing, or suspected to share, the same
435 mechanism of action. A next step would be to identify compounds the populations are
436 frequently exposed to, and then define corresponding cumulative assessment groups, such it
437 has already be done for certain environments (Fournier et al. 2014b), or regulatory processes
438 (Boobis et al. 2008).

439

440 Acknowledgments: Maud Pelletier's work was funded with a doctoral grant from the
441 Fondation de France and the Agence de l'Environnement et de la maîtrise de l'énergie
442 (Ademe).

REFERENCES

- Armstrong, B.; Hutchinson, E.; Unwin, J.; Fletcher, T. Lung cancer risk after exposure to polycyclic aromatic hydrocarbons: A review and meta-analysis. *Environ. Health Perspect.* **2004**, *112* (9), 970–978.
- Beamer, P. I.; Canales, R. A.; Ferguson, A. C.; Leckie, J. O.; Bradman, A. Relative pesticide and exposure route contribution to aggregate and cumulative dose in young farmworker children. *Int. J. Environ. Res. Public. Health* **2012**, *9* (1), 73–96.
- Beko, G.; Weschler, C. J.; Langer, S.; Callesen, M.; Toftum, J.; Clausen, G. Children's Phthalate Intakes and Resultant Cumulative Exposures Estimated from Urine Compared with Estimates from Dust Ingestion, Inhalation and Dermal Absorption in Their Homes and Daycare Centers. *Plos One* **2013**, *8* (4), e62442.
- Benson, R. Hazard to the developing male reproductive system from cumulative exposure to phthalate esters—dibutyl phthalate, diisobutyl phthalate, butylbenzyl phthalate, diethylhexyl phthalate, dipentyl phthalate, and diisononyl phthalate. *Regul. Toxicol. Pharmacol.* **2009**, *53* (2), 90–101.
- Blanchard, O.; Glorennec, P.; Mercier, F.; Bonvallot, N.; Chevrier, C.; Ramalho, O.; Mandin, C.; Le Bot, B. Semivolatile Organic Compounds in Indoor Air and Settled Dust in 30 French Dwellings. *Environ. Sci. Technol.* **2014**, *48* (7), 3959–3969.
- Blaznik, U.; Yngve, A.; Eržen, I.; Ribič, C. H. Consumption of fruits and vegetables and probabilistic assessment of the cumulative acute exposure to organophosphorus and carbamate pesticides of schoolchildren in Slovenia. *Public Health Nutr.* **2015**, *19* (3), 557–563.

465 Boobis, A. R.; Ossendorp, B. C.; Banasiak, U.; Hamey, P. Y.; Sebestyen, I.; Moretto, A.
 466 Cumulative risk assessment of pesticide residues in food. *Toxicol. Lett.* **2008**, *180*(2),
 467 137-150.

468 Boobis, A.; Budinsky, R.; Collie, S.; Crofton, K.; Embry, M.; Felter, S.; Hertzberg, R.; Kopp,
 469 D.; Mihlan, G.; Mumtaz, M.; Price, P.; Solomon, K.; Teuschler, L.; Yang, R.; Zaleski,
 470 R. Critical analysis of literature on low-dose synergy for use in screening chemical
 471 mixtures for risk assessment. *Crit. Rev. Toxicol.* **2011**, *41* (5), 369–383.

472 Boon, P. E.; Van der Voet, H.; Van Raaij, M. T. M.; Van Klaveren, J. D. Cumulative risk
 473 assessment of the exposure to organophosphorus and carbamate insecticides in the
 474 Dutch diet. *Food Chem. Toxicol.* **2008**, *46* (9), 3090–3098.

475 Borg, D.; Lund, B.O.; Lindquist, N.G.; Håkansson, H. Cumulative health risk assessment of
 476 17 perfluoroalkylated and polyfluoroalkylated substances (PFASs) in the Swedish
 477 population. *Environ. Int.* **2013**, *59*, 112–123.

478 Bosgra, S.; Voet, H. van der; Boon, P. E.; Slob, W. An integrated probabilistic framework for
 479 cumulative risk assessment of common mechanism chemicals in food: An example
 480 with organophosphorus pesticides. *Regul. Toxicol. Pharmacol.* **2009**, *54* (2), 124–133.

481 Calabrese, E. J. *Multiple chemical interactions*; CRC Press, **1990**.

482 Caldas, E. D.; Boon, P. E.; Tressou, J. Probabilistic assessment of the cumulative acute
 483 exposure to organophosphorus and carbamate insecticides in the Brazilian diet.
 484 *Toxicology* **2006**, *222* (1–2), 132–142.

485 Castorina, R.; Bradman, A.; McKone, T. E.; Barr, D. B.; Harnly, M. E.; Eskenazi, B.
 486 Cumulative organophosphate pesticide exposure and risk assessment among pregnant

487 women living in an agricultural community: a case study from the CHAMACOS
 488 cohort. *Environ. Health Perspect.* **2003**, *111* (13), 1640–1648.

489 Chang, J. W.; Yan, B. R.; Chang, M. H.; Tseng, S. H.; Kao, Y. M.; Chen, J. C.; Lee, C. C.
 490 Cumulative risk assessment for plasticizer-contaminated food using the hazard index
 491 approach. *Environ. Pollut.* **2014**, *189*, 77–84.

492 Chauhan, V.; Kuo, B.; McNamee, J. P.; Wilkins, R. C.; Yauk, C. L. Transcriptional
 493 benchmark dose modeling: Exploring how advances in chemical risk assessment may
 494 be applied to the radiation field. *Environ. Mol. Mutagen.* **2016**, *57* (8), 589-604.

495 Christensen, K. L. Y.; Makris, S. L.; Lorber, M. Generation of hazard indices for cumulative
 496 exposure to phthalates for use in cumulative risk assessment. *Regul. Toxicol.*
 497 *Pharmacol.* **2014**, *69* (3), 380–389.

498 Costa, L. G.; de Laat, R.; Tagliaferri, S.; Pellacani, C. A mechanistic view of polybrominated
 499 diphenyl ether (PBDE) developmental neurotoxicity. *Toxicol. Lett.* **2014**, *230* (2),
 500 282–294.

501 Delmaar, J. E.; van Engelen, J. G. M. Aggregating human exposure to chemicals: An
 502 overview of tools and methodologies. The Netherlands: National Institute for Public
 503 Health and the Environment (RIVM). Report no. 630700001/2006, **2006**.

504 Dewalque, L.; Charlier, C.; Pirard, C. Estimated daily intake and cumulative risk assessment
 505 of phthalate diesters in a Belgian general population. *Toxicol. Lett.* **2014**, *231* (2),
 506 161–168.

507 Duggan, A.; Charnley, G.; Chen, W.; Chukwudebe, A.; Hawk, R.; Krieger, R. I.; Ross, J.;
 508 Yarborough, C. Di-alkyl phosphate biomonitoring data: assessing cumulative

509 exposure to organophosphate pesticides. *Regul. Toxicol. Pharmacol.* **2003**, 37 (3),
510 382–395.

511 Durmusoglu, E.; Aslan, S.; Can, E.; Bulut, Z. Health Risk Assessment of Workers' Exposure
512 to Organic Compounds in a Tire Factory. *Hum. Ecol. Risk Assess. Int. J.* **2007**, 13 (1),
513 209–222.

514 Dybing, E.; Doe, J.; Groten, J.; Kleiner, J.; O'brien, J.; Renwick, A. G.; Schlatter, J.;
515 Steinberg, P.; Tritscher, A.; Walker, R.; Younes, M. Hazard characterisation of
516 chemicals in food and diet: dose response, mechanisms and extrapolation issues. *Food*
517 *Chem. Toxicol.* **2002**, 40 (2), 237-282.

518 EFSA Panel on Plant Protection Products and their Residues (PPR) Scientific Opinion on a
519 request from the EFSA evaluate the suitability of existing methodologies and, if
520 appropriate, the identification of new approaches to assess cumulative and synergic
521 risks from pesticides to human health with a view to set MRLs for those pesticides in
522 the frame of Regulation (EC) 396/2005. *EFSA J.* **2008**, 704, 1-85.
523 <https://www.efsa.europa.eu/en/efsajournal/pub/705>.

524 Fournier, K.; Glorennec, P.; Bonvallot, N. Construction de valeurs toxicologiques de
525 référence adaptées à la prise en compte des mélanges en évaluation des risques
526 sanitaires: méthodes existantes et applications récentes. *Environ. Risques Santé* **2014a**,
527 13 (3), 203–221.

528 Fournier, K.; Glorennec, P.; Bonvallot, N. An exposure-based framework for grouping
529 pollutants for a cumulative risk assessment approach: Case study of indoor semi-
530 volatile organic compounds. *Environ. Res.* **2014b**, 130, 20–28.

531 Gao, C.-J.; Liu, L.-Y.; Ma, W.-L.; Ren, N.-Q.; Guo, Y.; Zhu, N.-Z.; Jiang, L.; Li, Y.-F.;
532 Kannan, K. Phthalate metabolites in urine of Chinese young adults: Concentration,
533 profile, exposure and cumulative risk assessment. *Sci. Total Environ.* **2016**, *543*, 19–
534 27.

535 Gaspar, F. W.; Castorina, R.; Maddalena, R. L.; Nishioka, M. G.; McKone, T. E.; Bradman,
536 A. Phthalate Exposure and Risk Assessment in California Child Care Facilities.
537 *Environ. Sci. Technol.* **2014**, *48* (13), 7593–7601.

538 Ginsberg, G.; Ginsberg, J.; Foos, B. Approaches to Children’s Exposure Assessment: Case
539 Study with Diethylhexylphthalate (DEHP). *Int. J. Environ. Res. Public. Health* **2016**,
540 *13* (7), 670.

541 Gong, M.; Zhang, Y.; Weschler, C. J. Predicting dermal absorption of gas-phase chemicals:
542 transient model development, evaluation, and application. *Indoor Air* **2014**, *24* (3),
543 292-306.

544 Gosens, I.; Delmaar, C. J. E.; ter Burg, W.; de Heer, C.; Schuur, A. G. Aggregate exposure
545 approaches for parabens in personal care products: a case assessment for children
546 between 0 and 3 years old. *J Expos Sci Env. Epidemiol* **2014**, *24* (2), 208–214.

547 Hannas, B. R.; Lambright, C. S.; Furr, J.; Evans, N.; Foster, P. M.; Gray, L. E.; Wilson, V. S.
548 Genomic biomarkers of phthalate-induced male reproductive developmental toxicity:
549 A targeted rtPCR array approach for defining relative potency. *Toxicol. Sci.* **2011**,
550 kfr315.

551 Hays, S. M.; Aylward, L. L. Using biomonitoring equivalents to interpret human
552 biomonitoring data in a public health risk context. *J. Appl. Toxicol.* **2009**, *29* (4), 275-
553 288.

554 IARC. *Polychlorinated biphenyls and polybrominated biphenyls*. IARC Monogr. Eval.
 555 Carcinog; Risk Chem. Hum. **2015a**, 107.

556 Jardim, A. N. O.; Mello, D. C.; Goes, F. C. S.; Frota Junior, E. F.; Caldas, E. D. Pesticide
 557 residues in cashew apple, guava, kaki and peach: GC- μ ECD, GC-FPD and LC-
 558 MS/MS multiresidue method validation, analysis and cumulative acute risk
 559 assessment. *Food Chem.* **2014**, *164*, 195–204.

560 Jensen, B. H.; Petersen, A.; Nielsen, E.; Christensen, T.; Poulsen, M. E.; Andersen, J. H.
 561 Cumulative dietary exposure of the population of Denmark to pesticides. *Food Chem.*
 562 *Toxicol.* **2015**, *83*, 300–307.

563 Joensen, U. N.; Frederiksen, H.; Jensen, M. B.; Lauritsen, M. P.; Olesen, I. A.; Lassen, T. H.;
 564 Andersson, A.-M.; Jørgensen, N. Phthalate excretion pattern and testicular function: a
 565 study of 881 healthy Danish men. *Environ. Health Perspect.* **2012**, *120* (10), 1397.

566 Kalantari, F.; Bergkvist, C.; Berglund, M.; Fattore, E.; Glynn, A.; Håkansson, H.; Sand, S.
 567 Establishment of the cumulative margin of exposure for a group of polychlorinated
 568 biphenyl (PCB) congeners using an improved approach that accounts for both
 569 variability and uncertainty. *Regul. Toxicol. Pharmacol.* **2013**, *65* (3), 325–333.

570 Kavlock, R.; Chandler, K.; Houck, K.; Hunter, S.; Judson, R.; Kleinstreuer, N.; Knudsen, T.;
 571 Richard, A.; Martin, M.; Padilla, S.; Reif, D.; Richard, A.; Rotroff, D.; Sipes, N.; Dix,
 572 D. Update on EPA's ToxCast program: providing high throughput decision support
 573 tools for chemical risk management. *Chem. Res. Toxicol.* **2012**, *25* (7), 1287–1302.

574 Kay, V. R.; Bloom, M. S.; Foster, W. G. Reproductive and developmental effects of phthalate
 575 diesters in males. *Crit. Rev. Toxicol.* **2014**, *44* (6), 467–498.

576 Koch, H. M.; Wittassek, M.; Brüning, T.; Angerer, J.; Heudorf, U. Exposure to phthalates in
577 5–6 years old primary school starters in Germany—A human biomonitoring study and
578 a cumulative risk assessment. *Int. J. Hyg. Environ. Health* **2011**, *214* (3), 188–195.

579 Kortenkamp, A.; Faust, M. Combined exposures to anti-androgenic chemicals: steps towards
580 cumulative risk assessment. *Int. J. Androl.* **2010**, *33* (2), 463–474.

581 Kranich, S. K.; Frederiksen, H.; Andersson, A.-M.; Jørgensen, N. Estimated Daily Intake and
582 Hazard Quotients and Indices of Phthalate Diesters for Young Danish Men. *Environ.*
583 *Sci. Technol.* **2014**, *48* (1), 706–712.

584 Labib, S.; Williams, A.; Kuo, B.; Yauk, C. L.; White, P. A.; Halappanavar, S. A framework
585 for the use of single-chemical transcriptomics data in predicting the hazards associated
586 with complex mixtures of polycyclic aromatic hydrocarbons. *Arch. Toxicol.* **2016**, 1–
587 18.

588 Leeman, W. R.; Krul, L.; Houben, G. F. Complex mixtures: Relevance of combined exposure
589 to substances at low dose levels. *Food Chem. Toxicol.* **2013**, *58*, 141–148.

590 Li, Z.; Nie, J.; Lu, Z.; Xie, H.; Kang, L.; Chen, Q.; Li, A.; Zhao, X.; Xu, G.; Yan, Z.
591 Cumulative risk assessment of the exposure to pyrethroids through fruits consumption
592 in China – Based on a 3-year investigation. *Food Chem. Toxicol.* **2016**, *96*, 234–243.

593 Linares, V.; Perelló, G.; Nadal, M.; Gómez-Catalán, J.; Llobet, J. M.; Domingo, J. L.
594 Environmental versus dietary exposure to POPs and metals: A probabilistic
595 assessment of human health risks. *J. Env. Monit* **2010**, *12* (3), 681–688.

596 Lorber, M. Exposure of Americans to polybrominated diphenyl ethers. *J. Expo. Sci. Environ.*
597 *Epidemiol.* **2007**, *18* (1), 2–19.

598 Lozowicka, B.; Mojsak, P.; Jankowska, M.; Kaczynski, P.; Hryenko, I.; Rutkowska, E.;
 599 Szabunko, J.; Borusiewicz, A. Toxicological studies for adults and children of
 600 insecticide residues with common mode of action (MoA) in pome, stone, berries and
 601 other small fruit. *Sci. Total Environ.* **2016**, 566–567, 144–156.

602 Lu, C.; Holbrook, C. M.; Andres, L. M. The implications of using a physiologically based
 603 pharmacokinetic (PBPK) model for pesticide risk assessment. *Environ. Health*
 604 *Perspect.* **2010**, 125–130.

605 Meek, M. E.; Boobis, A. R.; Crofton, K. M.; Heinemeyer, G.; Van Raaij, M.; Vickers, C. Risk
 606 assessment of combined exposure to multiple chemicals: A WHO/IPCS framework.
 607 *Regul Toxicol Pharmacol* **2011**, 60 (2 suppl 1), S1–S14.

608 Mileson, B. E.; Chambers, J. E.; Chen, W. L.; Dettbarn, W.; Ehrich, M.; Eldefrawi, A. T.;
 609 Gaylor, D. W.; Hamernik, K.; Hodgson, E.; Karczmar, A. G.; et al. Common
 610 mechanism of toxicity: a case study of organophosphorus pesticides. *Toxicol. Sci.*
 611 **1998**, 41 (1), 8–20.

612 Mitro, S. D.; Dodson, R. E.; Singla, V.; Adamkiewicz, G.; Elmi, A. F.; Tilly, M. K.; Zota, A.
 613 R. Consumer Product Chemicals in Indoor Dust: A Quantitative Meta-analysis of U.S.
 614 Studies. *Environ. Sci. Technol.* **2016**, 50 (19), 10661–10672.

615 Morrison, G. C., Weschler, C. J. and Bekö, G. Dermal uptake directly from air under transient
 616 conditions: advances in modeling and comparisons with experimental results for
 617 human subjects. *Indoor Air*, **2016**, 26: 913–924.

618 Muñoz-Quezada, M. T.; Lucero, B. A.; Barr, D. B.; Steenland, K.; Levy, K.; Ryan, P. B.;
 619 Iglesias, V.; Alvarado, S.; Concha, C.; Rojas, E.; Vega, C. Neurodevelopmental

620 effects in children associated with exposure to organophosphate pesticides: a
 621 systematic review. *Neurotoxicology* **2013**, 39, 158–168.

622 NHANES. Fourth National Report on Human Exposure to Environmental Chemicals. US
 623 Department of Health and Human and Service. **2015**.

624 Ogbeide, O.; Tongo, I.; Enuneku, A.; Ogbomida, E.; Ezemonye, L. Human Health Risk
 625 Associated with Dietary and Non-Dietary Intake of Organochlorine Pesticide Residues
 626 from Rice Fields in Edo State Nigeria. *Expo. Health* **2016**, 8 (1), 53–66.

627 Ortiz, R. H.; Maître, A.; Barbeau, D.; Lafontaine, M.; Bouchard, M. Use of physiologically-
 628 based pharmacokinetic modeling to simulate the profiles of 3-hydroxybenzo (a)
 629 pyrene in workers exposed to polycyclic aromatic hydrocarbons. *PloS One* **2014**, 9
 630 (7), e102570.

631 Peluso, F.; Othax, N.; Castelain, J. G.; Dubny, S. Applying Health Risk Analysis to Assess
 632 the Chemical Quality of Water for Recreational Bathing: The Case of Tres Arroyos
 633 Creek, Buenos Aires, Argentina. *Hum. Ecol. Risk Assess. Int. J.* **2014**, 20 (1), 45–68.

634 Peretz, J.; Vrooman, L.; Ricke, W. A.; Hunt, P. A.; Ehrlich, S.; Hauser, R.; Padmanabhan, V.;
 635 Taylor, H. S.; Swan, S. H.; VandeVoort, C. A.; et al. Bisphenol a and reproductive
 636 health: update of experimental and human evidence, 2007-2013. *Environ. Health*
 637 *Perspect. Online* **2014**, 122 (8), 775.

638 Poet, T. S.; Timchalk, C.; Hotchkiss, J. A.; Bartels, M. J. Chlorpyrifos PBPK/PD model for
 639 multiple routes of exposure. *Xenobiotica* **2014**, 44 (10), 868–881.

640 Quijano, L.; Yusà, V.; Font, G.; Pardo, O. Chronic cumulative risk assessment of the
 641 exposure to organophosphorus, carbamate and pyrethroid and pyrethrin pesticides

642 through fruit and vegetables consumption in the region of Valencia (Spain). *Food*
643 *Chem. Toxicol.* **2016**.

644 Ragas, A. M. J.; Oldenkamp, R.; Preeker, N. L.; Wernicke, J.; Schlink, U. Cumulative risk
645 assessment of chemical exposures in urban environments. *Environ. Int.* **2011**, *37* (5),
646 872–881.

647 Van Ravenzwaay, B.; Montoya, G. A.; Fabian, E.; Herold, M.; Krennrich, G.; Looser, R.;
648 Mellert, W.; Peter, E.; Strauss, V.; Walk, T.; Kamp, H. The sensitivity of
649 metabolomics versus classical regulatory toxicology from a NOAEL perspective.
650 *Toxicol. Lett.* **2014**, *227* (1), 20-28.

651 Roosens, L.; Cornelis, C.; D'Hollander, W.; Bervoets, L.; Reynders, H.; Van Campenhout,
652 K.; Van Den Heuvel, R.; Neels, H.; Covaci, A. Exposure of the Flemish population to
653 brominated flame retardants: Model and risk assessment. *Environ. Int.* **2010**, *36* (4),
654 368–376.

655 Rudel, R.A.; Camann, D. E.; Spengler, J.D.; Korn, L.R.; Brody, J.G. Phthalates,
656 Alkylphenols, Pesticides, Polybrominated Diphenyl Ethers, and Other Endocrine-
657 Disrupting Compounds in Indoor Air and Dust. *Environ. Sci. Technol.* **2003**, *37* (20),
658 4543–4553.

659 Sarigiannis, D. A.; Hansen, U. Considering the cumulative risk of mixtures of chemicals—A
660 challenge for policy makers. *Environ. Health* **2012**, *11* (Suppl 1), S18.

661 Sarigiannis, D. A.; Karakitsios, S. P.; Gotti, A. A tiered approach for aggregate exposure
662 assessment—the case of bisphenol-A. In *9th Panhellenic Conference in Chemical*
663 *Engineering, Athens, Greece* **2013**.

664 Shi, S., & Zhao, B. Modeled exposure assessment via inhalation and dermal pathways to
 665 airborne semivolatile organic compounds (SVOCs) in residences. *Environmental*
 666 *science & technology*, **2014**, 48(10), 5691-5699.

667 Shurdut, B. A.; Barraj, L.; Francis, M. Aggregate exposures under the Food Quality
 668 Protection Act: an approach using chlorpyrifos. *Regul. Toxicol. Pharmacol.* **1998**, 28
 669 (2), 165–177.

670 Søbørg, T.; Frederiksen, H.; Andersson, A. M. Cumulative risk assessment of phthalate
 671 exposure of Danish children and adolescents using the hazard index approach. *Int. J.*
 672 *Androl.* **2012**, 35 (3), 245–252.

673 Thomas, R. S.; Clewell, H. J.; Allen, B. C.; Wesselkamper, S. C.; Wang, N. C. Y.; Lambert, J.
 674 C.; Hess-Wilson, J. K.; Jay Zhao, Q., Andersen, M. E. Application of transcriptional
 675 benchmark dose values in quantitative cancer and noncancer risk assessment. *Toxicol.*
 676 *Sci.* **2010**, kfq355.

677 Thomas, R. S.; Clewell, H. J.; Allen, B. C.; Yang, L.; Healy, E.; Andersen, M. E. Integrating
 678 pathway-based transcriptomic data into quantitative chemical risk assessment: a five
 679 chemical case study. *Mut. Res./Gen. Tox. EN* **2012**, 746 (2), 135-143.

680 Trudel, D.; Scheringer, M.; von Goetz, N.; Hungerbühler, K. Total Consumer Exposure to
 681 Polybrominated Diphenyl Ethers in North America and Europe. *Environ. Sci. Technol.*
 682 **2011**, 45 (6), 2391–2397.

683 U.S. EPA. *Risk Assessment Guidance for Superfund Volume 1 Human Health Evaluation*
 684 *Manual (Part A)*. US EPA: Washington, DC, **1989**. EPA/540/1-89/002.

685 U.S. EPA. *Guidelines for Exposure Assessment. Risk Assessment Forum*; US EPA:
 686 Washington, DC, **1992**. EPA/600/Z-92/001.

687 U.S. EPA. *Supplementary guidance for conducting health risk assessment of chemical*
688 *mixtures*. US EPA: Washington, DC, **2000**. EPA/630/R-00/002.

689 U.S. EPA. *Organophosphorus Cumulative Risk Assessment*; US EPA: Washington, DC, **2006**.
690 [https://www.epa.gov/pesticide-science-and-assessing-pesticide-risks/cumulative-](https://www.epa.gov/pesticide-science-and-assessing-pesticide-risks/cumulative-assessment-risk-pesticides)
691 [assessment-risk-pesticides](https://www.epa.gov/pesticide-science-and-assessing-pesticide-risks/cumulative-assessment-risk-pesticides).

692 U.S. EPA. *Revised N-methyl Carbamate Cumulative Risk Assessment*; US EPA: Washington,
693 DC, **2007**.

694 U.S. EPA. *Pyrethrins/pyrethroid Cumulative Risk Assessment*; US EPA: Washington, DC,
695 **2011**.

696 Von Goetz, N.; Wormuth, M.; Scheringer, M.; Hungerbühler, K. Bisphenol A: How the Most
697 Relevant Exposure Sources Contribute to Total Consumer Exposure. *Risk Anal.* **2010**,
698 *30* (3), 473–487.

699 Wason, S. C.; Smith, T. J.; Perry, M. J.; Levy, J. I. Using physiologically-based
700 pharmacokinetic models to incorporate chemical and non-chemical stressors into
701 cumulative risk assessment: A case study of pesticide exposures. *Int. J. Environ. Res.*
702 *Public. Health* **2012**, *9* (5), 1971–1983.

703 Wei, B.; Isukapalli, S. S.; Weisel, C. P. Studying permethrin exposure in flight attendants
704 using a physiologically based pharmacokinetic model. *J. Expo. Sci. Environ.*
705 *Epidemiol.* **2013**, *23* (4), 416–427.

706 Weschler, C. J.; Nazaroff, W. W. Semivolatile organic compounds in indoor environments.
707 *Atmos. Environ.* **2008**, *42* (40), 9018–9040.

708 Weschler, C. J.; Nazaroff, W. W. SVOC exposure indoors: fresh look at dermal pathways.
 709 *Indoor Air* **2012**, 22 (5), 356–377.

710 Weschler, C. J., Bekö, G., Koch, H. M., Salthammer, T., Schripp, T., Toftum, J., Clausen, G.
 711 (2015). Transdermal uptake of diethyl phthalate and di (n-butyl) phthalate directly
 712 from air: experimental verification. *Environ. Health Perspect.* **2015**, 123 (10), 928.

713 Wilson, N. K.; Chuang, J. C.; Lyu, C.; Menton, R.; Morgan, M. K. Aggregate exposures of
 714 nine preschool children to persistent organic pollutants at day care and at home. *J.*
 715 *Expo. Anal. Environ. Epidemiol.* **2003**, 13 (3), 187–202.

716 Wormuth, M.; Scheringer, M.; Vollenweider, M.; Hungerbühler, K. What Are the Sources of
 717 Exposure to Eight Frequently Used Phthalic Acid Esters in Europeans? *Risk Anal.*
 718 **2006**, 26 (3), 803–824.

719 Xue, J.; Zartarian, V.; Tornero-Velez, R.; Tolve, N. S. EPA’s SHEDS-multimedia model:
 720 Children’s cumulative pyrethroid exposure estimates and evaluation against NHANES
 721 biomarker data. *Environ. Int.* **2014**, 73, 304–311.

722 Zartarian, V.; Xue, J.; Glen, G.; Smith, L.; Tolve, N.; Tornero-Velez, R. Quantifying
 723 children’s aggregate (dietary and residential) exposure and dose to permethrin:
 724 application and evaluation of EPA’s probabilistic SHEDS-Multimedia model. *J Expos*
 725 *Sci Env. Epidemiol* **2012**, 22 (3), 267–273.

726 Zentai, A.; Szabó, I. J.; Kerekes, K.; Ambrus, Á. Risk assessment of the cumulative acute
 727 exposure of Hungarian population to organophosphorus pesticide residues with regard
 728 to consumers of plant based foods. *Food Chem. Toxicol.* **2016**.

Table 1: Publications including exposure aggregation on semivolatile organic compounds

References (year of publication)	Chemical class (nb. of publications)	Exposure pathways			Exposure modeling		Aggregation			Comparison with biomonitoring data
		Source	Media	Route	Source-to-external dose	External-to-internal dose	Approach	Type of aggregated dose	Aggregation level	
Ginsberg and Foos (2016)	Phthalates (1)	Dietary Residential	Food Breast milk Drinking water Indoor air Indoor dust Indoor objects Clothing	Dermal Ingestion Inhalation	External dose equation	Absorption factors	Deterministic	Internal	Average individual	Yes
Mitro et al. (2016)	Musks (1) Parabens (3) PFAAs (11) Phenols (7) Phthalates (8) RFRs (15)	Residential	Indoor air Indoor dust	Dermal Ingestion Inhalation	External dose equation	Dermal absorption factors	Deterministic	External (ingestion, inhalation) Internal (dermal)	Average individual	No
Gosens et al. (2014)	Parabens (4)	Personal care products	Personal care products	Dermal Ingestion	External dose equation and PACEM	Absorption factors	Deterministic Probabilistic	Internal	Individual	No
Gaspar et al. (2014)	Phthalates (2)	Residential	Indoor air Indoor dust	Dermal Ingestion Inhalation	External dose equation	Dermal absorption factors	Probabilistic	Internal	Average individual	No
Poet et al. (2014)	OPs (1)	Residential	Indoor air Indoor surface (carpet treated)	Dermal Inhalation	PBPK/PD model		Deterministic	Internal	Individual	Yes
Ortiz et al. (2014)	PAHs (1)	Occupational	Indoor air	Dermal Inhalation	PBPK model and simple one-compartment PK model		Deterministic	Internal	Individual	Yes
Xue et al. (2014)	Pyrethroids (7)	Dietary Residential	Food Indoor air	Dermal Ingestion Inhalation	SHEDS-Multimedia	PK model	Probabilistic	Internal	Average individual	Yes
Beko et al. (2013)	Phthalates (5)	Residential	Gas phase Particle phase Indoor dust	Dermal Ingestion Inhalation	External dose equation	Dermal absorption factor	Probabilistic	External (ingestion, inhalation) Internal (dermal)	Average individual	Yes
Wei et al. (2013)	Pyrethroids (1)	Occupational (aircraft disinfection)	Indoor air	Dermal Ingestion Inhalation	PBPK model		Probabilistic	Internal	Average individual	Yes
Wason et al. (2013)	Pyrethroids (3) OPs (2)	Residential	Indoor dust Indoor surfaces Soil	Dermal Ingestion	SHEDS-Multimedia		Probabilistic	Internal	Average individual	No

Beamer et al. (2012)	OPs (2)	Dietary Residential	Food Indoor air Indoor dust Indoor surfaces and toy	Dermal Ingestion Inhalation	PBPK model		Probabilistic	Internal	Individual	Yes
Zartarian et al. (2012)	Pyrethroids (1)	Dietary Residential	Food Indoor air	Dermal Ingestion Inhalation	SHEDS-Multimedia	PK model	Probabilistic	Internal	Average individual	Yes
Trudel et al. (2011)	PBDEs (8)	Dietary Residential	Food Indoor air Indoor dust Indoor surfaces Soil	Dermal Ingestion Inhalation	External dose equation	Absorption factors	Probabilistic	Internal	Average individual	Yes
Roosens et al. (2010)	PBDEs (6)	Dietary Residential	Food Breast milk Indoor air Indoor dust Soil	Ingestion Inhalation	External dose equation	No	Probabilistic	External	Average individual	No
Von Goetz et al. (2010)	Phenols (1)	Dietary Residential	Food Breast milk Indoor air Indoor dust Dental sealings	Ingestion Inhalation	External dose equation	Absorption factors	Deterministic	Internal	Average individual	Yes
Linares et al. (2010)	PAHs (16) PCBs (7) PCNs (5)	Dietary Occupational (petrochemistry)	Food Drinking water Air Soil	Dermal Ingestion Inhalation	External dose equation	Dermal absorption factor	Probabilistic	External (ingestion, inhalation) Internal (dermal)	Average individual	No
Lu et al. (2010)	OPs (1)	Dietary Residential	Food Drinking water Indoor air Indoor dust Toy surfaces Soil	Ingestion Inhalation	PBPK model		Deterministic	Internal	Individual	Yes
Lorber (2007)	PBDEs (9)	Dietary Residential	Food Indoor air Indoor dust Soil	Dermal Ingestion Inhalation	Simple one-compartment PK model		Deterministic	Internal	Average individual	Yes
Wormuth et al. (2006)	Phthalates (8)	Dietary Residential Personal care products	Food Indoor air Outdoor air Indoor dust Clothing Personal care products	Dermal Ingestion Inhalation	External dose equation	Absorption factors	Deterministic	Internal	Average individual	Yes
Duggan et al. (2003)	OPs (31)	Dietary Residential	Food Drinking water Non-dietary	Dermal Ingestion Inhalation	External dose equation	Dermal absorption factor	Deterministic	External (ingestion, inhalation) Internal (dermal)	Average individual	Yes
Wilson et al. (2003)	OCs (9)	Dietary	Food	Ingestion	External dose	No	Deterministic	External	Individual	Yes

	OPs (2) PAHs (20) PCBs (18) Phenols (3) Phthalates (2)	Residential	Indoor air Outdoor air Indoor dust Soil	Inhalation	equation					
Shurdut et al. (1998)	OPs (1)	Dietary Residential	Food Indoor air Indoor surfaces	Dermal Ingestion Inhalation	External dose equation	Dermal absorption factor	Probabilistic	External (ingestion, inhalation) Internal (dermal)	Average individual	Yes

OCs = Organochlorines; OPs = Organophosphorus; PACEM = Probabilistic aggregate consumer exposure model; PAHs = Polycyclic aromatic hydrocarbons; PBDEs = Polybrominated diphenyl ethers; PCBs = Polychlorinated biphenyls; PCNs = Polychlorinated naphthalenes; PFAAs = Perfluoroalkyl acids; RFRs = Replacement flame retardants; SHEDS-multimedia = Stochastic human exposure and dose simulation model for multimedia.

Table 2: Publications performing cumulative risk assessment (CRA) on semivolatile organic compounds

References (year of publication)	Number of chemicals cumulated	SVOCs selection criteria		CRA			Toxicity indicators			
		Occurrence	Toxicity	Grouping by chemical class	Dose calculation basis	Method	Type	Source	Type of toxicological endpoint	POD
Li et al. (2016)	Pyrethroids (7)	Dietary contamination	AChE inhibitors	Yes	Contamination data	HI	ADI RfD	na	na	na
Lozowicka et al. (2016)	CBs (1) OPs (3) Pyrethroids (3)	Dietary contamination	AChE inhibitors Sodium channel modulators	Yes	Contamination data	HI	ADI	EFSA	na	na
Quijano et al. (2016)	CBs (7) OPs (12) Pyrethroids (7)	Dietary contamination	AChE inhibitors	Yes	Contamination data	RPF	RPF ADI	U.S. EPA	AChE activity	NOAEL BMD
Zentai et al. (2016)	OPs (24)	Dietary contamination	AChE inhibitors	Yes	Contamination data	RPF	RPF	Boon et al. (2008) Caldas et al. (2006)	AChE activity	NOAEL BMD ₁₀
							RfD	JMPR		
Gao et al. (2016)	Phthalates (3)	Urinary measurement	Multiple Anti-androgenic	Yes	Biomonitoring data	HI	TDI	EFSA	Effects on reproduction and development	NOAEL LOAEL BMD
							RfD	Kortenkamp and Faust (2010)	Anti-Androgenic	
Ogbeide et al. (2016)	OCs (13)	Water, sediment and biota contamination	-	Yes	Contamination data	HI	RfD	U.S. EPA	na	na
Jensen et al. (2015)	Different classes of pesticides (20)	Dietary contamination	Multiple	No	Contamination data	HI	ADI	EC EFSA JMPR	na	na
Blaznik et al. (2015)	CBs (4) OPs (10)	Dietary contamination	AChE inhibitors	No	Contamination data	RPF	RPF RfD	Boon et al. (2008) European commission	AChE activity	NOAEL LOAEL BMD ₁₀
Kranich et al. (2014)	Phthalates (5)	-	Anti-androgenic	Yes	Biomonitoring data	HI	TDI	EFSA	Effects on reproduction and development	NOAEL LOAEL BMD
							RfD	Kortenkamp and Faust (2010)	Anti-Androgenic	
Peluso et al. (2014)	OCs (9)	Water contamination	Multiple	Yes	Contamination data	HI	RfD	U.S. EPA IRIS	na	na
Xue et al. (2014)	Pyrethroids (7)	Residential (consumer use) Urinary measurement	-	Yes	Contamination data	RPF	RPF	U.S. EPA	na	na
							RfD			
Chang et al. (2014)	Phthalates (7)	-	Anti-androgenic	Yes	Contamination data	HI	TDI	EFSA WHO	Hepatotoxicity	na
							RfD	U.S. EPA	Anti-Androgenic	

Dewalque et al. (2014)	Phthalates (4)	Urinary measurement	Anti-androgenic	Yes	Biomonitoring data	HI	TDI	EFSA	Effects on reproduction and development	NOAEL LOAEL BMD
							RfD	Kortenkamp and Faust (2010)	Anti-Androgenic	
Christensen et al. (2014)	Phthalates (5)	Dietary contamination Urinary measurement	Anti-androgenic	Yes	Biomonitoring data	HI	TDI	EFSA	Effects on reproduction and development	NOAEL LOAEL
							RfD	Denmark EPA	Anti-Androgenic	
Jardim et al. (2014)	OPs (7) Pyrethroids (4)	Dietary contamination	Multiple	Yes	Contamination data	RPF	RPF	Caldas et al. (2006) U.S. EPA	AChE activity	NOAEL BMD
							RfD	JMPR U.S. EPA		
Leeman et al. (2013)	CBs (7) OPs (26) Phthalates (6)	Dietary contamination	AChE inhibitors AChE inhibitors Anti-androgenic	Yes	Contamination data	RPF	RPF	Calculated	na	na
							ADIRfD	EU Pesticide Database		
Borg et al. (2013)	PFASs (17)	Dietary contamination Urinary measurement	Multiple	Yes	Biomonitoring data	HI	RfD	Calculated	Hepatotoxicity Reproductive toxicity	NOAEL LOAEL BMD
Beko et al. (2013)	Phthalates (3)	Residential (dust contamination) Urinary measurement	Anti-androgenic	Yes	Biomonitoring data Contamination data	HI	TDI	EFSA	Effects on reproduction and development	NOAEL LOAEL
Kalantari et al. (2013)	PCBs (5)	Dietary contamination	Reduction of hepatic retinoid	Yes	Contamination data	RPF Combined MOE	RPF	Kalantari et al. (2012)	Hepatic retinoids	na
							RfD	Calculated	Body/liver weights	BMD ₀₅
Beamer et al. (2012)	OPs (2)	Dietary contamination Residential (media contamination)	AChE inhibitors	Yes	Contamination data	RPF HI	RPF RfD	U.S. EPA	ChE inhibition (plasma, RBC) ChE inhibition (brain)	BMD ₁₀ NOAEL
Søeborg et al. (2012)	Phthalates (4)	Urinary measurement	Anti-androgenic	Yes	Biomonitoring data	HI	TDI	EFSA	Effects on reproduction and development	NOAEL LOAEL
							RfD	Kortenkamp and Faust (2010)	Anti-Androgenic	
Ragas et al. (2011)	CBs (1) Neonicotinoids (1) OCs (1) OPs (2) Pyrethroids (1)	Dietary contamination	Nervous system Liver	Yes	Contamination data	HI	ADI	EC WHO	na	na
Koch et al. (2011)	Phthalates (3)	Urinary measurement	Anti-androgenic	Yes	Biomonitoring data	HI	TDI	EFSA	Effects on reproduction and development	NOAEL LOAEL
Kortenkamp and Faust (2010)	OCs (1) OPs (1) Parabens (2) Phenols (1)	-	Anti-androgenic	No	Contamination data	HI	RfD	Calculated	Anti-androgenic	NOAEL LOAEL BMDL

	Phthalates (5) PBDEs (1) Other classes of pesticides (4)						ADI	EFSA		
Linares et al. (2010)	PAHs (3)	Chemical and petrochemical industries emissions	-	Yes	Contamination data	HI	RfD	U.S. EPA	na	na
Bosgra et al. (2009)	OPs (31)	Dietary contamination	AChE inhibitors	Yes	Contamination data	RPF Combine d MOE	RPF	Calculated	AChE activity	BMD ₂₀
Benson (2009)	Phthalates (5)	-	Anti-androgenic	Yes	Biomonitoring data	RPF HI	RfD	Calculated	Anti-Androgenic	NOAEL LOAEL BMDL
Boon et al. (2008)	OPs (26) CBs (8)	Dietary contamination	AChE inhibitors	Yes	Contamination data	RPF	RPF	Calculated	AChE activity	NOAEL BMD ₁₀
							RfD	JMPR		
Durmusoglu et al. (2007)	Phenols (1) PAHs (4)	Tire industries emissions	Respiratory irritation	No	Contamination data	HI	RfD	U.S. EPA	Not specific	na
Caldas et al. (2006)	OPs (22) CBs (3)	Dietary contamination	AChE inhibitors	No	Contamination data	RPF	RPF	Calculated	AChE activity	NOAEL BMD ₁₀
							RfD	JMPR		
Castorina et al. (2003)	OPs (11)	OPs commonly applied that metabolize dialkyl compounds	AChE inhibitors	Yes	Biomonitoring data	RPF Combine d MOE	RPF	Calculated	AChE activity	BMD ₁₀

AChE = acetylcholinesterase; ADI = Acceptable daily intake; CBs = Carbamates; DNEL = Derived no effect level; EC = European Commission; HI = Hazard index; HQ = Hazard quotient; JMPR = Joint FAO/WHO Meeting on Pesticide Residues; MOE = margin of exposure; na = not available; OCs = Organochlorines; OPs = Organophosphorus; PAHs = Polycyclic aromatic hydrocarbons; PCBs = Polychlorinated biphenyls; PFAs = Perfluoroalkoxy alkanes; POD = Point of departure; POPs = Persistent organic pollutants; RBC = Red blood cell; RfD = Reference dose.