



OPE for XXX

Philippe Di Francesco, Fedor Smirnov

► To cite this version:

Philippe Di Francesco, Fedor Smirnov. OPE for XXX. Reviews in Mathematical Physics, 2018, 30 (6), pp.1840006. 10.1142/S0129055X18400068 . hal-01639490

HAL Id: hal-01639490

<https://hal.science/hal-01639490>

Submitted on 11 Apr 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

OPE FOR XXX

PHILIPPE DI FRANCESCO AND FEDOR SMIRNOV

In memory of Ludwig Dmitrievich Faddeev

ABSTRACT. We explain a new method for finding the correlation functions for the XXX model which is based on the concepts of Operator Product Expansion of Quantum Field Theory on one hand and of fermionic bases for the XXX spin chain on the other. With this method we are able to perform computations for up to 11 lattice sites. We show that these “experimental” data allow to guess exact formulae for the OPE coefficients.

1. INTRODUCTION

This paper deals with the isotropic Heisenberg spin chain. However, even dealing with a lattice model, it is important to have in mind certain general ideas coming from Quantum Field Theory (QFT). One of them is the operator product expansion (OPE).

Suppose for a given QFT we know a complete set of local operators $\mathcal{O}_i(x)$. Then for small x we have an expansion

$$\mathcal{O}_i(x)\mathcal{O}_j(0) = \sum_k C_{i,j}^k(x)\mathcal{O}_k(0).$$

We take the OPE quite symbolically leaving aside the necessity of time ordering, the convergence issues *etc.* An important feature of the OPE is that it is defined only by the short distance, ultra-violet (UV) nature of the theory. The infra-red (IR) environment becomes relevant when the correlation functions are computed. In the OPE approach, it enters through the one-point functions only:

$$(1.1) \quad \langle \mathcal{O}_i(x)\mathcal{O}_j(0) \rangle_{\text{env}} = \sum_k C_{i,j}^k(x) \langle \mathcal{O}_k(0) \rangle_{\text{env}}.$$

A first serious problem concerning the application of OPE consists in finding a convenient way to enumerate the local operators. It is well known that in a two-dimensional conformal field theory (CFT) they correspond to representations of the Virasoro algebra. A less known example is provided by the sine(h)-Gordon model for which there exists a fermionic basis for the local operators where all the one-point functions are expressed as determinants [1, 2]. In the UV limit the relation of the fermionic basis to the Virasoro basis is known, and this allows in principle to apply Perturbed CFT methods to the computation of the two-point functions.

On the other hand, let us imagine a situation when for a large number of IR environments the one-point functions and two-point functions are known. Then the relations (1.1) provide equations for the coefficients of the OPE. One can dream that these equations could be sufficient to entirely fix all the coefficients of the OPE. For the moment this program does not look realistic for the QFT. However, we shall show that it is applicable to integrable lattice models.

Generally, the OPE does not depend on the IR environment. However, one can consider classes of IR environments possessing certain symmetries. For example, they can respect

translational invariance. In this case the spatial derivatives of one-point functions vanish, and we can drop them in the right hand side of the OPE. We shall deal with much wider symmetries of this kind, and write the OPE modulo symmetries as

$$\mathcal{O}_i(x)\mathcal{O}_j(0) \equiv \sum_k C_{i,j}^k(x)\mathcal{O}_k(0).$$

Certainly, the symmetries in question will have to be carefully implemented. They can be not only continuous, but also discrete ones, like C -symmetry for instance.

In this paper we shall consider the celebrated isotropic Heisenberg magnet (XXX spin chain) with the Hamiltonian

$$H = \sum_{j=-\infty}^{\infty} h_{j,j+1}, \quad h_{j,j+1} = \frac{1}{2}(\sigma_j^a \sigma_{j+1}^a - 1),$$

with usual notations for Pauli matrices σ_j^a at site j , and where summation over repeated indices is always implied. For some technical reasons which will be explained later we consider the two-point functions of \mathfrak{sl}_2 -invariant operators. The examples will be $\sigma_1^a \sigma_N^a$ and $h_{1,2} h_{N-1,N}$.

We consider the correlation functions with arbitrary Matsubara data which will be introduced in the next section. This is a quite general situation which includes important physical applications: zero-temperature anti-ferromagnet, (anti)-ferromagnet with non-zero temperature and magnetic field or even the case of generalised Gibbs ensemble. All these cases correspond to choosing certain special Matsubara data and taking limits for them.

The set of local operators in the lattice analogue of OPE is expressed via the fermionic basis. Crucial to our study are the cases with simple Matsubara data. For these the one-point functions are computed as determinants and there is an efficient procedure for computing the two-point functions using the quantum inverse scattering method (QISM) techniques [3, 4, 5]. This allows to obtain an over-determined system of equations for the OPE coefficients. We check that these systems have solutions up to $N = 11$ and compute all the OPE coefficients. These coefficients can now be used for any Matsubara data.

It will be very interesting to guess general formulae for the OPE coefficients from our "experimental" data. This part of the work is still under way, but we shall present here some preliminary results.

The paper is organised as follows. In Section 2 we formulate the problem of computing correlation functions on a cylinder (Matsubara environment). In Section 3 the fermionic basis and its restriction to our problem are explained. In Section 4 we present an efficient computation procedure for small Matsubara chains. In Section 5 we further restrict the fermionic basis. The results of computations and formulae for some elements of OPE are given in Section 6.

2. FORMULATION OF THE PROBLEM

Consider the following square lattice wrapped onto a cylinder $\mathbb{R} \times S^1$

We associate the space \mathbb{C}^2 to each horizontal lattice site (along each of the parallel circles), and the space $\mathbb{C}^{2s_{\mathbf{m}}+1}$ together with the complex parameter $\tau_{\mathbf{m}}$ to the \mathbf{m} -th site of the compact Matsubara lattice (along horizontal lines). Denote by π_{λ}^{2s} the evaluation representation of the Yangian with the dimension $2s + 1$ and evaluation parameter λ . Every crossing corresponds to the \mathfrak{sl}_2 Yangian R -matrix in the tensor product of two evaluation representations: π_0^1 and $\pi_{\tau_{\mathbf{m}}-1/2}^{2s_{\mathbf{m}}}$. Summation is implied over all indices except




FIGURE 1.

the four fixed: $\epsilon_1, \epsilon_2, \epsilon'_1, \epsilon'_2$. We denote the partition function defined by the picture of Fig.1 by $Z_{\epsilon_1, \epsilon_2}^{\epsilon'_1, \epsilon'_2}$, then the expectation value of $\sigma_1^a \sigma_N^a$ (correlation function of spins) with the Matsubara data $\{s_{\mathbf{m}}, \tau_{\mathbf{m}}\}$ is defined by

$$G^{\sigma\sigma}(N) = 1 + \frac{2}{Z} (Z_{+-}^{-+} + Z_{-+}^{+-} - 2Z_{+-}^{+-}) ,$$

where Z is the usual partition function (without insertions). A similar formula based on the partition function with four operator insertions (at the sites $1, 2, N-1, N$) is easy to write for the expectation value $G^{hh}(N)$ of $h_{1,2} h_{N-1,N}$ (correlation function of densities of the Hamiltonian). The Matsubara direction has finite size \mathbf{n} while the horizontal (space) direction is infinite. The latter infinity may cause trouble, so we regularise the horizontal space by assuming it has finite length equal to $2L$ and imposing periodic boundary conditions (this amounts to putting our lattice on a torus), and then by finally sending $L \rightarrow \infty$.

Let us be more formal. Consider the representation of Yangian $\pi_{\mathbf{S}}$ which is the tensor product of $2L$ representations π_0^1 . Consider in addition the representation $\pi_{\mathbf{M}}$ which is the tensor product of the \mathbf{n} representations $\pi_{\tau_{\mathbf{m}}-1/2}^{2s_{\mathbf{m}}}$, $\mathbf{m} = 1, 2, \dots, \mathbf{n}$. Define

$$T_{\mathbf{S}, \mathbf{M}} = (\pi_{\mathbf{S}} \otimes \pi_{\mathbf{M}})(\mathcal{R}) ,$$

with \mathcal{R} being the universal R -matrix. Then for any \mathcal{O} acting on a small (compared to L) number of tensor components of the representation space of $\pi_{\mathbf{S}}$ define

$$Z(\mathcal{O}) = \lim_{L \rightarrow \infty} \frac{\text{Tr}_{\mathbf{M}} \text{Tr}_{\mathbf{S}} (T_{\mathbf{S}, \mathbf{M}} \mathcal{O})}{\text{Tr}_{\mathbf{M}} \text{Tr}_{\mathbf{S}} (T_{\mathbf{S}, \mathbf{M}})} .$$

This is the more general definition of the functional Z which was introduced earlier in a particular case.

To the left and to the right of the insertion (see *Fig.1*) we have products of Matsubara transfer-matrices

$$T_{\mathbf{M}} = T_{\mathbf{M}}(0), \quad T_{\mathbf{M}}(\lambda) = \text{Tr} (\pi_{\lambda}^1 \otimes \pi_{\mathbf{M}}(\mathcal{R})) ,$$

where the trace is taken with respect to the first tensor component. Generically the transfer-matrix $T_{\mathbf{M}}$ has a single eigenvector $|\text{max}\rangle$ corresponding to an eigenvalue T of maximal absolute value. Suppose the operator \mathcal{O} is localised on the interval $[1, N]$ (like on *Fig.1*). Clearly, because of the limit $L \rightarrow \infty$

$$Z(\mathcal{O}) = \frac{\langle \text{max} | \text{Tr}_{[1, N]} (T_{[1, N], \mathbf{M}} \mathcal{O}) | \text{max} \rangle}{T^N \langle \text{max} | \text{max} \rangle} .$$

This quantity was actually computed using the fermionic basis in [6]. The computation is purely algebraic and uses only the fact that $|\max\rangle$ is an eigenvector. For that reason the computation extends to more general quantities of the form

$$(2.1) \quad Z(\mathcal{O}, \Psi) = \frac{\langle \Psi | \text{Tr}_{[1, N]} (T_{[1, N], \mathbf{M}} \mathcal{O}) | \Psi \rangle}{T^N \langle \Psi | \Psi \rangle},$$

with an arbitrary eigenstate $|\Psi\rangle$ with eigenvalue T of the transfer-matrix $T_{\mathbf{M}}$.

3. FERMIONIC BASIS

The fermionic basis is defined in [7] for the anisotropic XXZ model with the anisotropy parameter $\Delta = \frac{1}{2}(q + q^{-1})$. Taking the limit $q \rightarrow 1$ is not a difficult problem. More importantly, Ref. [7] deals with the quasi-local operators of the form $q^\alpha \sum_{j=-\infty}^0 \sigma_j^3 \mathcal{O}$. Taking the limit $\alpha \rightarrow 0$ in this expression is non trivial, and requires some explanation.

In the XXZ case the basic algebraic tool is the quantum affine algebra $U_q(\widehat{\mathfrak{sl}}_2)$ which replaces the Yangian used for the XXX case. The main result of [7] is the existence of creation operators \mathbf{t}_j^* (boson), \mathbf{b}_j^* , \mathbf{c}_j^* (fermions) for $j = 1, 2, \dots$. By acting on the primary field $\Phi_\alpha = q^\alpha \sum_{j=-\infty}^0 \sigma_j^3$ these operators create a basis in the space of quasi-local operators for which the expectation values of the type considered in the previous section are easily computed. It should be noticed that contrary to the fermions the operators \mathbf{t}_j^* are of rather simple origin, however they make computations much more involved.

In the limit $\alpha \rightarrow 0$ the operators \mathbf{b}_j^* , \mathbf{c}_j^* acting on Φ_α develop simple poles in α . As a consequence, computing expectation values involves using L'Hospital's rules. This is doable, but leads eventually to the doubling of fermions: one has to consider for every fermion its regular part and residue at $\alpha = 0$. However, there is a notable exception of this general picture. The algebra $U_q(\widehat{\mathfrak{sl}}_2)$ contains two finite-dimensional subalgebras isomorphic to $U_q(\mathfrak{sl}_2)$. Consider the operators \mathcal{O} commuting with the action of one of them. In a *weak* sense (when inserted in expectation values) it can be shown that in the limit $\alpha \rightarrow 0$ these operators are created by the action of regular parts of fermions only. This is a great simplification. Another nice point is that the operators \mathbf{t}^* do not contribute for $\alpha = 0$.

Passing to the XXX case with $\alpha = 0$ we assert that the local operators are created by \mathbf{b}_j^* , \mathbf{c}_j^* . We use the multi-index notations: for $I = \{i_1, \dots, i_n\}$ we define $\mathbf{b}_I^* = \mathbf{b}_{i_1}^* \cdots \mathbf{b}_{i_n}^*$, $\mathbf{c}_I^* = \mathbf{c}_{i_1}^* \cdots \mathbf{c}_{i_n}^*$. We denote by $\#(I) = n$ the cardinality of I , and by $|I| = \sum_{\ell=1}^n i_\ell$. Then according to previous considerations the invariant operators are contained in the space \mathfrak{H}_0 of operators with the basis

$$(3.1) \quad \mathbf{b}_I^* \mathbf{c}_J^* \cdot \mathbf{I}, \quad \#(I) = \#(J),$$

\mathbf{I} standing for the unit operator. The operator (3.1) is supported on an interval of length at most $|I| + |J|$.

It is important to identify among these the operators which are localised on an interval of length N . The following first two conditions follow for instance from [8]:

$$(3.2) \quad \#(I) \leq [N/2], \quad \max(I \cup J) \leq N.$$

Moreover, we consider C -invariant operators (invariant under $\sigma_j^a \rightarrow -\sigma_j^a$). For them it can be shown that the following additional condition can be imposed:

$$(3.3) \quad |I| + |J| \equiv 0 \pmod{2}.$$

Let us denote the subspace of \mathfrak{H}_0 satisfying (3.2), (3.3) by $\mathfrak{H}_0^{(N)}$.

More sophisticated conditions result from the homogeneous limit of the results of [9]. Consider the annihilation operators $\mathbf{b}_j, \mathbf{c}_j$ canonically conjugated to $\mathbf{b}_j^*, \mathbf{c}_j^*$, and annihilating the vacuum. Introduce the operators

$$Q_m = \sum_{j=1}^{m-1} c_j b_{m-j}, \quad m = 1, 2, \dots$$

We state that the operators localised on $[1, N]$ are linear combinations of (3.1) satisfying

$$(3.4) \quad Q_m \mathcal{O} = 0, \quad m > N.$$

Due to the second condition in (3.2) for given N we can restrict the values of m in the above to $m \leq 2N + 1$.

The main theorem of [6] states that for $\#(I) = \#(J)$

$$(3.5) \quad Z(\mathbf{b}_I^* \mathbf{c}_J^* \cdot \mathbf{I}, \Psi) = \det(\omega_{i_p, j_q})_{p, q=1, \dots, \#(I)},$$

where $\omega_{i,j}$ is an infinite matrix depending on

$$\text{Matsubara data} = \{\{s_{\mathbf{m}}, \tau_{\mathbf{m}}\}_{\mathbf{m}=1}^{\mathbf{n}}, |\Psi\rangle\}.$$

The exact definition of $\omega_{i,j}$ will be given in the next section.

In our case ($\alpha = 0$) the matrix $\omega_{i,j}$ is symmetric. Since we decided to work in a weak sense, this imposes additional restrictions. The first of them is obvious: we can consider the space \mathfrak{H}_0/ι , with ι being the anti-homomorphism $\mathbf{b}_j^* \leftrightarrow \mathbf{c}_j^*$. For example, we can require that $I \leq J$ in lexicographical order. Another consequence is less trivial. The following simple example explains that there are linear relations between minors of any given symmetric matrix:

$$\begin{vmatrix} \omega_{1,3} & \omega_{1,4} \\ \omega_{2,3} & \omega_{2,4} \end{vmatrix} - \begin{vmatrix} \omega_{1,2} & \omega_{1,4} \\ \omega_{2,3} & \omega_{3,4} \end{vmatrix} + \begin{vmatrix} \omega_{1,2} & \omega_{1,3} \\ \omega_{2,4} & \omega_{3,4} \end{vmatrix} = 0.$$

This identity implies that working in a weak sense we can factor out the null-vector

$$(b_1^* b_2^* c_4^* c_3^* - b_1^* b_3^* c_4^* c_2^* + b_2^* b_3^* c_4^* c_1^*) \cdot \mathbf{I}.$$

We look for a general formula for such null-vectors. Consider the operator

$$C = \sum_{i=1}^{\infty} c_i^* b_i,$$

and the space \mathfrak{H}_2 spanned by $\mathbf{b}_I^* \mathbf{c}_J^* \cdot \mathbf{I}$ with $\#(I) = \#(J) + 2$. Then the subspace $\mathfrak{N} = C\mathfrak{H}_2$ of the space \mathfrak{H}_2 consists of null-vectors. Indeed, by taking linear combinations, the formula (3.5) can be rewritten as

$$Z(\mathcal{O}, \Psi) = I^* e^{\Omega} \mathcal{O},$$

where $\Omega = \sum_{i,j} \omega_{i,j} b_i c_j$ and the left vacuum I^* is annihilated by the right action of creation operators. The fact that \mathfrak{N} consists of null-vectors follows from the commutation of C with Ω . We do not have a formal proof that belonging to \mathfrak{N} is necessary for a vector to be a null-vector, but since the computer does not contradict this, we shall assume it is true.

Let us summarise what we have for the moment. In the weak sense for any local operator \mathcal{O} acting on N sites we have

$$(3.6) \quad \mathcal{O} \equiv \sum_{\alpha} C_{\alpha} v_{\alpha},$$

where v_α is a basis of the space

$$(3.7) \quad \mathfrak{V}^{(N)} = \bigcap_{i=N+1}^{2N+1} \text{Ker} \left(Q_i|_{\mathfrak{H}_0^{(N)}/\iota/\mathfrak{M}} \right).$$

the dimension of this space is much smaller than the dimension of $\mathfrak{H}_0^{(N)}/\iota$. We shall see that it is possible to reduce the space further, but with $\mathfrak{V}^{(N)}$ we can already start working.

4. EXPLICIT COMPUTATIONS FOR SMALL MATSUBARA CHAIN

The main goal of this section is to show how to produce a large number of Matsubara data, and subsequently compute the matrix $(\omega_{i,j})$ and expectation values using them.

Suppose we are given some Matsubara parameters $\tau_1, s_1, \dots, \tau_n, s_n$. The eigenvectors $|\Psi\rangle$ are parametrised by Bethe roots β_1, \dots, β_m . They satisfy the Bethe equations which are conveniently written in terms of the Baxter function $Q(\lambda) = \prod_{j=1}^m (\lambda - \beta_j)$:

$$(4.1) \quad a(\beta_j)Q(\beta_j + 1) + d(\beta_j)Q(\beta_j - 1) = 0, \quad j = 1, \dots, m,$$

where

$$a(\lambda) = \prod_{\mathbf{m}=1}^{\mathbf{n}} (\lambda - \tau_{\mathbf{m}} - s_{\mathbf{m}}), \quad d(\lambda) = \prod_{\mathbf{m}=1}^{\mathbf{n}} (\lambda - \tau_{\mathbf{m}} + s_{\mathbf{m}}).$$

Recall $s_{\mathbf{m}}$ are given integers, however the subsequent formulae are all analytical in $s_{\mathbf{m}}$, hence we may consider them as arbitrary complex parameters. Then $a(\lambda), d(\lambda)$ will be considered as arbitrary monic polynomials of degree \mathbf{n} :

$$(4.2) \quad a(\lambda) = \lambda^{\mathbf{n}} + \sum_{j=1}^{\mathbf{n}} a_j \lambda^{\mathbf{n}-j}, \quad d(\lambda) = \lambda^{\mathbf{n}} + \sum_{j=1}^{\mathbf{n}} d_j \lambda^{\mathbf{n}-j}.$$

The usual problem of the theory of quantum integrable models is to find solutions to Bethe equations for the input data

$$(4.3) \quad \text{input} = \{a_1, \dots, a_{\mathbf{n}}, d_1, \dots, d_{\mathbf{n}}\}.$$

Now solving the Bethe equations we would obtain an equivalent set of parameters to the previously introduced

$$(4.4) \quad \text{Matsubara data} = \{\{\tau_{\mathbf{m}}\}, \{s_{\mathbf{m}}\}, \Psi\} \equiv \{\beta_1, \dots, \beta_m, a_1, \dots, a_{\mathbf{n}}, d_1, \dots, d_{\mathbf{n}}\}.$$

For certain well-known reason we set $m \leq [\mathbf{n}/2]$.

The main problem here is that the Bethe equations constitute a complicated system of algebraic equations for β_1, \dots, β_m . But we have to realise that our goal is different from the usual one. We do not need to find the spectrum for a given chain. Rather we need a stock of solutions of the Bethe equations in order to obtain equations for the OPE coefficients. That is why we make the following new choice of input data

$$(4.5) \quad \text{input} = \{\beta_1, \dots, \beta_m, a_{m+1}, \dots, a_{\mathbf{n}}, d_1, \dots, d_{\mathbf{n}}\}.$$

The number of continuous parameters is the same, but the unknowns are now a_1, \dots, a_m . Note that for them the Bethe equations are *linear*. Solving them is easy, and we obtain as a result some Matsubara data (4.4).

Given some Matsubara data, we need first of all to construct the infinite matrix $\omega_{i,j}$. The latter is conveniently coded by the generating function

$$(4.6) \quad \omega(\lambda, \mu) := \sum_{i,j=1}^{\infty} \lambda^{i-1} \mu^{j-1} \omega_{i,j}.$$

Let us define the latter in the spirit of [10]. Introduce the kernel and “half-kernel” functions:

$$K(\lambda) = -\frac{2}{\lambda^2 - 1}, \quad H(\lambda) = \frac{1}{(\lambda - 1)\lambda}.$$

and the measure

$$dm(\lambda) = \frac{d\lambda}{1 + \mathfrak{a}(\lambda)}, \quad \mathfrak{a}(\lambda) = \frac{a(\lambda)Q(\lambda + 1)}{d(\lambda)Q(\lambda - 1)}.$$

We need an auxiliary function defined by the integral equation

$$G(\eta, \mu) = H(\eta - \lambda) - \frac{1}{2\pi i} \oint_{\Gamma} K(\eta - \sigma)G(\sigma, \mu)dm(\sigma),$$

where the contour Γ goes around the Bethe roots β_1, \dots, β_m and the point $\sigma = \mu$. For a finite Matsubara chain we have a finite number of Bethe roots for which the equation above reduces to a linear system for $G(\beta_j, \mu)$. The function $\omega(\lambda, \mu)$ is given by

$$\omega(\lambda, \mu) = \frac{1}{2\pi i} \oint_{\Gamma'} H(\eta - \lambda)G(\eta, \mu)dm(\eta) - \frac{1}{4}K(\lambda - \mu),$$

with Γ' containing one more point: $\eta = \lambda$. It is easy to see that this function is symmetric. Finally, extracting the coefficients of $\omega(\lambda, \mu)$ from (4.6) allows to compute the right hand side of

$$(4.7) \quad \langle \mathcal{O} \rangle_{\text{Md}} = \sum_{\alpha} C_{\alpha} \langle v_{\alpha} \rangle_{\text{Md}},$$

where Md stands for Matsubara data and $\{v_{\alpha}\}$ is the reduced fermion basis (3.6).

The left hand side of (4.7) is computed by QISM techniques. Let us begin by introducing the main character of QISM, the *monodromy matrix*. We use the block decomposition with respect to the first tensor component

$$(\pi_{\lambda}^1 \otimes \pi_{\mathbf{M}})(\mathcal{R}) = \begin{pmatrix} A(\lambda) & B(\lambda) \\ C(\lambda) & D(\lambda) \end{pmatrix},$$

with $A(\lambda), B(\lambda), C(\lambda), D(\lambda)$ acting in the Matsubara space. We shall denote $A = A(0)$, etc. We want to compute

$$\langle \Psi | \text{Tr}_{[1, N]} (T_{[1, N], \mathbf{M}} \mathcal{O}) | \Psi \rangle,$$

for \mathcal{O} located on sites $[1, N]$. For given Matsubara data we write

$$\begin{aligned} \langle \Psi | &= \langle \beta_1, \dots, \beta_m | = \langle \downarrow | B(\beta_1) \cdots B(\beta_m), \\ | \Psi \rangle &= | \beta_1, \dots, \beta_m \rangle = C(\beta_1) \cdots C(\beta_m) | \downarrow \rangle. \end{aligned}$$

The normalisation is provided by the Gaudin formula below. Obviously, for given \mathcal{O} the expression $\text{Tr}_{[1, N]} (T_{[1, N], \mathbf{M}} \mathcal{O})$ is certain linear combination of monomials of A, B, C, D of total degree N . So, we have to learn how to compute their averages in an efficient way.

We start with the Slavnov formula for the scalar product of the Bethe covector $\langle \beta_1, \dots, \beta_m |$ with an off-shell vector $|\mu_1, \dots, \mu_m\rangle = C(\mu_1) \cdots C(\mu_m) | \downarrow \rangle$, μ_j being arbitrary. The formula is

$$(4.8) \quad \langle \beta_1, \dots, \beta_m | \mu_1, \dots, \mu_m \rangle = \prod_{j=1}^m d(\beta_j) d(\mu_j) \frac{\prod_{i,j=1}^m (\beta_j - \mu_i + 1)}{\prod_{i < j} (\mu_i - \mu_j) \prod_{i < j} (\beta_j - \beta_i)} \det(\mathcal{N}),$$

where the matrix \mathcal{N} has entries

$$N_{i,j} = \frac{1}{(\mu_i - \beta_j)} \left(\frac{1}{(\mu_i - \beta_j - 1)} - \frac{1}{(\mu_i - \beta_j + 1)} \mathfrak{a}(\mu_i) \right).$$

The left hand side of (4.8) is by definition a symmetric polynomial of μ_1, \dots, μ_m , hence the singularities in the right hand side cancel. We shall return to this point later.

The normalisation of the Bethe vectors is given by the Gaudin formula, expressible via L'Hospital rules from (4.8). Explicitly, we have

$$(4.9) \quad \langle \beta_1, \dots, \beta_m | \beta_1, \dots, \beta_m \rangle = \prod_{j=1}^m a(\beta_j) d(\beta_j) \prod_{i \neq j} \frac{\beta_i - \beta_j + 1}{\beta_i - \beta_j} \det(\mathcal{G}),$$

where the matrix \mathcal{G} has entries

$$G_{k,l} = \frac{\partial}{\partial \beta_l} \log \mathfrak{a}(\beta_k), \quad k, l = 1, \dots, m.$$

We want to compute

$$\langle \beta_1, \dots, \beta_m | X_1 X_2 \dots X_N | \beta_1, \dots, \beta_m \rangle,$$

where X_j is one of A, B, C, D . We proceed as follows. Start with the Slavnov formula. Use the following formulae which are easily obtained by methods of QISM. Consider any Bethe covector $\langle \Phi |$. Introduce the notations $v(\lambda) = 1/\lambda$, $u(\lambda) = v(\lambda) + 1$. Provided

$$f(\mu_1, \dots, \mu_m) = \langle \Phi | \mu_1, \dots, \mu_m \rangle,$$

is known for arbitrary μ_1, \dots, μ_m we have

$$(4.10) \quad \begin{aligned} \langle \Phi | A | \mu_1, \dots, \mu_m \rangle &= \prod_{j=1}^m u(-\mu_j) a(0) f(\mu_1, \dots, \mu_m) \\ &- \sum_{j=1}^k v(-\mu_j) \prod_{r \neq j}^k u(\mu_j - \mu_r) a(\mu_j) f(\mu_1, \dots, \widehat{\mu}_j, \dots, \mu_m, 0), \end{aligned}$$

$$(4.11) \quad \begin{aligned} \langle \Phi | D | \mu_1, \dots, \mu_m \rangle &= \prod_{j=1}^k u(\mu_j) d(0) \prod_{j=1}^k C(\mu_j) | \downarrow \rangle \\ &- \sum_{j=1}^m v(\mu_j) \prod_{r \neq j}^m u(\mu_r - \mu_j) d(\mu_j) f(\mu_1, \dots, \widehat{\mu}_j, \dots, \mu_m, 0), \end{aligned}$$

$$\begin{aligned}
(4.12) \quad \langle \Phi | C | \mu_1, \dots, \mu_m \rangle &= \left(\sum_{j=1}^k v(-\mu_j) \prod_{r \neq j}^k u(-\mu_r) (\mu_r - \mu_j) a(0) d(\mu_j) \right. \\
&\quad \left. + v(\mu_j) \prod_{r \neq j}^k u(\mu_r) (\mu_j - \mu_r) d(0) a(\mu_j) \right) f(\mu_1, \dots, \widehat{\mu_j}, \dots, \mu_m) \\
&\quad + \sum_{j>i} \left(d(\mu_i) a(\mu_j) v(-\mu_i) v(-\lambda) u(\mu_j - \mu_i) \prod_{r \neq i, j}^k u(\mu_r - \mu_i) u(\mu_j - \mu_r) \right. \\
&\quad \left. + a(\mu_i) d(\mu_j) v(-\mu_j) v(\mu_i) u(\mu_i - \mu_j) \prod_{r \neq i, j}^k u(\mu_i - \mu_r) u(\mu_r - \mu_j) \right) \\
&\quad \times f(\mu_1, \dots, \widehat{\mu_i}, \dots, \dots, \widehat{\mu_j}, \dots, \mu_m, 0).
\end{aligned}$$

$$(4.13) \quad \langle \Phi | B | \mu_1, \dots, \mu_m \rangle = f(\mu_1, \dots, \mu_m, 0).$$

Notice that in the last two formulae the number of arguments changes. Using these formulae we compute inductively

$$\langle \beta_1, \dots, \beta_m | X_1 X_2 \dots X_N | \mu_1, \dots, \mu_m \rangle,$$

and then set $\mu_j = \beta_j$, $j = 1, \dots, m$.

Our goal is to implement the formulae above on a computer. To proceed fast we have to avoid symbolic computations as much as possible. The input data are chosen numerically as a set of random integers from 1 to 10. Still in the procedure described above we cannot immediately exclude the symbolic variables μ_j . It has been said that $\langle \beta_1, \dots, \beta_m | \mu_1, \dots, \mu_m \rangle$ is a symmetric polynomial in μ_1, \dots, μ_m . So are the expressions (4.10), (4.13), (4.12), (4.11). However, to arrive at the polynomial form we have to cancel some singularities by factorizing the expressions in the right hand side. Factorisation is a time consuming operation, it should be ultimately avoided. Actually, looking at the Slavnov formula one realises that it can be easily rewritten as a sum of Schur polynomials. Then with some effort one rewrites the right hand sides of (4.10), (4.13), (4.12), (4.11) as operators acting in the basis of Schur polynomials (indexed by Young diagrams). In this way we completely eliminate time-consuming symbolic computations. In fact acting in the space of Young diagrams can be made very fast.

5. FINAL RESTRICTION OF THE NUMBER OF VARIABLES

Now for any given input data we construct the full Matsubara data which allow to compute, as explained in the previous section, the quantities $\langle \mathcal{O} \rangle_{\text{Md}}$ and $\langle v_\alpha \rangle_{\text{Md}}$ in the main formula (4.7).

We start with $\mathcal{O} = \sigma_1^a \sigma_N^a$. It is interesting that in order to fix C_α up to $N = 4$ we do not need to resort to non-trivial excited states Ψ in the Matsubara data to fix the coefficients. Thus for $N = 4$, absolutely trivial computations give results allowing access to the correlation functions in any environment. For $N = 5$ we need one one-particle state Ψ . Starting from $N = 8$ a few two-particle states are needed. We always take more Matsubara data than the number of unknowns, so that the systems of equations to solve for C_α are overdetermined. The compatibility of all these equations is an important check of the validity of our procedure.

We computed up to $N = 9$, and going further was very hard. However, analysing the results we made the following important observation. Decompose the operators v_α in the

fermionic basis to rewrite

$$\mathcal{O} \equiv \sum_{I,J} D_{I,J} b_I^* c_J^* \cdot \mathbf{I}.$$

A priori I, J are only lexicographically ordered, but we observe that the coefficients $D_{I,J}$ vanish unless

$$i_p \leq j_p, \quad p = 1, \dots, \#(I).$$

We shall write $I \preccurlyeq J$.

We want to reduce the number of unknowns using this observation. The naïve idea consists in repeating the procedure of Section 3 starting with the subspace $\widetilde{\mathfrak{H}}_0^{(N)}$ spanned by $b_I^* c_J^* \cdot \mathbf{I}$ satisfying (3.3), (3.2) and $I \preccurlyeq J$. However, the resulting space (3.7) would be too small, there would be no solutions to the equations for C_α . More careful consideration and further experiments show that the correct choice is the subspace $\widetilde{\mathfrak{V}}^{(N)}$ of $\widetilde{\mathfrak{H}}_0^{(N)}$ such that

$$\bigcap_{i=N+1}^{2N+1} \text{Im}(Q_i|_{\widetilde{\mathfrak{V}}^{(N)}}) \subset \mathfrak{N}.$$

This results in a drastic reduction of the dimension of the space. The dimensions up to $N = 12$ are given in the table below.

N	2	3	4	5	6	7	8	9	10	11	12
$\dim(\widetilde{\mathfrak{H}}_0^{(N)})$	1	3	11	26	99	253	1038	2816	12041	34062	148630
$\dim(\widetilde{\mathfrak{V}}^{(N)})$	1	2	6	12	31	79	178	434	1141	2946	7888

6. RESULTS AND CONJECTURES

We computed the OPE coefficients for $\sigma_1^a \sigma_N^a$ and $h_{1,2} h_{N-1,N}$ up to $N = 11$. Starting from $N = 7$ the data becomes too large to be presented here, they are available upon request.

With the OPE coefficients at hand we can compute the correlation functions for any Matsubara data, as well as the corresponding matrix $\omega_{i,j}$ or equivalently the function $\omega(\lambda, \mu)$. In particular, for the anti-ferromagnet at zero temperature and zero magnetic field we have

$$\omega(\lambda, \mu) = -\frac{1}{2} + 2 \log(2) + \sum_{k=1}^{\infty} (\lambda - \mu)^{2k} \left(2\zeta(2k+1)(1 - 2^{-2k}) - \frac{1}{2} \right).$$

Let us present the numerical values for $G^{\sigma\sigma}$. Up to $N = 8$ they are known from [11], so, we use the normalisation of this paper $G^{\sigma\sigma}(N) \rightarrow G^{\sigma\sigma}(N)/12$ to simplify the comparison.

N	$G^{\sigma\sigma}(N)/12$
2	-0.147715726853315103139077
3	0.0606797699564353014934941
4	-0.0502486272572352479593931
5	0.0346527769827281656556596
6	-0.0308903666476093257628751
7	0.0244467383279589065417695
8	-0.0224982227633722183770986
9	0.0189734169587321977494075
10	-0.0177751064604679461357958
11	0.0155478493216075886422179

The last two values are already in a good agreement with the asymptotic formula of [12].

We can proceed with computations for finite temperature and other Matsubara environments, however, the most interesting applications of our “experimental” data consists in trying to guess general structure of the OPE coefficients. We have some incomplete results in this direction which we expose now.

Consider the decompositions

$$(6.1) \quad \begin{aligned} \sigma_1^1 \sigma_N^a &\equiv \sum_{I \preccurlyeq J} D_{I,J}^{\sigma\sigma}(N) \mathbf{b}_I^* \mathbf{c}_J^* \cdot \mathbf{I}, \\ h_1 h_{N-1} &\equiv \sum_{I \preccurlyeq J} D_{I,J}^{hh}(N) \mathbf{b}_I^* \mathbf{c}_J^* \cdot \mathbf{I}. \end{aligned}$$

A first question which we would like to address is the behaviour of the coefficients as functions of N . The experimental data show that they are polynomial in N of degrees

$$(6.2) \quad d_{I,J}^{\sigma\sigma} = \frac{1}{2}(|I| + |J|) + \#(I) - 1, \quad d_{I,J}^{hh} = \frac{1}{2}(|I| + |J|) + \#(I) - 4.$$

Moreover, in these polynomials not all the coefficients are independent, we find that they are of the form

$$(6.3) \quad D_{I,J}^{\sigma\sigma}(N) = \sum_{s=l_{I,J}^{\sigma\sigma}}^{d_{I,J}^{\sigma\sigma}} X_{s,I,J}^{\sigma\sigma} \binom{N-2}{s}, \quad D_{I,J}^{hh}(N) = \sum_{s=l_{I,J}^{hh}}^{d_{I,J}^{hh}} X_{s,I,J}^{hh} \binom{N-4}{s}.$$

The lower limits $l_{I,J}^{\sigma\sigma}$, $l_{I,J}^{hh}$ are rather subtle. Lower bounds for them are

$$l_{I,J}^{\sigma\sigma} \geq \max(J) - 2, \quad l_{I,J}^{hh} \geq \max(J) - 4.$$

These estimates show that the coefficients vanish if $\max(J)$ goes far from the rest of elements of $I \cup J$. However, if two elements of J become large no vanishing happens. There are some additional simple reason for the coefficients to vanish which will be clear from examples below. In certain cases we were even able to make general conjectures for the coefficients. Let us present them.

We begin with the simple case of $X_{s,I,J}^{\sigma\sigma}$ (coefficients in (6.3)) for the two-fermion case with $I = \{k-p\}$ and $J = \{k+p\}$. In that case we have $3-p \leq 3$ terms in our formula. Since the space \mathfrak{N} is empty for the zero fermion case we have for any s

$$(6.4) \quad \sum_p X_{s,\{k-p\},\{k+p\}}^{\sigma\sigma} = 0.$$

In particular, for $s = k-2$ there is only one term in this sum. Hence $X_{k-2,\{k\},\{k\}}^{\sigma\sigma} = 0$, this is the above mentioned additional reason for coefficients to vanish. For nontrivial coefficients it is easy to guess from our data that

$$X_{k,\{k-p\},\{k+p\}}^{\sigma\sigma} = 2(-1)^{p-1} \left(\binom{2}{p} + 2 \binom{1}{p} \right), \quad X_{k-1,\{k-p\},\{k+p\}}^{\sigma\sigma} = 2(-1)^{p-1} \binom{1}{p}.$$

Both coefficients vanish for $p > 2$ in agreement with the previous explanation.

It so happens that all the two fermions contribution to $G^{hh}(N)$ vanish. The next simplest case is that of four fermions contribution. For that case we use for the sets $I = \{k-p, i\}$, $J = \{k+p, j\}$ for $i+j$ even and $I = \{k-p, i\}$, $J = \{k+p+1, j\}$ for $i+j$ odd. For fixed i, j we shall vary k and p within the obvious limits. Non-vanishing coefficients X^{hh} for some (k, p) correspond to boxes in a triangular array (see *Tab. 1* for $i = 1, j = 1$). The vertical and horizontal sizes of the array are equal to $S = \left\lceil \frac{i+j}{2} \right\rceil + 3$. Counting the rows by $p = 0, \dots, S-1$, and the columns by $r = 0, \dots, S-1$, then let $s = r + k - 4$ for $i+j$ even and $s = k + r - 3$ for $i+j$ odd. For given i, j we shall

simplify notations: define $X_{r,p}(k) := X_{s,\{k-p,1\},\{k+p,1\}}^{hh}$ with r and s related as explained. Sums over columns equal zero for the same reason as explained above, so, we do not write down $X_{r,r}(k)$. Notice, in particular, that $X_{0,0}(k) = 0$.

We start from the simplest case $i = j = 1$. Non-vanishing coefficients correspond to boxes in the table

•			
	•		
		•	
			•

Tab. 1

So, we shall not write formulae for the boxes with bullets for the reasons which have been explained. The coefficients grow rather fast with k , so, at the first glance there is no hope to find a formula for them. However, looking at them attentively, we observed that the simplest one (the second from the left in the first row) is just

$$X_{1,0}(k) = \frac{1}{6}(3^{k-2} - 1).$$

Based on this observation and the experimental data we guessed further

$$\begin{aligned} X_{2,0}(k) &= \frac{5}{8} \cdot 3^{k-1} + \frac{(2k^2 + 4k - 45)}{24}, \\ X_{3,0}(k) &= \frac{5}{4} \cdot 3^{k-1} + \frac{2k^2 + 8k - 49}{12}, \\ X_{2,1}(k) &= -\frac{5}{2} \cdot (3^{k-2} - 1), \\ X_{3,1}(k) &= -\frac{5}{8} \cdot 3^k - \frac{2k^2 + 8k - 159}{24}, \\ X_{3,2}(k) &= \frac{27}{4} \cdot (3^{k-3} - 1) - \frac{(k-3)(k+7)}{6}. \end{aligned}$$

For certain reason $X_{3,2}(3) = 0$. This is made explicit in the last line. We shall continue making such vanishings explicit.

Let us consider now the case $i = 1$, $j = 2$. The table is the same as for the previous case. From experimental data we guess the formulae

$$\begin{aligned} X_{1,0}(k) &:= -\frac{k-2}{24} \cdot (3^{k-1} - 1) - \frac{(k-1)(k+2)}{12}, \\ X_{2,0}(k) &= -\frac{5(k-1)}{16} \cdot 3^{k-1} - \frac{2k^3 + 18k^2 + 23k + 21}{48}, \\ X_{3,0}(k) &:= -\frac{5k}{32} \cdot 3^k - \frac{6k^3 + 52k^2 + 89k + 128}{96}, \\ X_{2,1}(k) &= \frac{5(k-1)}{32} \cdot 3^k + \frac{2k^3 + 38k^2 + 57k + 63}{96}, \\ X_{3,1}(k) &:= \frac{k}{32} \cdot 3^{k+2} + \frac{2k^3 + 20k^2 + 31k + 64}{32}, \\ X_{3,2}(k) &= -\frac{5}{32}k \cdot 3^k + \frac{2k^3 + 4k^2 + 23k - 64}{96}. \end{aligned}$$

At this point one can think that the general formula consists of combinations with polynomial coefficients of 1 and 3^k . Actually, this is not the case: we were able to fix several additional coefficients observing that 6^k starts to appear.

Consider $i = 2, j = 2$. In this case our data allow to define only three coefficients:

$$\begin{aligned} X_{1,0}(k) &= \frac{1}{125} \cdot (6^{k-1} - 1) - \frac{(2k+1)}{144} \cdot (3^{k-1} - 1) + \frac{(k-1)(45k-37)}{1800}, \\ X_{2,0}(k) &= \frac{49}{250} \cdot (6^{k-1} - 1) + \frac{(k^2 - 19k - 41)}{96} \cdot (3^{k-1} - 1) \\ &\quad + \frac{(k-1)(25k^2 + 210k + 179)}{1200}, \\ X_{2,1}(k) &= -\frac{98}{375} \cdot (6^{k-1} - 1) + \frac{(6k+13)}{24} \cdot (3^{k-1} - 1) - \frac{(45k+43)(k-1)}{300}. \end{aligned}$$

Similarly for $i = 1, j = 3$:

$$\begin{aligned} X_{1,0}(k) &= -\frac{48}{625} \cdot (6^{k-2} - 1) + \frac{(7k-11)}{60} \cdot (3^{k-2} - 1) - \frac{(k-2)(5k-53)}{750}, \\ X_{2,0}(k) &= -\frac{144 \cdot 49}{625} \cdot (6^{k-3} - 1) - \frac{27(2k^2 - 15k - 18)}{720} \cdot (3^{k-3} - 1) \\ &\quad + \frac{(k-3)(25k^2 - 465k + 10054)}{3000}, \\ X_{2,1}(k) &= \frac{32 \cdot 49}{625} \cdot 6^{k-2} - 2k \cdot 3^{k-2} + \frac{75k^2 + 1105k - 964}{1875}. \end{aligned}$$

Finally, we were able to guess one coefficient for each $i = 2, j = 3$ and $i = 1, j = 4$. They are respectively

$$\begin{aligned} X_{1,0}(k) &= -\frac{6(k-4)}{625} \cdot (6^{k-1} - 1) - \frac{(4k^2 - 9k - 8)}{240} \cdot (3^{k-1} - 1) \\ &\quad - \frac{(k-1)(65k^2 + 91k + 96)}{3000}, \\ X_{1,0}(k) &= \frac{12(k-4)}{625} \cdot (6^{k-1} - 1) + \frac{k^2 + 9k - 40}{120} \cdot (3^{k-1} - 1) \\ &\quad + \frac{(k-1)(5k^2 + 47k + 132)}{500}. \end{aligned}$$

An important question is that of the general structure. Is it true that the next exponent to appear will be 9^k ? This should be possible to answer considering the cases $i+j = 6$ for which we observe experimentally faster growth than 6^k . Unfortunately, our experimental data do not allow to guess a general formula for this case.

For four fermions contributions to $G^{\sigma\sigma}$ the coefficients are more complicated, we shall not present them because we know only very few. We would mention, however, that the structure looks similar, but 6^k starts to appear from the very beginning.

7. CONCLUSION

We have demonstrated the power of the method to fix the OPE coefficients for the XXX model based on the fermionic basis. Somebody with better computer skills than ours may probably produce some more data. For example $N = 12$ should be reachable.

The main problem which we could not solve for the moment is finding general formulae for the coefficients even for the four fermion contributions to G^{hh} . For example, the coefficient $X_{r,p}(k)$ is known for $\{i, j\} = \{1, 1\}, \{1, 2\}, \{1, 3\}, \{1, 4\}, \{2, 2\}, \{2, 3\}$, but still

we were unable to find a conjecture for general i, j . Probably, an independent look is needed?

ACKNOWLEDGEMENTS

PDF is partially supported by the Morris and Gertrude Fine endowment. FS would like to thank H. Boos, M. Jimbo and T. Miwa for stimulating discussions.

REFERENCES

- [1] M. Jimbo, T. Miwa, F. Smirnov, Hidden Grassmann structure in the XXZ model V: sine-Gordon model. *Lett. Math. Phys.*, **96** (2011) 325–365
- [2] S. Negro, F. Smirnov On one-point functions for sinh-Gordon model at finite temperature *Nuclear Physics B* **875(FS)** (2013) 166–185
- [3] L.D. Faddeev, E.K. Sklyanin, L.A. takhtajan The quantum inverse problem method *Theoretical and Mathematical Physics* **40** (1980) 688–711
- [4] V.E. Korepin, N.M. Bogoliubov, A.G. Izergin Quantum inverse scattering method and correlation functions *Cambridge University Press* (1993)
- [5] N.A. Slavnov Calculation of scalar products of wave functions and form factors in the framework of the alcebraic Bethe ansatz *Theoretical and Mathematical Physics* **79** (1989) 502–508
- [6] M. Jimbo, T. Miwa, and F. Smirnov. Hidden Grassmann structure in the XXZ model III: Introducing Matsubara direction. *J. Phys. A* **42** (2009) 304018 (31pp)
- [7] H. Boos, M. Jimbo, T. Miwa, F. Smirnov, and Y. Takeyama. Hidden Grassmann structure in the XXZ model II : Creation operators. *Commun. Math. Phys.* **286** (2009) 875–932.
- [8] H. Boos, M. Jimbo, T. Miwa, F. Smirnov. Completeness of a fermionic basis in the homogeneous XXZ model. *J. Math. Phys.*, **50** (2009) 095206
- [9] H. Boos, M. Jimbo, T. Miwa, F. Smirnov, and Y. Takeyama. Fermionic basis for space of operators in the XXZ model. *SISSA Proceedings of Science* (2007), Paper 015, 34
- [10] H. Boos, F. Göhmann, A. Klümper and J. Suzuki, Factorization of the finite temperature correlation functions of the XXZ chain in a magnetic field, *J. Phys. A* **40** (2007) 10699–10727.
- [11] J. Sato, M. Shiroishi, M. Takahashi Exact evaluation of density matrix elements for the Heisenberg chain *J.Stat.Mech.* **0612** (2006) P12017
- [12] S. Lukyanov, V. Terras. Long-distance asymptotics of spinDspin correlation functions for the XXZ spin chain *Nuclear Physics B* **654** (2003), 323–356

PHDF: IPHT, UNIV. PARIS SACLAY, CNRS, CEA,, F-91191 GIF-SUR-YVETTE, FRANCE, AND DEPT OF MATHEMATICS, UNIV. OF ILLINOIS MC-382,, URBANA, IL 61821, U.S.A.

E-mail address: philippe.difrancesco@gmail.com

FS¹ SORBONNE UNIVERSITÉ, UPMC UNIV. PARIS 06, CNRS, UMR 7589, LPTHE, F-75005, PARIS, FRANCE

E-mail address: smirnov@lpthe.jussieu.fr

¹Membre du CNRS