Inhibition of P-glycoprotein-mediated multidrug efflux by aminomethylene and ketomethylene analogs of reversins

Ali Koubeissi, Imad Raad, Laurent Ettouati, David Guilet, Charles Dumontet, Joëlle Paris

► To cite this version:

Ali Koubeissi, Imad Raad, Laurent Ettouati, David Guilet, Charles Dumontet, et al.. Inhibition of P-glycoprotein-mediated multidrug efflux by aminomethylene and ketomethylene analogs of reversins. *Bioorganic and Medicinal Chemistry Letters*, 2006, 16 (21), pp.5700 - 5703. 10.1016/j.bmcl.2006.07.059 . hal-01653673

HAL Id: hal-01653673

<https://hal.science/hal-01653673>

Submitted on 15 Feb 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Inhibition of P-glycoprotein-mediated multidrug efflux by aminomethylene and ketomethylene analogs of reversins

Ali Koubeissi,^a Imad Raad,^a Laurent Ettouati,^a David Guilet,^a Charles Dumontet^b and Joelle Paris^{a*}

^a *Université Claude Bernard Lyon 1, Institut des Sciences Pharmaceutiques et Biologiques, EA 3741 Écosystèmes et Molécules bioactives, 69373 Lyon cedex 08, France*

^b *Université Claude Bernard Lyon 1, Faculté de Médecine, Laboratoire de Cytologie analytique, INSERM U590, 69373 Lyon cedex 08, France*

This is where the receipt/accepted dates will go; Received Month XX, 2000; Accepted Month XX, 2000 [BMCL RECEIPT]

Abstract— Several aminomethylene analogs and a ketomethylene analog of reversins were synthesized in order to evaluate their ability to inhibit P-glycoprotein-mediated drug efflux in K562/R7 human leukemic cells overexpressing P-glycoprotein. These analogs retained good activity compared to cyclosporin A and the original reversins. ©2000 Elsevier Science Ltd. All rights reserved.

Multidrug resistance (MDR) to anticancer agents remains a major cause of treatment failure in cancer chemotherapy. MDR describes the cross-resistance of tumor cell lines to several structurally unrelated chemotherapeutic agents after exposure to a single cytotoxic drug. This phenomenon is often associated with overexpression of several proteins.¹ Among them P-glycoprotein (Pgp) is the most important one that belongs to the ABC superfamily of transporters which acts as a drug efflux pump.^{2,3}

Numerous molecules have shown some activity on Pgp.⁴ Among them short linear hydrophobic peptides were described as chemosensitizers.^{5a} Seprödi and coll. showed that small hydrophobic peptide derivatives modulate Pgp-ATPase activity and inhibited the drug extrusion function of Pgp.^{5b,c} These compounds are a family of di- and tripeptide derivatives sharing some common physico-chemical and structural features. Some of them are dimerized aminoacids by diacid derivatives. The enhanced affinity to Pgp of these chemosensitizers finally coined reversins is ascribed to the hydrophobic nature of the side chains protected with bulky aromatic or alkyl groups.⁶ Among them reversin

121 **1a** showed the highest affinity and specificity for Pgp.

1a $n = 1$, $R^1 = \text{Bn}$

1b $n = 2$, $R^1 = \text{Dmp}$

1c $n = 2$, $R^1 = \text{cHex}$

At 1-2 μM this reversin was more effective than cyclosporin A for blocking colchicine transport in isolated membranes and reconstituted systems. In order to improve proteolytic stability and bioavailability of reversins we planned to synthesize aminomethylene and ketomethylene analogs of typical reversin representatives. We chose reversin 121 **1a** as our working model for dipeptide-type reversins and reversin 213 **2a** as a dimerized aminoacid containing succinyl unit. The reduced analogs **3a-b** were synthesized using the classical reductive amination

* Corresponding authors : fax: +33 4 78 77 75 49; e-mail: laurent.ettouati@univ-lyon1.fr, joelle.paris@univ-lyon1.fr

Scheme 1. Reagents and conditions: (a) NaBH₃CN, MeOH, AcOH, rt 1 h, 52% (n = 1) and 57% (n = 2).

Scheme 2. Reagents and conditions: (a) *N*-Hydroxysuccinimide, DMAP, DCC, THF, 0°C 10 min then rt 48 h, 94%; (b) DIEA, DMF, rt 24 h, quant yield; (c) OsO₄ 2.5% in *t*-BuOH, NMO, THF, H₂O, rt 24 h, 86%; (d) NaIO₄, THF, H₂O, 3 h, quant yield; (e) NaBH₃CN, MeOH, AcOH, rt 1-2 h.

strategy⁷ starting from chiral pool derived aminoaldehydes according to Scheme 1.

As far as reduced analogs of reversin 213 **2a** were concerned we started from 4-pentenoic acid which was activated as succinimidyl ester and then coupled to L-glutamic acid dibenzyl ester tosylate salt to obtain the amide **4** (Scheme 2). The corresponding aldehyde **6** was obtained by a two-step reaction with osmium tetroxide/4-methylmorpholine *N*-oxide⁹ followed by oxidative cleavage of the diol **5** with sodium periodate. It is noteworthy that aldehyde **6** exists in equilibrium with the corresponding hemiaminals **6'** as previously described in similar reactions.^{10a,b} Structure of the diastereoisomeric hemiaminals **6'** was assessed by ESIMS and HSQC and HMBC experiments.^{10c} The last step consisted in obtaining the reduced analogs **8a-d** of reversin 213 **2a**. However in the standard reductive amination conditions, the reaction between diprotected L-glutamic acid **7b** and aldehyde **6** did not afford the corresponding secondary amine **8b** but instead the pyrrolidinone **9** in 38% yield (Figure 1). This result is explained by nucleophilic attack of the generated secondary amine on the adjacent benzyl ester side chain as observed for similar derivatives.¹¹ So as to avoid

cyclization several protected derivatives of L-glutamic acid such as the 2,4-dimethyl-3-pentyl^{12a} and cyclohexyl^{12b} esters known to prevent aspartimide formation, were used to obtain respectively analogs **8c** and **8d**.^{12c} Moreover by using an excess of aldehyde **6** a double addition product **10** was obtained in 47 % yield in the case of reductive amination of L-glutamic acid derivative **7c** (Figure 1).¹³ Derivatives **1b** and **1c** along with compound **2b** were also synthesized by standard procedures for sake of comparison.

We thought it would be worthwhile to test the introduction of a spacer between the Asp and Lys residue of reversin 121 **1**. Thus, we chose to synthesize the protected Asp-ψ(CO-CH₂)Gly-Lys ketomethylene analog **16** as outlined in Scheme 3. The synthesis started from L-homoserine which was protected on the side chain as a *tert*-butyl dimethyl silyl ether followed by protection of the free α-amine as *tert*-butyloxycarbonyl derivative and finally transformed in Weinreb amide **11** in 40% overall yield over three steps. The alkene **12** was then obtained from Weinreb amide **11** by alkylation with butenyl magnesium bromide as described¹⁴ in 55% yield. Transformation of the protected hydroxyl of **12** in benzyl ester **13** was carried out by oxidation of the deprotected hydroxyl with PDC and then esterification of the cesium salt in 53% overall yield as described.^{15a,15b} Oxidation of alkene **13** with RuCl₃·xH₂O/NaIO₄ afforded free acid **14** in 96% yield. The ketomethylene analog **16** was finally obtained by coupling the succinimidyl derivative **15** and diprotected L-lysine hydrochloride in 94% yield.

Figure 1. Structures of by-products **9** and **10** obtained respectively in the reductive amination of **7b** and **7c** with aldehyde **6**.

Scheme 3. Reagents and conditions: (a) TBDMSCl, DBU, CH₃CN, 0°C then rt 24 h, 82%; (b) (Boc)₂O, Et₃N, acetone, H₂O, rt 24 h, 80%; (c) *N,O*-dimethylhydroxylamine hydrochloride, TBTU, HOBt, DIEA, CH₂Cl₂, rt 12 h, 61%; (d) Butenyl magnesium bromide, THF, -78°C then 3.5 h rt 30 min, 55%; (e) TBAF, THF, rt 1 h, 87%; (f) PDC, DMF, rt 3 h, 71%; (g) i. Cs₂CO₃, MeOH, H₂O then reduced pressure ii. Benzyl bromide, DMF; rt 4.5 h, 86%; (h) RuCl₃·xH₂O NaIO₄, H₂O, CH₃CN, rt 1 h, 96%; (i) *N*-Hydroxysuccinimide, DMAP, DCC, THF, 0°C 10 min then rt 48 h, 49%; (j) DIEA, DMF, rt 24 h, 94%.

The efficiency of reversin analogs to inhibit Pgp-mediated daunorubicin efflux was investigated by monitoring the intracellular accumulation of this drug in K562/R7 human leukemic cells overexpressing Pgp in the presence of daunorubicin.^{16a} Cyclosporin A was used as a positive control (Table 1). These analogs showed strong inhibitory activity comparable to that of reversin 121 **1a** except in the presence of the tertiary amine **10** which caused an important decrease in activity. It was noteworthy that the replacement of the amide bond with an amino methylene isostere did not impair the capacity to inhibit Pgp-mediated drug efflux (**1a** versus **3a** and **2b** versus **8d**). Analogs with cyclohexyl ester presented an increased inhibitory activity when compared to 2,4-dimethyl-3-pentyl ester analogs (**1b** versus **1c** and **8c** versus **8d**). Moreover, exchanging benzyl ester by cyclohexyl ester was tolerated as shown in Table 1 by comparing activity for compounds **2a** and **2b**. Finally, the inhibitory activity of ketomethylene analog **16** was conserved indicating that inserting a succinyl moiety into reversin 121 **1a** was not detrimental to the Pgp-mediated drug efflux inhibitory activity.

Table 1. Mean inhibitory activity (%) of reversins and synthesized analogs on human leukemic cells K562/R7.^{16b}

Compds ^a	Mean inhibiting activity % ^b
Reversin 121 1a	72.73 (± 5.27)
1b	52.70 (± 5.27)
1c	85.43 (± 5.69)
Reversin 213 2a	93.74 (± 2.17)
2b	94.45 (± 1.36)
3a	74.14 (± 4.06)
3b	79.08 (± 5.13)
8a	82.22 (± 4.57)
8c	54.71 (± 3.63)
8d	98.33 (± 1.35)
10	3.35 (± 3.44)
16	86.62 (± 4.86)

^aCompounds were tested at a 10 μM concentration.

^bCyclosporin A was used as positive control (mean inhibitory activity of 100%) at a final concentration of 2 μM. Standard deviation is given in parentheses.

We have designed aminomethylene and ketomethylene analogs with inhibitory activity of Pgp-mediated drug efflux comparable to that of reversins 121 **1** and 213 **2a**. We envisage now the synthesis of ketomethylene analogs of reversin 121 and modified side chain analogs to confirm and refine our first results.

References and Notes

- Gottesman, M. M.; Fojo, T.; Bates S. E. *Nature Rev. Cancer* **2002**, 2, 48.
- Gottesman, M. M.; Pastan, I.; Ambudkar, S. V. *Curr. Opin. Genet. Dev.*, **1996**, 6, 610.
- Szakács, G.; Paterson, J. K.; Ludwig, J. A.; Booth-Genthe, C.; Gottesman, M. M. *Nature Rev. Drug Discovery*, **2006**, 5, 219.
- Teodori, E.; Dei, S.; Scapecchi, S.; Gualtieri, F. *II Farmaco*, **2002**, 57, 385.
- (a) Sharom, F. J.; DiDiodato, G.; Yu, X.; Ashbourne, K. J. D. *J. Biol. Chem.* **1995**, 270, 10334. (b) Seprödi, J.; Mezö, I.; Vadász, Zs.; Szabó, K.; Sarkadi, B.; Teplán, I. In *Peptides 1996*; Proceedings of the 24th European Symposium, 1996, Edinburgh; Ramage, R., Epton, R., Eds; Mayflower Scientific Ltd: Kingswinford, 1998; pp 801-802. (c) Vadász, Zs.; Szabó, K.; Sarkadi, B.; Teplán, I.; Mák, M.; Miklós, I.; Györffy, E.; Seprödi, J. In *Peptides 1998*; Proceedings of the 25th European Symposium, 1998, Budapest; Bajusz, S., Hudecz, F., Eds; Akademiai Kiado Ed: Budapest, 1999; pp 640-641.
- Sharom, F. J.; Yu, X.; Lu, P.; Liu, R.; Chu, J. W. K.; Szabó, K.; Müller, M.; Hose, C. D.; Monks, A.; Várad, A.; E.; Seprödi, J.; Sarkadi, B. *Biochem. Pharmacol.*, **1999**, 58, 571.
- Martinez, J.; Bali, J.-P.; Rodriguez, M.; Castro, B.; Magous, R.; Laur, J.; Lignon, M.-F. *J. Med. Chem.*, **1985**, 28, 1874.
- The aminoaldehydes were synthesized by reduction of aminoacids derived mixed anhydrides with NaBH₄ and then oxidized by Swern oxidation.
- VanRheenen, V.; Kelly, R. C.; Cha, D. Y. *Tetrahedron Lett.*, **1976**, 17, 1973.
- (a) Baldwin, J. E.; Hulme, C.; Edwards, A. J.; Schofield, C. J.; Parkes, K. E. B. *Tetrahedron Lett.*, **1993**, 34, 1665. (b) Delcros, J.-G.; Tomasi, S.; Carrington, S.; Martin, B.; Renault, J.; Blagbrough, I. S.; Uriac, P. *J. Med. Chem.*, **2002**, 45, 5098. (c) Spectral data of diastereoisomeric hemiaminals **6'**: IR (film, ν cm⁻¹): 3412 (OH and NH), 3066 (arom CH), 2954 (al CH), 1738 (C=O ester), 1674 (CO amide). ESIMS (negative mode, NaCl): 856,9 (2M+Cl⁻), 446,0 (M+Cl⁻). ¹H NMR (300 MHz, CDCl₃, δ ppm): 7.36 (20H, m, 4 Phe), 5.31 (2H, m, 2 CH(OH)), 5.18 (4H, m, 2 $\text{CH}_2\text{O Bn}$), 5.12 (4H, m, 2 $\text{CH}_2\text{O Bn}$), 4.77 (1H, dd, J = 4.5 and 9.8 Hz, $\text{CH}\alpha$), 4.64 (1H, t, J = 7.1 Hz, $\text{CH}\alpha$), 4.32 (1H, d, J = 4.5 Hz, exch OH), 3.41 (1H, d, J = 7.8 Hz, exch OH), 1.8-2.8 (16H, m, 2 $-\text{CH}_2\text{CH}_2-$ and 2 $-\text{CH}_2\text{CH}_2-$ Glu). ¹³C NMR (75 MHz, CDCl₃, δ ppm): 176.3 (C=O), 175.8 (C=O), 173.7 (C=O), 173.1 (C=O), 172.8 (C=O), 171.0 (C=O), 136.0 (Cq Phe), 135.9 (Cq Phe), 135.5 (Cq Phe), 135.2 (Cq Phe), 129.1 (CH Phe), 129.07 (CH Phe), 129.02 (CH Phe), 128.9 (CH Phe), 128.8 (CH Phe), 128.7 (CH Phe), 128.6 (2 CH Phe), 84.4 (CH(OH)), 82 (CH(OH)), 68.3 (CH_2O), 67.9 (CH_2O), 67.1 (CH_2O), 67.0 (CH_2O), 54.7 ($\text{CH}\alpha$), 54.3 ($\text{CH}\alpha$), 31.3 (CH_2), 30.0 (CH_2), 29.5 (CH_2), 29.3 (CH_2), 29.0 (CH_2), 28.8 (CH_2), 25.7 (CH_2), 24.6 (CH_2).
- Gareau, Y.; Zamboni, R.; Wong, A.W. *J. Org. Chem.*, **1993**, 58, 1582.
- (a) Karlström, A. H.; Undén, A. E. *Tetrahedron Lett.*, **1995**, 36, 3909. (b) Tam, J. P.; Wong, T.-W.; Riemen, M. W.; Tjoeng, F.-S.; Merrifield, R. B. *Tetrahedron Lett.*, **1979**, 42, 4033. (c) H-Glu(ODmp)-OBn **7c** was synthesized from commercial *N*-Boc-Glu-OBn by side chain esterification with DCC/DMAP and 2,4-dimethyl-3-pentanol in dichloromethane in 70% yield and then Boc deprotected with conc H₂SO₄ in AcOEt in 95% yield. H-Glu(OcHex)-OBn **7d** was obtained from commercial *N*-Boc-Glu(OcHex)-OH by esterification of the α -carboxylic acid with benzyl bromide and cesium carbonate in MeOH/H₂O in 70% yield and then Boc deprotected in the same conditions as previously in 96% yield.
- See Salvi, J. P.; Walchshofer, N.; Paris, J. *Tetrahedron Lett.*, **1994**, 35, 1181 for similar formation of double condensation product in reductive amination reactions.
- Kaiser, M.; Siciliano, C.; Assfalg-Machleidt, I.; Groll, M.; Milbradt, A. G.; Moroder, L. *Org. Lett.*, **2003**, 5, 3435.
- (a) Våbenø, J.; Nielsen, U.; Ingebrigtsen, T.; Lejon, T.; Steffansen, B.; Luthman, K. *J. Med. Chem.*, **2004**, 47, 4755. (b) As racemisation of a similar benzylated ester was noted in the cited reference 15a the enantiomeric purity of **13** was checked and found enantiomerically pure by C18 RP-HPLC analysis after deprotection (TFA/CH₂Cl₂) and derivatisation with GITC.
- (a) Comte, G.; Daskiewicz, J.-P.; Bayet, C.; Conseil, G.; Viorner-Vanier A.; Dumontet, C.; Di Pietro, A.; Barron, D. *J. Med. Chem.*, **2001**, 44, 763. (b) One million K562/R7 human leukemic cells expressing high levels of P-glycoprotein were incubated for 1 hour at 37°C in 1mL of RPMI 1640 medium containing a final concentration of 10 μ M daunorubicin, in the presence or absence of inhibitor. The cells were then washed twice with ice-cold phosphate buffer saline (PBSt), and kept on ice until analysis by flow cytometry on a FACS-II. Assays were performed in duplicate, with at least three separate experiments.