



HAL
open science

New Insights into the Anti-pathogenic Potential of *Lactococcus garvieae* against *Staphylococcus aureus* Based on RNA Sequencing Profiling

Pierre Delpech, Etienne Rifa, Graham Ball, Sabine Nidelet, Emeric Dubois,
Geneviève Gagne, Marie-Christine Montel, Céline Delbes, Stephanie Bornes

► **To cite this version:**

Pierre Delpech, Etienne Rifa, Graham Ball, Sabine Nidelet, Emeric Dubois, et al.. New Insights into the Anti-pathogenic Potential of *Lactococcus garvieae* against *Staphylococcus aureus* Based on RNA Sequencing Profiling. *Frontiers in Microbiology*, 2017, 8, pp.359. 10.3389/fmicb.2017.00359 . hal-01661796

HAL Id: hal-01661796

<https://hal.science/hal-01661796>

Submitted on 12 Dec 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License



New Insights into the Anti-pathogenic Potential of *Lactococcus garvieae* against *Staphylococcus aureus* Based on RNA Sequencing Profiling

OPEN ACCESS

Edited by:

Fliss Ismail,
Laval University, Canada

Reviewed by:

Luis Augusto Nero,
Universidade Federal de Viçosa, Brazil
Young Min Kwon,
University of Arkansas, USA

*Correspondence:

Céline Delbès
celine.delbes@inra.fr

† Present address:

Sabine Nidelet,
Institut National de la Recherche
Agronomique, UMR 1062, Centre
de Biologie pour la Gestion des
Populations, Montferrier-sur-Lez,
France

Pierre Delpech,
Department of Health and Life
Sciences, Glasgow Caledonian
University, Cowcaddens Road,
Glasgow, UK

‡ These authors have contributed
equally to this work.

Specialty section:

This article was submitted to
Food Microbiology,
a section of the journal
Frontiers in Microbiology

Received: 28 September 2016

Accepted: 21 February 2017

Published: 08 March 2017

Citation:

Delpech P, Rifa E, Ball G, Nidelet S,
Dubois E, Gagne G, Montel M-C,
Delbès C and Bornes S (2017) New
Insights into the Anti-pathogenic
Potential of *Lactococcus garvieae*
against *Staphylococcus aureus*
Based on RNA Sequencing Profiling.
Front. Microbiol. 8:359.
doi: 10.3389/fmicb.2017.00359

Pierre Delpech^{1†}, Etienne Rifa¹, Graham Ball², Sabine Nidelet^{3†}, Emeric Dubois³, Geneviève Gagne¹, Marie-Christine Montel¹, Céline Delbès^{1*‡} and Stéphanie Bornes^{1‡}

¹ Université Clermont Auvergne, INRA, UMRF, Aurillac, France, ² John van Geest Cancer Research Centre, School of Science and Technology, Nottingham Trent University, Nottingham, UK, ³ Montpellier GenomiX, Institut de Génomique Fonctionnelle, Montpellier, France

The bio-preservation potential of *Lactococcus garvieae* lies in its capacity to inhibit the growth of *staphylococci*, especially *Staphylococcus aureus*, in dairy products and *in vitro*. *In vitro*, inhibition is modulated by the level of aeration, owing to hydrogen peroxide (H₂O₂) production by *L. garvieae* under aeration. The *S. aureus* response to this inhibition has already been studied. However, the molecular mechanisms of *L. garvieae* underlying the antagonism against *S. aureus* have never been explored. This study provides evidence of the presence of another extracellular inhibition effector *in vitro*. This effector was neither a protein, nor a lipid, nor a polysaccharide, nor related to an L-threonine deficiency. To better understand the H₂O₂-related inhibition mechanism at the transcriptome level and to identify other mechanisms potentially involved, we used RNA sequencing to determine the transcriptome response of *L. garvieae* to different aeration levels and to the presence or absence of *S. aureus*. The *L. garvieae* transcriptome differed radically between different aeration levels mainly in biological processes related to fundamental functions and nutritional adaptation. The transcriptomic response of *L. garvieae* to aeration level differed according to the presence or absence of *S. aureus*. The higher concentration of H₂O₂ with high aeration was not associated with a higher expression of *L. garvieae* H₂O₂-synthesis genes (*pox*, *sodA*, and *spxA1*) but rather with a repression of *L. garvieae* H₂O₂-degradation genes (*trxB1*, *ahpC*, *ahpF*, and *gpx*). We showed that *L. garvieae* displayed an original, previously undiscovered, H₂O₂ production regulation mechanism among bacteria. In addition to the key factor H₂O₂, the involvement of another extracellular effector in the antagonism against *S. aureus* was shown. Future studies should explore the relation between H₂O₂-metabolism, H₂O₂-producing LAB and the pathogen they inhibit. The nature of the other extracellular effector should also be determined.

Keywords: *Lactococcus garvieae*, *Staphylococcus aureus*, hydrogen peroxide, transcriptome, growth inhibition, anti-pathogenic interactions

INTRODUCTION

Interest in Lactic Acid Bacteria (LAB) as bio-preservation agents against foodborne pathogens has been growing for the last 15 years (Ito et al., 2003; Batdorj et al., 2007; Charlier et al., 2009; Adesokan et al., 2010). Considering their beneficial properties, the dairy industry could find it useful to employ *Lactococcus garvieae* strains as starter or adjunct cultures, provided the strains are safe (Fernández et al., 2010). Aquatic strains of *L. garvieae* are frequently referenced as fish pathogens (Eldar and Ghittino, 1999; Vendrell et al., 2006) but other strains are regarded as opportunistic pathogens for humans (Aguado-Urda et al., 2011; Ortiz et al., 2014). Despite its ubiquity in foods such as milk (Devriese et al., 1999; Villani et al., 2001; Callon et al., 2007) and dairy products of various origins (Flórez and Mayo, 2006; Fortina et al., 2007; Alomar et al., 2008b; El-Baradei et al., 2008; Jokovic et al., 2008; Alegria et al., 2009; Sip et al., 2009; Monfredini et al., 2012; Pangallo et al., 2014; Morandi et al., 2015), to our knowledge, *L. garvieae* has never been involved in a foodborne disease outbreak. *L. garvieae* is of interest for bio-preservation owing to its ability to inhibit the growth of staphylococci, particularly *Staphylococcus aureus*, as has been observed in milk, in cheese and *in vitro* (Alomar et al., 2008a,b; Delbes-Paus et al., 2010; Delpéch et al., 2015). The transcriptome response of *S. aureus* to this inhibition has already been explored (Delpéch et al., 2015). It is associated with a repression of the stress response (especially H₂O₂ response) and of cell division genes and with modulation of the expression of virulence genes (particularly *agrA*, *hld*, and enterotoxin-encoding genes) genes. However, the molecular mechanisms of *L. garvieae* underlying the antagonism against *S. aureus* have never been explored.

Since *L. garvieae* produces low amounts of acetic and lactic acid in milk (Alomar et al., 2008a,b; Nouaille et al., 2009; Sip et al., 2009; Delbes-Paus et al., 2010; Delpéch et al., 2015; Morandi et al., 2015), its antagonism against *S. aureus* is not associated with acidification. It is also unlikely to be associated with nutritional competition (Alomar et al., 2008a). With high aeration, *S. aureus* inhibition is mainly associated with hydrogen peroxide (H₂O₂) production by *L. garvieae*, as already observed *in vitro* (Delbes-Paus et al., 2010; Delpéch et al., 2015). With low aeration, the inhibition is weaker and H₂O₂ is not detected. The addition of catalase (an H₂O₂-degrading enzyme) partly suppressed the inhibition (Delbes-Paus et al., 2010). Delbes-Paus et al. (2010) therefore suggested that at least one other molecule from *L. garvieae* must be involved in the residual inhibition of *S. aureus* in the absence of H₂O₂. The role and nature of this molecule have not yet been determined.

In Gram-positive bacteria, the aeration level and the presence of H₂O₂ generally strongly affect the expression of H₂O₂ metabolism genes. The expression of major H₂O₂ degradation genes (*ahpC*, *ahpF*, *gshR*, *gpo*, *trxA*, and *trxB*) and their related proteins is strongly induced in *L. lactis* (Pedersen et al., 2008) under aeration and in *Bacillus subtilis* in the presence of H₂O₂ (Mostertz et al., 2004). As regards the major H₂O₂ synthesis genes, *sodA* is generally regulated in the same way as H₂O₂-degradation genes while the pyruvate oxidase (*pox*) gene is not affected by these parameters. Biological functions of *L. lactis*

related to aerobiosis, i.e., O₂ response, menaquinone metabolism and stress response, are variably affected by the presence of *S. aureus* (Nouaille et al., 2009). However, little is known about the H₂O₂-metabolism of *L. garvieae* as an anti-pathogenic process.

In this study, we aimed to characterize the antagonism of *L. garvieae* against *S. aureus* *in vitro* in greater depth. Firstly, we investigated the presence of potential antagonist molecules in H₂O₂-free *L. garvieae* supernatants and their impact on *S. aureus* growth. Secondly, the transcriptome of *L. garvieae* with different aeration levels and in different biotic environments (absence or presence of *S. aureus*) was determined through RNA sequencing, an accurate and efficient method for revealing bacterial transcriptome profiles (Pinto et al., 2011). The resulting data led us to a better understanding of *L. garvieae* hydrogen peroxide metabolism and to some initial hypotheses on the nature of a new inhibition effector.

MATERIALS AND METHODS

Strains and Culture Conditions

Lactococcus garvieae N201 and *S. aureus* SA15, isolated from raw milk, were obtained from the INRA UR545 collection (Alomar et al., 2008a). Both strains were aerobically grown in Brain-Heart Infusion broth ("BHI", Biokar Diagnostic, Pantin, France) for 20 h, at 30°C for *L. garvieae* and at 37°C for *S. aureus*. They were then inoculated separately or in co-culture at 10⁶ cells.mL⁻¹ for *S. aureus* and 10⁷ cells.mL⁻¹ for *L. garvieae* into BHI buffered at pH = 7 with phosphate buffer KH₂PO₄, 3H₂O/K₂HPO₄ at 0.1 mol.L⁻¹ (KH₂PO₄, 3H₂O, Riedel-de-Haen, Honeywell GmbH, Seelze, Germany; K₂HPO₄, Merck KGaA, Darmstadt, Germany) previously equilibrated at 30°C. Pure cultures and co-cultures of both strains were performed with either a high or a low aeration depending on the experiment. Low aeration cultures were set in static, fully filled and sealed, 50-mL Nunc EZ Flip conical centrifuge tubes (Sigma-Aldrich, St. Louis, MO, USA). High aeration was obtained by a mechanical shaking at 150 rpm on 50-mL cultures in 250-mL Erlenmeyers. All cultures were incubated at 30°C for 24 h in an Infors HT Minitron (Infors AG, Bottmingen, Switzerland). The cultivable cell counts were determined after plating for each sampling time as described by Delpéch et al. (2015).

Impact of H₂O₂-Free Co-culture Supernatants on *S. aureus* Growth

The potential presence of other antistaphylococcal molecules in *L. garvieae* N201 and *S. aureus* SA15 co-culture supernatant and their impact on *S. aureus* planktonic growth were investigated. After adding catalase at 400 U.mL⁻¹, a pure culture of *S. aureus* SA15, a co-culture of *S. aureus* SA15 and *L. garvieae* N201 and plain BHI were incubated with low aeration as described above. After 6 h of incubation, 40 mL of each tube were centrifuged at 9,600 × g for 10 min at 4°C and supernatants stored at 4°C until utilization. Four milliliters of the remaining *S. aureus* culture were centrifuged at 7,500 × g for 10 min. The cell pellet was resuspended at a concentration of ~10⁶ cells.mL⁻¹ in 20 mL of

fresh 2X-concentrated BHI buffered at pH = 7. The supernatants were filtered through a cellulose acetate membrane (pore size 0.45 μm ; GVS S.p.A., Zola Predosa, Italy). They were either treated with proteinase K (AMRESCO LLC, Solon, OH, USA, ref: 0706-100MG) and pronase E (Merck KGaA, ref: 537088), i.e., with two proteases, or treated with lipase (Sigma-Aldrich, ref: L3126-100G), or treated with α -amylase (Sigma-Aldrich, ref: A3176-500KU), or not treated at all. Each enzyme was added at 0.2 $\text{mg}\cdot\text{mL}^{-1}$ from stock solutions at 10 $\text{mg}\cdot\text{mL}^{-1}$ prepared in 200 mM of phosphate buffer at pH = 7. Treatment with proteases consisted of a first incubation of the supernatants with proteinase K at 50°C for 2 h 15 and a second incubation with pronase E at 37°C for 2 h 15. Treatments with lipase and α -amylase consisted of an incubation of the supernatants with the enzyme at 37°C for 5 h. After each incubation step, the enzyme was inactivated by heating (95°C during 10 min) and all supernatants were filtered again through a cellulose acetate membrane (pore size 0.45 μm ; GVS S.p.A.). Prepared supernatants and *S. aureus* culture in 2X-BHI were then distributed at 1:1 volume ratio (total volume = 1.5 mL) in a CytoOne 24-well cell plate covered with a lid (ref: CC7672-7524; STARLAB, Hambourg, Germany). The cell plate was incubated for 25 h in a SAFAS Xenius XC spectrophotometer (SAFAS Monaco, Monaco) and thermostated at 30°C with a Julabo CryoStat (JULABO GmbH, Seelbach, Germany). *S. aureus* growth in each well was determined by measuring the OD₆₀₀ every 15 min for 1 h and then every 30 min for 24 h. Before each OD₆₀₀ measurement, the cell plate was shaken at 5 Hz for 20 s with an orbital diameter of 6 mm. The whole experimental design was repeated three times. To identify sample means that were significantly different from each other, statistical analyses were performed on values at 9, 12, 15, 18, 21, and 24 h, using Statistica software (StatSoft) with a single-factor analysis of variance (ANOVA) followed by a Newman–Keuls *post hoc* test.

Sample Preparation for RNA Analysis

Pure cultures and co-cultures of *L. garvieae* N201 and *S. aureus* SA15 were grown with high or low aeration, as specified above. After 3, 6, 9, and 24 h of incubation, 40 mL of each culture were centrifuged at 9,600 $\times g$ for 10 min at 4°C. Hydrogen peroxide concentrations and pH values were determined on the supernatants by enzymatic reaction and spectrophotometry as described by Delbes-Paus et al. (2010). The cell pellets were immediately frozen in an ethanol bath and stored at -80°C . Extraction of total RNA from the frozen cell pellets was performed as described by Delpéch et al. (2015). For each sample of total RNA obtained with one culture condition, a first part of the aliquot was used for RNA sequencing after rRNA depletion and a second part of the aliquot was used for RT-qPCR analyses. The whole experimental design was repeated three times.

Determination of *L. garvieae* Transcriptome Changes by RNA Sequencing

Ribosomal RNAs were removed from the total RNA ($2 \times 5 \mu\text{g}$ of RNA by sample) using a RiboZero Magnetic Kit for

Gram Positive Bacteria (Illumina Inc., San Diego, CA, USA) according to the manufacturer's instructions. The quality and concentration of RNA in each sample were assessed using a RNA 6000 pico kit (Agilent Technologies, Santa Clara, CA, USA). The following steps were performed by the MGX Platform (Montpellier GenomiX, CNRS, Montpellier, France) using Illumina kits and devices (Illumina Inc.): construction of the mRNA library using a TruSeq Stranded mRNA Sample Preparation Kit; cluster generation with the cBot system using a Cluster Generation Kit; hybridization of the sequencing primer on the flow-cell; 50-bp single-read sequencing using a HiSeq 2000 device with SBS technology; informatic pretreatments, i.e., image analysis with the HiSeq Control Software and Real-Time Analysis component, base-calling with the RTA software and demultiplexing with CASAVA (Illumina). The RNAseq data are available from NCBI GEO datasets under the accession number GSE74030.

The quality scores across all bases of all reads and the N (non-attributed bases) content across all bases were determined for each condition with the FastQC software from the Babraham Institute¹. Both analyses showed good quality values (data not shown). The reads were next aligned simultaneously on reference genomes, i.e., *L. garvieae* N201 (unpublished, GOLD project ID = Gp0034836 and NCBI BioProject ID = 184287) and *S. aureus* MW2 (RefSeq number = NC_003923.1), using the BWA package (Li and Durbin, 2009) with a seed of 32 bases and a maximum of two mismatches tolerated on the seed. Using the Samtools suite (Li et al., 2009) we excluded from further analyses those reads with low alignment quality scores (MAPQ index < 20). Reads which mapped on multiple sites (between 0.3 and 3.2% depending on the sample) and reads which did not map on any site considering the stringency applied (between 8.7 and 15.5% depending on the sample) were excluded from further analysis (see Supplementary Table 1). Reads overlapping genes were counted with the HTSeq Count software in Union mode (Anders et al., 2015). Differentially expressed genes were identified using the Bioconductor R² packages EdgeR, DESeq and DESeq2 (Anders and Huber, 2010; Robinson et al., 2010; Love et al., 2014). Genes with less than 15 reads (from the three biological replicates of two compared samples) were filtered and thus removed from the analysis. Data were normalized using the Relative Log Expression (RLE) normalization factor for EdgeR and the DESeq normalization factor for DESeq and DESeq2. Gene expression changes with adjusted *p*-value of less than 0.05 (by the FDR method from Benjamini–Hochberg) were declared differentially expressed. Differentially expressed genes highlighted by at least one of the three packages were considered for further investigation.

Genes were sorted into two lists: one of genes differentially expressed in both pure culture and co-culture and one of genes differentially expressed only in pure culture or only in co-culture. These lists were separately subjected to a

¹<http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>

²<http://www.R-project.org/>

Blast2Go analysis (Conesa et al., 2005). Blast2Go analysis consisted first of a BlastX³ on the nr database with 20 hits and a maximum *E*-value of $1.0E^{-15}$. The resulting data were enriched with an InterPro scan analysis (Zdobnov and Apweiler, 2001). Blast hits of each sequence were then mapped with Gene Ontology terms (Gene Ontology Consortium, 2008) annotated with a maximum *E*-value of $1.0E^{-6}$, a cut-off of 55 and a GO weight of 5. The number of genes involved in each Biological Process GO category was calculated from combined graphs with no filter and a score alpha of 0.6. Since more than 450 different biological processes were identified for each category of genes, we excluded biological processes involving less than nine genes. When several biological processes involved the same genes and were associated with comparable functions, we considered only the one most relevant to our scientific hypotheses.

For a deeper analysis of *S. aureus* direct effect on *L. garvieae* gene expression regardless of aeration level, data mining was undertaken using the regression functions in Microsoft excel. Expression values were systematically plotted for each pair of samples (co-culture versus pure culture), a standard residual value was determined from the regression analysis for each gene across multiple pair wise comparisons and the mean value was determined. This mean standard residual value was used to rank the genes and identify the strongest gene expression differences. A standard residual cut-off of 5 was used as the threshold for significance to account for false discovery using a Bonferroni correction and approximating the standard residual to an equivalent *p*-value for the number of comparisons.

Determination of Gene Expression by RT-qPCR

Total RNA was retro-transcribed using a High Capacity cDNA Reverse Transcription kit (Invitrogen, Life Technologies, Carlsbad, CA, USA) following the supplier's instructions. Ct of genes of interest (see Supplementary Table 2) were determined by RT-qPCR as described by Delpéch et al. (2015). All primers were designed using PrimerExpress® software (Applied Biosystems®, Life Technologies). By comparison with the Ct of the *tufB* reference gene stable in our conditions (data not shown), expression of a gene of interest ("goi") was calculated using the formula introduced by Pfaffl (2001).

The influence of two experimental factors was studied: the presence of *S. aureus* (with high or low aeration, gene expression in co-culture divided by gene expression in pure culture) and aeration level (in presence or absence of *S. aureus*, gene expression in shaken condition divided by gene expression in static condition). Statistical analyses were performed using Statistica software (StatSoft, Inc., StatSoft France, Maisons-Alfort, France) by single-factor ANOVA followed by a Newman-Keuls *post hoc* test.

³<http://blast.ncbi.nlm.nih.gov/Blast.cgi>

RESULTS

Effect of Enzyme-Treated Co-culture Supernatants on *S. aureus* Planktonic Growth

Since the inhibition of *S. aureus* may not be related solely to hydrogen peroxide, we looked for another possible anti-staphylococcal molecule produced by *L. garvieae* in the culture supernatants. We monitored *S. aureus* growth (OD₆₀₀) over 24 h in either non-inoculated BHI or in supernatants prepared from H₂O₂-free cultures (presence of catalase) with low aeration (either pure culture of *S. aureus* or co-culture of *L. garvieae* and *S. aureus*). In order to avoid any nutritional competition, BHI at a final concentration of 1X was added to each culture.

Growth of *S. aureus* in supernatant from *S. aureus* pure culture was comparable to that in plain BHI (data not shown). During the stationary phase, the OD₆₀₀ measured in *S. aureus* cultures in supernatant from co-culture was lower than that in supernatant from *S. aureus* pure culture (see Data Sheet 1). This inhibition was still observed when these supernatants were treated with proteases, lipase or α -amylase, the main macromolecule-degrading enzymes. pH values remained between 6.9 and 7.1 in all cultures (data not shown).

Determination of Aeration and *S. aureus* Effects on *L. garvieae* Transcriptome by RNA Sequencing

To identify the best conditions for studying the transcriptome of *L. garvieae* N201 with respect to its capacity to inhibit *S. aureus* SA15, we followed the growth of the two bacteria in pure cultures and co-cultures in BHI for 24 h under high or low aeration conditions (Table 1).

The growth of *L. garvieae* was not affected by *S. aureus*. The growth of *S. aureus* was inhibited by *L. garvieae* at both aeration levels. With high aeration, maximal inhibition was observed from 9 to 24 h (difference with pure culture of 4 and 4.5 log CFU/ml, respectively, at 9 and 24 h). At 9 h, H₂O₂ concentration reached a peak concomitant with the lowest *S. aureus* concentration. With low aeration, inhibition was weaker than with high aeration and was observed later, at 24 h. Concomitantly, H₂O₂ was not detected in these cultures. pH values remained between 6.9 and 7.1 in all cultures (data not shown).

Considering these data, we analyzed the *L. garvieae* transcriptome in pure culture and in co-culture with *S. aureus* after 9 h of incubation with high and the low aerations. RNA sequencing generated a number of reads per sample, ranging from 12,365,133 to 15,670,131 depending on the sample after the initial quality filter (see Supplementary Table 1). Between 83.1 and 91.0% of these reads mapped correctly onto the reference genomes. Considering the direct effect of *S. aureus* on *L. garvieae* gene expression, analysis of the mapped reads using the Bioconductor R packages failed to identify expression differences (data not shown). Additional regression analyses highlighted 39 genes differentially

TABLE 1 | Cell counts and H₂O₂ concentration over 24 h in cultures of *Lactococcus garvieae* N201 and *Staphylococcus aureus* SA15 with high or low aeration levels.

Aeration level	Culture	<i>L. garvieae</i> cell concentration (log [CFU.mL ⁻¹])					<i>S. aureus</i> cell concentration (log [CFU.mL ⁻¹])					Hydrogen peroxide concentration (mM)				
		0 h	3 h	6 h	9 h	24 h	0 h	3 h	6 h	9 h	24 h	0 h	3 h	6 h	9 h	24 h
High	N201	6.6 ^a	7.3 ^a	8.4 ^a	9.0 ^a	8.6 ^a	NT	NT	NT	NT	NT	ND	NT	0.5 ^a	1.7 ^a	1.5 ^a
	SA15	NT	NT	NT	NT	NT	5.6 ^a	5.2 ^{ab}	6.1 ^a	7.4 ^a	9.0 ^a	ND	NT	ND	ND	ND
	N201 + SA15	6.5 ^a	7.4 ^a	8.5 ^a	8.6 ^a	8.7 ^a	5.6 ^a	4.8 ^a	3.9 ^b	3.4 ^b	4.5 ^b	ND	NT	0.5 ^a	1.6 ^a	1.5 ^a
Low	N201	6.6 ^a	7.4 ^a	8.4 ^a	8.7 ^a	8.9 ^a	NT	NT	NT	NT	NT	ND	NT	ND	ND	ND
	SA15	NT	NT	NT	NT	NT	5.6 ^a	6.1 ^c	7.0 ^a	7.3 ^a	8.0 ^c	ND	NT	ND	ND	ND
	N201 + SA15	6.5 ^a	7.3 ^a	8.8 ^a	8.8 ^a	8.9 ^a	5.6 ^a	5.6 ^{bc}	6.2 ^a	6.5 ^a	6.2 ^d	ND	NT	ND	ND	ND

In the same column, letters (a, b, and c) indicate homogeneous statistical groupings (p -value < 0.05 by Newman-Keuls method). ND, non-detectable values (below the spectrometry detection limit). NT, not tested.

expressed in presence of *S. aureus*, of which 4 genes under low aeration and 35 genes under high aeration conditions (see Supplementary Table 3). The expression of ~18% of *L. garvieae* genes, i.e., 358 genes, differed between the two aeration levels (see Supplementary Table 4). Among these 358 genes, the expression of 181 genes differed regardless of the presence or absence of *S. aureus* (i.e., similarly in pure culture and in co-culture). The expression of 177 *L. garvieae* genes responded differently to different aeration levels, depending on the presence or absence of *S. aureus*: 88 gene expressions differed only in pure culture and 89 gene expressions differed only in co-culture.

Effect of Aeration Level on *L. garvieae* Biological Processes

After RNA sequencing data treatments, 22 biological processes (named according to Gene Ontology termes) related to *L. garvieae* genes differentially expressed depending on aeration level in both pure culture and co-culture were identified using Blast2Go (Figure 1). Changes in gene expression are shown in Supplementary Table 4.

Most of the biological processes affected were related to fundamental growth functions (Figures 1A,B) and nutrition (Figure 1C). With high aeration, two genes (*ilvA*, LCGN_1922) related to “threonine metabolism” were repressed while two genes (LCGN_1919, LCGN_1920) were over-expressed.

Fourteen genes related to “transport” processes were differently expressed depending on aeration level both in pure culture and co-culture, including genes involved in the transport of metals (lead, cadmium, zinc, copper and/or mercury) and vitamins (riboflavin and folate).

Genes and GO biological processes related to O₂ and H₂O₂ metabolism were also affected. “Oxidation-reduction process” and “response to stress” biological processes involved more repressed genes (17 and 8, respectively) than over-expressed genes (6 and 1, respectively). Three genes related to H₂O₂ metabolism (*ahpF*, *pox*, and *spxA1*) and one gene related to O₂ consumption (*lox*) were repressed with the high aeration level. Several stress response genes (*hrcA*, *groES*, *groEL*, *dnaK*, *dnaJ*, *grpE*, *clpB*, and five genes belonging to the universal stress protein family) were repressed. Genes related

to peroxide resistance (*ohrA* and *ohrR*) were strongly over-expressed under high aeration conditions. Electron Transport Chain (ETC) genes (*cydB*, *menH*, and *ubiE*) were over-expressed.


Modulation by *S. aureus* of *L. garvieae* Response to Aeration

We identified 27 Gene Ontology biological processes related to genes differentially expressed depending on aeration level either in pure culture or in co-culture (Figure 2).

Changes in gene expressions are shown in Supplementary Table 4. Most of the biological processes affected were related to fundamental growth functions (Figures 2A,B) or nutrition (Figure 2C). Under high aeration conditions, four lysine metabolism genes (LCGN_0575, LCGN_0576, LCGN_0577, LCGN_0578) and two threonine metabolism genes (LCGN_0576 and LCGN_0577) were over-expressed exclusively in pure culture while four galactose metabolism genes were over-expressed exclusively in co-culture (LCGN_1809, LCGN_1810, LCGN_1811, and LCGN_1812). Eight genes involved in pyruvate and carbohydrate metabolism and the citrate cycle (*glmS*, *glmM*, *galE*, PTS-Man-EIIC and EIID *pdhA*, *pdhB*, DLAT) were repressed by *S. aureus*. With low aeration, the enolase encoding gene was repressed by *S. aureus*.

Four biological processes related to transport functions were affected (Figure 2D): “transport,” “transmembrane transport,” “organic substance transport” and “ion transport.” In pure culture, genes involved in the transport of cobalt/zinc/cadmium (LCGN_1867) and an unspecified monosaccharide (LCGN_0332) were over-expressed with high aeration while a gene involved in copper transport (LCGN_1427) was repressed. Two genes of the *fur* regulon (ferrous iron transport) were over-expressed with high aeration, one of them in pure culture (*feoA*) and one in co-culture (*feoB*).

With low aeration, the H₂O₂ synthesis gene *sodA* was up-regulated by *S. aureus*. The H₂O₂ degradation gene *ahpC* was repressed by high aeration in co-culture but not in pure culture. In pure culture, two genes related to O₂ consumption, *noxE* and LCGN_0208, were repressed and six genes related to ETC, *cydA*, *cydC*, *menB*, *menC*, *menD*, and LCGN_0364, were over-expressed.



Impact of Aeration on the Expression of *L. garvieae* Genes Potentially Involved in the Antagonism against *S. aureus*

To complete the RNA-seq data obtained only at 9 h, we determined the expression of ten genes from the cultures previously used for RNA-seq analyses, at 6, 9, and 24 h by RT-qPCR. These genes, potentially involved in the antagonism mechanisms, were involved in O₂ consumption (*noxE* and *lox*), H₂O₂ synthesis (*pox* and *sodA*), H₂O₂ degradation (*ahpC*, *ahpF*, *gpx*, and *trxB1*) and resistance to other peroxides (*ohrA* and *ohrR*).

The presence of *S. aureus* induced only slight changes in their expression (data not shown). With high aeration, *ahpC* and *pox* were over-expressed 2.4-fold and 2.0-fold in the presence of *S. aureus* at 6 h. With low aeration, *S. aureus* induced a 3.1-fold over-expression of *trxB1* at 9 h and a 3.3-fold repression of *gpx* at 24 h.


Conversely, RT-qPCR results showed major differences in H₂O₂-metabolism gene expression between the two aeration levels (Table 2). The expression of *ohrA* was strongly induced under high aeration conditions in both pure culture and co-culture at 9 h. While RNA sequencing identified *trxA2* as up-regulated with high aeration in both pure culture and co-culture at 9 h (Supplementary Table 4), RT-qPCR identified other H₂O₂-degradation genes (*ahpC*, *ahpF*, *gpx*, and *trxB1*) as

repressed at 6, 9, or 24 h. The lactate mono-oxygenase gene *lox* seemed slightly repressed with high aeration but this modification was significant only at 9 h in co-culture. In pure culture with high aeration, the expression of *noxE* reached a peak at 6 h when it was 11.0 times higher than with low aeration. With low aeration, *noxE* expression gradually increased over time. No significant difference in the expression of H₂O₂-synthesis genes (*pox* and *sodA*) according to aeration level was observed.

DISCUSSION

This study aimed to improve understanding of the mechanisms underlying the antagonism of *L. garvieae* against *S. aureus*, especially as regards H₂O₂-related pathways.

The high aeration level and the resulting high H₂O₂ concentration were associated with stronger inhibition of *S. aureus*, confirming previous observations under the same conditions (Alomar et al., 2008a; Delbes-Paus et al., 2010; Delpéch et al., 2015). It was already known that this inhibition is associated with a modulation of the expression of *S. aureus* virulence genes and a repression of the H₂O₂ response, stress response and cell division genes of *S. aureus* by *L. garvieae* and with high aeration (Delpéch et al., 2015). However, the transcriptome adaptation of *L. garvieae* to prevailing aeration conditions during this interaction had not been explored until



this study. It is known that difference in aeration is associated with drastic modifications of the *L. lactis* transcriptome (Pedersen et al., 2008; Dijkstra et al., 2014) and consequent metabolic adaptations (Jensen et al., 2001; Nordkvist et al., 2003; Pedersen et al., 2008; Dijkstra et al., 2014; Larsen et al., 2016a,b). In accordance with these findings, we found that different aeration levels were associated with significant differences in the *L. garvieae* transcriptome in biological processes related to fundamental growth functions, nutritional and metabolic adaptations and transport functions. Moreover, *L. garvieae* genes

involved in fundamental growth functions were slightly repressed by *S. aureus* under both aeration levels suggesting an impact of *S. aureus* on *L. garvieae* metabolism. While *L. garvieae* is a catalase-negative bacterium, the main *S. aureus* enzyme involved in H₂O₂ dismutation is the O₂-forming catalase KatA (Cosgrove et al., 2007). The *S. aureus* catalase may affect the transcriptomic response of *L. garvieae* to different aeration conditions by modulating O₂ and H₂O₂ concentrations. It is known that the *katA* gene of *S. aureus* SA15 is repressed by *L. garvieae* in high aeration conditions but not in low aeration conditions (Delpech

TABLE 2 | Effect of high aeration level on the expression of *L. garvieae* genes in pure culture and co-culture, as determined by RT-qPCR.

	Gene	In pure culture ^a			In co-culture ^a		
		6 h	9 h	24 h	6 h	9 h	24 h
Degradation of H ₂ O ₂	<i>ahpC</i>		0.6*		0.6*	0.4**	
	<i>ahpF</i>		0.3**	NT		0.4*	NT
	<i>gpx</i>			0.2*			0.4*
	<i>trxB1</i>			0.2**			0.2**
Degradation of other peroxides	<i>ohrA</i>		16.7**	NT	7.0*	20.2*	NT
O ₂ consumption/H ₂ O synthesis	<i>noxE</i>	11.0**	0.4*	0.1*			
	<i>lox</i>			NT		0.4**	NT

^aIndicated values, at 6, 9, or 24 h in pure culture or in co-culture, corresponds to the ratios of gene expressions with high aeration to gene expressions with low aeration. Only ratios that are significant according to the Newman-Keuls test are shown (*p-value < 0.1, **p-value < 0.05). NT, not tested.

et al., 2015). This may explain why the presence of *S. aureus* (via its catalase production) modulated the *L. garvieae* transcriptome response under different aeration conditions. Indeed, with low aeration, the *L. garvieae* H₂O₂ synthesis gene *sodA* expression was slightly higher in presence of *S. aureus*. With high aeration, the main enzymes involved in O₂ consumption of *L. lactis* (NADH-oxidase NoxE and ETC enzymes (Tachon et al., 2010), were repressed only in *L. garvieae* pure culture. The expression of the *L. garvieae* *noxE* reached a peak at 6 h (beginning of exponential growth) and then decreased until 24 h. This suggests that *L. garvieae* NoxE is the main enzyme responsible for O₂ consumption at the beginning of exponential growth with high aeration, as observed for *L. lactis* NoxE (Lopez de Felipe and Hugenholz, 2001; Tachon et al., 2010). The induction of most of the *L. garvieae* ETC genes (*cydA*, *cydB*, *cydC*, *menB*, *menC*, *menD*, *menH*, and LCGN_0364) under high aeration conditions at 9 h (beginning of stationary phase) suggested that ETC may be involved in O₂ consumption by *L. garvieae* during the stationary phase, as already shown with *L. lactis* (Tachon et al., 2009, 2010).

Previous studies have showed that hydrogen peroxide production by *L. garvieae* depends on aeration level and plays a key role in *S. aureus* inhibition. As regards *S. aureus*, Delpéché et al. (2015) suggested that the stronger inhibition with a high aeration level may be caused by the higher concentration of H₂O₂ associated with the *L. garvieae*-induced repression of *S. aureus* genes involved in the H₂O₂ response (*katA* and *sodA* at 6 and 9 h) and cell division (*mraZ*, *mraW* and potentially the *dcw* cluster). As regards *L. garvieae*, our RNA-sequencing and RT-qPCR analyses showed an overexpression of the H₂O₂-degradation genes *ahpC*, *ahpF*, *gpx*, and *trxB1* under low aeration conditions compared to high aeration, while the expression of H₂O₂ synthesis genes *pox* and *sodA* remained stable. The expression of the main H₂O₂ degradation genes of Gram-positive bacteria is generally induced more under aeration and in the presence of H₂O₂ (Mostertz et al., 2004; Pedersen et al., 2008). Also, H₂O₂ metabolism may be very different in *L. garvieae* compared to other Gram-positive bacteria. Although the difference in H₂O₂ concentration between the two aeration levels was probably primarily conditioned by the availability of O₂, our transcriptome results suggest that it was also associated with a control of H₂O₂ degradation by *L. garvieae* rather than with a control of H₂O₂ synthesis. Since the AhpCF peroxy-redoxin system was repressed, the OhrAR system was probably essential for the resistance of *L. garvieae* to ROS (other than H₂O₂) under the high aeration conditions. The widespread organic hydroperoxide detoxifying system *ohrAR*, known to be over-expressed under high O₂ conditions (Mongkolsuk et al., 1998; Fuangthong et al., 2001; Chuchue et al., 2006; Oh et al., 2007; Atichartpongkul et al., 2010; da Silva Neto et al., 2012; Clair et al., 2013), was consistently induced in *L. garvieae* at the high aeration level.

The fact that *S. aureus* population levels were lower in the stationary phase in H₂O₂-free supernatant from a co-culture of *S. aureus* and *L. garvieae* revealed the presence of a new molecule involved in this inhibition. This effector is extracellular, is produced by *L. garvieae* during its exponential growth phase and can reduce the population level of *S. aureus*

during its stationary growth phase. In view of the results of our enzymatic treatments on supernatants, the inhibitory gap during the stationary phase was probably caused neither by hydrogen peroxide, nor by a protein, nor by a lipid, nor by a polysaccharide. The only putative bacteriocin identified in the *L. garvieae* N201 genome was homologous to garvieacin Q (GarQ, data not shown), a class IId bacteriocin (Tosukhowong et al., 2012). Class IId bacteriocins are generally sensitive to protease treatments as stringent as the one we used in this study (Kuo et al., 2013; Song et al., 2014), suggesting that garvieacin Q is unlikely to be the effector we are seeking. This effector may instead be related to several genes identified by RNA seq as being regulated by aeration level, such as genes involved in metal homeostasis (e.g., siderophores (Hannauer et al., 2015), chemical and ionic equilibrium (Doyle et al., 1975; Chudobova et al., 2015), transport of vitamin-related compounds (Schlievert et al., 2013) and export of unknown proteins. For example, the differential expression of ferrous ion transport encoding genes has already been observed in *L. lactis* under oxidative stress in milk (Larsen et al., 2016b). It may also be related to signaling molecules (stress, quorum sensing). It is known that *L. garvieae* can modify the expression of several *S. aureus* genes involved in environment-sensing systems like the *agr* system, CodY or two-component systems SaeRS and SrrAB (Delpéché et al., 2015).

RNA-seq revealed variations in the expression of the *codY* gene (involved in nutritional adaptation (Guédon et al., 2001; Ercan et al., 2015), and of several nutritional-related metabolisms (lysine, threonine, mannitol, aspartate, ribose, fructose, and galactose). This suggests that *L. garvieae* adapts its nutritional behavior to the prevailing aeration level and the presence or absence of *S. aureus*. In a rich medium like BHI, there should be little nutritional competition. It is known that *L. garvieae* can consume all the L-threonine in micro-filtered milk in less than 3 h (Alomar et al., 2008b) and that *S. aureus* growth could be inhibited by L-threonine depletion (Pohl et al., 2009). However, we showed that the antagonism of *L. garvieae* against *S. aureus* was not associated with nutritional competition for L-threonine in micro-filtered milk (see Supplementary Table 5).

CONCLUSION

RNA sequencing analyses revealed a *L. garvieae* transcriptome adaptation to aeration level. This adaptation differed depending on the presence or absence of *S. aureus*. Our findings show that the control of autogenic H₂O₂ levels by *L. garvieae* was probably carried out by H₂O₂ degradation genes rather than H₂O₂ synthesis genes. Our study also leads us to suggest that an unidentified effector was involved in the inhibition of *S. aureus* in the stationary phase. The potential inhibitory role of metals, siderophores and signal molecules (e.g., stress signal, quorum sensing) generated by *L. garvieae* should be investigated. In order to promote the use of H₂O₂-producing bacteria as bio-preservation agents, future studies should explore the relation between H₂O₂-metabolism, H₂O₂-producing LAB and the pathogen they inhibit.

AUTHOR CONTRIBUTIONS

PD carried out all the experiments, excluding preparation of cDNA libraries and RNA sequencing, and drafted the manuscript helped by CD and SB. ER, ED, and GB analyzed RNA sequencing data. SN performed the RNA sequencing (preparation of libraries and the sequencing itself). GG, M-CM, CD and SB conceived the study. All authors read and approved the final manuscript.

FUNDING

PD received a Ph.D. fellowship from the FEDER (*Fond Européen de Développement Economique et Régional*) and the *Conseil Régional* of the Auvergne region in the framework of the Pharmabiotic Research Institute (PRI) cluster.

REFERENCES

- Adesokan, I. A., Ekanola, Y. A., and Okanlawon, B. M. (2010). Influence of cultural conditions on hydrogen peroxide production by lactic acid bacteria isolated from some Nigerian traditional fermented foods. *Afr. J. Microbiol. Res.* 4, 1991–1996.
- Aguado-Urda, M., López-Campos, G. H., Blanco, M. M., Fernández-Garayzábal, J. F., Cutuli, M. T., Aspiroz, C., et al. (2011). Genome sequence of *Lactococcus garvieae* 21881, isolated in a case of human septicemia. *J. Bacteriol.* 193, 4033–4034. doi: 10.1128/JB.05090-11
- Alegria, A., Alvarez-Martín, P., Sacristán, N., Fernández, E., Delgado, S., and Mayo, B. (2009). Diversity and evolution of the microbial populations during manufacture and ripening of Casin, a traditional Spanish, starter-free cheese made from cow's milk. *Int. J. Food Microbiol.* 136, 44–51. doi: 10.1016/j.ijfoodmicro.2009.09.023
- Alomar, J., Lebert, A., and Montel, M.-C. (2008a). Effect of temperature and pH on growth of *Staphylococcus aureus* in co-culture with *Lactococcus garvieae*. *Curr. Microbiol.* 56, 408–412. doi: 10.1007/s00284-007-9079-3
- Alomar, J., Loubiere, P., Delbes, C., Nouaille, S., and Montel, M.-C. (2008b). Effect of *Lactococcus garvieae*, *Lactococcus lactis* and *Enterococcus faecalis* on the behaviour of *Staphylococcus aureus* in microfiltered milk. *Food Microbiol.* 25, 502–508. doi: 10.1016/j.fm.2008.01.005
- Anders, S., and Huber, W. (2010). Differential expression analysis for sequence count data. *Genome Biol.* 11:R106. doi: 10.1186/gb-2010-11-10-r106
- Anders, S., Pyl, P. T., and Huber, W. (2015). HTSeq—a Python framework to work with high-throughput sequencing data. *Bioinformatics* 31, 166–169. doi: 10.1093/bioinformatics/btu638
- Atichartpongkul, S., Fuangthong, M., Vattanaviboon, P., and Mongkolsuk, S. (2010). Analyses of the regulatory mechanism and physiological roles of *Pseudomonas aeruginosa* OhrR, a transcription regulator and a sensor of organic hydroperoxides. *J. Bacteriol.* 192, 2093–2101. doi: 10.1128/JB.01510-09
- Batdorj, B., Trinetta, V., Dalgalarondo, M., Prévost, H., Dousset, X., Ivanova, I., et al. (2007). Isolation, taxonomic identification and hydrogen peroxide production by *Lactobacillus delbrueckii* subsp. *lactis* T31, isolated from Mongolian yoghurt: inhibitory activity on food-borne pathogens. *J. Appl. Microbiol.* 103, 584–593. doi: 10.1111/j.1365-2672.2007.03279.x
- Callon, C., Duthoit, F., Delbès, C., Ferrand, M., Le Frileux, Y., De Crémoux, R., et al. (2007). Stability of microbial communities in goat milk during a lactation year: molecular approaches. *Syst. Appl. Microbiol.* 30, 547–560. doi: 10.1016/j.syapm.2007.05.004
- Charlier, C., Cretenet, M., Even, S., and Le Loir, Y. (2009). Interactions between *Staphylococcus aureus* and lactic acid bacteria: an old story with new perspectives. *Int. J. Food Microbiol.* 131, 30–39. doi: 10.1016/j.ijfoodmicro.2008.06.032
- Chuchue, T., Tanboon, W., Prapagdee, B., Dubbs, J. M., Vattanaviboon, P., and Mongkolsuk, S. (2006). ohrR and ohr are the primary sensor/regulator

ACKNOWLEDGMENTS

The authors would like to thank Adrien Nivoliez for his help in determining the quality of our RNAs in the ProBionov laboratory, Magali Cordaillat-Simmons, David Tropel, Laurent Rios, Monique Zagorec, Yves Le Loir, Pierre Renault, Hélène Falentin, and Christophe Chassard for their helpful advice and Maryline Bornes and Harriet Coleman for English proofreading.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fmicb.2017.00359/full#supplementary-material>

- and protective genes against organic hydroperoxide stress in *Agrobacterium tumefaciens*. *J. Bacteriol.* 188, 842–851. doi: 10.1128/JB.188.3.842-851.2006
- Chudobova, D., Dostalova, S., Ruttikay-Nedecky, B., Guran, R., Rodrigo, M. A. M., Tmejova, K., et al. (2015). The effect of metal ions on *Staphylococcus aureus* revealed by biochemical and mass spectrometric analyses. *Microbiol. Res.* 170, 147–156. doi: 10.1016/j.micres.2014.08.003
- Clair, G., Lorphelin, A., Armengaud, J., and Dupont, C. (2013). OhrRA functions as a redox-responsive system controlling toxinogenesis in *Bacillus cereus*. *J. Proteomics* 94, 527–539. doi: 10.1016/j.jprot.2013.10.024
- Conesa, A., Götz, S., García-Gómez, J. M., Terol, J., Talón, M., and Robles, M. (2005). Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics* 21, 3674–3676. doi: 10.1093/bioinformatics/bti610
- Cosgrove, K., Coutts, G., Jonsson, I.-M., Tarkowski, A., Kokai-Kun, J. F., Mond, J. J., et al. (2007). Catalase (KatA) and alkyl hydroperoxide reductase (AhpC) have compensatory roles in peroxide stress resistance and are required for survival, persistence, and nasal colonization in *Staphylococcus aureus*. *J. Bacteriol.* 189, 1025–1035. doi: 10.1128/JB.01524-06
- da Silva Neto, J. F., Negretto, C. C., and Netto, L. E. S. (2012). Analysis of the organic hydroperoxide response of chromobacterium violaceum reveals that OhrR is a cys-based redox sensor regulated by thioredoxin. *PLoS ONE* 7:e47090. doi: 10.1371/journal.pone.0047090
- Delbes-Paus, C., Dorchies, G., Chaabna, Z., Callon, C., and Montel, M.-C. (2010). Contribution of hydrogen peroxide to the inhibition of *Staphylococcus aureus* by *Lactococcus garvieae* in interaction with raw milk microbial community. *Food Microbiol.* 27, 924–932. doi: 10.1016/j.fm.2010.05.031
- Delpéch, P., Bornes, S., Alaterre, E., Bonnet, M., Gagne, G., Montel, M.-C., et al. (2015). *Staphylococcus aureus* transcriptomic response to inhibition by H₂O₂-producing *Lactococcus garvieae*. *Food Microbiol.* 51, 163–170. doi: 10.1016/j.fm.2015.05.014
- Devriese, L. A., Homme, J., Laevens, H., Pot, B., Vandamme, P., and Haesebrouck, F. (1999). Identification of aesculin-hydrolyzing streptococci, lactococci, aerococci and enterococci from subclinical intramammary infections in dairy cows. *Vet. Microbiol.* 70, 87–94. doi: 10.1016/S0378-1135(99)00124-8
- Dijkstra, A. R., Alkema, W., Starrenburg, M. J., Hugenholtz, J., van Hijum, S. A., and Bron, P. A. (2014). Fermentation-induced variation in heat and oxidative stress phenotypes of *Lactococcus lactis* MG1363 reveals transcriptome signatures for robustness. *Microb. Cell Fact.* 13, 148. doi: 10.1186/s12934-014-0148-6
- Doyle, J. J., Marshall, R. T., and Pfander, W. H. (1975). Effects of cadmium on the growth and uptake of cadmium by microorganisms. *Appl. Microbiol.* 29, 562–564.
- El-Baradei, G., Delacroix-Buchet, A., and Ogier, J. C. (2008). Bacterial biodiversity of traditional Zabady fermented milk. *Int. J. Food Microbiol.* 121, 295–301. doi: 10.1016/j.ijfoodmicro.2007.11.014

- Eldar, A., and Ghittino, C. (1999). *Lactococcus garvieae* and *Streptococcus iniae* infections in rainbow trout *Oncorhynchus mykiss*: similar, but different diseases. *Dis. Aquat. Org.* 36, 227–231. doi: 10.3354/dao036227
- Ercan, O., Wels, M., Smid, E. J., and Kleerebezem, M. (2015). Genome-wide transcriptional responses to carbon starvation in nongrowing *Lactococcus lactis*. *Appl. Environ. Microbiol.* 81, 2554–2561. doi: 10.1128/AEM.03748-14
- Fernández, E., Alegría, Á., Delgado, S., and Mayo, B. (2010). Phenotypic, genetic and technological characterization of *Lactococcus garvieae* strains isolated from a raw milk cheese. *Int. Dairy J.* 20, 142–148. doi: 10.1016/j.idairyj.2009.11.004
- Flórez, A. B., and Mayo, B. (2006). Microbial diversity and succession during the manufacture and ripening of traditional, Spanish, blue-veined Cabrales cheese, as determined by PCR-DGGE. *Int. J. Food Microbiol.* 110, 165–171. doi: 10.1016/j.ijfoodmicro.2006.04.016
- Fortina, M. G., Ricci, G., Foschino, R., Picozzi, C., Dolci, P., Zeppa, G., et al. (2007). Phenotypic typing, technological properties and safety aspects of *Lactococcus garvieae* strains from dairy environments. *J. Appl. Microbiol.* 103, 445–453. doi: 10.1111/j.1365-2672.2006.03265.x
- Fuangthong, M., Atichartpongkul, S., Mongkolsuk, S., and Helmann, J. D. (2001). OhrR is a repressor of ohrA, a key organic hydroperoxide resistance determinant in *Bacillus subtilis*. *J. Bacteriol.* 183, 4134–4141. doi: 10.1128/JB.183.14.4134-4141.2001
- Gene Ontology Consortium (2008). The gene ontology project in 2008. *Nucleic Acids Res.* 36, D440–D444.
- Guédon, E., Serror, P., Ehrlich, S. D., Renault, P., and Delorme, C. (2001). Pleiotropic transcriptional repressor CodY senses the intracellular pool of branched-chain amino acids in *Lactococcus lactis*. *Mol. Microbiol.* 40, 1227–1239. doi: 10.1046/j.1365-2958.2001.02470.x
- Hannauer, M., Sheldon, J. R., and Heinrichs, D. E. (2015). Involvement of major facilitator superfamily proteins SfaA and SbnD in staphyloferrin secretion in *Staphylococcus aureus*. *FEBS Lett.* 589, 730–737. doi: 10.1016/j.febslet.2015.02.002
- Ito, A., Sato, Y., Kudo, S., Sato, S., Nakajima, H., and Toba, T. (2003). The screening of hydrogen peroxide-producing lactic acid bacteria and their application to inactivating psychrotrophic food-borne pathogens. *Curr. Microbiol.* 47, 0231–236. doi: 10.1007/s00284-002-3993-1
- Jensen, N. B. S., Melchiorsen, C. R., Jokumsen, K. V., and Villadsen, J. (2001). Metabolic behavior of *Lactococcus lactis* MG1363 in microaerobic continuous cultivation at a low dilution rate. *Appl. Environ. Microbiol.* 67, 2677–2682. doi: 10.1128/AEM.67.6.2677-2682.2001
- Jokovic, N., Nikolic, M., Begovic, J., Jovic, B., Savic, D., and Topisirovic, L. (2008). A survey of the lactic acid bacteria isolated from Serbian artisanal dairy product kajmak. *Int. J. Food Microbiol.* 127, 305–311. doi: 10.1016/j.ijfoodmicro.2008.07.006
- Kuo, Y.-C., Liu, C.-F., Lin, J.-F., Li, A.-C., Lo, T.-C., and Lin, T.-H. (2013). Characterization of putative class II bacteriocins identified from a non-bacteriocin-producing strain *Lactobacillus casei* ATCC 334. *Appl. Microbiol. Biotechnol.* 97, 237–246. doi: 10.1007/s00253-012-4149-2
- Larsen, N., Brøsted Werner, B., and Jespersen, L. (2016a). Transcriptional responses in *Lactococcus lactis* subsp. *cremoris* to the changes in oxygen and redox potential during milk acidification. *Let. Appl. Microbiol.* 63, 117–123. doi: 10.1111/lam.12596
- Larsen, N., Moslehi-Jenabian, S., Werner, B. B., Jensen, M. L., Garrigues, C., Vogensen, F. K., et al. (2016b). Transcriptome analysis of *Lactococcus lactis* subsp. *lactis* during milk acidification as affected by dissolved oxygen and the redox potential. *Int. J. Food Microbiol.* 226, 5–12. doi: 10.1016/j.ijfoodmicro.2016.03.002
- Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with burrows-wheeler transform. *Bioinformatics* 25, 1754–1760. doi: 10.1093/bioinformatics/btp324
- Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., et al. (2009). The sequence alignment/map format and SAMtools. *Bioinformatics* 25, 2078–2079. doi: 10.1093/bioinformatics/btp352
- Lopez de Felipe, F., and Hugenholtz, J. (2001). Purification and characterisation of the water forming NADH-oxidase from *Lactococcus lactis*. *Int. Dairy J.* 11, 37–44. doi: 10.1016/S0958-6946(01)00031-0
- Love, M. I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* 15, 550. doi: 10.1186/s13059-014-0550-8
- Monfredini, L., Settanni, L., Poznanski, E., Cavazza, A., and Franciosi, E. (2012). The spatial distribution of bacteria in Grana-cheese during ripening. *Syst. Appl. Microbiol.* 35, 54–63. doi: 10.1016/j.syapm.2011.07.002
- Mongkolsuk, S., Praituan, W., Loprasert, S., Fuangthong, M., and Chamnongpol, S. (1998). Identification and characterization of a new organic hydroperoxide resistance (ohr) gene with a novel pattern of oxidative stress regulation from *Xanthomonas campestris* pv. *phaseoli*. *J. Bacteriol.* 180, 2636–2643.
- Morandi, S., Silveti, T., Miranda Lopez, J. M., and Brasca, M. (2015). Antimicrobial activity, antibiotic resistance and the safety of lactic acid bacteria in raw milk valtellina casera cheese. *J. Food Saf.* 35, 193–205. doi: 10.1111/jfs.12171
- Mostertz, J., Scharf, C., Hecker, M., and Homuth, G. (2004). Transcriptome and proteome analysis of *Bacillus subtilis* gene expression in response to superoxide and peroxide stress. *Microbiology* 150, 497–512. doi: 10.1099/mic.0.26665-0
- Nordkvist, M., Jensen, N. B. S., and Villadsen, J. (2003). Glucose metabolism in *Lactococcus lactis* MG1363 under different aeration conditions: requirement of acetate to sustain growth under microaerobic conditions. *Appl. Environ. Microbiol.* 69, 3462–3468. doi: 10.1128/AEM.69.6.3462-3468.2003
- Nouaille, S., Even, S., Charlier, C., Le Loir, Y., Coccagn-Bousquet, M., and Loubière, P. (2009). Transcriptomic response of *Lactococcus lactis* in mixed culture with *Staphylococcus aureus*. *Appl. Environ. Microbiol.* 75, 4473–4482. doi: 10.1128/AEM.02653-08
- Oh, S.-Y., Shin, J.-H., and Roe, J.-H. (2007). Dual role of OhrR as a repressor and an activator in response to organic hydroperoxides in *Streptomyces coelicolor*. *J. Bacteriol.* 189, 6284–6292. doi: 10.1128/JB.00632-07
- Ortiz, C., López, J., Del Amo, E., Sevilla, T., García, P. E., and San Román, J. A. (2014). *Lactococcus garvieae* infective endocarditis: report of 2 cases and review of the literature. *Rev. Esp. Cardiol.* 67, 776–778. doi: 10.1016/j.recesp.2014.04.010
- Pangallo, D., Saková, N., Korešová, J., Puškárová, A., Kraková, L., Valík, L., et al. (2014). Microbial diversity and dynamics during the production of May bryndza cheese. *Int. J. Food Microbiol.* 170, 38–43. doi: 10.1016/j.ijfoodmicro.2013.10.015
- Pedersen, M. B., Garrigues, C., Tuphile, K., Brun, C., Vido, K., Bennedsen, M., et al. (2008). Impact of aeration and heme-activated respiration on *Lactococcus lactis* gene expression: identification of a heme-responsive operon. *J. Bacteriol.* 190, 4903–4911. doi: 10.1128/JB.00447-08
- Pfaffl, M. W. (2001). A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res.* 29:e45. doi: 10.1093/nar/29.9.e45
- Pinto, A. C., Melo-Barbosa, H. P., Miyoshi, A., Silva, A., and Azevedo, V. (2011). Application of RNA-seq to reveal the transcript profile in bacteria. *Genet. Mol. Res.* 10, 1707–1718. doi: 10.4238/vol10-3gmr1554
- Pohl, K., Francois, P., Stenz, L., Schlink, F., Geiger, T., Herbert, S., et al. (2009). CodY in *Staphylococcus aureus*: a regulatory link between metabolism and virulence gene expression. *J. Bacteriol.* 191, 2953–2963. doi: 10.1128/JB.01492-08
- Robinson, M. D., McCarthy, D. J., and Smyth, G. K. (2010). edgeR: a bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 26, 139–140. doi: 10.1093/bioinformatics/btp616
- Schlievert, P. M., Merriman, J. A., Salgado-Pabón, W., Mueller, E. A., Spaulding, A. R., Vu, B. G., et al. (2013). Menaquinone analogs inhibit growth of bacterial pathogens. *Antimicrob. Agents Chemother.* 57, 5432–5437. doi: 10.1128/AAC.01279-13
- Sip, A., Wiecekiewicz, M., Olejnik-Schmidt, A., Gardo, A., Gorlas, R., and Grajek, W. (2009). Occurrence of lactic acid bacteria with activity against *Listeria* in Polish regional cheeses produced in the Tatrzan'Sko-Beskidzki District. *Acta Sci. Pol.* 8, 27–44.
- Song, D.-F., Zhu, M.-Y., and Gu, Q. (2014). Purification and characterization of Plantaricin ZJ5, a new bacteriocin produced by *Lactobacillus plantarum* ZJ5. *PLoS ONE* 9:e105549. doi: 10.1371/journal.pone.0105549
- Tachon, S., Brandsma, J. B., and Yvon, M. (2010). NoxE NADH oxidase and the electron transport chain are responsible for the ability of *Lactococcus lactis* to decrease the redox potential of milk. *Appl. Environ. Microbiol.* 76, 1311–1319. doi: 10.1128/AEM.02120-09
- Tachon, S., Michelon, D., Chambellon, E., Cantonnet, M., Mezange, C., Henno, L., et al. (2009). Experimental conditions affect the site of tetrazolium violet

- reduction in the electron transport chain of *Lactococcus lactis*. *Microbiology* 155, 2941–2948. doi: 10.1099/mic.0.029678-0
- Tosukhowong, A., Zendo, T., Visessanguan, W., Roytrakul, S., Pumpuang, L., Jaresitthikunchai, J., et al. (2012). ieacin Q, a novel class II bacteriocin from *Lactococcus garvieae* BCC 43578. *Appl. Environ. Microbiol.* 78, 1619–1623. doi: 10.1128/AEM.06891-11
- Vendrell, D., Balcázar, J. L., Ruiz-Zarzuela, I., de Blas, I., Gironés, O., and Múzquiz, J. L. (2006). *Lactococcus garvieae* in fish: a review. *Comp. Immunol. Microbiol. Infect. Dis.* 29, 177–198. doi: 10.1016/j.cimid.2006.06.003
- Villani, F., Aponte, M., Blaiotta, G., Mauriello, G., Pepe, O., and Moschetti, G. (2001). Detection and characterization of a bacteriocin, garviecin L1-5, produced by *Lactococcus garvieae* isolated from raw cow's milk. *J. Appl. Microbiol.* 90, 430–439. doi: 10.1046/j.1365-2672.2001.01261.x
- Zdobnov, E. M., and Apweiler, R. (2001). InterProScan—an integration platform for the signature-recognition methods in InterPro. *Bioinformatics* 17, 847–848. doi: 10.1093/bioinformatics/17.9.847

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Delpéché, Rifa, Ball, Nidelet, Dubois, Gagne, Montel, Delbès and Bornes. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.