

# Positive solutions for a second order multi-point boundary value problem with delay

Abdelkader Lakmeche, Horiya Kouadri Habbaze, Ahmed Lakmeche

### ▶ To cite this version:

Abdelkader Lakmeche, Horiya Kouadri Habbaze, Ahmed Lakmeche. Positive solutions for a second order multi-point boundary value problem with delay. 2018. hal-01677056

## HAL Id: hal-01677056 https://hal.science/hal-01677056

Preprint submitted on 7 Jan 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. 

### Positive solutions for a second order

#### multi-point boundary value problem with delay

Horiya Kouadri Habbaze, Ahmed Lakmeche and Abdelkader Lakmeche

Biomathematics Laboratory Univ Sidi-Bel-Abbès Sidi-Bel-Abbès Algeria lakahmed2000@yahoo.fr and lakmeche@yahoo.fr

**RÉSUMÉ.** Dans ce travail, nous étudions l'existence des solutions positives pour un problème aux limites en plusieurs points pour une équation différentielle du second ordre avec retard. Sous certaines conditions de croissance sur la non-linéarité, et moyennant le théorème du point fixe de Leray-Schauder, on obtient des conditions suffisantes pour l'existence d'une solution non triviale, ce qui améliorent les résultats de J. Chen *et al.* [3].

**ABSTRACT.** In this work, we investigate the existence of positive solutions for a multi-point boundary value problem for a second order delay differential equation. Under certain growth conditions on the nonlinearity, and by the mean of Leray-Schauder fixed point theorem, sufficient conditions for the existence of nontrivial solution are obtained, which improve the results of J. Chen *et al.* [3].

**MOTS-CLÉS** : Solution positive, Equation différentielle à retard, Problème aux limites en plusieurs points, Théorème de point fixe de Leray-Schauder.

**KEYWORDS**: Positive solution, Delay differential equation, Multi-point boundary value problem, Leray-Schauder fixed point theorem.

Volume 26 - 2016, pages 1 à 11 - Revue

#### 1. Introduction

,,

The boundary value problems for delay differential equations arise in a variety of areas of applied mathematics, physics and variational problems of control theory [4]. Recently, many researchers have done a great deal of research works upon boundary value problems of lower order differential equations with delay, and some interesting results were produced, see for example [1], [2] and [6]-[9].

In this work, we study the existence of positive solutions of the following nonlinear multipoint boundary value problem with delay

$$u''(t) + \lambda a(t) f(t, u(t - \tau)) = 0, \qquad t \in [0, 1], u(t) = \beta u(\eta), \qquad -\tau \le t \le 0, u(1) = \alpha u(\eta)$$
[1]

where  $0 < \tau < 1$ ,  $0 < \eta < 1$ ,  $0 < \alpha < \frac{1}{\eta}$  and  $0 < \beta < \frac{1-\alpha\eta}{1-\eta}$  are constants, and  $\lambda$  is a positive real parameter.

The paper is organized as follows, in section tow we give definitions and preliminaries, and in section there we give our main results.

#### 2. Preliminaries

In this section we give some preliminary results.

#### **Definition 1**

u(t) is called a positive solution of (1) if  $u \in C[-\tau, 1] \cap C^2(0, 1)$ ,  $u(t) \ge 0$  for  $t \in (0, 1)$  and satisfies (1).

#### Lemma 1

Let  $\beta \neq \frac{1-\alpha\eta}{1-\eta}$ . Then for  $y \in C([0,T], \mathbf{R})$ , the boundary value problem

$$u''(t) + y(t) = 0, \quad t \in [0, T],$$
[2]

$$u(0) = \beta u(\eta), \quad u(1) = \alpha u(\eta)$$
[3]

has a unique solution

$$u(t) = \int_0^1 G(t,s)y(s)ds$$
[4]

where

$$G(t,s) = g(t,s) + \frac{\beta + (\alpha - \beta)t}{(1 - \alpha\eta) - \beta(1 - \eta)}g(\eta,s)$$
<sup>[5]</sup>

and

$$g(t,s) = \left\{ \begin{array}{ll} s(1-t), & 0 \leq s \leq t \leq 1, \\ t(1-s), & 0 \leq t \leq s \leq 1. \end{array} \right.$$

PROOF. — From equation (2), we have

$$u(t) = u(0) + u'(0)t - \int_0^t (t-s)y(s)ds := A + Bt - \int_0^t (t-s)y(s)ds$$

with

$$u(0) = A,$$
  
$$u(\eta) = A + B\eta - \int_0^{\eta} (\eta - s)y(s)ds$$

and

$$u(1) = A + B - \int_0^1 (1 - s)y(s)ds.$$

From  $u(0) = \beta u(\eta)$ , we have

$$(1-\beta)A - B\beta\eta = -\beta \int_0^\eta (\eta - s)y(s)ds.$$

From  $u(1) = \alpha u(\eta)$ , we have

$$(1 - \alpha)A + B(1 - \alpha\eta) = \int_0^1 (1 - s)y(s)ds - \alpha \int_0^\eta (\eta - s)y(s)ds.$$

Therefore,

$$A = \frac{\beta\eta}{(1-\alpha\eta) - \beta(1-\eta)} \int_0^1 (1-s)y(s)ds - \frac{\beta}{(1-\alpha\eta) - \beta(1-\eta)} \int_0^\eta (\eta-s)y(s)ds$$
  
and

$$B = \frac{1-\beta}{(1-\alpha\eta) - \beta(1-\eta)} \int_0^1 (1-s)y(s)ds - \frac{\alpha-\beta}{(1-\alpha\eta) - \beta(1-\eta)} \int_0^\eta (\eta-s)y(s)ds.$$
 From which it follows that

From which it follows that  $\beta n$ 

$$\begin{split} u(t) &= \frac{\beta\eta}{(1-\alpha\eta) - \beta(1-\eta)} \int_0^1 (1-s)y(s)ds - \frac{\beta}{(1-\alpha\eta) - \beta(1-\eta)} \int_0^\eta (\eta-s)y(s)ds \\ &+ \frac{(1-\beta)t}{(1-\alpha\eta) - \beta(1-\eta)} \int_0^1 (1-s)y(s)ds - \frac{(\alpha-\beta)t}{(1-\alpha\eta) - \beta(1-\eta)} \int_0^\eta (\eta-s)y(s)ds \\ &- \int_0^t (t-s)y(s)ds \\ &= -\int_0^t (t-s)y(s)ds + \frac{(\beta-\alpha)t - \beta}{(1-\alpha\eta) - \beta(1-\eta)} \int_0^\eta (\eta-s)y(s)ds \\ &+ \frac{(1-\beta)t + \beta\eta}{(1-\alpha\eta) - \beta(1-\eta)} \int_0^1 (1-s)y(s)ds \\ &= \int_0^1 g(t,s)y(s)ds + \frac{\beta + (\alpha-\beta)t}{(1-\alpha\eta) - \beta(1-\eta)} \int_0^1 g(\eta,s)y(s)ds. \end{split}$$

Then,  $u(t) = \int_0^1 G(t, s)y(s)ds$ . The function u presented above is the unique solution to the problem (2), (3).

#### Lemma 2

Let  $0 < \alpha < \frac{1}{\eta}$  and  $0 \le \beta < \frac{1-\alpha\eta}{1-\eta}$ . If  $y \in C([0,1], [0,\infty))$ , then the unique solution u of the problem (2), (3) satisfies

$$u(t) \ge 0, \quad t \in [0,1].$$

PROOF. — We know that if  $u''(t) = -y(t) \le 0$  for  $t \in (0, 1)$ ,  $u(0) \ge 0$  and  $u(1) \ge 0$ , then  $u(t) \ge 0$  for  $t \in [0, 1]$ . We have

$$\begin{split} u(0) &= \frac{-\beta}{(1-\alpha\eta)-\beta(1-\eta)} \int_{0}^{\eta} (\eta-s)y(s)ds + \frac{\beta\eta}{(1-\alpha\eta)-\beta(1-\eta)} \int_{0}^{1} (1-s)y(s)ds \\ &= \frac{\beta}{(1-\alpha\eta)-\beta(1-\eta)} [-\int_{0}^{\eta} (\eta-s)y(s)ds + \eta \int_{0}^{\eta} (1-s)y(s)ds] \\ &+ \frac{\beta\eta}{(1-\alpha\eta)-\beta(1-\eta)} \int_{\eta}^{1} (1-s)y(s)ds \\ &= \frac{\beta}{(1-\alpha\eta)-\beta(1-\eta)} [\int_{0}^{\eta} s(1-\eta)y(s)ds] + \frac{\beta\eta}{(1-\alpha\eta)-\beta(1-\eta)} \int_{\eta}^{1} (1-s)y(s)ds \ge 0 \end{split}$$

and

$$u(1) = -\int_{0}^{1} (1-s)y(s)ds + \frac{(\beta-\alpha)-\beta}{(1-\alpha\eta)-\beta(1-\eta)} \int_{0}^{\eta} (\eta-s)y(s)ds + \frac{(1-\beta)+\beta\eta}{(1-\alpha\eta)-\beta(1-\eta)} \int_{0}^{1} (1-s)y(s)ds$$
  
$$= \frac{\alpha}{(1-\alpha\eta)-\beta(1-\eta)} [\eta \int_{0}^{1} (1-s)y(s)ds + \int_{0}^{\eta} (\eta-s)y(s)ds]$$
  
$$\geq \frac{\alpha}{(1-\alpha\eta)-\beta(1-\eta)} [\eta \int_{0}^{\eta} (1-s)y(s)ds + \int_{0}^{\eta} (\eta-s)y(s)ds]$$
  
$$= \frac{\alpha}{(1-\alpha\eta)-\beta(1-\eta)} \int_{0}^{\eta} s(1-\eta)y(s)ds \ge 0.$$

Then,  $u(t) \ge 0 \ \forall t \in [0, 1].$ 

Lemma 3 The function g has the following properties

 $\begin{array}{ll} (i) \ \ 0 \leq g(t,s) \leq s(1-s) = g(s,s) & \forall t,s \in [0,1]. \\ (ii) \ \ \text{Let} \ \theta \in [0,\frac{1}{2}]. \ \text{Then, for} \ t \in [\theta,1-\theta] \ \text{and} \ s \in [0,1], \ \text{we have} \\ & g(t,s) \geq \min\{t,1-t\}g(s,s) \geq \theta g(s,s). \end{array}$ 

PROOF. — For  $0 \le s \le t \le 1$ , we have

$$0 \le g(t,s) = s(1-t) \le s(1-s) = g(s,s).$$

And for  $0 \le t \le s \le 1$ , we have

$$g(t,s) = t(1-s) \le s(1-s) = g(s,s).$$

Thus (i) holds.

If s = 0 or s = 1, we show that (ii) holds. For  $0 < s \le t \le 1$  and  $s \ne 1$  we have

$$\frac{g(t,s)}{g(s,s)} = \frac{t(1-s)}{s(1-s)} = \frac{t}{s} \ge t \quad \forall t \in [0,1].$$

For  $0 \le t \le s < 1$  and  $s \ne 0$  we have

$$\frac{g(t,s)}{g(s,s)} = \frac{s(1-t)}{s(1-s)} = \frac{(1-t)}{(1-s)} \ge (1-t) \quad \forall t \in [0,1].$$

Then

$$g(t,s) \ge \min\{t, 1-t\}g(s,s).$$

Thus, there exist  $\theta \in ]0, \frac{1}{2}]$  such that

$$\frac{g(t,s)}{g(s,s)} \ge \theta, \quad \forall t \in [\theta, 1-\theta]$$

Thus (ii) holds.

Lemma 4 The function G has the following properties

 $\begin{array}{ll} (i) \ \ G(t,s) \geq 0 & \forall t,s \in [0,1], \\ (ii) \ \ G(t,s) \leq k_1 g(s,s) & \forall t,s \in [0,1] \ and \ k_1 = 1 + \frac{\max\{\alpha,\beta\}}{(1-\alpha\eta) - \beta(1-\eta)}, \\ (iii) \ \ \min_{\theta \leq t \leq 1-\theta} G(t,s) \geq k_2 g(s,s) & \forall t,s \in [0,1] \ where \ \theta \in (0,\frac{1}{2}) \ and \\ k_2 = \theta \left[ 1 + \frac{\beta + \min\{(\alpha - \beta)\theta, (\alpha - \beta)(1-\theta)\}}{(1-\alpha\eta) - \beta(1-\eta)} \right] \end{array}$ 

Proof. —

(i) From equation (5) and (i) of Lemma 3, we get

$$G(t,s) \ge 0 \ \forall t,s \in [0,1].$$

(ii) By equation (5) and (i) of Lemma 3, we have

$$G(t,s) = g(t,s) + \frac{\beta + (\alpha - \beta)t}{(1 - \alpha\eta) - \beta(1 - \eta)}g(\eta,s)$$
  
$$\leq g(s,s) + \frac{\max(\alpha,\beta)}{(1 - \alpha\eta) - \beta(1 - \eta)}g(s,s) = k_1g(s,s).$$

(*iii*) From (*ii*) of Lemma 3, for  $t \in [\theta, 1 - \theta]$  we have

$$\begin{aligned} G(t,s) &= g(t,s) + \frac{\beta + (\alpha - \beta)t}{(1 - \alpha \eta) - \beta(1 - \eta)}g(\eta,s) \\ &\geq \theta g(s,s) + \frac{\beta + \min\{(\alpha - \beta)\theta, (\alpha - \beta)(1 - \theta)\}}{(1 - \alpha \eta) - \beta(1 - \eta)}\theta g(s,s) \\ &\geq \theta \left[1 + \frac{\beta + \min\{(\alpha - \beta)\theta, (\alpha - \beta)(1 - \theta)\}}{(1 - \alpha \eta) - \beta(1 - \eta)}\right]g(s,s) = k_2g(s,s). \end{aligned}$$

**Lemma 5** If  $y \in C([0,1])$  and  $y \ge 0$ , then the unique solution u of the boundary value problem (2), (3) satisfies  $\min_{\substack{\theta \le t \le 1-\theta}} u(t) \ge \gamma ||u||_1$  where  $||u||_1 := \sup\{|u(t)|; 0 \le t \le 1\}$  and  $\gamma := \frac{k_2}{k_1}$ .

**PROOF.** — For any  $t \in [0, 1]$ , by Lemma 4 we have

$$u(t) = \int_0^1 G(t, s) y(s) ds \le k_1 \int_0^1 g(s, s) y(s) ds,$$

thus  $||u||_1 \le k_1 \int_0^1 g(s,s)y(s)ds$ . Moreover, from (*iii*) of Lemma 4 for  $t \in [\theta, 1-\theta]$ , we have

$$u(t) = \int_0^1 G(t,s)y(s)ds \ge k_2 \int_0^1 g(s,s)y(s)ds \ge \frac{k_2}{k_1} ||u||_1.$$

Therefore  $\min_{\theta \le t \le 1-\theta} u(t) \ge \gamma \|u\|_1.$ 

By Lemma 1, we can show that the BVP (2), (3) has a solution u = u(t) if and only if u is a solution of the operator equation u = Tu, where

$$Tu(t) = \begin{cases} \beta u(\eta), & -\tau \le t \le 0, \\ \lambda \int_0^1 G(t,s)a(s)f(s,u(s-\tau))ds, & 0 \le t \le 1. \end{cases}$$

We assume the following hypothesis :

 $(H_1) f \in C([0,1] \times [0,\infty); [0,\infty)),$ 

 $(H_2) \ a \in C([0,1];[0,\infty))$  and there exists  $t_0 \in (0,1)$  such that  $a(t_0) > 0$ , Let define,

$$\begin{split} f^{0} &:= \limsup_{u \to 0} \max_{t \in [0,1]} \frac{f(t,u)}{u}, \quad f^{\infty} := \limsup_{u \to \infty} \max_{t \in [0,1]} \frac{f(t,u)}{u}, \\ M_{1} &:= \beta \int_{0}^{\tau} g(s,s)a(s)ds + \int_{\tau}^{1} g(s,s)a(s)ds \text{ and } M_{2} := \int_{0}^{1} g(s,s)a(s)ds \end{split}$$

The proof of our main results is based upon an application of the following Leray-Schauder fixed point theorem.

#### **Theorem 2.1** ([5])

Let  $\Omega$  be a convex subset of a Banach space  $X, 0 \in \Omega$  and  $\Phi : \Omega \to \Omega$  be a completely continuous operator. Then either

- 1)  $\Phi$  has at least one fixed point in  $\Omega$ , or
- 2) the set  $\{x \in \Omega | x = \mu \Phi x, 0 < \mu < 1\}$  is unbounded.

#### 3. Main results

Let  $X = C[-\tau, 1]$  be a Banach space with norm  $||u|| = \sup\{|u(t)| : -\tau \le t \le 1\}$ .

#### Theorem 3.1

Assume  $(H_1)$  and  $(H_2)$  hold. If  $f^0 < \infty$ , then the boundary value problem (1) has at least one positive solution.

PROOF. — Choose  $\epsilon > 0$  such that  $(f^0 + \epsilon)\lambda k_1 M_1 \leq 1$ . Since  $f^0 < \infty$ , then there exists constant B > 0, such that  $f(s, u) < (f^0 + \epsilon)u$  for  $0 < u \leq B$ . Let

$$\Omega = \{ u \ / u \in C([-\tau, 1]), u \ge 0, \|u\| \le B, \min_{\theta \le t \le 1 - \theta} u(t) \ge \gamma \|u\| \}.$$

Then  $\Omega$  is a convex subset of X.

For  $u \in \Omega$ , by Lemmas 2 and 5, we know that  $Tu(t) \ge 0$  and  $\min_{\theta \le t \le 1-\theta} (Tu)(t) \ge \gamma ||Tu||$ . Moreover

$$Tu \leq \lambda k_1 \int_0^1 g(s,s)a(s)f(s,u(s-\tau))ds$$
  
$$\leq \lambda (f^0 + \epsilon)k_1 \int_0^1 g(s,s)a(s)u(s-\tau)$$
  
$$= \lambda (f^0 + \epsilon)k_1 \left( \int_0^\tau g(s,s)a(s)\beta u(\eta)ds + \int_\tau^1 g(s,s)a(s)u(s-\tau)ds \right)$$

$$\leq \lambda(f^0 + \epsilon)k_1 \left(\beta \int_0^\tau g(s, s)a(s)ds + \int_\tau^1 g(s, s)a(s)ds\right) \|u\|$$
  
$$\leq \|u\| \leq B.$$

Thus,  $||Tu|| \leq B$ . Hence,  $T\Omega \subset \Omega$ .

We shall show that T is completely continuous. Suppose  $u_n \to u$   $(n \to \infty)$  and  $u_n \in \Omega \ \forall n \in \mathbb{N}$ , then there exists M > 0 such that  $||u_n|| \leq M$ .

Since f is continuous on  $[0,1] \times [0,M]$ , it is uniformly continuous.

Therefore,  $\forall \varepsilon > 0$  there exists  $\delta > 0$  such that  $|x - y| < \delta$  implies  $|f(s, x) - f(s, y)| < \epsilon \ \forall s \in [0, 1], x, y \in [0, M]$  and there exists N such that  $||u_n - u|| < \delta$  for n > N, so  $|f(s, u_n(s - \tau)) - f(s, u(s - \tau))| < \varepsilon$ , for n > N and  $s \in [0, 1]$ . This implies

$$\begin{aligned} |Tu_n(t) - Tu(t)| &\leq \lambda k_1 \int_0^1 g(s,s)a(s)|f(s,u_n(s-\tau)) - f(s,u(s-\tau))|ds\\ &\leq \lambda \epsilon k_1 \int_0^1 g(s,s)a(s)ds. \end{aligned}$$

Therefore T is continuous.

Let D be any bounded subset of  $\Omega$ , then there exists  $\gamma > 0$  such that  $||u|| \leq \gamma$  for all  $u \in D$ .

Since f is continuous on  $[0, 1] \times [0, \gamma]$  there exists L > 0 such that  $|f(t, v)| < L \forall (t, v) \in [0, 1] \times [0, \gamma]$ .

Consequently, for all  $u \in D$  and  $t \in [0.1]$  we have

$$\begin{aligned} |Tu(t)| &\leq \left| \lambda k_1 \int_0^1 g(s,s) a(s) f(s,u(s-\tau)) ds \right| \\ &\leq \left| \lambda k_1 L \int_0^1 g(s,s) a(s) ds. \end{aligned}$$

Which implies the boundedness of TD.

Since G is continuous on  $[0, 1] \times [0, 1]$ , it is uniformly continuous.

Then 
$$\forall \epsilon > 0$$
 there exists  $\delta > 0$  such that  $|t_1 - t_2| < \delta$  implies that  $|G(t_1, s) - G(t_2, s)| < \epsilon \ \forall s \in [0, 1]$ . So, if  $u \in D$ ,  $|Tu(t_1) - Tu(t_2)| \leq \lambda \int_0^1 |G(t_1, s) - G(t_2, s)|a(s)f(s, u_n(s - \tau))ds \leq \lambda L\epsilon \int_0^1 a(s)ds$ .

From the arbitrariness of  $\epsilon$ , we get the equicontinuity of TD.

The operator T is completely continuous by the mean of the Ascoli-Arzela theorem. For  $u \in \Omega$  and  $u = \mu T u$ ,  $0 < \mu < 1$ , we have  $u(t) = \mu T u(t) < T u(t) < B$ , which implies  $||u|| \le B$ . So,  $\{x \in \Omega/x = \mu \Phi x, 0 < \mu < 1\}$  is bounded.

By theorem 2.1, we deduce that operator T has at least one fixed point in  $\Omega$ . Thus the

boundary value problem (1) has at least one positive solution.

#### Remark. —

The conditions of Theorem 3.1 are weaker than those of Theorem 3.1 in [3].

#### Theorem 3.2

Assume  $(H_1) - (H_2)$  hold. If  $f^{\infty} < \infty$  is satisfied, then the boundary value problem (1) has at least one positive solution.

PROOF. — Choose  $\epsilon > 0$  such that  $(f^{\infty} + \epsilon)\lambda k_1 M_1 \leq \frac{1}{2}$ . Since  $f^{\infty} < \infty$ , then there exists constant N > 0, such that  $f(s, u) < (f^{\infty} + \epsilon)u$  for u > N. Let B > 0 such that

$$B \ge N + 1 + 2\lambda k_1 M_2 \quad \max_{\substack{0 \le s \le 1\\ 0 \le u \le N}} f(s, u).$$

Let

$$\Omega = \{ u/u \in C[-\tau, 1], \ u \ge 0, \ \|u\| \le B, \min_{\theta \le t \le 1-\theta} u(t) \ge \gamma \|u\| \}.$$

Then  $\Omega$  is a convex subset of X.

For  $u \in \Omega$ , by Lemmas 2 and 5, we have  $Tu(t) \ge 0$  and  $\min_{\theta \le t \le 1-\theta} (Tu)(t) \ge \gamma ||Tu||$ .

Moreover, for  $u \in \Omega$ , we have

$$\begin{aligned} Tu(t) &= \lambda \int_{0}^{1} G(t,s)a(s)f(s,u(s-\tau))ds \\ &\leq \lambda k_{1} \int_{0}^{1} g(s,s)a(s) f(s,u(s-\tau))ds \\ &= \lambda k_{1} \left( \int_{J_{1}=\{s\in[0,1]/u>N\}} g(s,s)a(s)f(s,u(s-\tau))ds \right) \\ &+ \int_{J_{2}=\{s\in[0,1]/u\le N\}} g(s,s)a(s)f(s,u(s-\tau))ds \right) \\ &\leq \lambda k_{1} \left( \int_{0}^{1} g(s,s)a(s)(f^{\infty}+\epsilon)u(s-\tau)ds + \int_{0}^{1} g(s,s)a(s) \max_{\substack{0 \le s \le 1 \\ 0 \le u \le N}} f(s,u(s-\tau))ds \right) \\ &\leq \lambda k_{1} \left( (f^{\infty}+\epsilon) \left[ \beta \int_{0}^{\tau} g(s,s)a(s)ds + \int_{\tau}^{1} g(s,s)a(s)ds \right] \|u\| \\ &+ \int_{0}^{1} g(s,s)a(s) \max_{\substack{0 \le s \le 1 \\ 0 \le u \le N}} f(s,u(s-\tau))ds \right) \end{aligned}$$

$$\leq \lambda(f^{\infty} + \epsilon)k_1M_1B + \lambda k_1M_2 \max_{\substack{0 \leq s \leq 1\\ 0 \leq u \leq N}} f(s, u(s - \tau)) \leq \frac{B}{2} + \frac{B}{2} = B.$$

Thus,  $||Tu|| \leq B$ . Hence,  $T\Omega \subset \Omega$ . We can show that  $T : \Omega \to \Omega$  is completely continuous. For  $u \in \Omega$  and  $u = \mu Tu$ ,  $0 < \mu < 1$ , we have  $u(t) = \mu Tu(t) < Tu(t) < B$ , which implies  $||u|| \leq B$ . So,  $\{x \in \Omega/x = \mu \Phi x, 0 < \mu < 1\}$  is bounded. By theorem 2.1, we show that the operator T has at least one fixed point in  $\Omega$ . Thus, the boundary value problem (1) has at least one positive solution.

Remark. —

The conditions of Theorem 3.2 are weaker than those of Theorem 3.2 in [3].

#### 4. References

- D. BAI AND Y. XU, « Existence of positive solutions for boundary-value problems of second order delay differential equations », *Appl. Math. Lett.*, vol. 18, 2005, 621-630.
- A. CARVALHO, L. A. LADEIRA AND M. MARTELLI, « Forbidden periods in delay differential equation », *Portugal. Math.*, vol. 57, n° 3, 2000, 259-271
- J. CHEN, J. TARIBOON AND S. KOONPRASERT, « Existence of positive solutions to a second order multi point boundary value problem with delay », *Thai Journal of Mathematics*, 2010, 21-32.
- L. H. ERBE AND Q. K. KONG, « Boundary value problems for singular second-order functional differential equations », *J. Comput. Appl. Math.*, vol. 53, 1994, 377–388.
- A. GRANAS AND J. DUGUNDJI, « Fixed point theory », Springer-Verlag, New York, 2003.
- T. JANKOWSKI, « Solvability of three point boundary value problems for second order ordinary differential equations with deviating arguments », *J. Math. Anal. Appl.*, vol. 312, 2005, 620-636.
- D. Q. JIANG, « Multiple positive solutions for boundary value problems of second-order delay differential equations », *Appl. Math. Lett.*, vol. 15, 2002, 575-583.
- J. W. LEE AND D. O'REGAN, « Existence results for differential delay equations-I », J. Differential Equations, vol. 102, 1993, 342-359.
- W. B. WANG AND J. H. SHEN, « Positive solutions to a multi-point boundary value problem with delay », *Appl. Math. Comput.*, vol. 188, 2007, 96-102.
- Z. YAO, « New results of positive solutions for second-order nonlinear three-point integral boundary value problems », *J. Nonlinear Sci. Appl.*, vol. 8, 2015, 93-98.