(2 S ,3 S)-2-(N , N -Dibenzylamino)butane-1,3-diol refined using a multipolar atom model

Krzysztof Ejsmont, Jean-Pierre Joly, Emmanuel Wenger, Christian Jelsch

► To cite this version:

Krzysztof Ejsmont, Jean-Pierre Joly, Emmanuel Wenger, Christian Jelsch. (2 S ,3 S)-2-(N , N -Dibenzylamino)butane-1,3-diol refined using a multipolar atom model. Acta Crystallographica Section C: Crystal Structure Communications [1968-2013], 2008, 64 (1), pp.o18 - o20. 10.1107/S0108270107043296 . hal-01713024

HAL Id: hal-01713024

<https://hal.science/hal-01713024>

Submitted on 20 Feb 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

(2*S*,3*S*)-2-(*N,N*-Dibenzylamino)butane-1,3-diol refined using a multipolar atom model

Krzysztof Ejsmont, Jean-Pierre Joly, Emmanuel Wenger and Christian Jelsch

Acta Cryst. (2008). **C64**, o18–o20

Copyright © International Union of Crystallography

Author(s) of this paper may load this reprint on their own web site or institutional repository provided that this cover page is retained. Republication of this article or its storage in electronic databases other than as specified above is not permitted without prior permission in writing from the IUCr.

For further information see <http://journals.iucr.org/services/authorrights.html>

Acta Crystallographica Section C: Crystal Structure Communications specializes in the rapid dissemination of high-quality studies of crystal and molecular structures of interest in fields such as chemistry, biochemistry, mineralogy, pharmacology, physics and materials science. The numerical and text descriptions of each structure are submitted to the journal electronically as a Crystallographic Information File (CIF) and are checked and typeset automatically prior to peer review. The journal is well known for its high standards of structural reliability and presentation. *Section C* publishes approximately 1000 structures per year; readers have access to an archive that includes high-quality structural data for over 10000 compounds.

Crystallography Journals **Online** is available from journals.iucr.org

**(2*S*,3*S*)-2-(*N,N*-Dibenzylamino)-
butane-1,3-diol refined using a
multipolar atom model**Krzysztof Ejsmont,^a Jean-Pierre Joly,^b Emmanuel Wenger^c
and Christian Jelsch^{c*}

^aInstitute of Chemistry, University of Opole, Oleska 48, 45-052 Opole, Poland,
^bUMR 7565 (SRSMC) – Structure et Réactivité des Systèmes Moléculaires Complexes,
 Université Henri Poincaré Nancy 1, Faculté des Sciences, BP 239, 54506
 Vandoeuvre lès Nancy Cedex, France, and ^cLaboratoire de Cristallographie et
 Modélisation des Matériaux Minéraux et Biologiques (LCM3B), CNRS, UMR 7036,
 Université Henri Poincaré Nancy 1, Faculté des Sciences, BP 239, 54506
 Vandoeuvre lès Nancy Cedex, France
 Correspondence e-mail: christian.jelsch@lcm3b.uhp-nancy.fr

Received 27 June 2007

Accepted 4 September 2007

Online 14 December 2007

The crystal structure of the title compound, C₁₈H₂₃NO₂, was determined using the experimental library multipolar atom model. The refinement showed a significant improvement of crystallographic statistical indices when compared with a conventional spherical neutral atom refinement.

Comment

Enantiomerically pure amino acids and their derivatives play an important role as indispensable intermediates in the pharmaceutical industry and agrochemistry (Breuer *et al.*, 2004, and references therein). Among these optically active amino-alcohols, L- and/or D-threoninol have recently received much attention as chiral parts of promising drugs against chemokine-mediated diseases (Brough & McNally, 2004), for the synthesis of acridine–DNA conjugates for site-selective RNA scissors (Shi *et al.*, 2005), as linkers for Methyl Red incorporation into DNA (Kashida, Tanaka *et al.*, 2006), as asymmetric *cis*-platinum(II) complexes for cancer therapy (van Rijn *et al.*, 2006), as intercalators between pyrene moieties and an oligodeoxyribonucleotide for the detection of deletion polymorphisms (Kashida, Asanuma *et al.*, 2006) or between 10–23 DNAzyme and a binding arm for RNA cleavage activity enhancement (Asanuma *et al.*, 2006), and for the synthesis of inhibitors of protein kinase C for the treatment of inflammation (Peng *et al.*, 2006) or of a small combinatorial library of dihydroceramide analogs (Villorbin *et al.*, 2007).

As part of a programme for the evaluation of new asymmetric catalysts from the chiral pool (Waykole *et al.*, 2007) (*i.e.* amino acids, sugars, *etc.*), we have synthesized the new optically pure α -amino alcohol (II) in two steps from readily available D-threonine. Diol (II) is a convenient source of D-threoninol by catalytic hydrogenolysis under mild condi-

tions with toluene as the sole by-product (Yoshida *et al.*, 1988). We report here the X-ray structure of (II), which was prepared in good yield by reduction of the intermediate ester, (I).

The molecular structure of (II) is depicted in Fig. 1. An intramolecular O1–HO1···N1 hydrogen bond (Table 2) forms a five-membered ring. Atom N1 lies about 0.407 Å out of the plane defined by the three neighbouring C atoms (C1, C5 and C12), towards atom HO1 (Fig. 1). The sum of the valence angles around the N atom is 338.9°, indicating *sp*³-hybridization (328° for *sp*³ and 360° for *sp*²). The crystal structure of (II) is presented in Fig. 2. Chains along the *b* axis are formed by O2–HO2···O1ⁱ hydrogen bonds [symmetry code: (i) *x*, *y* + 1, *z*; Table 2].

Initially, in the independent atom model (IAM) refinement, a conventional spherical neutral atom model was applied. Scale factors, atomic positions and displacement parameters for all atoms were refined using the *MoPro* program (Guillot *et al.*, 2001; Jelsch *et al.*, 2005) until convergence. In the experimental library multipolar atom model (ELMAM) refinement, the same parameters were varied but a multipolar charged atom model was applied. The electron-density parameters were transferred from the ELMAM library (Pichon-Pesme *et al.*, 2004; Zarychta *et al.*, 2007) and subsequently kept fixed. Riding restraints on H-atom *B* factors were applied similarly in both refinements, which were carried out with the same intensity data and cut-off criterion [*I*/ σ (*I*) ≥ 0]. The H–X distances were restricted to standard values in X-ray and

Figure 1

The molecular structure of (II), showing the intramolecular hydrogen bond (dashed line). Displacement ellipsoids are drawn at the 50% probability level and H atoms are shown as small spheres of arbitrary radii.

Figure 2
The packing of (II), showing the arrangement of the chains.

neutron diffraction studies (Allen, 1986) in the IAM and ELMAM refinements, respectively (0.002 Å distance σ). The ELMAM refinement shows a good improvement in statistical indexes when compared with the IAM refinement; the $R(F)$ factor is reduced from 4.22 to 3.02% and $wR(R)$ from 5.12 to 3.37%. The minimum and maximum peaks in the residual electron density are -0.036 and 0.056 e Å^{-3} , respectively, after the IAM refinement and -0.028 and 0.039 e Å^{-3} after the ELMAN refinement. The most significant differences in the geometry of compound (II) are, as expected, the bond distances involving H atoms, which had different targets in the ELMAN and IAM refinements. When H atoms are not considered, the r.m.s. deviation between the two structures is 0.0067 Å , while the r.m.s. discrepancy for the bond lengths is 0.0055 Å . The differences do not exceed 0.015 Å for bond distances and 0.5° for bond angles when H atoms are excluded. The H—C—H angles show the strongest (2.9°) r.m.s. discrepancy, the X—Y—H angles differ by 1.8° , while the X—Y—Z angles are less affected, with a low (0.2°) r.m.s. difference. The s.u. values of the bond distances and angles are smaller with the ELMAM refinement owing to better R factor values (Table 1).

The largest effect of the multipoles transfer on the crystallographic structure is observed in the atomic thermal motion. The r.m.s. difference over all the U^{ij} parameters reaches 16% between the two refinement models. The U^{ij} values derived from the IAM refinement have an r.m.s. value 10% larger on average than those derived from the ELMAM refinement. With the independent spherical atom model, the displacement parameters are incorrect as they incorporate some significant deformation electron density due to improper deconvolution between these two features (Jelsch *et al.*, 1998).

Experimental

For the preparation of (2*S*,3*S*)-benzyl 2-(*N,N*-dibenzylamino)-3-hydroxybutanoate, (I), benzyl bromide (6.6 ml, 55 mmol, 3.3 equivalents) was dropped over a period of 30 min on to a magneti-

cally stirred dispersion of 98% D-threonine (2.02 g, 16.6 mmol) and Na_2CO_3 (3.875 g, 19.5 mmol, ~ 2.2 equivalents) in 75% aqueous EtOH (50 ml) at room temperature. The resulting mixture was refluxed for 2.5 h [the formation of (I) being monitored by SiO_2 thin-layer chromatography (TLC); $R_F = 0.43$; AcOEt–cyclohexane 1:4], cooled to room temperature, concentrated under reduced pressure, and partitioned between dichloromethane and water ($2 \times 50 \text{ ml}$). The aqueous phase was extracted with dichloromethane ($3 \times 50 \text{ ml}$). The combined organic phases were washed with saturated aqueous NaHCO_3 (50 ml) and water ($3 \times 50 \text{ ml}$), dried over MgSO_4 , and finally concentrated *in vacuo* to afford the crude ester (I) as a pale-yellow syrup, which was used for the next step without further purification (yield 5.8 g, 14.9 mmol, $\sim 90\%$). An analytical sample was isolated by TLC (SiO_2 , AcOEt–hexane 1:9) as a colourless gum. $[\alpha]_D^{25} = +148.9^\circ$ ($c = 2.6$, CHCl_3). MS (70 eV, EI^+) calculated for $\text{C}_{25}\text{H}_{27}\text{NO}_3 = 389$; found m/z (%) = 390 (23) $[M + \text{H}]^+$, 344 (32) $[M - \text{C}_2\text{H}_5\text{O}]^+$. IR (film): 1730 cm^{-1} . ^1H NMR (250 MHz, CDCl_3 , 298 K): δ 7.15–7.55 (15H, *m*, aromatic), 5.35 (1H, *d*, $J = 12.4 \text{ Hz}$, CHHPh), 5.22 (1H, *d*, CHHPh), 4.07 (1H, *m*, H- β), 4.07 (2H, *d*, $J = 13.2 \text{ Hz}$, NCHHPh), 3.5 (1H, *s*, OH), 3.39 (2H, *d*, NCHHPh), 3.11 (1H, *d*, $J = 9.5 \text{ Hz}$, H- α), 1.1 (3H, *d*, $J = 5.8 \text{ Hz}$, CH_3). ^{13}C NMR (100 MHz, CDCl_3): δ 139.45 (C aromatic), 135.7 (C aromatic), 129.1, 128.7, 128.6, 128.5, 127.4 (CH aromatic), 67.2 (C- β), 66.3 (PhCH $_2$ O), 63.2 (C- α), 54.8 (PhCH $_2$ N), 19.2 (CH_3). Elemental analysis calculated for $\text{C}_{25}\text{H}_{27}\text{NO}_3$: C 77.09, H 6.99, N 3.60%; found: C 76.91, H 6.81, N 3.67%.

For the preparation of (II), a solution of crude ester (I) (5.453 g, 14.0 mmol) in absolute ether (50 ml) was dropped over a period of 30 min on to a magnetically stirred suspension of 95% LiAlH_4 (0.78 g, ~ 19.5 mmol, 1.15 equivalents) in absolute tetrahydrofuran (THF, 50 ml) under argon cooled below 277 K. The mixture was stirred for 25 h at room temperature and refluxed for an additional hour. After completion of the reaction (checked by SiO_2 TLC; ethyl acetate–*n*-hexane 1:1; $R_F \approx 0.29$), the mixture was cooled to room temperature, quenched by slow addition of ethyl acetate (5 ml) and saturated aqueous Na_2SO_4 (5 ml), and stirred overnight in air. The suspension was filtered through sintered glass and the remaining salts thoroughly washed with a dichloromethane–ethanol mixture (1:1, 100 ml). The filtrates were concentrated under reduced pressure and the residue partitioned in dichloromethane–water ($2 \times 50 \text{ ml}$), washed with water ($2 \times 20 \text{ ml}$), dried over MgSO_4 , concentrated under reduced pressure, and finally stored at 277 K. The resulting solids were isolated by filtration and crystallized twice from ethyl acetate–hexanes (1:19) to yield pure alcohol (II) (3.2 g, $\sim 73\%$) as white crystals suitable for X-ray diffraction (m.p. 363–364 K). $[\alpha]_D^{25} = +54.0^\circ$ ($c = 1$, CHCl_3). MS (70 eV, EI^+): m/z 286.1 $[M + \text{H}]^+$. ^1H NMR (250 MHz, $\text{CDCl}_3/\text{D}_2\text{O}$): δ 7.2–7.42 (*m*, 10H, aromatic), 3.99 (2H, *d*, $J = 13.2 \text{ Hz}$, NCHHPh), 3.85 (1H, *m*, H- β), 3.8 (2H, *d*, $J = 13.2 \text{ Hz}$, $J = 5.8 \text{ Hz}$, CH_2OD), 3.72 (2H, *d*, NCHHPh), 2.61 (1H, *d*, $J = 8.8 \text{ Hz}$, H- α), 1.15 (3H, *d*, $J = 5.8 \text{ Hz}$, CH_3). ^{13}C NMR (100 MHz, CDCl_3): δ 139.5 (C aromatic), 129.4, 128.7, 127.5 (CH aromatic), 65.5 (C- β), 64.8 (C- α), 59.1 (PhCH $_2$), 54.7 ($\text{CH}_2\text{—OH}$), 20.4 (CH_3). Elemental analysis calculated for $\text{C}_{18}\text{H}_{23}\text{NO}_2$: C 75.76, H 8.12, N 4.91%; found: C 75.79, H 8.09, N 4.90%.

Crystal data

$\text{C}_{18}\text{H}_{23}\text{NO}_2$
 $M_r = 285.37$
Monoclinic, C_2
 $a = 18.6701$ (4) Å
 $b = 6.652$ (1) Å
 $c = 16.060$ (2) Å
 $\beta = 123.126$ (8)°

$V = 1670.4$ (4) Å 3
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.07 \text{ mm}^{-1}$
 $T = 100$ (2) K
 $0.35 \times 0.25 \times 0.20 \text{ mm}$

Data collection

Oxford Diffraction Xcalibur2 diffractometer	2016 independent reflections
8273 measured reflections	1966 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.022$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$	46 restraints
$wR(F^2) = 0.034$	H-atom parameters constrained
$S = 1.04$	$\Delta\rho_{\text{max}} = 0.05 \text{ e } \text{\AA}^{-3}$
2016 reflections	$\Delta\rho_{\text{min}} = -0.03 \text{ e } \text{\AA}^{-3}$
199 parameters	

Table 1
Hydrogen-bond geometry ($\text{\AA}$, $^\circ$).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$O1-HO1\cdots N1$	0.97	2.06	2.729 (3)	125
$O2-HO2\cdots O1^i$	0.97	1.87	2.789 (2)	158

Symmetry code: (i) $x, y + 1, z$.

Table 2
Selected geometric data for (II) ($\text{\AA}$, $^\circ$).

ELMAM is the experimental library multipolar atom model and IAM is the independent atom model.

Distance/angle	ELMAM	IAM
$O1-C2$	1.465 (2)	1.461 (3)
$O2-C4$	1.436 (2)	1.435 (3)
$N1-C1$	1.502 (2)	1.502 (3)
$N1-C5$	1.506 (1)	1.504 (1)
$N1-C12$	1.496 (2)	1.500 (4)
$O1-C2-C1$	107.3 (2)	107.3 (4)
$O1-C2-C3$	108.2 (2)	108.4 (4)
$O2-C4-C1$	112.5 (3)	112.5 (5)
$C1-N1-C5$	112.6 (3)	112.8 (5)
$C1-N1-C12$	115.8 (3)	115.7 (4)
$C5-N1-C12$	110.5 (3)	110.4 (5)

At first, a least-squares refinement, based on $|F^2|$, was carried out using the program *SHELXL97* (Sheldrick, 1997), the H atoms being constrained according to standard crystallography stereochemistry. A least-squares refinement, based on $|F|$, was then carried out with the program *MoPro* (Guillot *et al.*, 2001; Jelsch *et al.*, 2005) using the ELMAM approach (Zarychta *et al.*, 2007). The reflection weights were set equal to $1/\sigma^2(F_o)$. The $U_{\text{iso}}(\text{H})$ values were restrained to be $1.2U_{\text{eq}}$ of the attached atom, with a standard deviation of $0.01 \text{ \AA}$. In the absence of suitable anomalous scattering, the refinement of the Flack (1983) parameter led to inconclusive values; therefore Friedel equivalent reflections were merged prior to the final refinements. The absolute structure was set by reference to the known chirality of the enantiomerically pure D-threonine used in the chemical synthesis.

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2006); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2006); data reduc-

tion: *CrysAlis RED*; program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1993); program(s) used to refine structure: *MoPro* (Guillot *et al.*, 2001; Jelsch *et al.*, 2005); molecular graphics: *SHELXTL* (Sheldrick, 1990); software used to prepare material for publication: *publCIF* (Westrip, 2007).

This work was partially supported by the Agence Nationale de la Recherche, programme Blanc.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: DN3056). Services for accessing these data are described at the back of the journal.

References

- Allen, F. H. (1986). *Acta Cryst.* **B42**, 515–522.
- Altomare, A., Casciarano, G., Giacovazzo, C. & Guagliardi, A. (1993). *J. Appl. Cryst.* **26**, 343–350.
- Asanuma, H., Hayashi, H., Zhao, J., Liang, X., Yamazawa, A., Kuramochi, T., Matsunaga, D., Aiba, Y., Kashida, H. & Komiyama, M. (2006). *Chem. Commun.* **48**, 5062–5064.
- Breuer, M., Ditrich, K., Habicher, T., Hauer, B., Kessler, M., Stürmer, R. & Zelinski, T. (2004). *Angew. Chem. Int. Ed.* **43**, 788–824.
- Brough, S. J. & McNally, T. (2004). Patent No. WO2003-GB400020030916, 1–24.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Guillot, B., Viry, L., Guillot, R., Lecomte, C. & Jelsch, C. (2001). *J. Appl. Cryst.* **34**, 214–223.
- Jelsch, C., Guillot, B., Lagoutte, A. & Lecomte, C. (2005). *J. Appl. Cryst.* **38**, 38–54.
- Jelsch, C., Pichon-Pesme, V., Lecomte, C. & Aubry, A. (1998). *Acta Cryst.* **D54**, 1306–1318.
- Kashida, H., Asanuma, H. & Komiyama, M. (2006). *Chem. Commun.* **26**, 2768–2770.
- Kashida, H., Tanaka, M., Baba, S., Sakamoto, T., Kawai, G., Asanuma, H. & Komiyama, M. (2006). *Chem. Eur. J.* **12**, 777–784.
- Oxford Diffraction (2006). *CrysAlis CCD* and *CrysAlis RED*. Versions 1.171.31.7. Oxford Diffraction Ltd, Abingdon, Oxfordshire, England.
- Peng, S., Zhao, M., Wang, C. & Gu, K. (2006). Chinese Patent CN1785995 A 20060614, 1–38.
- Pichon-Pesme, V., Jelsch, C., Guillot, B. & Lecomte, C. (2004). *Acta Cryst.* **A60**, 204–208.
- Rijt, S. van, van Zutphen, S., den Dulk, H., Brouwer, J. & Reedijk, J. (2006). *Inorg. Chim. Acta*, **359**, 4125–4129.
- Sheldrick, G. M. (1990). *SHELXTL*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (1997). *SHELXL97*. University of Göttingen, Germany.
- Shi, Y., Machida, K., Kuzuya, A. & Komiyama, M. (2005). *Bioconjug. Chem.* **16**, 306–311.
- Villorina, G., Canals, D., Carde, L., Grijalvo, S., Pascual, R., Rabal, O., Teixido, J., Fabrias, G., Llebaria, A., Casas, J. & Delgado, A. (2007). *Bioorg. Med. Chem.* **15**, 50–62.
- Waykole, L. M., McKenna, J. J., Bach, A., Prashad, M., Repic, O. & Blacklock, T. J. (2007). *Synth. Commun.* **37**, 1445–1454.
- Westrip, S. P. (2007). *publCIF*. In preparation.
- Yoshida, K., Nakajima, S., Wakamatsu, T., Ban, Y. & Shibasaki, M. (1988). *Heterocycles*, **27**, 1167–1168.
- Zarychta, B., Pichon-Pesme, V., Guillot, B., Lecomte, C. & Jelsch, C. (2007). *Acta Cryst.* **A63**, 108–125.