Explicit expressions related to degenerate Cauchy numbers and their generating function

Feng Qi, Ai-Qi Liu, Dongkyu Lim

► To cite this version:

Feng Qi, Ai-Qi Liu, Dongkyu Lim. Explicit expressions related to degenerate Cauchy numbers and their generating function: Explicit expressions related to degenerate Cauchy numbers. International Workshop of Mathematical Modelling, Applied Analysis and Computation, Jul 2018, Jaipur, India. pp.41–52, 10.1007/978-981-13-9608-3_2 . hal-01725045v2

HAL Id: hal-01725045

<https://hal.science/hal-01725045v2>

Submitted on 19 Jan 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EXPLICIT EXPRESSIONS RELATED TO DEGENERATE CAUCHY NUMBERS AND THEIR GENERATING FUNCTION

FENG QI, AI-QI LIU, AND DONGKYU LIM

ABSTRACT. In the paper, by virtue of the Faà di Bruno formula and two identities for the Bell polynomials of the second kind, the authors establish an explicit expression for degenerate Cauchy numbers and find explicit, meaningful, and significant expressions for coefficients in a family of nonlinear differential equations for the generating function of degenerate Cauchy numbers.

1. MOTIVATIONS AND MAIN RESULTS

It is well known [4, 7, 8, 34, 39] that the Cauchy numbers of the first kind C_n can be generated by

$$\frac{t}{\ln(1+t)} = \sum_{n=0}^{\infty} C_n \frac{t^n}{n!}.$$

In [2], degenerate Cauchy numbers $C_n(\lambda)$ were defined by

$$\frac{\lambda[e^{[(1+t)^\lambda-1]/\lambda} - 1]}{(1+t)^\lambda - 1} = \sum_{n=0}^{\infty} C_n(\lambda) \frac{t^n}{n!}.$$

Because

$$\lim_{\lambda \rightarrow 0} \frac{(1+t)^\lambda - 1}{\lambda} = \ln(1+t) \quad \text{or} \quad \lim_{\lambda \rightarrow 0} e^{[(1+t)^\lambda-1]/\lambda} = 1+t,$$

it follows that

$$\lim_{\lambda \rightarrow 0} C_n(\lambda) = C_n, \quad n \geq 0. \quad (1)$$

In [2, Theorem 2.1], it was established that the family of nonlinear differential equations

$$(1+t)^n [(1+t)^\lambda - 1]^n F_\lambda^{(n)}(t) = F_\lambda(t) \sum_{i=1}^{2n} a_i(n, \lambda) (1+t)^{i\lambda} + \sum_{i=1}^{2n-1} b_i(n, \lambda) (1+t)^{i\lambda} \quad (2)$$

2010 *Mathematics Subject Classification.* Primary 11B68; Secondary 11B83, 33B10, 33B99, 34A05, 34A34.

Key words and phrases. explicit expression; degenerate Cauchy number; coefficient; nonlinear differential equation; generating function; Bell polynomial of the second kind; Faà di Bruno formula.

Please cite this article as “Feng Qi, Ai-Qi Liu, and Dongkyu Lim, *Explicit expressions related to degenerate Cauchy numbers and their generating function*, In: J. Singh, D. Kumar, H. Dutta, D. Baleanu, and S. Purohit (eds), *Mathematical Modelling, Applied Analysis and Computation*, ICMMAAC 2018 (Jaipur, India, July 6-8, 2018), Springer Proceedings in Mathematics & Statistics, vol. 272, Chapter 2, pp. 41–52, Springer, Singapore, 2019; available online at https://doi.org/10.1007/978-981-13-9608-3_2.”

for $n \in \mathbb{N}$ has the same solution

$$F_\lambda(t) = \frac{e^{[(1+t)^\lambda - 1]/\lambda} - 1}{(1+t)^\lambda - 1}, \quad (3)$$

where $a_i(n, \lambda)$ for $1 \leq i \leq 2n$ and $b_i(n, \lambda)$ for $1 \leq i \leq 2n-1$ are uniquely determined by

$$\begin{aligned} a_1(n, \lambda) &= -\frac{1}{\lambda} \langle n-1-\lambda \rangle_{n+1}, \quad a_2(n, \lambda) = \langle n-1-2\lambda \rangle_{n-1} \\ &\quad - \frac{1}{\lambda} \sum_{i=0}^{n-2} [\lambda - (\lambda+1)(n-i)] \langle n-i-2-\lambda \rangle_{n-i} \langle n-1-2\lambda \rangle_i, \\ a_i(n, \lambda) &= [(i-1)\lambda - (\lambda+1)n] a_i(n-1, \lambda) \\ &\quad + a_{i-2}(n-1, \lambda) + (n-1-i\lambda) a_i(n-1, \lambda), \quad 3 \leq i \leq 2n-2, \\ a_{2n-1}(n, \lambda) &= \frac{1}{2} n [(\lambda-1)(n-1) - 2(\lambda+1)], \\ a_{2n}(n, \lambda) &= 1, \quad b_1(n, \lambda) = \langle n-1-\lambda \rangle_{n-1}, \\ b_i(n, \lambda) &= [(i-1)\lambda - (\lambda+1)(n-1)] b_{i-1}(n-1, \lambda) \\ &\quad + a_{i-1}(n-1, \lambda) + (n-1-i\lambda) b_i(n-1, \lambda), \quad 2 \leq i \leq 2n-3, \\ b_{2n-2}(n, \lambda) &= (\lambda-1) \binom{n-1}{2} - 2(n-1) - \lambda, \quad b_{2n-1}(n, \lambda) = 1 \end{aligned}$$

in terms of the falling factorials

$$\langle x \rangle_n = \prod_{k=0}^{n-1} (x-k) = \begin{cases} x(x-1) \cdots (x-n+1), & n \geq 1; \\ 1, & n = 0. \end{cases}$$

It is clear that the generating function $F_\lambda(t)$ defined by (3) satisfies

$$\lim_{\lambda \rightarrow 0} [\lambda F_\lambda(t)] = \frac{t}{\ln(1+t)}.$$

It is obvious that

- (1) the above expressions for $a_i(n, \lambda)$ and $b_i(n, \lambda)$ are recursive and can not be computed easily;
- (2) the original proof of [2, Theorem 2.1] is inductive, recursive, and long;
- (3) there was no any application given in [2].

In this paper, by virtue of the Faà di Bruno formula (9) and two identities (10) and (11) for the Bell polynomials of the second kind $B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$, we will establish an explicit expression (4) for degenerate Cauchy numbers $C_n(\lambda)$ and find explicit, meaningful, and significant expressions (7) and (8) for coefficients $a_i(n, \lambda)$ and $b_i(n, \lambda)$ in the family of nonlinear differential equations (2) related to the generating function $F_\lambda(t)$ of degenerate Cauchy numbers $C_n(\lambda)$.

Our main results can be stated as the following theorems.

Theorem 1. *For $n \geq 0$, degenerate Cauchy numbers $C_n(\lambda)$ and the Cauchy numbers C_n can be explicitly and respectively computed by*

$$C_n(\lambda) = \sum_{k=0}^n \frac{(-1)^k}{(k+1)! \lambda^k} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \langle \ell \lambda \rangle_n \quad (4)$$

and

$$C_n = \sum_{k=0}^n \frac{s(n, k)}{k+1}. \quad (5)$$

Theorem 2. For $n \in \mathbb{N}$, the generating function $F_\lambda(t)$ and its derivatives of degenerate Cauchy numbers $C_n(\lambda)$ satisfy

$$F_\lambda^{(n)}(t) = \frac{F_\lambda(t)}{(1+t)^n [1 - (1+t)^\lambda]^n} \sum_{i=1}^{2n} \alpha_i(n, \lambda) (1+t)^{i\lambda} + \sum_{i=1}^{2n-1} \beta_i(n, \lambda) (1+t)^{i\lambda} \quad (6)$$

with

$$\alpha_i(n, \lambda) = \sum_{\substack{k+m=i \\ 1 \leq k \leq n \\ 0 \leq m \leq n}} (-1)^m A_k(n, \lambda) \sum_{\ell=0}^{\min\{n-m, k\}} \frac{\lambda^\ell}{(k-\ell)!} \binom{n-\ell}{m} \quad (7)$$

for $1 \leq i \leq 2n$ and

$$\beta_i(n, \lambda) = \sum_{\substack{k+m=i \\ 1 \leq k \leq n \\ 0 \leq m \leq n-1}} (-1)^{m+1} A_k(n, \lambda) \sum_{\ell=0}^{\min\{k-1, n-m-1\}} \frac{\lambda^\ell}{(k-\ell)!} \binom{n-\ell-1}{m} \quad (8)$$

for $1 \leq i \leq 2n-1$, where

$$A_k(n, \lambda) = \frac{(-1)^k}{\lambda^k} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \prod_{q=0}^{n-1} (\ell\lambda - q).$$

2. LEMMAS

In order to obtain our main results, we need the following lemmas.

Lemma 1 ([1, pp. 134 and 139]). For $n \geq k \geq 0$, the Bell polynomials of the second kind, denoted by $B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$, are defined by

$$B_{n,k}(x_1, x_2, \dots, x_{n-k+1}) = \sum_{\substack{1 \leq i \leq n-k+1 \\ \ell_i \in \{0\} \cup \mathbb{N} \\ \sum_{i=1}^{n-k+1} i\ell_i = n \\ \sum_{i=1}^{n-k+1} \ell_i = k}} \frac{n!}{\prod_{i=1}^{n-k+1} \ell_i!} \prod_{i=1}^{n-k+1} \left(\frac{x_i}{i!} \right)^{\ell_i}.$$

The Faà di Bruno formula can be described in terms of the Bell polynomials of the second kind $B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$ by

$$\frac{d^n}{dt^n} f \circ h(t) = \sum_{k=0}^n f^{(k)}(h(t)) B_{n,k}(h'(t), h''(t), \dots, h^{(n-k+1)}(t)). \quad (9)$$

Lemma 2 ([1, p. 135]). For $n \geq k \geq 0$, we have

$$B_{n,k}(abx_1, ab^2x_2, \dots, ab^{n-k+1}x_{n-k+1}) = a^k b^n B_{n,k}(x_1, x_2, \dots, x_{n-k+1}), \quad (10)$$

where a and b are any complex numbers.

Lemma 3. For $n \geq k \geq 0$ and $\lambda, \alpha \in \mathbb{C}$, we have

$$B_{n,k} \left(1, 1 - \lambda, (1 - \lambda)(1 - 2\lambda), \dots, \prod_{\ell=0}^{n-k} (1 - \ell\lambda) \right) = \frac{(-1)^k}{k!} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \prod_{q=0}^{n-1} (\ell - q\lambda) \quad (11)$$

or, equivalently,

$$B_{n,k}(\langle \alpha \rangle_1, \langle \alpha \rangle_2, \dots, \langle \alpha \rangle_{n-k+1}) = \frac{(-1)^k}{k!} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \langle \alpha \ell \rangle_n. \quad (12)$$

Proof. This explicit formula (11) was first established in [25, Remark 1] and then was applied in [16, Section 2], [17, First proof of Theorem 2], [19, Lemma 2.2], [22, Remark 6.1], [23, Lemma 4], and [32, Lemma 2.6]. The formula (12) and the equivalence were presented in [33, Theorems 2.1 and 4.1]. $\square$

3. PROOFS OF THEOREMS 1 AND 2

We are now in a position to prove our main results.

Proof of Theorem 1. For $n \geq 0$, applying $u = h(t) = \frac{(1+t)^\lambda - 1}{\lambda}$ and $f(u) = \frac{e^u - 1}{u}$ to (9) and making use of (10) and (11) in sequence arrive at

$$\begin{aligned} \frac{d^n[\lambda F_\lambda(t)]}{d t^n} &= \sum_{k=0}^n \frac{d^k}{d u^k} \left(\frac{e^u - 1}{u} \right) B_{n,k} \left(\frac{\lambda(1+t)^{\lambda-1}}{\lambda}, \frac{\lambda(\lambda-1)(1+t)^{\lambda-2}}{\lambda}, \right. \\ &\quad \left. \dots, \frac{\lambda(\lambda-1) \cdots [\lambda - (n-k)](1+t)^{\lambda-(n-k+1)}}{\lambda} \right) \\ &= \sum_{k=0}^n \frac{d^k}{d u^k} \left(\sum_{\ell=1}^{\infty} \frac{u^{\ell-1}}{\ell!} \right) (1+t)^{k\lambda-n} B_{n,k}(1, \lambda-1, \dots, (\lambda-1) \cdots [\lambda - (n-k)]) \\ &= \sum_{k=0}^n \frac{d^k}{d u^k} \left(\sum_{\ell=0}^{\infty} \frac{u^\ell}{(\ell+1)!} \right) (1+t)^{k\lambda-n} \\ &\quad \times B_{n,k} \left(1, \lambda \left(1 - \frac{1}{\lambda} \right), \dots, \lambda^{n-k} \left(1 - \frac{1}{\lambda} \right) \cdots \left(1 - \frac{n-k}{\lambda} \right) \right) \\ &= \sum_{k=0}^n \left[\sum_{\ell=0}^{\infty} \frac{d^k}{d u^k} \left(\frac{u^\ell}{(\ell+1)!} \right) \right] (1+t)^{k\lambda-n} \\ &\quad \times \lambda^{n-k} B_{n,k} \left(1, 1 - \frac{1}{\lambda}, \dots, \left(1 - \frac{1}{\lambda} \right) \cdots \left(1 - \frac{n-k}{\lambda} \right) \right) \\ &= \sum_{k=0}^n \left[\sum_{\ell=k}^{\infty} \frac{\ell!}{(\ell-k)!} \frac{u^{\ell-k}}{(\ell+1)!} \right] (1+t)^{k\lambda-n} \lambda^{n-k} \frac{(-1)^k}{k!} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \prod_{q=0}^{n-1} \left(\ell - \frac{q}{\lambda} \right) \\ &= \sum_{k=0}^n \left[\sum_{\ell=k}^{\infty} \frac{1}{(\ell-k)!} \frac{u^{\ell-k}}{\ell+1} \right] (1+t)^{k\lambda-n} \frac{1}{\lambda^k} \frac{(-1)^k}{k!} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \prod_{q=0}^{n-1} (\ell\lambda - q) \\ &\rightarrow \sum_{k=0}^n \frac{1}{k+1} \frac{1}{\lambda^k} \frac{(-1)^k}{k!} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \prod_{q=0}^{n-1} (\ell\lambda - q) \end{aligned}$$

$$= \sum_{k=0}^n \frac{(-1)^k}{(k+1)! \lambda^k} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \langle \ell \lambda \rangle_n$$

as $t \rightarrow 0$ and, consequently, $u \rightarrow 0$. This implies that

$$C_n(\lambda) = \lambda \lim_{t \rightarrow 0} F_\lambda^{(n)}(t) = \sum_{k=0}^n \frac{(-1)^k}{(k+1)! \lambda^k} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \langle \ell \lambda \rangle_n.$$

The explicit formula (4) is thus proved.

It is well known [1, Theorem A] that the Stirling numbers of the first kind $s(n, k)$ can be generated by

$$\langle x \rangle_n = \sum_{k=0}^n s(n, k) x^k.$$

Hence, by the L'Hôpital rule, we have

$$\begin{aligned} \lim_{\lambda \rightarrow 0} \frac{1}{\lambda^k} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \langle \ell \lambda \rangle_n &= \lim_{\lambda \rightarrow 0} \frac{1}{\lambda^k} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \sum_{m=0}^n s(n, m) (\ell \lambda)^m \\ &= \frac{1}{k!} \lim_{\lambda \rightarrow 0} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \sum_{m=0}^n s(n, m) \langle m \rangle_k \ell^k (\ell \lambda)^{m-k} \\ &= \frac{1}{k!} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} s(n, k) \langle k \rangle_k \ell^k = s(n, k) \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \ell^k = (-1)^k k! s(n, k). \end{aligned}$$

Combining this with (1) and (4) gives

$$C_n = \lim_{\lambda \rightarrow 0} C_n(\lambda) = \sum_{k=0}^n \frac{(-1)^k}{(k+1)!} \lim_{\lambda \rightarrow 0} \frac{1}{\lambda^k} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \langle \ell \lambda \rangle_n = \sum_{k=0}^n \frac{s(n, k)}{k+1}.$$

The explicit formula (5) follows. The proof of Theorem 1 is complete. $\square$

Proof of Theorem 2. For $n \in \mathbb{N}$, applying $u = h(t) = \frac{(1+t)^\lambda - 1}{\lambda}$ and $f(u) = \frac{e^u - 1}{u}$ to (9) and making use of (10) and (11) in sequence arrive at

$$\begin{aligned} \frac{d^n [\lambda F_\lambda(t)]}{d t^n} &= \sum_{k=1}^n \frac{d^k}{d u^k} \left(\frac{e^u - 1}{u} \right) B_{n,k} \left(\frac{\lambda(1+t)^{\lambda-1}}{\lambda}, \frac{\lambda(\lambda-1)(1+t)^{\lambda-2}}{\lambda}, \right. \\ &\quad \left. \dots, \frac{\lambda(\lambda-1) \cdots [\lambda - (n-k)](1+t)^{\lambda-(n-k+1)}}{\lambda} \right) \\ &= \sum_{k=1}^n \left[\sum_{\ell=0}^k \binom{k}{\ell} (e^u - 1)^{(\ell)} \left(\frac{1}{u} \right)^{(k-\ell)} \right] (1+t)^{k\lambda-n} \\ &\quad \times B_{n,k}(1, \lambda-1, \dots, (\lambda-1) \cdots [\lambda - (n-k)]) \\ &= \sum_{k=1}^n \left[\frac{(-1)^k k! (e^u - 1)}{u^{k+1}} + \sum_{\ell=1}^k \binom{k}{\ell} e^u \frac{(-1)^{k-\ell} (k-\ell)!}{u^{k-\ell+1}} \right] (1+t)^{k\lambda-n} \\ &\quad \times B_{n,k} \left(1, \lambda \left(1 - \frac{1}{\lambda} \right), \dots, \lambda^{n-k} \left(1 - \frac{1}{\lambda} \right) \cdots \left(1 - \frac{n-k}{\lambda} \right) \right) \\ &= \sum_{k=1}^n \left[\lambda F_\lambda(t) \sum_{\ell=0}^k \binom{k}{\ell} \frac{(-1)^{k-\ell} (k-\ell)!}{u^{k-\ell}} + \sum_{\ell=1}^k \binom{k}{\ell} \frac{(-1)^{k-\ell} (k-\ell)!}{u^{k-\ell+1}} \right] (1+t)^{k\lambda-n} \end{aligned}$$

$$\begin{aligned}
& \times \lambda^{n-k} B_{n,k} \left(1, 1 - \frac{1}{\lambda}, \dots, \left(1 - \frac{1}{\lambda} \right) \cdots \left(1 - \frac{n-k}{\lambda} \right) \right) \\
& = \sum_{k=1}^n \left[\lambda F_{\lambda}(t) \sum_{\ell=0}^k \binom{k}{\ell} \frac{(-1)^{k-\ell} \lambda^{k-\ell} (k-\ell)!}{[(1+t)^{\lambda} - 1]^{k-\ell}} + \sum_{\ell=1}^k \binom{k}{\ell} \frac{(-1)^{k-\ell} \lambda^{k-\ell+1} (k-\ell)!}{[(1+t)^{\lambda} - 1]^{k-\ell+1}} \right] \\
& \quad \times (1+t)^{k\lambda-n} \lambda^{n-k} \frac{(-1)^k}{k!} \sum_{\ell=0}^k (-1)^{\ell} \binom{k}{\ell} \prod_{q=0}^{n-1} \left(\ell - \frac{q}{\lambda} \right) \\
& = \frac{1}{(1+t)^n} \sum_{k=1}^n \left[\lambda F_{\lambda}(t) \sum_{\ell=0}^k \binom{k}{\ell} \frac{(-1)^{\ell} \lambda^{\ell} \ell!}{[(1+t)^{\lambda} - 1]^{\ell}} + \sum_{\ell=0}^{k-1} \binom{k}{\ell} \frac{(-1)^{\ell} \lambda^{\ell+1} \ell!}{[(1+t)^{\lambda} - 1]^{\ell+1}} \right] \\
& \quad \times \frac{(1+t)^{k\lambda}}{\lambda^k} \frac{(-1)^k}{k!} \sum_{\ell=0}^k (-1)^{\ell} \binom{k}{\ell} \prod_{q=0}^{n-1} (\ell\lambda - q).
\end{aligned}$$

Accordingly, we have

$$\begin{aligned}
(1+t)^n [(1+t)^{\lambda} - 1]^n F_{\lambda}^{(n)}(t) & = \sum_{k=1}^n \left[\frac{(1+t)^{k\lambda}}{\lambda^k} \frac{(-1)^k}{k!} \sum_{\ell=0}^k (-1)^{\ell} \binom{k}{\ell} \prod_{q=0}^{n-1} (\ell\lambda - q) \right] \\
& \times \left[F_{\lambda}(t) \sum_{\ell=0}^k \binom{k}{\ell} (-1)^{\ell} \lambda^{\ell} \ell! [(1+t)^{\lambda} - 1]^{n-\ell} + \sum_{\ell=0}^{k-1} \binom{k}{\ell} (-1)^{\ell} \lambda^{\ell} \ell! [(1+t)^{\lambda} - 1]^{n-\ell-1} \right] \\
& = \sum_{k=1}^n \left[\frac{(-1)^k}{\lambda^k} \sum_{\ell=0}^k (-1)^{\ell} \binom{k}{\ell} \prod_{q=0}^{n-1} (\ell\lambda - q) \right] (1+t)^{k\lambda} \\
& \quad \times \left[F_{\lambda}(t) \sum_{\ell=0}^k \frac{(-1)^{\ell} \lambda^{\ell}}{(k-\ell)!} \sum_{m=0}^{n-\ell} \binom{n-\ell}{m} (1+t)^{m\lambda} (-1)^{n-\ell-m} \right. \\
& \quad \left. + \sum_{\ell=0}^{k-1} \frac{(-1)^{\ell} \lambda^{\ell}}{(k-\ell)!} \sum_{m=0}^{n-\ell-1} \binom{n-\ell-1}{m} (1+t)^{m\lambda} (-1)^{n-\ell-1-m} \right] \\
& = (-1)^n \sum_{k=1}^n \left[\frac{(-1)^k}{\lambda^k} \sum_{\ell=0}^k (-1)^{\ell} \binom{k}{\ell} \prod_{q=0}^{n-1} (\ell\lambda - q) \right] (1+t)^{k\lambda} \\
& \quad \times \left[F_{\lambda}(t) \sum_{\ell=0}^k \frac{\lambda^{\ell}}{(k-\ell)!} \sum_{m=0}^{n-\ell} (-1)^m \binom{n-\ell}{m} (1+t)^{m\lambda} \right. \\
& \quad \left. + \sum_{\ell=0}^{k-1} \frac{\lambda^{\ell}}{(k-\ell)!} \sum_{m=0}^{n-\ell-1} (-1)^{m+1} \binom{n-\ell-1}{m} (1+t)^{m\lambda} \right] \\
& = (-1)^n F_{\lambda}(t) \sum_{k=1}^n A_k(n, \lambda) \sum_{\ell=0}^k \frac{\lambda^{\ell}}{(k-\ell)!} \sum_{m=0}^{n-\ell} (-1)^m \binom{n-\ell}{m} (1+t)^{(k+m)\lambda} \\
& \quad + (-1)^n \sum_{k=1}^n A_k(n, \lambda) \sum_{\ell=0}^{k-1} \frac{\lambda^{\ell}}{(k-\ell)!} \sum_{m=0}^{n-\ell-1} (-1)^{m+1} \binom{n-\ell-1}{m} (1+t)^{(k+m)\lambda}.
\end{aligned}$$

Since

$$\begin{aligned}
& \sum_{k=1}^n A_k(n, \lambda) \sum_{\ell=0}^k \frac{\lambda^\ell}{(k-\ell)!} \sum_{m=0}^{n-\ell} (-1)^m \binom{n-\ell}{m} (1+t)^{(k+m)\lambda} \\
&= \sum_{k=1}^n A_k(n, \lambda) \sum_{\ell=0}^k \frac{\lambda^\ell}{(k-\ell)!} \sum_{m=0}^n (-1)^m \binom{n-\ell}{m} (1+t)^{(k+m)\lambda} \\
&= \sum_{k=1}^n A_k(n, \lambda) (1+t)^{k\lambda} \sum_{m=0}^n (-1)^m \left[\sum_{\ell=0}^k \frac{\lambda^\ell}{(k-\ell)!} \binom{n-\ell}{m} \right] (1+t)^{m\lambda} \\
&= \sum_{k=1}^n A_k(n, \lambda) (1+t)^{k\lambda} \sum_{m=0}^n (-1)^m \left[\sum_{\ell=0}^{\min\{n-m, k\}} \frac{\lambda^\ell}{(k-\ell)!} \binom{n-\ell}{m} \right] (1+t)^{m\lambda} \\
&= \sum_{k=1}^n A_k(n, \lambda) \sum_{m=0}^n (-1)^m \left[\sum_{\ell=0}^{\min\{n-m, k\}} \frac{\lambda^\ell}{(k-\ell)!} \binom{n-\ell}{m} \right] (1+t)^{(k+m)\lambda} \\
&= \sum_{i=1}^{2n} \left[\sum_{\substack{k+m=i \\ 1 \leq k \leq n \\ 0 \leq m \leq n}} (-1)^m A_k(n, \lambda) \sum_{\ell=0}^{\min\{n-m, k\}} \frac{\lambda^\ell}{(k-\ell)!} \binom{n-\ell}{m} \right] (1+t)^{i\lambda}
\end{aligned}$$

and

$$\begin{aligned}
& \sum_{k=1}^n A_k(n, \lambda) \sum_{\ell=0}^{k-1} \frac{\lambda^\ell}{(k-\ell)!} \sum_{m=0}^{n-\ell-1} (-1)^{m+1} \binom{n-\ell-1}{m} (1+t)^{(k+m)\lambda} \\
&= \sum_{k=1}^n A_k(n, \lambda) \sum_{\ell=0}^{k-1} \frac{\lambda^\ell}{(k-\ell)!} \sum_{m=0}^{n-1} (-1)^{m+1} \binom{n-\ell-1}{m} (1+t)^{(k+m)\lambda} \\
&= \sum_{k=1}^n A_k(n, \lambda) \sum_{m=0}^{n-1} (-1)^{m+1} \left[\sum_{\ell=0}^{k-1} \frac{\lambda^\ell}{(k-\ell)!} \binom{n-\ell-1}{m} \right] (1+t)^{(k+m)\lambda} \\
&= \sum_{k=1}^n A_k(n, \lambda) \sum_{m=0}^{n-1} (-1)^{m+1} \left[\sum_{\ell=0}^{\min\{k-1, n-m-1\}} \frac{\lambda^\ell}{(k-\ell)!} \binom{n-\ell-1}{m} \right] (1+t)^{(k+m)\lambda} \\
&= \sum_{i=1}^{2n-1} \left[\sum_{\substack{k+m=i \\ 1 \leq k \leq n \\ 0 \leq m \leq n-1}} (-1)^{m+1} A_k(n, \lambda) \sum_{\ell=0}^{\min\{k-1, n-m-1\}} \frac{\lambda^\ell}{(k-\ell)!} \binom{n-\ell-1}{m} \right] (1+t)^{i\lambda},
\end{aligned}$$

where an empty sum is understood to be 0 and $\binom{p}{q} = 0$ for $q > p \geq 0$, the equation (6) and the formulas (7) and 8 are thus proved. The proof of Theorem 2 is complete. $\square$

4. REMARKS

Finally, we list several remarks on our main results and closely related things.

Remark 1. Comparing (6) with (2) reveals that

$$a_i(n, \lambda) = (-1)^n \alpha_i(n, \lambda) \quad \text{and} \quad b_i(n, \lambda) = (-1)^n \beta_i(n, \lambda).$$

Remark 2. It is easy to see that explicit expressions (7) and (8) for $\alpha_i(n, \lambda)$ and $\beta_i(n, \lambda)$ are more meaningful and more significant than those in [2, Theorem 2.1] for $a_i(n, \lambda)$ and $b_i(n, \lambda)$ mentioned above.

Remark 3. The formula (5) was derived in [3] and mentioned in [4, 8].

Remark 4. Per requests of anonymous referees, the preprint [17] is split into and simplified as two formally published papers [18, 29].

Remark 5. The motivations in the papers [5, 6, 9, 10, 11, 12, 13, 14, 15, 16, 20, 21, 24, 25, 26, 27, 30, 31, 32, 38, 35, 36, 37, 40, 41, 42] are same as the one in this paper.

Remark 6. This paper is a slightly revised version of the preprint [28].

Acknowledgements. The authors are thankful to Professor Taekyun Kim for his sending a doi link for his paper [2] through an e-mail on 28 February 2018.

The third author was supported by the National Research Foundation of Korea (Grant No. 2018R1D1A1B07041846).

REFERENCES

- [1] L. Comtet, *Advanced Combinatorics: The Art of Finite and Infinite Expansions*, Revised and Enlarged Edition, D. Reidel Publishing Co., Dordrecht and Boston, 1974; available online at <https://doi.org/10.1007/978-94-010-2196-8>.
- [2] T. Kim, D. S. Kim, and G.-W. Jang, *Differential equations associated with degenerate Cauchy numbers*, Iran. J. Sci. Technol. Trans. Sci. **43** (2019), no. 3, 1021–1025; available online at <https://doi.org/10.1007/s40995-018-0531-y>.
- [3] G. Nemes, *An asymptotic expansion for the Bernoulli numbers of the second kind*, J. Integer Seq. **14** (2011), Article 11.4.8.
- [4] F. Qi, *A new formula for the Bernoulli numbers of the second kind in terms of the Stirling numbers of the first kind*, Publ. Inst. Math. (Beograd) (N.S.) **100(114)** (2016), 243–249; available online at <https://doi.org/10.2298/PIM150501028Q>.
- [5] F. Qi, *A simple form for coefficients in a family of nonlinear ordinary differential equations*, Adv. Appl. Math. Sci. **17** (2018), no. 8, 555–561.
- [6] F. Qi, *A simple form for coefficients in a family of ordinary differential equations related to the generating function of the Legendre polynomials*, Adv. Appl. Math. Sci. **17** (2018), no. 11, 693–700.
- [7] F. Qi, *An integral representation, complete monotonicity, and inequalities of Cauchy numbers of the second kind*, J. Number Theory **144** (2014), 244–255; available online at <https://doi.org/10.1016/j.jnt.2014.05.009>.
- [8] F. Qi, *Explicit formulas for computing Bernoulli numbers of the second kind and Stirling numbers of the first kind*, Filomat **28** (2014), no. 2, 319–327; available online at <https://doi.org/10.2298/FIL1402319Q>.
- [9] F. Qi, *Explicit formulas for the convolved Fibonacci numbers*, ResearchGate Working Paper (2016), available online at <https://doi.org/10.13140/RG.2.2.36768.17927>.
- [10] F. Qi, *Notes on several families of differential equations related to the generating function for the Bernoulli numbers of the second kind*, Turkish J. Anal. Number Theory **6** (2018), no. 2, 40–42; available online at <https://doi.org/10.12691/tjant-6-2-1>.
- [11] F. Qi, *Simple forms for coefficients in two families of ordinary differential equations*, Glob. J. Math. Anal. **6** (2018), no. 1, 7–9; available online at <https://doi.org/10.14419/gjma.v6i1.9778>.
- [12] F. Qi, *Simplification of coefficients in two families of nonlinear ordinary differential equations*, Turkish J. Anal. Number Theory **6** (2018), no. 4, 116–119; available online at <https://doi.org/10.12691/tjant-6-4-2>.

- [13] F. Qi, *Simplifying coefficients in a family of nonlinear ordinary differential equations*, Acta Comment. Univ. Tartu. Math. **22** (2018), no. 2, 293–297; available online at <https://doi.org/10.12697/ACUTM.2018.22.24>.
- [14] F. Qi, *Simplifying coefficients in a family of ordinary differential equations related to the generating function of the Laguerre polynomials*, Appl. Appl. Math. **13** (2018), no. 2, 750–755.
- [15] F. Qi, *Simplifying coefficients in a family of ordinary differential equations related to the generating function of the Mittag-Leffler polynomials*, Korean J. Math. **27** (2019), no. 2, 417–423; available online at <https://doi.org/10.11568/kjm.2019.27.2.417>.
- [16] F. Qi, *Simplifying coefficients in differential equations related to generating functions of reverse Bessel and partially degenerate Bell polynomials*, Bol. Soc. Paran. Mat. **39** (2021), no. 4, in press; available online at <http://dx.doi.org/10.5269/bspm.41758>.
- [17] F. Qi, V. Čerňanová, and Y. S. Semenov, *On tridiagonal determinants and the Cauchy product of central Delannoy numbers*, ResearchGate Working Paper (2016), available online at <https://doi.org/10.13140/RG.2.1.3772.6967>.
- [18] F. Qi, V. Čerňanová, and Y. S. Semenov, *Some tridiagonal determinants related to central Delannoy numbers, the Chebyshev polynomials, and the Fibonacci polynomials*, Politehn. Univ. Bucharest Sci. Bull. Ser. A Appl. Math. Phys. **81** (2019), no. 1, 123–136.
- [19] F. Qi, V. Čerňanová, X.-T. Shi, and B.-N. Guo, *Some properties of central Delannoy numbers*, J. Comput. Appl. Math. **328** (2018), 101–115; available online at <https://doi.org/10.1016/j.cam.2017.07.013>.
- [20] F. Qi and B.-N. Guo, *A diagonal recurrence relation for the Stirling numbers of the first kind*, Appl. Anal. Discrete Math. **12** (2018), no. 1, 153–165; available online at <https://doi.org/10.2298/AADM170405004Q>.
- [21] F. Qi and B.-N. Guo, *Explicit formulas and recurrence relations for higher order Eulerian polynomials*, Indag. Math. **28** (2017), no. 4, 884–891; available online at <https://doi.org/10.1016/j.indag.2017.06.010>.
- [22] F. Qi and B.-N. Guo, *Explicit formulas for special values of the Bell polynomials of the second kind and for the Euler numbers and polynomials*, Mediterr. J. Math. **14** (2017), no. 3, Article 140, 14 pages; available online at <https://doi.org/10.1007/s00009-017-0939-1>.
- [23] F. Qi and B.-N. Guo, *Several explicit and recursive formulas for generalized Motzkin numbers*, AIMS Mathematics **5** (2020), no. 3, in press; available online at <https://doi.org/10.20944/preprints201703.0200.v1>.
- [24] F. Qi and B.-N. Guo, *Some properties of the Hermite polynomials*, Georgian Math. J. **27** (2020), in press; available online at <https://doi.org/10.20944/preprints201611.0145.v1>.
- [25] F. Qi and B.-N. Guo, *Viewing some ordinary differential equations from the angle of derivative polynomials*, Iran. J. Math. Sci. Inform. **15** (2020), no. 2, in press; available online at <https://doi.org/10.20944/preprints201610.0043.v1>.
- [26] F. Qi, D. Lim, and B.-N. Guo, *Explicit formulas and identities for the Bell polynomials and a sequence of polynomials applied to differential equations*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM **113** (2019), no. 1, 1–9; available online at <https://doi.org/10.1007/s13398-017-0427-2>.
- [27] F. Qi, D. Lim, and B.-N. Guo, *Some identities related to Eulerian polynomials and involving the Stirling numbers*, Appl. Anal. Discrete Math. **12** (2018), no. 2, 467–480; available online at <https://doi.org/10.2298/AADM171008014Q>.
- [28] F. Qi, D. Lim, and A.-Q. Liu, *Explicit expressions related to degenerate Cauchy numbers and their generating function*, HAL preprint (2018), available online at <https://hal.archives-ouvertes.fr/hal-01725045>.
- [29] F. Qi and A.-Q. Liu, *Alternative proofs of some formulas for two tridiagonal determinants*, Acta Univ. Sapientiae Math. **10** (2018), no. 2, 287–297; available online at <https://doi.org/10.2478/ausm-2018-0022>.
- [30] F. Qi, D.-W. Niu, and B.-N. Guo, *Simplification of coefficients in differential equations associated with higher order Frobenius–Euler numbers*, Tatra Mt. Math. Publ. **72** (2018), 67–76; available online at <https://doi.org/10.2478/tmmp-2018-0022>.
- [31] F. Qi, D.-W. Niu, and B.-N. Guo, *Simplifying coefficients in differential equations associated with higher order Bernoulli numbers of the second kind*, AIMS Math. **4** (2019), no. 2, 170–175; available online at <https://doi.org/10.3934/Math.2019.2.170>.
- [32] F. Qi, D.-W. Niu, and B.-N. Guo, *Some identities for a sequence of unnamed polynomials connected with the Bell polynomials*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A

- Math. RACSAM **113** (2019), no. 2, 557–567; available online at <https://doi.org/10.1007/s13398-018-0494-z>.
- [33] F. Qi, D.-W. Niu, D. Lim, and B.-N. Guo, *Closed formulas and identities for the Bell polynomials and falling factorials*, Contrib. Discrete Math. **14** (2019), no. 2, in press; available online at <https://doi.org/10.13140/RG.2.2.34679.52640>.
- [34] F. Qi, X.-T. Shi, and F.-F. Liu, *Several identities involving the falling and rising factorials and the Cauchy, Lah, and Stirling numbers*, Acta Univ. Sapientiae Math. **8** (2016), no. 2, 282–297; available online at <https://doi.org/10.1515/ausm-2016-0019>.
- [35] F. Qi, J.-L. Wang, and B.-N. Guo, *Notes on a family of inhomogeneous linear ordinary differential equations*, Adv. Appl. Math. Sci. **17** (2018), no. 4, 361–368.
- [36] F. Qi, J.-L. Wang, and B.-N. Guo, *Simplifying and finding ordinary differential equations in terms of the Stirling numbers*, Korean J. Math. **26** (2018), no. 4, 675–681; available online at <https://doi.org/10.11568/kjm.2018.26.4.675>.
- [37] F. Qi, J.-L. Wang, and B.-N. Guo, *Simplifying differential equations concerning degenerate Bernoulli and Euler numbers*, Trans. A. Razmadze Math. Inst. **172** (2018), no. 1, 90–94; available online at <https://doi.org/10.1016/j.trmi.2017.08.001>.
- [38] F. Qi and Y.-H. Yao, *Simplifying coefficients in differential equations for generating function of Catalan numbers*, J. Taibah Univ. Sci. **13** (2019), no. 1, 947–950; available online at <https://doi.org/10.1080/16583655.2019.1663782>.
- [39] F. Qi and X.-J. Zhang, *An integral representation, some inequalities, and complete monotonicity of the Bernoulli numbers of the second kind*, Bull. Korean Math. Soc. **52** (2015), no. 3, 987–998; available online at <https://doi.org/10.4134/BKMS.2015.52.3.987>.
- [40] F. Qi and J.-L. Zhao, *Some properties of the Bernoulli numbers of the second kind and their generating function*, Bull. Korean Math. Soc. **55** (2018), no. 6, 1909–1920; available online at <https://doi.org/10.4134/BKMS.b180039>.
- [41] F. Qi, Q. Zou, and B.-N. Guo, *The inverse of a triangular matrix and several identities of the Catalan numbers*, Appl. Anal. Discrete Math. **13** (2019), no. 2, 518–541; available online at <https://doi.org/10.2298/AADM190118018Q>.
- [42] J.-L. Zhao, J.-L. Wang, and F. Qi, *Derivative polynomials of a function related to the Apostol–Euler and Frobenius–Euler numbers*, J. Nonlinear Sci. Appl. **10** (2017), no. 4, 1345–1349; available online at <https://doi.org/10.22436/jnsa.010.04.06>.

(Qi) COLLEGE OF MATHEMATICS, INNER MONGOLIA UNIVERSITY FOR NATIONALITIES, TONGLIAO 028043, INNER MONGOLIA, CHINA; SCHOOL OF MATHEMATICAL SCIENCES, TIANJIN POLYTECHNIC UNIVERSITY, TIANJIN 300387, CHINA; INSTITUTE OF MATHEMATICS, HENAN POLYTECHNIC UNIVERSITY, JIAOZUO 454010, HENAN, CHINA

URL: <https://qifeng618.wordpress.com>, <http://orcid.org/0000-0001-6239-2968>

Email address: qifeng618@gmail.com, qifeng618@hotmail.com, qifeng618@qq.com

(Liu) DEPARTMENT OF MATHEMATICS, SANMENXIA POLYTECHNIC, SANMENXIA 472000, HENAN, CHINA

Email address: lady_smx@126.com, 740297962@qq.com

(Lim) DEPARTMENT OF MATHEMATICS EDUCATION, ANDONG NATIONAL UNIVERSITY, ANDONG 36729, REPUBLIC OF KOREA

URL: <http://orcid.org/0000-0002-0928-8480>

Email address: dgrim84@gmail.com, dklim@andong.ac.kr