



HAL
open science

The determination of new stable zeolite frameworks using a parallel hybrid genetic algorithm

Omar Abdelkafi, Lhassane Idoumghar, Julien Lepagnot, Jean-Louis Paillaud

► **To cite this version:**

Omar Abdelkafi, Lhassane Idoumghar, Julien Lepagnot, Jean-Louis Paillaud. The determination of new stable zeolite frameworks using a parallel hybrid genetic algorithm. OLA 2018 - International Workshop on Optimization and Learning: Challenges and Applications, Feb 2018, Alicante, Spain. pp.1-2. hal-01726493

HAL Id: hal-01726493

<https://hal.science/hal-01726493>

Submitted on 9 Mar 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The determination of new stable zeolite frameworks using a parallel hybrid genetic algorithm

Omar Abdelkafi¹, Lhassane Idoumghar², Julien Lepagnot², and Jean-Louis Paillaud³

¹ Université Lille 1, CRISTAL/UMR CNRS 9189 - INRIA Lille Nord Europe, 59655 - Villeneuve d'Ascq cedex, France

`omar.abdelkafi@univ-lille1.fr`

² Université de Haute-Alsace, LMIA, EA 3993, F-68093 Mulhouse, France

`{lhassane.idoumghar, julien.lepagnot}@uha.fr`

³ Université de Haute-Alsace, CNRS, IS2M UMR 7361, F-68093 Mulhouse, France

`jean-louis.paillaud@uha.fr`

1 Introduction

Zeolites were originally discovered by the swedish mineralogist Axel Fredrik Cronstedt in 1756 [1]. However, it is only from the 1930s with Barrer pioneering work on the synthesis and adsorption properties of zeolites that the science of zeolites has really taken off [2]-[3]. Over the years 1950s the first processes for the synthesis of zeolites A, X and Y are developed before their commercialization by Union Carbide in 1954 [2]. The zeolites are crystalline microporous materials with outstanding characteristics of catalysis and adsorption. Zeolites are composed mainly of silicon (Si) and aluminum (Al) atoms which represent two of the most abundant elements on earth. The Zeolite Structures Problem (ZSP), so called by Falcioni and Deem [4], can be classified in the area of the computational chemistry [5]. The aim of this problem is to find optimal zeolite structures which depend on initial atoms positions.

Metaheuristics [6] are powerful heuristic algorithms capable of solving, among other problems, combinatorial and continuous optimization problems. The common characteristic of metaheuristics is to use a random mechanism to explore the space of solutions. Beyond the topology based methods, two major metaheuristics have been successfully exploited in the zeolite structures prediction: the former one is simulated annealing (SA) [7], used in particular for the crystal structure resolution from experimentally available data. The second, more recent, is the genetic algorithm (GA) [8].

2 Methods

Our proposition to solve the ZSP is a genetic algorithm hybridized with a heuristic based on the memorization and the prediction of the atoms positions. We have named this algorithm MEGA-HZ (from *ME*emory *GE*netic *AL*gorithm *HY*bridized for *ZE*olites). The MEGA-HZ generates a random population. The oxygen atoms are added at the end of the algorithm to the center of two atoms T that can be connected. The aim of this algorithm is to find atoms positions to generate potential zeolite structures. The evolutionary algorithms are naturally adapted to propose original structures [9]. In this work, a distributed and massively parallel genetic hybrid algorithm is developed.

To create a viable zeolites by the MEGA-HZ, the algorithm needs to put the atoms at the right positions inside the AU. Each atom position of the Assymmetric Unit (AU) is defined by 3 atomic coordinates (x,y,z) . This step creates the AU. The symmetric operations defined in the space group are applied to the AU to generate the UC (Unit Cell). This UC is the smallest volume unit that contains all of the structural and symmetry information. Then, in order to form the bulk atoms arrangement of the crystals, the unit cells are stacked in the three-dimensional space. The objective is to generate viable zeolites frameworks taking into account the constraints determined by the user input.

3 Results

The MEGA-HZ algorithm found an unknown stable zeolite structure not recognized by the Structural Commission of the International Zeolite Association and the Atlas of Prospective Zeolite Structures [10, 11].

The unknown topology $HZM\#4$ has a three-dimensional pore system with 8 and 10 T atoms. One type of Composite Building Units (CBU) is observable ($[4^55^2]$ or $d5r$), these CBUs are connected with 4 tetrahedres to form pairs (Figure 1a). The arrangement of these pairs, separated from each other in space, gives rise to large cages ($[4^{12}5^46^28^410^2]$, Figure 1b), each large cage is linked to 8 pairs of CBUs $d5r$ by pooling their cycles to 4 and 5 tetrahedron as shown in Figure 1c. Similarly, each pair of CBUs $d5r$ is linked to 8 large cages ($[4^{12}5^46^28^410^2]$).




Fig. 1. 'Tiling' of the structure $HZM\#4$ showing (a) the spatial arrangement of CBUs pairs $d5r$, (b) a large cage $[4^{12}5^46^28^410^2]$ and (c) surrounded by 8 pairs of CBUs $d5r$. Oxygen atoms have been omitted for clarity.

References

1. A. Cronstedt, "Ron och beskrifning om en obekant barg art, som kallas zeolites," *Kungliga Vetenskapsakademiens Handlingar*, vol. 17, pp. 120–123, 1756.
2. R. M. Barrer, *Hydrothermal chemistry of zeolites*. London, New York: Academic Press, 1982.
3. M. Guisnet and J. Gilson, *Introduction to zeolite science and technology*, ser. Catalytic Science Series. Imperial College Press, 2002, ch. 1, pp. 1–28.
4. M. Falcioni and M. W. Deem, "A biased Monte Carlo scheme for zeolite structure solution," *J. Chem. Phys.*, vol. 110, no. 3, pp. 1754–1766, 1999.
5. B. Archibald, O. Brümmer, M. Devenney, D. M. Giaquinta, B. Jandeleit, W. H. Weinberg, and T. Weskamp, *Combinatorial Aspects of Materials Science*. Wiley-VCH Verlag GmbH & Co. KGaA, 2005, pp. 1017–1062.
6. C. Blum and A. Roli, "Metaheuristics in combinatorial optimization: Overview and conceptual comparison," *ACM Computing Surveys*, vol. 35, no. 3, pp. 268–308, 2003.
7. J. Pannetier, J. Bassas-Alsina, J. Rodriguez-Carvajal, and V. Caignaert, "Prediction of crystal structures from crystal chemistry rules by simulated annealing," *Nature*, vol. 346, no. 6282, pp. 343–345, 1990.
8. D. A. Coley, *An Introduction to genetic algorithms for scientists and engineers*. World Scientific Publishing Co., Inc., 1998.
9. G. Jones, P. Willett, and R. C. Glen, "Molecular recognition of receptor sites using a genetic algorithm with a description of desolvation," *Molecular Biology*, vol. 245, no. 1, pp. 43–53, 1995.
10. C. Baerlocher and L. B. McCusker, "Database of zeolite structures," online since 1996. [Online]. Available: <http://www.iza-structure.org/databases/>
11. M. D. Foster and M. M. J. Treacy, "A database of hypothetical zeolite structures," online since April 2004. [Online]. Available: <http://www.hypotheticalzeolites.net>